



HAL
open science

Semi-industrial production of a minimally processed infant formula powder using membrane filtration

X. Yu, Nadine Leconte, S. Méjean, Gilles Garric, Sergine Even, G. Henry, Frederic J. Tessier, M. Howsam, Thomas Croguennec, Geneviève Gésan-Guiziou, et al.

► To cite this version:

X. Yu, Nadine Leconte, S. Méjean, Gilles Garric, Sergine Even, et al.. Semi-industrial production of a minimally processed infant formula powder using membrane filtration. *Journal of Dairy Science*, 2021, 104 (5), pp.5265-5278. 10.3168/jds.2020-19529 . hal-03176288

HAL Id: hal-03176288

<https://hal.inrae.fr/hal-03176288v1>

Submitted on 22 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Semi-industrial production of a minimally processed infant formula powder using membrane filtration

X. Yu,¹ N. Leconte,¹ S. Méjean,¹ G. Garric,¹ S. Even,¹ G. Henry,¹ F. J. Tessier,² M. Howsam,² T. Croguennec,¹ G. Gésan-Guiziuou,¹ D. Dupont,¹ R. Jeantet,¹ and A. Deglaire^{1*}

¹STLO, INRAE, Institut Agro, 35042, Rennes, France

²University of Lille, Inserm, CHU Lille, Pasteur Institute of Lille, U1167 – RID-AGE, F-59000 Lille, France

ABSTRACT

Infant formula (IF) is submitted to several heat treatments during production, which can lead to denaturation or aggregation of proteins and promote Maillard reaction. The objective of this study was to investigate innovative minimal processing routes for the production of first-age IF powder, thus ensuring microbial safety with minimal level of protein denaturation. Three nutritionally complete IF powders were produced at a semi-industrial scale based on ingredients obtained by fresh bovine milk microfiltration (0.8 and 0.1- μm pore size membranes). Low-temperature vacuum evaporation (50°C) and spray-drying (inlet and outlet temperatures of 160 and 70°C, respectively) were conducted to produce the T- formula with no additional heat treatment. The T+ formula was produced with a moderate heat treatment (75°C for 2 min) applied before spray-drying, whereas the T+++ formula received successive heat treatments (72°C for 30 s on the milk; 90°C for 2–3 s before evaporation; 85°C for 2 min before spray-drying), thus mimicking commercial powdered IF. Protein denaturation and Maillard reaction products were followed throughout the production steps and the physicochemical properties of the powders were characterized. The 3 IF powders presented satisfactory physical properties in terms of a_w , free fat content, glass transition temperature, and solubility index, as well as satisfactory bacteriological quality with a total flora $<10^3$ cfu/g and an absence of pathogens when a high level of bacteriological quality of the ingredients was ensured. Protein denaturation occurred mostly during the heat treatments of T+ and T+++ and was limited during the spray-drying process. The IF powder produced without heat treatment (T-) presented a protein denaturation extent ($6 \pm 4\%$) significantly lower than that in T+++ ($58 \pm 0\%$), but not significantly differ-

ent from that in T+ ($10 \pm 4\%$). Although T- tended to contain less Maillard reaction products than T+ and T+++ , the Maillard reaction products did not significantly discriminate the infant formulas in the frame of this work. The present study demonstrated the feasibility of producing at a semi-industrial scale an infant formula being bacteriologically safe and containing a high content of native proteins. Application of a moderate heat treatment before spray-drying could further guarantee the microbiological quality of the IF powders while maintaining a low protein denaturation extent. This study opens up new avenues for the production of minimally processed IF powders.

Key words: infant formula, filtration, minimally processed, protein denaturation

INTRODUCTION

According to UNICEF (2020), 56% of infants 0 to 5 mo of age receive a human milk substitute, which is in most circumstances an infant formula (IF), most often available in a powder form. The IF powders are generally manufactured by blending skim milk with other dried ingredients, including whey proteins (WP) originating from cheese whey or less commonly from skim milk microfiltration (MF) dried permeate, which are all together rehydrated and subjected to a succession of thermal processes (pasteurization, evaporation, drying) to ensure the microbial safety and shelf life stability of the product. These successive heat treatments induce numerous physicochemical modifications of the components, among them the denaturation or aggregation of WP (Brodkorb et al., 2016). The latter can affect WP functionality during IF processing, in particular when they form aggregates that increase viscosity, reduce the emulsion stability, and decrease the overall processing performance (Joyce et al., 2017). In addition, the heat-induced protein denaturation may have nutritional consequences. It has been previously reported to affect the in vitro digestion kinetics of caseins (CN; Dupont et al., 2010; Halabi et al., 2020a) and of β -LG (Takagi

Received August 26, 2020.

Accepted January 8, 2021.

*Corresponding author: amelie.deglair@agrocampus-ouest.fr

et al., 2003; Peram et al., 2013), as well as the microstructure of the gastric digesta (Halabi et al., 2020a). It could further affect the dietary nitrogen retention by the body, such as reported in adults fed UHT-treated milk versus pasteurized or microfiltered milk (Lacroix et al., 2008). In addition, the heat-induced protein denaturation can modulate the protein allergenicity, by either masking or, on the contrary, exposing epitopes, depending on the intensity of the heat treatment (Golkar et al., 2019). However, some authors have reported an increased allergic sensitization in pasteurized versus raw milk (Roth-Walter et al., 2008) and a loss of immunologically active WP due their denaturation, thus decreasing the allergy-protective capacity of raw cow's milk (Abbring et al., 2020). This can be further modulated by the Maillard reaction products, resulting from the reaction between lysine-rich proteins and reducing sugar (lactose) induced by heat treatment. This results in the formation in the first stage of Amadori products, such as lactulosyllysine, up to more advanced glycation end products, such as N^ε-carboxymethyl-lysine (CML). Consequently, the lysine bioavailability can be affected (Nunes et al., 2019), in addition to other potential health effects, such as pro-inflammatory and pro-oxidant effects for CML (Elmhiri et al., 2015) and the promotion of allergies in infants (Baskara et al., 2017). Overall, maintaining as much as possible the proteins in their native state may improve the nutritional quality of IF, even though bovine and human WP differ by nature.

Integrating some MF steps to fresh milk into the IF production process could limit the heat treatment effect on IF constituents while maintaining the bacteriological safety, and is to date an understudied strategy in the area of IF production. Several studies have demonstrated that MF with 1.4- μm pore size ceramic membranes can substantially retain bacteria in skim milk, with an overall decimal reduction above 4 log₁₀ units (Trouvé et al., 1991; Fauquant et al., 2009, 2014; Tomasula et al., 2011; Schmidt et al., 2012). More specifically, Madec et al. (1992) have reported a decimal reduction around 2 log₁₀ units for *Listeria* and *Salmonella*, respectively, using nonvirulent strains, and Trouvé et al. (1991) have shown a decimal reduction close to 3 log₁₀ units for bacteria with various morphologies and cellular volumes, representative of raw milk contamination. Therefore, MF appears as an alternative to usually applied thermal processes such as pasteurization for reducing microbial load of fresh milk, while maintaining proteins in their native state. In addition, MF with 0.1- μm pore size membrane can be used to separate CN micelles from the other components of smaller molecular weight (WP, lactose, vitamins, and minerals; Gésan-Guiziou et al., 1999; Saboya and Maubois,

2000). After concentration by ultrafiltration, the liquid WP-rich permeate can be mixed with the microfiltered milk to achieve the human milk protein CN:WP ratio (40:60), while ensuring bacteriological safety without heat treatments. Notwithstanding significant development of filtration in the dairy industry over the past decade, few studies have used its potential for limiting thermal processes, and thus, maintaining proteins in their native state in IF powder. The main challenge is to integrate such nonthermal process in an overall low temperature processing route, allowing the production of a first-age (0–6 mo) IF powder.

The objective of this study was to investigate whether a minimal processing route, based on filtration of fresh milk and other unit operations (evaporation, homogenization, spray-drying), could allow the production, at a semi-industrial scale, of a qualitative IF powder presenting a limited level of protein denaturation and being bacteriologically safe. The microfiltration for skim milk debacterization was decreased to a pore size of 0.8 μm to improve the bacteria retention (Tomasula et al., 2011), particularly, regarding *Cronobacter spp.* Three IF powders were produced: T–, without heat treatment; T+, with pasteurization of the concentrate alone (75°C for 2 min); T+++ , with successive heat treatments including one on the concentrate at 85°C for 2 min before spray-drying. The extent of WP denaturation and the formation of Maillard reaction products were evaluated throughout the process. The physicochemical properties of IF powders before and after reconstitution were characterized. To the best of our knowledge, this is the first time that such a minimal processing technological route, integrating membrane filtration and spray-drying operations, is proposed and characterized for the production of a first-age IF.

MATERIALS AND METHODS

Chemicals

Unless stated otherwise, chemicals are from commercial origin (Sigma-Aldrich).

IF Ingredients

Raw cow milk (900 kg) was purchased from a local dairy cooperative (SODIAAL Entremont). Liquid WP isolate (**WPI**), obtained by low temperature (<10°C) MF 0.1 μm of fresh skim milk, followed by ultrafiltration, was purchased from a dairy company (confidential source). The WPI had a DM of 26.6% (wt/wt) of which 92% were proteins. The WP accounted for 93% (wt/wt) of the protein fraction, such as determined by the Kjeldahl method. Lactose was supplied by Lactalis In-

redients. Oil blend and vitamin and mineral premixes, adapted to IF, were purchased from Cargill (Schiphol) and Fimalis (Levallois-Perret), respectively.

IF Processing

Three types of IF powders, namely T⁻, T⁺, and T⁺⁺⁺ according to the heat treatments received during manufacture, were produced according to Figure 1. Fresh raw whole milk was skimmed by centrifugal cream separation at 50°C in rising phase and then subjected to MF using 0.8- μm pore size graded permeability (GP) Pall mineral membranes (Pall Corporation, Bazel, France) with a total filtration area of 1.68 m². The observed retention of total proteins, as well as of caseins, was 97.5%. The crossflow velocity was maintained at 7 m·s⁻¹ (retentate flowrate of 43 m³·h⁻¹) and the retentate pressure drop was maintained constant at 2 bar. The temperature was maintained at 50 \pm 2°C with continuous cooling of the filtrate (microfiltered skim milk, MSM) using plate heat exchangers. The filtrate and the retentate fluxes were adjusted to 630 L/h (375 L·h⁻¹/m) and 33 L/h, respectively, to maintain a volume concentration factor of 20. For T⁻, the MSM was directly subjected to the mixing step. For T⁺, the MSM was stored overnight at 4°C before mixing. For T⁺⁺⁺, the MSM was pasteurized at 72°C for 30 s, stored overnight at 4°C, and repasteurized at 72°C for 30 s before mixing. For the mixing step, MSM, WPI (with a DM of 26.6% wt/wt), lactose [solubilized in sterilized osmosis water (130°C, 20 s) at 50% (wt/wt) and heated at 60°C] and minerals were mixed at 50°C under stirring. The mixed solutions were then concentrated to approximately 40% (wt/wt) of DM in a 2-stage semi-industrial scale falling film vacuum evaporator (GEA Process Engineering) with an evaporation capacity of 270 L·h⁻¹ at 50°C. For T⁺⁺⁺, the mixed solution received a heat treatment at 90°C for 2 to 3 s before concentration. After oil blend and vitamin addition at 50°C, and pH adjustment to 6.6 \pm 0.2 (KOH, 45% wt/wt), the mixes were homogenized (50°C, 140/40 bar). For T⁻, the homogenized concentrate was directly spray-dried using a semi-industrial-scale 2-stage spray-dryer (Niro Atomizer, GEA-PE, Saint Quentin en Yvelines, France) at Bionov (Rennes, France), for which maximum theoretical evaporation capacity is approximately 90 kg·h⁻¹. The inlet air temperature was set at 160°C and the outlet air temperature was set at 70°C. The homogenized concentrate flow rate was 100 \pm 10 L·h⁻¹ and the major airflow rate was 2,400 \pm 50 kg/h. Fine particles were recycled at the top of the chamber. For T⁺, the homogenized concentrate received a heat treatment at 75°C for 2 min before spray-drying. For T⁺⁺⁺, the homogenized

concentrate received a heat treatment at 85°C for 2 min before spray-drying. The whole process for producing T⁻, T⁺, and T⁺⁺⁺ was duplicated. An average of 100 kg of IF powder was produced for each formula and replicate. Throughout the IF productions, samples were collected in the middle of mixing (Mix-MID), after vacuum evaporation (Mix-EV), after the addition of oil blend and homogenization (MixL-HO), after heat treatment and before spray-drying (only for T⁺ and T⁺⁺⁺; MixL-HOHT), and after spray-drying (spray dried powder, SDP) at the beginning and at the end of the batch.

Physicochemical Characterization

Dry Matter and Ash Content. The DM and ash contents (g/100 g) were determined in duplicate as described previously (ISO, 1987; Schuck, Dolivet and Jeantet, 2012).

Nitrogen and Protein Contents. Total nitrogen (TN) content, non-CN nitrogen content corresponding to the soluble fraction at pH 4.6, and nonprotein nitrogen (NPN) content corresponding to the soluble fraction after protein precipitation by trichloroacetic acid 12% (wt/vol) were determined by the Kjeldhal method (IDF, 1993). The protein content was determined as TN subtracted of NPN. The specific N conversion factor for bovine milk protein based IF of 6.38 was used, as reported previously (Maubois and Lorient, 2016). The denaturation extent of WP was calculated by comparison of its measured content (g/kg) in the samples collected throughout the IF production or in the reconstituted IF powders with the theoretical WP content (g/kg); the latter being calculated based on the WP content in the ingredients (MSM and WPI) and their mass used for the IF production.

Individual WP Contents. Quantification of individual WP, α -LA, and β -LG, was conducted after precipitation at pH 4.6 using the reversed-phase HPLC method adapted from Resmini et al. (1989), using a PLRP-S column (gel of polystyrene divinylbenzene, 300 Å, 8 μm , 150 \times 3.0 mm) connected to a HPLC System (Dionex UltiMate 3000, Thermo Fisher), equipped with a detector VWD3400RS (Thermo Fisher) operating at 214 nm for the quantification of eluted proteins. Chromatographs were analyzed by Chromeleon software (Version 7.2 SR4, Thermo Fisher). The denaturation extent of individual WP (α -LA or β -LG) was calculated as described above. All measurements were carried out in duplicate.

Total and Free Fat Contents. Total fat content was measured in IF powders by Gerber's acid-butyrometric method (AFNOR, 1990). Free fat content, extracted with petroleum ether, was determined gravimetrically




Figure 1. Semi-industrial process flow diagram for production of 3 types of infant formula powders. Solid arrow = T- (no heat treatment), round dot arrow = T+ (moderate heat treatment), hyphen arrow = T+++ (intense heat treatments).

after solvent evaporation. Total and free fat analyses were carried out in duplicate (Schuck et al., 2012).

Mineral Contents. Sodium, potassium, calcium, magnesium, iron, zinc, and copper in IF powders were quantified by atomic absorption spectrometry as described by Brulé et al. (1974). Chloride was quantified with the Sherwood Scientific chloride analyzer (Model 926, Sherwood Scientific Ltd.) as described by Berdagué et al. (1987). Inorganic phosphorus content was quantified by using the molybdenum blue method according to Jean (1969). All measurements were carried out in duplicate.

Water Activity and Glass Transition Temperature. Water activity (a_w) of IF powders was measured at $25 \pm 0.1^\circ\text{C}$ using the Novasina aw-meter (Novasina, Switzerland). The glass transition temperature (T_g) was determined using a modulated temperature differential scanning calorimetry method according to Schuck et al. (2012). Water activity and T_g measurements were carried out in triplicate.

Rehydration Properties. Dispersibility and solubility of IF powders were determined according to Schuck et al. (2012). All measurements were carried out in duplicate.

Color Parameters. Infant formula powder color parameters (L = brightness; a^* and b^* = chromaticity coordinates) were determined in triplicate with a chromameter (Konica Minolta Photo Imaging France SAS). Browning index (**BI**), an indicator of Maillard reaction extent (Martinez-Alvarenga et al., 2014), was calculated using Equations [1] and [2] (Maskan, 2001):

$$BI = \frac{100 \times (x - 0.31)}{0.17}, \quad [1]$$

with

$$x = \frac{a + 1.75 \times L}{5.645 \times L + a - 3.012 \times b}. \quad [2]$$

Particle Size Distribution. Size distribution of the IF powder particles was determined using a laser light-scattering granulometer (Mastersizer, Malvern Instruments Ltd.). The median diameter $d(0.5)$ was chosen to describe the particle size distribution of the IF powders.

Quantification of Maillard Reaction Products

Furosine. The levels of Amadori products (early Maillard reaction products, such as lactulosylsine) in samples collected throughout the IF production and in IF powders were measured after transformation into

furosine according to ISO (2004) and Giannetti et al. (2013) with some modifications. Samples adjusted to 3.1% (wt/wt) total protein were hydrolyzed with HCl 8 M at 110°C for 23 h under nitrogen. Hydrolyzed samples were purified on a Chromabond 3 mL/500 mg SPE cartridge (Macherey Nagel) using 3 mL of HCl 3 N, after a cartridge conditioning step with 5 mL of methanol and 10 mL of Milli-Q water. The eluate was filtered through a Chromafil xtra 13 mm, 0.2 μm filter and subjected to chromatographic separation. A 25- μL aliquot was injected into the HPLC column (Acclaim Trinity P1, 100 mm \times 3 mm i.d., 3- μm pore size) connected to a Dionex Ultimate 3000 HPLC System (Thermo Fisher), equipped with a detector VWD3400RS (Thermo Fisher) operating at 280 nm for the quantification of eluted furosine. The mobile phase composed of a mixture of acetonitrile (phase A), sodium acetate at 0.4% (vol/vol), pH 4.5 (phase B) and KCl 0.27% (wt/vol) in phase B (phase C) was delivered at a flow rate of 0.5 mL/min at 30°C . The gradient started with 60.0% of mobile phase A and 40.0% of mobile phase B and then continued with linear decrease to 10.0% of phase B and increase to 30.0% of phase C in 10 min. This gradient was maintained for 2 min before being brought back to 40% of phase B in 0.5 min. The gradient was then maintained for 5.5 min before the next injection. Furosine in hydrolyzed samples was quantified using a calibration curve established by direct injection of furosine samples at concentration ranging from 1.5 to 18 mg/L. Chromatographs were analyzed by Chromeleon software (Version 7.2 SR4). All measurements were carried out in triplicate after the hydrolysis step.

N^c-Carboxymethyl-Lysine. Sample preparation for CML analysis was adapted from Niquet-Léridon and Tessier (2011). Samples (of different weights, being equivalent to ~ 10 mg protein) were weighed into 2-mL polypropylene tubes and lyophilized. Their dry weight was then recorded, and the samples reduced by adding 500 μL of 0.1 M NaBH₄. This reduction step (2.5 h at 20°C) is considered necessary to block the Amadori product and avoid de novo formation of CML during subsequent acid hydrolysis. To each tube, 500 μL of 12 M HCl was then added and hydrolysis performed at 110°C for 21 h. Aliquots (200 μL) of hydrolysates were evaporated and resolubilized in 200 μL of 10 mM nonafluoropentanoic acid (**NFPA**). All samples were filtered (0.45 μm) before their analysis by tandem MS coupled to liquid chromatography.

Isotope dilution, HPLC with tandem MS detection analyses were performed in multiple reaction monitoring mode on a Waters Quattro Premier XE instrument with a Heated Electrospray Ionization source (HESI-Waters). A Hypercarb column (100 \times 2.1 mm, 5 μm ; Thermo) with a guard column (10 \times 2.1 mm, 5 μm ,

same phase) was used for chromatographic separation. An ion-pairing, binary mobile phase flow rate was 0.2 mL/min (A – aqueous 10 mM NFFPA, B – acetonitrile; percentage of A: 0–9 min, 100–75%; 9–11 min, 75–40%; 11–13 min, 40%; 13.1–21 min, 100%). The following specific transitions were monitored (in elution order): m/z 205.0 → 130.0 and m/z 207.0 → 130.0 for CML and its isotope, respectively. Quantification of all compounds used the ratio of the analyte peak areas or the peak area of its isotope and comparison with 9-point calibration curves. All measurements were carried out in duplicate from the first step of the protocol.

Bacteriological Analysis

Assessment of the bacteriological quality was performed according to the European regulation (Commission of European Communities, 2007), and completed by additional analyses. Numeration of the total aerobic flora at 30°C and of Enterobacteriaceae was conducted throughout the IF production based on AFNOR (2013, 2017a). Numeration of *Enterobacteriaceae*, *Bacillus cereus*, total coliforms, *Cronobacter* spp., *Listeria monocytogenes*, *Salmonella*, *Staphylococcus aureus*, total aerobic flora, yeasts and molds in IF powders was conducted by an external accredited laboratory (LAB-OCEA, Fougères, France) based on AFNOR (2003a,b, 2005, 2009, 2013, 2017a,b,c, 2018).

Statistical Analysis

Statistical analyses were conducted with R, version 3.3.1 (<https://www.r-project.org>). A nonparametric analysis for repeated measurements was conducted with treatment and production time (and their interaction) as factors, using the nparLD package (Noguchi et al., 2012). For a significant treatment effect ($P < 0.05$), the nparcomp package (Konietschke et al., 2012) was used. For a significant interaction effect, a linear mixed effect model taking into account the repeated measurements was performed and followed by the diffIsmmeans of the lmerTest package (Kuznetsova et al., 2017). Results are expressed as means \pm standard deviation.

RESULTS AND DISCUSSION

Physicochemical Properties of 3 IF Powders

The 3 IF powders, T–, T+ and T+++ , complied with the requirements of regulations (European Commission, 2016) in terms of total content of proteins, lipids, lactose, vitamins, and minerals (Table 1 and Supplemental Table S1, <https://doi.org/10.15454/DW23PE>). The average DM was 97.8 ± 0.1 g/100 g of powder

with an average a_w of 0.19 ± 0.01 (Table 2), close to the optimal value of 0.2 as defined by Efstathiou et al. (2002) to ensure the stability of powders during storage. Lipid stability during storage may also influence the nutritional value and flavor of IF powders. The free fat content of the present IF powders was equal to $2.9 \pm 0.1\%$ (wt/wt) of total fat content, regardless of the heat treatment intensity. This value is below the critical value (5%) reported by Vignolles et al. (2007) for whole milk powder and should be satisfactory for lipid stability. However, this parameter should be monitored during storage. The storage ability of milk powder also depends very much on its Tg (Schuck et al., 2007). The average Tg of the 3 IF powders ($a_w = 0.19$) was $61.2 \pm 0.4^\circ\text{C}$. Tham et al. (2017) reported comparable Tg values for IF at a_w of 0.18, which confirmed a good storage ability of IF powders at 25°C, which should be free from physical changes such as stickiness and caking. The median diameter of the powder grains $d(0.5)$ was $170 \pm 9 \mu\text{m}$, namely in the upper range of nongranulated powders. The 3 IF powders were considered as soluble with a solubility index (SI) higher than 99% (Schuck et al., 2012). With a dispersibility index (DI) ranging between $97.8 \pm 2.0\%$ and $98.9 \pm 0.3\%$, the 3 IF powders could be considered as dispersible. However, it was visually noticeable that T+++ encountered some slight dispersion impairments (flecking) when powder was rehydrated. Concomitantly, a small proportion ($2 \pm 0.4\%$) of large particles ($25\text{--}220 \mu\text{m}$) was observed by laser light diffraction solely for this reconstituted formula (data not shown), particles that were dissociated by addition of an anionic dissociating agent (sodium dodecyl sulfate). Thus, these flecks, although present in limited quantity, could be due to some insoluble protein aggregates, either by themselves or acting as bridges between fat droplets through hydrophobic interaction, thus resulting in fat droplet flocculation, such as previously reported (Schmidmeier et al., 2019; Toikkanen et al., 2018).

Bacteriological Quality Throughout the Production

Microfiltration with 0.8- μm pore size membranes has been used as a physical technique to selectively separate microorganisms from the other components of skim milk (i.e., proteins, lactose, minerals) based on size differences. Our choice of using MF membranes with a smaller pore diameter (0.8 μm instead of 1.4 μm commonly used in dairy industry) was based on previous studies reported by Fauquant et al. (2009) and Tomasula et al. (2011), which indicated that 0.8- μm membrane ensured a higher decimal reduction of microflora and both 2 membranes showed a near complete transmission of proteins. The GP concept applied in

Table 1. Average biochemical composition (\pm SD) of the 3 infant formula (IF) powders over the 2 productions

Item	Unit	Per 100 g of powder			Per 100 kcal			Per 100 kcal	
		T–	T+	T+++	T–	T+	T+++	UE 2016/127	
								Min	Max
Ash	g	3.1 \pm 0.6	3.1 \pm 0.7	3.3 \pm 0.8	0.6 \pm 0.1	0.6 \pm 0.1	0.6 \pm 0.2		
CP ¹	g	11.0 \pm 0.8	11.3 \pm 0.5	10.7 \pm 1.0	2.1 \pm 0.1	2.1 \pm 0.1	2.0 \pm 0.2	1.8	2.5
True protein ²	g	10.8 \pm 0.9	11.1 \pm 0.5	10.5 \pm 1.0	2.0 \pm 0.2	2.1 \pm 0.1	2.0 \pm 0.2		
CN	%	42 \pm 5	41 \pm 6	41 \pm 2					
WP	%	58 \pm 5	59 \pm 6	59 \pm 2					
Lipids	g	29.6 \pm 0.8	30.2 \pm 0.1	30.3 \pm 0.0	5.6 \pm 0.1	5.7 \pm 0.0	5.7 \pm 0.0	4.4	6.0
Lactose ³	g	54.5 \pm 2.2	53.5 \pm 1.1	53.8 \pm 1.8	10.3 \pm 0.5	10.1 \pm 0.1	10.2 \pm 0.3	4.5	14.0
Minerals									
Sodium	mg	167 \pm 10	161 \pm 16	161 \pm 15	32 \pm 2	30 \pm 3	30 \pm 3	25	60
Potassium	mg	705 \pm 4	663 \pm 34	664 \pm 17	134 \pm 1	125 \pm 7	125 \pm 4	80	160
Calcium	mg	355 \pm 19	373 \pm 15	374 \pm 13	67 \pm 2	70 \pm 2	71 \pm 2	50	140
Magnesium	mg	47 \pm 1	47 \pm 0	48 \pm 1	9 \pm 0	9 \pm 0	9 \pm 0	5	15
Chloride	mg	518 \pm 36	492 \pm 10	493 \pm 11	98 \pm 7	93 \pm 1	93 \pm 1	60	160
Phosphorus	mg	256 \pm 7	245 \pm 21	246 \pm 33	49 \pm 2	46 \pm 4	47 \pm 6	25	90
Iron	mg	4.1 \pm 0.1	4.1 \pm 0.2	4.0 \pm 0.3	0.8 \pm 0.0	0.8 \pm 0.0	0.8 \pm 0.0	0.3	1.3
Zinc	mg	4.1 \pm 0.8	4.1 \pm 0.9	4.4 \pm 0.8	0.8 \pm 0.2	0.8 \pm 0.2	0.8 \pm 0.1	0.5	1
Copper	μ g	440 \pm 80	449 \pm 99	410 \pm 81	83 \pm 15	85 \pm 19	78 \pm 16	60	100

¹CP content was expressed as required in the UE 2016/127 regulation (i.e., total N \times 6.25).

²True protein content was determined as total N content subtracted of nonprotein N content, and converted using the specific factor of 6.38.

³Lactose content was determined by calculation: Lactose (g/100 g) = DM (g/100 g) – Ash (g/100 g) – Proteins (g/100 g) – Lipids (g/100 g).

our study has several advantages compared with the uniform transmembrane pressure approach, such as low investment and operating costs due to the removal of a permeate recycle pump. As a consequence, no beads were present on the permeate side of the GP membrane, reducing contamination risks for permeate MSM stream. Moreover, GP membranes include a longitudinal permeability gradient built into the support structure without modification of the filtration layer. This design ensures a stable MF operation with uniform permeate flux (J) and transmembrane pressure (Tomasula et al., 2011; Zulewska and Barbano, 2014).

On the other hand, a liquid WPC obtained by MF with 0.1- μ m pore size membranes, combined with other

filtration techniques, was used instead of cheese whey, a by-product of cheese production, more generally used to re-equilibrate the CN:WP ratio in IF. Separation and purification of WP using membrane filtration preserve WP in their native state and allows to produce a more qualitative product. The production of cheese whey involves enzymatic coagulation, acidification, and, most of the time, heat treatment, which may alter the mineral equilibrium and protein integrity, and produce some by-products such as the glycomacropeptide. The latter is a soluble peptide containing 64 amino acid residues, which is released by the chymosin hydrolysis of κ -casein (Britten and Pouliot, 1996). Its presence in cheese whey alters its amino acid profile. Therefore,

Table 2. Physical properties of the 3 infant formula powders over the 2 productions (average \pm SD)¹

Item ²	T–	T+	T+++
Free fat (% wt/wt total fat)	2.8 \pm 0.3	3.0 \pm 0.6	2.8 \pm 0.3
d(0.5) (μ m)	172.12 \pm 15.43	177.69 \pm 24.82	160.05 \pm 6.53
Water activity	0.20 \pm 0.02	0.19 \pm 0.03	0.19 \pm 0.01
DM (g/100 g of powder)	97.9 \pm 0.1	97.8 \pm 0.0	97.8 \pm 0.1
Tg ($^{\circ}$ C)	61.47 \pm 3.44	61.31 \pm 7.51	60.74 \pm 8.12
Solubility index (%)	>99	>99	>99
Dispersibility index (%)	98.89 \pm 0.36	97.77 \pm 2.01	98.87 \pm 0.25
Color parameters			
L	70.5 \pm 0.4	71.0 \pm 0.4	72.2 \pm 0.0
a*	–3.1 \pm 0.5	–3.2 \pm 0.7	–3.6 \pm 0.7
b*	14.4 \pm 0.7	14.3 \pm 1.6	14.3 \pm 1.3
BI	19.0 \pm 0.8	18.5 \pm 2.0	17.7 \pm 1.4

¹T–: with no heat treatment; T+: with moderate heat treatment; T+++ : with intense heat treatments.

²d(0.5) = median diameter of powder grains; Tg = glass transition temperature; L = brightness; a* and b* = chromaticity coordinates; BI = browning index.

for sectors such as IF, which have stringent quality specifications, WPC produced by filtration is an ideal alternative to cheese whey, which also tends toward a global shortage (Henchion et al., 2017).

Over the 2 productions, the MF with 0.8- μm pore size membranes showed its sufficiency in allowing a reduction of bacterial counts of 3 up to 4 \log_{10} for the total aerobic flora, resulting in a microfiltered milk almost free of bacteria (<10 cfu·g $^{-1}$ for an initial population of $\sim 10^4$ cfu·g $^{-1}$) and totally free of *Enterobacteriaceae*. A more specific experiment of MF with 0.8- μm pore size membranes was conducted with nonpathogenic bacteria having morphological characteristics and cell volume close to *Cronobacter spp* (*Lactobacillus casei* and *Propionibacterium freudenreichii*). This confirmed its efficiency to retain these specific bacteria with a reduction of bacterial counts of 4 to 5 \log_{10} (data not shown).

Regarding the WPC obtained from MF 0.1 μm , its bacteriological quality differed between the 2 productions. Although the first production led to a total flora of 1.15×10^2 cfu·g $^{-1}$ and an *Enterobacteriaceae* population of 1 cfu·g $^{-1}$, higher values were obtained for the second production (1.8×10^4 and 12 cfu·g $^{-1}$ for total flora and *Enterobacteriaceae*, respectively). This was due to postfiltration contamination, which was more important in the second production.

Throughout the 2 IF productions, both heat treatments applied for T+ (75°C, 2 min on the concentrate) and for T+++ (90°C before evaporation, 85°C for 2 min on the concentrate) allowed a minimal reduction of bacteria counts of 2 \log_{10} . No *Enterobacteriaceae* was detected in the collected samples before spray-drying. The spray-drying step resulted in an average increase of 1 \log_{10} , mainly for T-. The T+ IF, spray-dried after T-, inherited the flora of T-, an effect that decreased along the spray-drying process. This effect was less noticeable for T+++; spray-dried after T+ but with a concentrate sent in the tower at a higher temperature (85°C instead of 75°C).

Overall, the 3 IF powders of the first production presented a satisfactory bacteriological quality with a total flora $<10^3$ cfu·g $^{-1}$ and an absence of pathogens, including *Salmonella* and *Cronobacter spp* (Table 3). Powder samples collected at the beginning of the spray-drying of T- presented higher but acceptable level of total flora and molds (1.07×10^3 and <40 cfu·g $^{-1}$, respectively) than those analyzed at the end of process (3.63×10^2 and <10 cfu·g $^{-1}$, respectively). This suggests a residual contamination of the spray-drying tower before drying of T-, which was eliminated with the first amount of T-.

During the second production, a total flora of 5.60×10^3 and 3.57×10^3 cfu·g $^{-1}$ were observed in the powders

Table 3. Bacteriological quality of ingredients and the 3 infant formula powders produced during the first production (T-: with no heat treatment; T+: with a moderate heat treatment; T+++; with intense heat treatments)¹

Item, cfu/g	Raw skim milk	MSM ²	Lactose	WP concentrate	T-		T+		T+++	
					Beginning	End	Beginning	End	Beginning	End
Total flora at 30°C	1.32×10^4 $\pm 5.18 \times 10^3$	6	0	1.15×10^2 ± 2.67	1.07×10^3 $\pm 4.00 \times 10^2$ <10	3.63×10^2 ± 32.1 <10	4.60×10^2 ± 28.3 <10	2.20×10^2 ± 0.00 <10	1.00×10^2 ± 28.3 <10	95 ± 77.8 <10
<i>Bacillus cereus</i>	NA	NA	NA	NA	<10	<10	<10	<10	<10	<10
Total coliforms	NA	NA	NA	NA	<10	<10	<10	<10	<10	<10
<i>Cronobacter spp.</i>	NA	NA	NA	NA	ND/10 g	ND/10 g	ND/10 g	ND/10 g	ND/10 g	ND/10 g
<i>Enterobacteriaceae</i>	1.38×10^2 ± 1.40	0	0	1	ND/10 g	ND/10 g	ND/10 g	ND/10 g	ND/10 g	ND/10 g
<i>Listeria monocytogenes</i>	NA	NA	NA	NA	abs/25 g	abs/25 g	abs/25 g	abs/25 g	abs/25 g	abs/25 g
<i>Salmonella</i>	NA	NA	NA	NA	abs/25 g	abs/25 g	abs/25 g	abs/25 g	abs/25 g	abs/25 g
<i>Staphylococcus aureus</i>	NA	NA	NA	NA	abs/1 g	abs/1 g	abs/1 g	abs/1 g	abs/1 g	abs/1 g
Yeasts	NA	NA	NA	NA	<10	<10	<10	<10	<10	<10
Molds	NA	NA	NA	NA	<40	<10	<10	<10	<10	<10

¹NA = not analyzed; ND/10 g = not detected in 10 g; abs/25 g. = absence in 25 g; abs/1 g. = absence in 1 g. MSM = microfiltered skim milk; WP = whey protein.

of T⁻ and T⁺ with a presence of *Enterobacteriaceae* (73.3 and 10 cfu·g⁻¹, respectively), due to the lower quality of the WP concentrate resulting from post-filtration contamination. The T⁺ powder could have reached a satisfactory quality if dried independently of T⁻, but this was not possible with the present experimental design. During this second production, only T⁺⁺⁺ showed a satisfactory bacteriological quality with a total flora <3.00 × 10² cfu·g⁻¹ and an absence of *Enterobacteriaceae*.

Overall, we can conclude that integrating MF of fresh milk into IF production can be efficient to ensure the bacteriological safety of the ingredients without additional heat treatments; however, a high microbiological quality of the ingredients (*Cronobacter*- and *Enterobacteriaceae*-free ingredients); of the processing water, which can be sterilized to ensure its microbiological safety; and of the production environment, by a thorough cleaning, is required. The present experiments were performed at a semi-industrial scale on a workshop pilot not specifically designed for the different successive steps involved, which were thus carried out in an open space. In industry, such production would be performed with a well-adapted continuous processing line, in a controlled environment and with the benefits of continuous improvement process. Such production frame should avoid any postfiltration contamination.

Denaturation Extent of WP

The denaturation extent of WP was followed throughout the production of the 3 types of IF powders by determining the proportion of whey proteins insoluble at pH 4.6, as shown in Figure 2. For T⁺⁺⁺, the effect of the double pasteurization (72°C, 30 s) of milk on protein integrity was minimal (denaturation extent <4%). Such HTST pasteurization is widely used in the dairy industry to inactivate pathogenic microorganisms and is proved to cause low levels of denaturation for major WP (α-LA and β-LG) in the milk (Rynne et al., 2004; Rohitha Prasantha and Wimalasiri, 2019; Halabi et al., 2020b). However, HTST can lead to denaturation of some minor proteins, such as lactoferrin, and does not ensure the inactivation of endogenous enzymes, such as plasmin (Rauh et al., 2014; Klotz et al., 2017; Wang et al., 2019). It is noteworthy that, unlike HTST alone, MF allows an efficient removal of somatic cells present in milk containing numerous thermo-resistant enzymes such as plasmin, which can cause proteolysis and thus, off-flavors (Wang et al., 2019).

Regarding vacuum evaporation (0.3 bar, 270 l/h, 50°C) of the mixed solution, this had no effect on the denaturation of WP (from Mix-MID to Mix-EV) in T⁻ and T⁺. The higher denaturation extent for T⁺⁺⁺ in

Mix-EV was therefore only due to the heat treatment at 90°C for 2 to 3 s between mixing and evaporation steps; the extent of WP denaturation increased 26 ± 6%. The oil and vitamin addition followed by the homogenization step (140/40 bar, 50°C) had, as expected, virtually no influence on the denaturation of WP (from Mix-EV to MixL-HO). The final heat treatment of the concentrated mix doubled the denaturation extent of WP in both T⁺ and T⁺⁺⁺ (75°C for 2 min and 85°C for 2 min, respectively), although this did not reach statistical significance due to a lack of statistical power. Similar results were reported by previous studies (Singh and Creamer, 1991; Morgan et al., 1999; Mehta and Deeth, 2016) showing that the key factor for protein denaturation during the production of powdered dairy products is the intensity of heat treatments rather than the drying process itself. In addition, it can be noticed that the highest heat treatment (85°C for 2 min applied to the concentrated mix) induced significant whey protein gelation, resulting in pipe clogging at this end of one of the replicates; this never occurred for the production of T⁻ and T⁺. The denaturation extent of WP tended to decrease for T⁺⁺⁺ after spray-drying, although not statistically significant. This can be due to a lack of representativeness of the T⁺⁺⁺ sample at




Figure 2. Average denaturation extent (±SD, n = 2) of whey proteins over the 2 productions of the infant formulas powders. T⁻: no additional heat treatment; T⁺: moderate heat treatment at 75°C for 2 min, applied before spray-drying; T⁺⁺⁺: successive heat treatments at 72°C for 30 s on the milk, 90°C for 2–3 s before evaporation, 85°C for 2 min before spray-drying; Mix-MID: middle point of mixing; Mix-EV: mixture after evaporation; MixL-HO: mixture with lipids after homogenization; MixL-HOHT: mixture with lipids after homogenization and heat treatment (only for T⁺ and T⁺⁺⁺); SDP: spray-dried powder. Statistics were calculated using a nonparametric analysis for repeated measurement. ***P < 0.001, **P < 0.01, NS: P ≥ 0.05. Different letters indicate statistically significant differences between treatments or between time (P < 0.05).

the MixL-HOHT stage or to some loss of protein aggregates during the process, although this was very limited during spray-drying. The final IF powders produced according to the present processing routes presented a low denaturation extent for T⁻ and T⁺ ($6 \pm 4\%$ and $10 \pm 4\%$, respectively) and a high denaturation extent for T⁺⁺⁺ ($58 \pm 0\%$). The latter value was in accordance with the denaturation extent determined in 4 commercial IF powders ($65.0 \pm 11.1\%$). Compared with T⁻ without any heat treatment, T⁺ might be of particular interest for food safety at the industrial level because the pasteurization step before spray-drying permitted to eliminate pathogens including potential coronaviruses, such as SARS CoV-2 (Duan et al., 2003) or tick-borne encephalitis, which can be present in bovine milk from Central and Eastern Europe and in Asia (Saier et al., 2015), thus further guaranteeing the microbiological quality of the powders without leading to a high protein denaturation extent.

First-age IF is administered to infants as a human milk substitute and thus should mimic this nutritional gold standard as much as possible. The present IF with a low denaturation extent contains mainly native proteins, such as in a raw human milk. Thus, such a minimally processing route can help to develop more biomimetic IF. To get even closer to human milk, the WP profile could also be improved, for instance by the addition of bioactive proteins such as lactoferrin. The present processing would be of particular interest as this protein needs to remain in its native form to preserve its bioactive properties.

Denaturation Extent of Individual WP (α -LA or β -LG)

The denaturation extent of individual WP (α -LA or β -LG) during the production of the 3 types of IF is presented in Figure 3. Heat treatment at 90°C for 2 to 3 s between mixing and evaporation steps induced a loss of native β -LG of $29 \pm 10\%$ for T⁺⁺⁺, whereas no significant loss of native α -LA was observed. A study by Halabi et al. (2020b) in the context of IF also showed that the decrease of the native WP after heat treatment at 90°C for 15 s was mainly due to the heat denaturation of β -LG. At temperatures above 68°C , β -LG undergoes conformational changes and partially unfolds, exposing its 2 disulfide bridges (Cys66-Cys160 and Cys106-Cys119) and a free sulfhydryl group (-SH group; Cys121). When heated for a short time, the reshuffling of intramolecular disulfide bonds may be responsible for the formation of irreversible non-native monomers of β -LG that can further aggregate (Croguenec et al., 2003; Kleber and Hinrichs, 2007; Wijayanti et al., 2014; Brodkorb et al., 2016; Nielsen et al., 2018), whereas α -LA, without free-SH group, is more heat sta-

ble at this point (Mulvihill and Donovan, 1987; Calvo et al., 1993). Heat treatment at 75°C for 2 min had very little influence on the denaturation of α -LA and β -LG for T⁺, whereas heat treatment at 85°C for 2 min for T⁺⁺⁺ induced a denaturation extent of $59 \pm 5\%$ for α -LA and $41 \pm 8\%$ for β -LG. When heated for a longer time (2 min), although intramolecular reshuffling cannot occur with α -LA because of absence of free -SH group, the intermolecular reshuffling is more active for α -LA than for β -LG because the former possesses more disulfide bonds, leading to higher denaturation extent of α -LA (Wijayanti et al., 2014). As observed for the overall denaturation extent, the denaturation extent of α -LA and β -LG decreased by $26 \pm 8\%$ and $14 \pm 3\%$ for T⁺⁺⁺ after spray-drying.

Effect of Heat Treatment on the Formation of Maillard Reaction Products

The formation of Amadori products, measured as furosine, during the production of the 3 types of IF is presented in Figure 4. The furosine concentration doubled after the addition of vitamins, oil, and pH adjustment




Figure 3. Average denaturation extent (\pm SD, $n = 2$) of α -lactalbumin (α -LA) and β -lactoglobulin (β -LG) over the 2 productions of the infant formula powders. T⁻: no additional heat treatment; T⁺: moderate heat treatment at 75°C for 2 min, applied before spray-drying; T⁺⁺⁺: successive heat treatments at 72°C for 30 s on the milk, 90°C for 2–3 s before evaporation, 85°C for 2 min before spray-drying; Mix-MID: middle point of mixing; Mix-EV: mixture after evaporation; MixL-HO: mixture with lipids after homogenization; MixL-HOHT: mixture with lipids after homogenization and heat treatment (only for T⁺ and T⁺⁺⁺); SDP: spray-dried powder. Statistics were calculated using a nonparametric analysis for repeated measurement. *** $P < 0.001$, ** $P < 0.01$, NS: $P \geq 0.05$. Different letters indicate statistically significant differences between treatments or between time ($P < 0.05$).

followed by homogenization (140/40 bar, 50°C) for all the 3 IF. Vitamin C can act as precursor or intermediate for the formation of Maillard reaction products (Pischetsrieder et al., 2005; Troise et al., 2016). After encapsulation of vitamin C, an increase of 45% of available lysine and a decrease of 10 to 53% in furosine and CML concentrations were observed in milk (Troise et al., 2016). In addition, pH adjustment of the mix from acidic pH after the addition of vitamins to more neutral values (pH from 6.02 ± 0.08 to 6.6 ± 0.2) after addition of potassium hydroxide can also accelerate the initial step of the Maillard reaction by increasing the nucleophilicity of protein amino groups (Lund and Ray, 2017; Nunes et al., 2019). Heat treatment at 75°C for 2 min for T+ and 85°C/2 min for T+++ made their furosine concentration doubled again from 255 ± 18 to 445 ± 208 mg/100 g MAT and from 249 ± 36 to 582 ± 170 mg/100 g MAT. In contrast, the 3 IF powders reached similar furosine concentrations after spray-drying, in accordance with previous studies, in which Henle et al. (1995) reported furosine concentration from 930 to 1,890 mg/100 g of proteins in commercial IF powders determined by isocratic ion-pair reversed-phase HPLC method. Lower values have been reported by Birlouez-Aragon et al. (2004) using ion-pair HPLC, ranging from 170 to 370 mg/100 g of proteins in 17 commercial IF powders, possibly due to a higher rate of transformation of the Amadori products into advanced products of the Maillard reaction.

Concomitantly to the evolution of furosine, the CML concentration doubled after the addition of vitamins, oil, pH adjustment followed by homogenization, and after heat treatments of the concentrate before spray-drying for T+ and T+++ (Figure 4). After spray-drying, the CML concentration was numerically lower in powder T- than in powder T+++ (8.8 ± 0.2 and 10.6 ± 0.0 mg/100 g MAT, respectively), but this did not reach statistical significance due to the variability observed along the entire production and due to the limited number of production replicate ($n = 2$). Contrary to the WP denaturation extent, which was more reproducible, a high variability was observed regarding furosine and CML between the 2 IF productions, particularly from the step of vitamin and oil addition, pH adjustment, and homogenization until spray-drying. The Maillard reaction products are likely to be more affected by some slight difference of the production parameters (pH, temperature) inherent to the semi-industrial level. Nevertheless, the present values are in the range of those previously reported by Delatour et al. (2009), who analyzed the CML concentrations in 7 commercial IF powders using the same technique as in the present study and indicated an average value of 7.6 mg/100 g MAT.




Figure 4. Average furosine (as a surrogate marker of the Amadori products) and *N*-ε-carboxymethyl-lysine (CML) concentration (\pm SD, $n = 2$) over the 2 productions of the infant formula powders. T-: no additional heat treatment; T+: moderate heat treatment at 75°C for 2 min, applied before spray-drying; T+++ : successive heat treatments at 72°C for 30 s on the milk, 90°C for 2–3 s before evaporation, 85°C for 2 min before spray-drying; Mix-MID: middle point of mixing; Mix-EV: mixture after evaporation; MixL-HO: mixture with lipids after homogenization; MixL-HOHT: mixture with lipids after homogenization and heat treatment (only for T+ and T+++); SDP: spray-dried powder. Statistics were calculated using a nonparametric analysis for repeated measurement. *** $P < 0.001$, ** $P < 0.01$, NS: $P \geq 0.05$. Different letters indicate statistically significant differences between treatments or between time ($P < 0.05$).

The concentration of the Maillard reaction products of the 3 IF powders was in accordance with their BI, one of the first indicators of the extent of the Maillard reaction. The latter was in the same range among the 3 IF powders, varying from 17.7 up to 19 (Table 2). This value was related to a pale yellow color, which was induced not only by the Maillard products but also by the yellow-color vitamin mix.

IF powders are products devoted to be stored for several months, during which the Maillard reaction products may increase. To this end, these compounds will be followed during the storage step and will be presented in a future study.

The present study presents some limitations. First, due to the scale of the production, only 2 replicates could be performed, which limits the statistical power. The present study has demonstrated that a native IF powder could be bacteriologically safe, however this does not guarantee viral decontamination, which, if suspected, could be performed by the moderate heat

treatment, depending on the virus. Further validation is required concomitantly to the development of viral analyses in dairy products. Finally, the viscosity of the product along IF processing could not be followed, however no problem of processability was encountered except for T+++ , such as stated above.

CONCLUSIONS

This study demonstrated the feasibility of producing, at semi-industrial scale, a bacteriologically safe IF powder with a high content of native proteins, based on membrane filtration of fresh milk, low temperature unit operations, and high level of bacteriological quality of the ingredients. Adding a moderate heat treatment step (75°C, 2 min) on the concentrate before spray-drying could further guarantee the microbiological quality of the IF powder, if dried independently, without leading to a high protein denaturation extent. The absence of heat treatment on the IF tended to limit the formation of advanced Maillard reaction products (CML); however, the extent of the difference was much lower on these products than on the WP denaturation extent. This study opens up new avenues for the production of minimally processed IF powders, although the nutritional and physiological impacts of such IF remain unknown. Such minimally processing route can be highly relevant for IF including heat-sensitive bioactive components, such as lactoferrin.

ACKNOWLEDGMENTS

This research project was funded by the institute Carnot Qualiment. We acknowledge A. Dolivet, F. Lambrouin, A. Leduc, J.-L. Thomas (STLO, INRAE, Institut Agro, Rennes, France), and G. Le Maout (Binov SA, Rennes, France) for their technical support. The authors have not stated any conflicts of interest.

REFERENCES

- Abbring, S., B. R. J. Blokhuis, J. L. Miltenburg, K. G. J. R. Olmedo, J. Garssen, F. A. Redegeld, and B. C. A. M. van Esch. 2020. Direct inhibition of the allergic effector response by raw cow's milk—An extensive in vitro assessment. *Cells* 9:1258. <https://doi.org/10.3390/cells9051258>.
- AFNOR. 1990. Milk – Determination of Fat Content – Acid-butyrometric Method. (NF V04-210). French Standardization Association.
- AFNOR. 2003a. Food Microbiology – Horizontal Method for the Enumeration of Coagulase-Positive Staphylococci (*Staphylococcus Aureus* and Other Species) – Part 3: Search and MPN Method for Low Numbers. (NF EN ISO 6888-3). French Standardization Association.
- AFNOR. 2003b. Food microbiology – Horizontal Method for the Enumeration of Yeasts and Molds Growing on a Medium with Low a_w. (NF V08-036). French Standardization Association.
- AFNOR. 2005. Food Microbiology – Horizontal Method for the Enumeration of Presumptive *Bacillus cereus* – Colony Counting Technique at 30°C. (NF EN ISO 7932). French Standardization Association.
- AFNOR. 2009. Food Microbiology – Enumeration of Suspected Coliforms by Counting the Colonies Obtained at 30°C. (NF V08-050). French Standardization Association.
- AFNOR. 2013. Food Microbiology – Horizontal Method for the Enumeration of Microorganisms – Part 1: Colony Count at 30°C by the Deep Sowing Technique. (NF EN ISO 4833-1). French Standardization Association.
- AFNOR. 2017a. Microbiology of the Food Chain – Horizontal Method for the Search and Enumeration of *Enterobacteriaceae* – Part 1: Search for *Enterobacteriaceae*. (NF EN ISO 21528-1). French Standardization Association.
- AFNOR. 2017b. Microbiology of the Food Chain – Horizontal method for the detection of Cronobacter spp. (NF EN ISO 22964). French Standardization Association.
- AFNOR. 2017c. ALOA One Day Method for the Detection of *Listeria monocytogenes* and *Listeria* spp in Food Products and Environmental Samples. (AES 10/3-09/00). French Standardization Association.
- AFNOR. 2018. RAPID/Salmonella: Validated for the Detection of *Salmonella* spp. (BRD 07/11-12/05). French Standardization Association.
- Baskara, I., C. Niquet-Leridon, P. M. Anton, and C. Delayre-Orthez. 2017. Neoformed compounds from the maillard reaction in infant formulas: A new risk factor for allergy? *EMJ Allergy Immunol.* 2:87-93.
- Berdagué, J. L., R. Grappin, and G. Duboz. 1987. Affinage et qualité du Gruyère de Comté. II. Influence de l'affinage sur l'évolution des caractéristiques physico-chimiques des fromages. *Le Lait.* INRA Ed. 67:237-247.
- Birlouez-Aragon, I., M. Pischetsrieder, J. Leclère, F. J. Morales, K. Hasenkopf, R. Kientsch-Engel, C. J. Ducauze, and D. Rutledge. 2004. Assessment of protein glycation markers in infant formulas. *Food Chem.* 87:253-259. <https://doi.org/10.1016/j.foodchem.2003.11.019>.
- Britten, M., and Y. Pouliot. 1996. Characterization of whey protein isolate obtained from milk microfiltration permeate. *Le Lait.* INRA Ed. 76:255-265.
- Brodkorb, A., T. Croguennec, S. Bouhallab, and J. J. Kehoe. 2016. Heat-induced denaturation, aggregation and gelation of whey proteins. Pages 155-178 in *Advanced Dairy Chemistry: Volume 1B: Proteins: Applied Aspects*. P. L. H. McSweeney and J. A. O'Mahony, ed. Springer.
- Brulé, G., J. L. Maubois, and J. Fauquant. 1974. Étude de la teneur en éléments minéraux des produits obtenus lors de l'ultrafiltration du lait sur membrane. *Lait* 54:600-615. <https://doi.org/10.1051/lait:1974539-54030>.
- Calvo, M. M., J. Leaver, and J. M. Banks. 1993. Influence of other whey proteins on the heat-induced aggregation of α-lactalbumin. *Int. Dairy J.* 3:719-727. [https://doi.org/10.1016/0958-6946\(93\)90085-E](https://doi.org/10.1016/0958-6946(93)90085-E).
- Commission of the European Communities. 2007. Commission Regulation (EC) No 1441/2007. Amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs. *Off. J. Eur. Union* 332:12-29.
- Croguennec, T., S. Bouhallab, D. Mollé, B. T. O'Kennedy, and R. Mehra. 2003. Stable monomeric intermediate with exposed Cys-119 is formed during heat denaturation of beta-lactoglobulin. *Biochem. Biophys. Res. Commun.* 301:465-471. [https://doi.org/10.1016/s0006-291x\(02\)02997-2](https://doi.org/10.1016/s0006-291x(02)02997-2).
- Delatour, T., J. Hegele, V. Parisod, J. Richo, S. Maurer, M. Steven, and T. Buetler. 2009. Analysis of advanced glycation endproducts in dairy products by isotope dilution liquid chromatography-electrospray tandem mass spectrometry. The particular case of carboxymethyllysine. *J. Chromatogr. A* 1216:2371-2381. <https://doi.org/10.1016/j.chroma.2009.01.011>.
- Duan, S.-M., X.-S. Zhao, R.-F. Wen, J.-J. Huang, G.-H. Pi, S.-X. Zhang, J. Han, S.-L. Bi, L. Ruan, and X. P. Dong. 2003. Stability

- of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomed. Environ. Sci.* 16:246–255.
- Dupont, D., G. Mandalari, D. Mollé, J. Jardin, O. Rolet-Répécaud, G. Duboz, J. Léonil, C. E. N. Mills, and A. R. Mackie. 2010. Food processing increases casein resistance to simulated infant digestion. *Mol. Nutr. Food Res.* 54:1677–1689. <https://doi.org/10.1002/mnfr.200900582>.
- Efstathiou, T., C. Feuadent, S. Méjean, and P. Schuck. 2002. The use of carbonyl analysis to follow the main reactions involved in the process of deterioration of dehydrated dairy products: Prediction of most favourable degree of dehydration. *Lait.* 82:423–439. <https://doi.org/10.1051/lait:2002021>.
- Elmhiri, G., D. F. D. Mahmood, C. Niquet-Leridon, P. Jacolot, S. Firmin, L. Guigand, F. J. Tessier, T. Larcher, and L. Abdennebi-Najar. 2015. Formula-derived advanced glycation end products are involved in the development of long-term inflammation and oxidative stress in kidney of IUGR piglets. *Mol. Nutr. Food Res.* 59:939–947. <https://doi.org/10.1002/mnfr.201400722>.
- European Commission. 2016. Commission directive 2016/127/EC of 25 September 2015 on infant formulas and follow-on formulas and completed regulation n° 609/2013 and amending directive 2006/141/EC. *Off. J. Eur. Union* 25:30–43.
- Fauquant, J., E. Beaucher, C. Sinet, B. Robert, and C. Lopez. 2014. Combination of homogenization and cross-flow microfiltration to remove microorganisms from industrial buttermilks with an efficient permeation of proteins and lipids. *Innov. Food Sci. Emerg. Technol.* 21:131–141. <https://doi.org/10.1016/j.ifset.2013.10.004>.
- Fauquant, J., B. Robert and C. Lopez. 2009. Procédé pour réduire la teneur bactérienne d'un milieu alimentaire et/ou biologique d'intérêt, contenant des gouttelettes lipidiques. Brevet INRA FR 2 953 686, 2009.
- Gésan-Guizou, G., E. Boyaval, and G. Daufin. 1999. Critical stability conditions in crossflow microfiltration of skimmed milk: Transition to irreversible deposition. *J. Membr. Sci.* 158:211–222. [https://doi.org/10.1016/S0376-7388\(99\)00017-4](https://doi.org/10.1016/S0376-7388(99)00017-4).
- Giannetti, V., M. B. Mariani, and P. Mannino. 2013. Furosine as a pasta quality marker: Evaluation by an innovative and fast chromatographic approach. *J. Food Sci.* 78:C994–C999. <https://doi.org/10.1111/1750-3841.12163>.
- Golkar, A., J. M. Milani, and T. Vasiljevic. 2019. Altering allergenicity of cow's milk by food processing for applications in infant formula. *Crit. Rev. Food Sci. Nutr.* 59:159–172. <https://doi.org/10.1080/10408398.2017.1363156>.
- Halabi, A., T. Croguennec, S. Bouhallab, D. Dupont, and A. Deglaire. 2020a. Modification of protein structures by altering the whey protein profile and heat treatment affects *in vitro* static digestion of model infant milk formulas. *Food Funct.* 11:6933–6945. <https://doi.org/10.1039/D0FO01362E>.
- Halabi, A., A. Deglaire, P. Hamon, S. Bouhallab, D. Dupont, and T. Croguennec. 2020b. Kinetics of heat-induced denaturation of proteins in model infant milk formulas as a function of whey protein composition. *Food Chem.* 302:125296. <https://doi.org/10.1016/j.foodchem.2019.125296>.
- Henchion, M., M. Hayes, A. M. Mullen, M. Fenelon, and B. Tiwari. 2017. Future protein supply and demand: Strategies and factors influencing a sustainable equilibrium. *Foods* 6:53. <https://doi.org/10.3390/foods6070053>.
- Henle, T., G. Zehetner, and H. Klostermeyer. 1995. Fast and sensitive determination of furosine. *Z. Lebensm. Unters. Forsch.* 200:235–237. <https://doi.org/10.1007/BF01190503>.
- IDF (International Dairy Federation). 1993. IDF Standard 20B:1993. Milk: Determination of nitrogen content. IDF.
- ISO (International Organization for Standardization). 1987. ISO 6731:1989. Milk, cream and evaporated milk—Determination of total solids content (reference method). ISO.
- ISO (International Organization for Standardization) 2004. ISO18329:2004 Lait et produits laitiers—Détermination de la teneur en furosine – Méthode par chromatographie liquide à haute performance en phase inverse par paire d'ions. ISO.
- Jean, P. 1969. Dosage du phosphore dans le lait. *Le Lait.* INRA Ed. 49:175–188.
- Joyce, A. M., A. Brodkorb, A. L. Kelly, and J. A. O'Mahony. 2017. Separation of the effects of denaturation and aggregation on whey-casein protein interactions during the manufacture of a model infant formula. *Dairy Sci. Technol.* 96:787–806.
- Kleber, N., and J. Hinrichs. 2007. Antigenic response of β -lactoglobulin in thermally treated bovine skim milk and sweet whey. *Milchwissenschaft* 62:121–124.
- Klotz, D., M. Joellenbeck, K. Winkler, M. Kunze, D. Huzly, and R. Hentschel. 2017. High-temperature short-time pasteurisation of human breastmilk is efficient in retaining protein and reducing the bacterial count. *Acta Paediatr.* 106:763–767. <https://doi.org/10.1111/apa.13768>.
- Konietschke, F., L. A. Hothorn, and E. Brunner. 2012. Rank-based multiple test procedures and simultaneous confidence intervals. *Electron. J. Stat.* 6:738–759. <https://doi.org/10.1214/12-EJS691>.
- Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in linear mixed effects models. *J. Stat. Softw.* 82:1–26. <https://doi.org/10.18637/jss.v082.i13>.
- Lacroix, M., C. Bon, C. Bos, J. Léonil, R. Benamouzig, C. Luengo, J. Fauquant, D. Tomé, and C. Gaudichon. 2008. Ultra high temperature treatment, but not pasteurization, affects the postprandial kinetics of milk proteins in humans. *J. Nutr.* 138:2342–2347. <https://doi.org/10.3945/jn.108.096990>.
- Lund, M. N., and C. A. Ray. 2017. Control of Maillard reactions in foods: Strategies and chemical mechanisms. *J. Agric. Food Chem.* 65:4537–4552. <https://doi.org/10.1021/acs.jafc.7b00882>.
- Madeç, M. N., S. Méjean, and J. L. Maubois. 1992. Retention of Listeria and Salmonella cells contaminating skim milk by tangential membrane microfiltration (“Bactocatch” process). *Dairy Sci. Technol. (Lait)* 72:327–332.
- Martinez-Alvarenga, M. S., E. Y. Martinez-Rodriguez, L. E. Garcia-Amezquita, G. I. Olivas, P. B. Zamudio-Flores, C. H. Acosta-Muniz, and D. R. Sepulveda. 2014. Effect of Maillard reaction conditions on the degree of glycation and functional properties of whey protein isolate – Maltodextrin conjugates. *Food Hydrocoll.* 38:110–118. <https://doi.org/10.1016/j.foodhyd.2013.11.006>.
- Maskan, M. 2001. Kinetics of colour change of kiwifruits during hot air and microwave drying. *J. Food Eng.* 48:169–175. [https://doi.org/10.1016/S0260-8774\(00\)00154-0](https://doi.org/10.1016/S0260-8774(00)00154-0).
- Maubois, J. L., and D. Lorient. 2016. Dairy proteins and soy proteins in infant foods nitrogen-to-protein conversion factors. *Dairy Sci. Technol.* 96:15–25. <https://doi.org/10.1007/s13594-015-0271-0>.
- Mehta, B. M., and H. C. Deeth. 2016. Blocked lysine in dairy products: Formation, occurrence, analysis, and nutritional implications. *Compr. Rev. Food Sci. Food Saf.* 15:206–218. <https://doi.org/10.1111/1541-4337.12178>.
- Morgan, F., J. Léonil, D. Mollé, and S. Bouhallab. 1999. Modification of bovine beta-lactoglobulin by glycation in a powdered state or in an aqueous solution: effect on association behavior and protein conformation. *J. Agric. Food Chem.* 47:83–91. <https://doi.org/10.1021/jf9804387>.
- Mulvihill, D. M., and M. Donovan. 1987. Whey proteins and their thermal denaturation - A review. *Irish J. Food Sci. Technol.* 11:43–75.
- Nielsen, L. R., M. N. Lund, M. J. Davies, J. H. Nielsen, and S. B. Nielsen. 2018. Effect of free cysteine on the denaturation and aggregation of holo α -lactalbumin. *Int. Dairy J.* 79:52–61. <https://doi.org/10.1016/j.idairyj.2017.11.014>.
- Niquet-Léridon, C., and F. J. Tessier. 2011. Quantification of N-carboxymethyl-lysine in selected chocolate-flavored drink mixes using high-performance liquid chromatography-linear ion trap tandem mass spectrometry. *Food Chem.* 126:655–663. <https://doi.org/10.1016/j.foodchem.2010.10.111>.
- Noguchi, K., Y. R. Gel, E. Brunner, and F. Konietschke. 2012. nparLD: An R Software Package for the Nonparametric Analysis of Longitudinal Data in Factorial Experiments. *J. Stat. Softw.* 50:1–23. <https://doi.org/10.18637/jss.v050.i12>.
- Nunes, L., E. Martins, I. Tuler Perrone, and A. Fernandes de Carvalho. 2019. The Maillard reaction in powdered infant formula. *J. Food Nutr. Res.* 7:33–40. <https://doi.org/10.12691/jfnr-7-1-5>.

- Peram, M. R., S. M. Loveday, A. Ye, and H. Singh. 2013. In vitro gastric digestion of heat-induced aggregates of β -lactoglobulin. *J. Dairy Sci.* 96:63–74. <https://doi.org/10.3168/jds.2012-5896>.
- Pischetsrieder, M., B. Larisch, and T. Severin. 2005. The Maillard reaction of ascorbic acid with amino acids and proteins – Identification of products. Pages 107–112 in *The Maillard Reaction in Foods and Medicine*. J. O'Brien, H. E. Nursten, M. J. C. Crabbe and J. M. Ames, ed. Woodhead Publishing. <https://doi.org/10.1533/9781845698447.2.107>.
- Rauh, V. M., L. B. Johansen, R. Ipsen, M. Paulsson, L. B. Larsen, and M. Hammershøj. 2014. Plasmin activity in UHT milk: Relationship between proteolysis, age gelation, and bitterness. *J. Agric. Food Chem.* 62:6852–6860. <https://doi.org/10.1021/jf502088u>.
- Resmini, P., L. Pellegrino, R. Andreini, and F. Prati. 1989. Determinazione delle sieroproteine solubili del latte per HPLC (cromatografia liquida ad alta prestazione) in fase inversa. *Sci. E. Technica Latterio-Casearia* 40:7–23.
- Rohitha Prasantha, B. D., and K. M. S. Wimalasiri. 2019. Effect of HTST thermal treatments on end-use quality characteristics of goat milk. *Int. J. Food Sci.* 2019:1801724. <https://doi.org/10.1155/2019/1801724>.
- Roth-Walter, F., M. C. Berin, P. Arnaboldi, C. R. Escalante, S. Dahan, J. Rauch, E. Jensen-Jarolim, and L. Mayer. 2008. Pasteurization of milk proteins promotes allergic sensitization by enhancing uptake through Peyer's patches. *Allergy* 63:882–890. <https://doi.org/10.1111/j.1398-9995.2008.01673.x>.
- Rynne, N. M., T. P. Beresford, A. L. Kelly, and T. P. Guinee. 2004. Effect of milk pasteurization temperature and in situ whey protein denaturation on the composition, texture and heat-induced functionality of half-fat cheddar cheese. *Int. Dairy J.* 14:989–1001. <https://doi.org/10.1016/j.idairyj.2004.03.010>.
- Saboya, L. V., and J.-L. Maubois. 2000. Current developments of microfiltration technology in the dairy industry. *Lait* 80:541–553. <https://doi.org/10.1051/lait:2000144>.
- Saier, R., G. Maier, Z. Atamer, and J. Hinrichs. 2015. Thermal inactivation of tickborne encephalitis virus in milk. *Int. J. Dairy Technol.* 68:366–373. <https://doi.org/10.1111/1471-0307.12230>.
- Schmidmeier, C., C. O'Gorman, K. P. Drapala, D. Waldron, and J. O'Mahony. 2019. Elucidation of factors responsible for formation of white flecks in reconstituted fat filled milk powders. *Colloids Surf. A Physicochem. Eng. Asp.* 575:245–255. <https://doi.org/10.1016/j.colsurfa.2019.03.034>.
- Schmidt, V. S. J., V. Kaufmann, U. Kulozik, S. Scherer, and M. Wenning. 2012. Microbial biodiversity, quality and shelf life of microfiltered and pasteurized extended shelf life (ESL) milk from Germany, Austria and Switzerland. *Int. J. Food Microbiol.* 154:1–9. <https://doi.org/10.1016/j.ijfoodmicro.2011.12.002>.
- Schuck, P., A. Dolivet, and R. Jeantet. 2012. *Analytical Methods for Food and Dairy Powders*. Wiley- Blackwell. <https://doi.org/10.1002/9781118307397>.
- Schuck, P., S. Mejean, A. Dolivet, R. Jeantet, and B. Bhandari. 2007. Keeping quality of dairy ingredients. *Lait* 87:481–488.
- Singh, H., and L. K. Creamer. 1991. Denaturation, aggregation and heat stability of milk protein during the manufacture of skim milk powder. *J. Dairy Res.* 58:269–283. <https://doi.org/10.1017/S002202990002985X>.
- Takagi, K., R. Teshima, H. Okunuki, and J. Sawada. 2003. Comparative study of in vitro digestibility of food proteins and effect of preheating on the digestion. *Biol. Pharm. Bull.* 26:969–973. <https://doi.org/10.1248/bpb.26.969>.
- Tham, T. W. Y., A. T. H. Yeoh, and W. Zhou. 2017. Characterisation of aged infant formulas and physicochemical changes. *Food Chem.* 219:117–125. <https://doi.org/10.1016/j.foodchem.2016.09.107>.
- Toikkanen, O., M. Outinen, L. Malafronte, and O. J. Rojas. 2018. Formation and structure of insoluble particles in reconstituted model infant formula powders. *Int. Dairy J.* 82:19–27. <https://doi.org/10.1016/j.idairyj.2018.03.001>.
- Tomasula, P. M., S. Mukhopadhyay, N. Datta, A. Porto-Fett, J. E. Call, J. B. Luchansky, J. Renye, and M. Tunick. 2011. Pilot-scale crossflow-microfiltration and pasteurization to remove spores of *Bacillus anthracis* (Sterne) from milk. *J. Dairy Sci.* 94:4277–4291. <https://doi.org/10.3168/jds.2010-3879>.
- Troise, A. D., D. Vitiello, C. Tsang, and A. Fiore. 2016. Encapsulation of ascorbic acid promotes the reduction of Maillard reaction products in UHT milk. *Food Funct.* 7:2591–2602. <https://doi.org/10.1039/C6FO00151C>.
- Trouvé, E., J. L. Maubois, M. Piot, M. N. Madec, J. Fauquant, A. Rouault, J. Tabard, and G. Brinkman. 1991. Rétention de différentes espèces microbiennes lors de l'épuration du lait par microfiltration en flux tangentiel. *Dairy Sci. Technol.* 71:1–13.
- UNICEF. 2020. Global UNICEF Global databases: Infant and young child feeding: Exclusive breastfeeding, predominant breastfeeding. UNICEF Division of Data, Analysis, Planning and Monitoring, New York.
- Vignolles, A. D., R. Jeantet, C. Lopez, and P. Schuck. 2007. Free fat, surface fat and dairy powders: interactions between process and product. A review. *Lait* 87:187–236. <https://doi.org/10.1051/lait:2007010>.
- Wang, D., J. Fritsch, and C. I. Moraru. 2019. Shelf life and quality of skim milk processed by cold microfiltration with a 1.4- μ m pore size membrane, with or without heat treatment. *J. Dairy Sci.* 102:8798–8806. <https://doi.org/10.3168/jds.2018-16050>.
- Wijayanti, H. B., N. Bansal, and H. C. Deeth. 2014. Stability of whey proteins during thermal processing: A review. *Compr. Rev. Food Sci. Food Saf.* 13:1235–1251. <https://doi.org/10.1111/1541-4337.12105>.
- Zulewska, J., and D. M. Barbano. 2014. The effect of linear velocity and flux on performance of ceramic graded permeability membranes when processing skim milk at 50°C. *J. Dairy Sci.* 97:2619–2632. <https://doi.org/10.3168/jds.2013-7635>.

ORCID

- F. J. Tessier  <https://orcid.org/0000-0001-8096-5715>
M. Howsam  <https://orcid.org/0000-0002-6344-8908>
T. Croguennec  <https://orcid.org/0000-0003-4720-3641>
G. Gésan-Guiziou  <https://orcid.org/0000-0001-7001-4684>
R. Jeantet  <https://orcid.org/0000-0001-5405-5056>
A. Deglaire  <https://orcid.org/0000-0001-8366-116X>