



**HAL**  
open science

## Linking Pélardon artisanal goat cheese microbial communities to aroma compounds during cheese-making and ripening

Marine Penland, Hélène Falentin, Sandrine Parayre, Audrey Pawtowski, Marie-Bernadette Maillard, Anne Thierry, Jérôme Mounier, Monika Coton, Stéphanie-Marie Deutsch

### ► To cite this version:

Marine Penland, Hélène Falentin, Sandrine Parayre, Audrey Pawtowski, Marie-Bernadette Maillard, et al.. Linking Pélardon artisanal goat cheese microbial communities to aroma compounds during cheese-making and ripening. *International Journal of Food Microbiology*, 2021, 345, pp.109130. 10.1016/j.ijfoodmicro.2021.109130 . hal-03181204

HAL Id: hal-03181204

<https://hal.inrae.fr/hal-03181204>

Submitted on 6 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 **Linking Pélardon artisanal goat cheese microbial communities to aroma**  
2 **compounds during cheese-making and ripening**

3 Marine Penland<sup>1,2</sup>, Hélène Falentin<sup>1</sup>, Sandrine Parayre<sup>1</sup>, Audrey Pawtowski<sup>2</sup>, Marie-  
4 Bernadette Maillard<sup>1</sup>, Anne Thierry<sup>1</sup>, Jérôme Mounier<sup>2</sup>, Monika Coton<sup>2</sup> & Stéphanie-Marie  
5 Deutsch<sup>1</sup>

6 <sup>1</sup> STLO, INRAE, Institut Agro, 35042, RENNES, FRANCE

7 <sup>2</sup> Univ Brest, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, F-29280  
8 Plouzané, France.

9 Marine Penland: marine.penland@inrae.fr

10 Hélène Falentin: helene.falentin@inrae.fr

11 Sandrine Parayre: sandrine.parayre@inrae.fr

12 Audrey Pawtowski: audrey.pawtowski@univ-brest.fr

13 Marie-Bernadette Maillard: marie-bernadette.maillard@inrae.fr

14 Anne Thierry: anne.thierry@inrae.fr

15 Jérôme Mounier: jerome.mounier@univ-brest.fr

16 Monika Coton: monika.coton@univ-brest.fr

17 Stéphanie-Marie Deutsch: stephanie-marie.deutsch@inrae.fr

18 **\*Corresponding author:** Hélène Falentin

19 E-mail: helene.falentin@inrae.fr

20 Phone: +33(0)223485334

## 21 **Abstract**

22 Pélardon is an artisanal French raw goat's milk cheese, produced using natural whey as a  
23 backslopping method. The aim of this study was to identify key microbial players involved in  
24 the acidification and aroma production of this Protected Designation of Origin cheese.  
25 Microbial diversity of samples, collected from the raw milk to 3-months cheese ripening, was  
26 determined by culture-dependent (MALDI-TOF analysis of 2877 isolates) and -independent  
27 (ITS2 and 16S metabarcoding) approaches and linked to changes in biochemical profiles  
28 (volatile compounds and acids). In parallel, potential dominant autochthonous microorganism  
29 reservoirs were also investigated by sampling the cheese-factory environment. Complex and  
30 increasing microbial diversity was observed by both approaches during ripening although  
31 major discrepancies were observed regarding *Lactococcus lactis* and *Lactiplantibacillus*  
32 *paracasei* fate. By correlating microbial shifts to biochemical changes, *Lactococcus lactis* was  
33 identified as the main acidifying bacterium, while *L. mesenteroides* and *Geotrichum*  
34 *candidum* were prevalent and associated with amino acids catabolism after the acidification  
35 step. The three species were dominant in the whey (backslop). In contrast, *L. paracasei*,  
36 *Enterococcus faecalis*, *Penicillium commune* and *Scopulariopsis brevicaulis*, which  
37 dominated during ripening, likely originated from the cheese-making environment. All these  
38 four species were positively correlated to major volatile compounds responsible for goaty and  
39 earthy Pélardon cheese aroma. Overall, this work highlighted the power of MALDI-TOF and  
40 molecular techniques combined with volatilome analyses to dynamically follow and identify  
41 microbial communities during cheese-making and successively identify the key-players  
42 involved in aroma production and contributing to the typicity of Pélardon cheese.

43 **Keywords: microbial dynamics; MALDI-TOF; GC-MS; metabarcoding; backslopping**

44

45 **1. Introduction**

46 Artisanal cheeses are highly appreciated worldwide because of their typical sensory  
47 attributes such as intense and complex flavors compared to industrial cheeses (Montel et al.  
48 2014; Van Hoorde, Vandamme and Huys 2008). They are often made from raw milk and rely  
49 on spontaneous fermentation, which implies that indigenous microorganisms (bacteria and  
50 fungi) naturally present in raw materials also contribute to the fermentation in addition to  
51 defined cultures sometimes added for acidification or ripening. In this context, the origin of  
52 these microorganisms has been investigated and milk is considered as a main microbial  
53 reservoir. However, traditional cheese-making practices such as manual salting, renneting and  
54 backslopping (i.e. the use of a small portion of milk or whey from a previous successful  
55 fermentation) have also been identified as microbial sources (Ercolini et al. 2008; Montel et  
56 al. 2014). Furthermore, some studies highlighted the impact of the production environment on  
57 the microbial communities involved in the fermentation process of some artisanal cheeses, in  
58 particular the milking environment equipment such as vats and trays (Calasso et al., 2016;  
59 Carpino et al., 2016), the cheese-making and ripening environment (equipments and surfaces).  
60 All these potential reservoirs enrich the microbial diversity of artisanal cheeses, which  
61 therefore harbor a complex microbiota (Delcenserie et al., 2014).

62 Cultural methods are classically used to characterize microbial diversity in fermented  
63 foods. They rely on diverse and more or less selective media to enumerate and isolate  
64 microorganisms, which constitute both the advantages and limits of the approach. While  
65 enumeration and isolation are useful to understand population shifts and preserve microbial  
66 diversity for further use, it only reveals the viable and cultivable fraction of the microbial  
67 communities and is particularly fastidious and time-consuming. In addition, the depth of the  
68 analysis depends on the number and selectivity of the media used. In the last decade, culture-  
69 independent approaches, based on DNA analyses, have been widely used to describe



70 microbial communities in cheese as they have the advantage to provide an overview of the  
71 microorganisms found in an ecosystem, cultivable or not. Different PCR-based techniques  
72 have been applied such as PCR-DDGE (Dolci et al., 2008) or more recently high-throughput  
73 sequencing techniques such as metabarcoding (De Filippis et al., 2018; Kergourlay et al.,  
74 2015). Using this approach, studies were conducted to characterize the bacterial communities  
75 of artisanal Irish cheeses based on their geographic production area (Quigley et al., 2012) or  
76 to determine the microbiota associated with specific cheese types (Dugat-Bony et al., 2015) or  
77 cheese rinds (Wolfe et al., 2014). Such methods revealed complex microbiota that were  
78 previously overlooked by cultural methods. However, most studies focused on microbial  
79 diversity and compared the microbial communities encountered in different cheese varieties  
80 thus rendering a static view of the microbial diversity rather than investigating the changes in  
81 the microbial community during cheese-making and ripening periods.

82 Cheese-making and ripening are complex phenomena during which the cheese develops  
83 its aroma and other sensory characteristics. Many successive or simultaneous enzymatic  
84 reactions occur throughout the process, which include lactose fermentation process,  
85 proteolysis and amino acid catabolism, and lipolysis. These reactions were shown to directly  
86 or indirectly involve bacteria and/or fungi depending on cheese type (Bertuzzi et al., 2018;  
87 Delgado et al., 2011). As a consequence, investigating the microbial dynamics and their  
88 correlation with the cheese aroma profile might help to better apprehend the key-players and  
89 their respective role throughout the process.

90 Pélardon cheese is a Protected Designation of Origin (PDO) raw goat milk artisanal  
91 cheese. It is mostly produced **in the mountainous Cevennes and Corbières areas and in**  
92 **Southern France** and is characterized by a soft creamy texture with a thin wrinkled white and  
93 blue mold rind. It is solely produced using an artisanal process by small farm-style

94 cheesemakers. Pélardon production requirements forbid the use of commercial starters but  
95 authorize the use of whey from a previous batch as a backslopping inoculum.  
96 The objective of this study was thus to understand the microbial dynamics associated with  
97 Pélardon cheese and how they could be linked with the cheese aroma characteristics. For this  
98 purpose, we first (i) described fungal and bacterial dynamics during Pélardon cheese-making  
99 and ripening by both culture-dependent and -independent approaches, (ii) **investigated**  
100 **potential microbial sources in the process environment and** (iii) quantified sugars, organic  
101 acids and aroma compounds throughout the entire process. Finally, we assessed how  
102 microbial diversity and dynamics influence the overall cheese aroma characteristics.

103

## 104 **2. Material and methods**

### 105 **2.1. Pélardon cheese sampling strategy**

#### 106 **2.1.1. Pélardon cheese-making and sample collection**

107 **Samples were collected from a local cheese-maker from the Southern France (Gard) in the**  
108 **Cevennes area.** Cheeses were prepared with the traditional process described by PDO  
109 requirements and in the same conditions as standard manufacturing. Briefly, raw goat milk  
110 from two successive milkings was supplemented with 2 % kid rennet (**chymosin: 180 mg/L**)  
111 and 2 % whey from the previous production batch. No starter cultures were added.  
112 Acidification was carried out for 18 to 24 h until the acidity reached 58 ° Dornic at 20°C with  
113 controlled relative humidity (RH=80 %). Curd was manually moulded with a ladle into  
114 individual jars and drained naturally **at room temperature (20°C)** for 24 h before salting by  
115 manually sprinkling salt on the surface. After demoulding, fresh cheeses were drained for 48  
116 h more prior to being placed in the drying room (T°=14°C; RH = 85 %) for 48 h. Finally,  
117 ripening was carried out in a cellar, at 12°C (RH = 90 %) for a minimum period of four days  
118 and up to three months.

119 Samples came from the same production batch (April 7, 2018) and sampling was done at  
120 seven steps during the cheese making process: raw materials (raw milk and whey used for  
121 backslopping inoculum), cheese samples after curdling (day 2), after salting and drying (day  
122 8), after two weeks (day 14), two months (day 62) and three months ripening (day 90).  
123 Altogether, the sampling steps provided 33 samples corresponding to the biological replicates  
124 collected throughout cheese-making and ripening. Whole cheeses were used for each analysis  
125 point. Samples were subjected to microbial enumeration, isolation and identification of  
126 bacteria and fungi, metabarcoding analyses and biochemical analyses to quantify sugars, acids  
127 and volatile compounds. General sampling strategy and experiments are summarized in Fig. 1  
128 and primers used in this study in Supplementary Table S1.

#### 129 **2.1.2. Production environment sample collection**

130 Sixteen samples were taken at day 2 from different areas in the production and ripening  
131 facilities in order to track potential environmental microbial sources. Airborne  
132 microorganisms from the main production, drying and ripening rooms were collected using an  
133 air sampler (Sampl'air Lite, AES) operated at flow rate of 100 L/min for 10 min while ten  
134 surfaces were also analysed by streaking swabs moistened with Tryptone Salt (TS) diluent  
135 (sodium chloride 8.5 g/L ; tryptone 1 g/L) over a 31.5 cm<sup>2</sup> area as well as three intrants  
136 (running water samples and salt) and **stored at 4°C until analysis**. All samples were analysed  
137 within 24 h and further subjected to microbial enumeration, bacterial and fungal isolations,  
138 identifications and metabarcoding analyses (except air samples).

#### 139 **2.2. Microbial counts during cheese-making process and in environmental samples**

140 For raw materials and cheese samples, microbial populations of interest were monitored  
141 at each sampling point. For cheese samples following the drying step (day 8), core and rind  
142 were analyzed separately. In each case, 10 g of cheese were mixed with 90 mL of 2 %

143 trisodium citrate buffer pre-heated to 42°C and then blended with a stomacher for 3 min at  
144 high speed. Serial dilutions were then prepared in TS diluent and plated on different media.  
145 The following microbial populations were enumerated: mesophilic aerobic bacteria on milk  
146 plate count agar (MPCA) (30°C, 72 h), mesophilic halotolerant bacterial populations on  
147 MPCA supplemented with 5 % NaCl (30°C, 72 h), enterococci on Kenner fecal agar  
148 supplemented with 1% (w/v) 2,3,5-triphenyltetrazolium chloride (KF) (37°C, 48 h),  
149 presumptive lactococci on M17 (30°C, 48 h), presumptive anaerobic lactobacilli on de Man  
150 Rogosa Sharpe (MRS) pH 5.4 (30°C, 48 h, anaerobiosis), presumptive aerobic lactobacilli on  
151 MRS pH 5.4 (30°C, 48 h, aerobiosis), total fungal populations on Yeast Glucose  
152 Chloramphenicol (YGC) agar (25°C, 5 days) and halotolerant fungi on YGC supplemented  
153 with 5 % NaCl (25°C, 5 days). All media targeting bacterial populations were supplemented  
154 with 0.1 % natamycin to inhibit fungal growth, except for KF medium.

155 For environment samples, enumerations were performed on three media: PCA (30°C, 72 h)  
156 for mesophilic aerobic microorganisms, MRS (37°C, 48h, anaerobiosis) for lactic acid  
157 bacteria and YGC (25°C, 5 days) for fungi after initial serial dilutions in TS diluent. Air  
158 samples were directly collected by impacting 1 m<sup>3</sup> of air on two media: PCA (30°C, 72 h) for  
159 mesophilic aerobic microorganisms and YGC (25°C, 5 days) for fungi. For those samples,  
160 analyses were performed on one replicate.

## 161 **2.3. Isolation and identification of microorganisms**

### 162 **2.3.1. Collection of microbial isolates**

163 For each biological replicate of raw materials (raw milk and whey used for backslopping)  
164 and cheese samples (core and rind analyzed separately from day 8 onwards), microorganisms  
165 were isolated from five media: bacteria from MPCA + natamycin, M17, MRS pH 5.4  
166 (anaerobiosis) and KF media and fungi from YGC medium. Twenty representative isolates (if

167 possible) were randomly selected from each medium from the Petri dish showing the highest  
168 morphological diversity. Regarding environmental samples, 15 isolates (if possible) were  
169 selected from PCA and YGC media.

### 170 2.3.2. Isolate dereplication using MALDI-TOF MS analysis and identification by 171 sequencing

172 A dereplication step using MALDI-TOF mass spectroscopy was performed for all  
173 bacterial and fungal isolates to identify clones prior to species level molecular identifications.  
174 Analyses were performed on a VITEK MS instrument (bioMérieux, Marcy l'Étoile, France)  
175 equipped with the Launchpad V2.8.4 acquisition software. Bacterial and fungal isolates were  
176 processed following the manufacturer's instructions. For filamentous fungi isolates, extraction  
177 and plate preparation were performed as described by Quéro et al. (2019) as well as spectra  
178 acquisition. Then, mass spectra of each isolate were first compared against the bioMérieux  
179 bacterial or fungal industrial-clinical VITEK MS databases (V3.2.0 for bacteria and new  
180 update of the VITEK MS database for fungi developed in the laboratory together with  
181 bioMérieux) for presumptive identifications (internal research tool). Then, the spectra of  
182 isolates identified as belonging to the same species were compared against one another to  
183 build clusters based on their similarities using SARAMIS software (bioMérieux, Marcy  
184 l'Étoile, France). A threshold of 65 % similarity was applied for bacteria and of 50 % for  
185 fungi, as described by Lindgren et al. (2018). Representative isolates (isolates with the lowest  
186 similarity within a species cluster) were then chosen within each cluster and for each  
187 presumptive species. For each presumptive species, the choice of representative isolates was  
188 based on the heterogeneity between their spectra and when discrepancies were observed  
189 between MALDI-TOF identification and position within the cluster. An example is given for  
190 *Staphylococcus* genus (Supplementary figure S1).

191 Species-level molecular identifications for filamentous fungi were done after amplification  
192 and sequencing of either the internal transcribed spacer (ITS) region using ITS4/ITS5 primers  
193 or the partial  $\beta$ -tubulin gene using Bt2a/Bt2b primers (Glass and Donaldson, 1995) depending  
194 on the presumptive genus (i.e. *Penicillium*, *Aspergillus*) provided by MALDI-TOF MS  
195 analysis. Yeasts were identified by targeting the D1-D2 domain using NL1/NL4 primers  
196 (Kurtzman and Robnett, 1998). Finally, bacteria were identified by sequencing the complete  
197 16S rRNA gene after amplification with fD1 and rP2 primers (Weisburg et al., 1991).  
198 Sequencing was performed using the same primers as those used for PCR amplifications.  
199 Sequences were assembled into contigs using Geneious software (<http://www.geneious.com>,  
200 Kearse et al. 2012) and compared with the GenBank databases using the “Basic Local  
201 Alignment Search Tool” (BLAST) (<http://www.ncbi.nlm.nih.gov/BLAST>) and “SILVA  
202 Incremental Aligner” (SINA) (<https://www.arb-silva.de/aligner/>). The obtained genus or  
203 species level identifications were then confronted with MALDI-TOF identifications and  
204 dendrogram analyses. When both identifications were consistent inside a given cluster, the  
205 identification was expanded to all the isolates within the cluster. When the identification was  
206 inconsistent, isolates were excluded.

## 207 **2.4. Metabarcoding analysis**

### 208 **2.4.1. Raw materials, cheese and environment sample preparation**

209 Metabarcoding analysis was performed on raw milk (triplicates), whey (triplicates), curd (3  
210 curds), cheeses at day 8, 14, 62 and 90 and 13 environmental samples. The same raw material  
211 and cheese samples (core and rind separately), diluted in citrate buffer, were used as described  
212 previously for culture-dependent analyses (see 2.2). Aliquots of 1 mL were centrifuged (9000  
213 g, 15 min, 4°C) then supernatants were removed and the cell pellets stored at -20°C until  
214 DNA extraction. Regarding environmental samples, pellets were obtained from liquid

215 samples by centrifuging 1 mL (9000 g, 15 min, 4°C), whereas for surface samples, swabs  
216 were placed in 1 mL of TS diluent prior to centrifugation (9000 g, 15 min, 4°C) to obtain cell  
217 pellets.

#### 218 **2.4.2.** Total DNA extraction from cheese and environment samples

219 Total DNA extractions were performed using the DNeasy Blood and tissue kit (Qiagen,  
220 Germany) with a supplementary initial enzymatic lysis. First, cell pellets were thawed at room  
221 temperature then resuspended in 400 µL of lysis buffer (Tris-HCl 20 mM at pH 8.0, EDTA 2  
222 mM, Triton X-100 1.2 %) supplemented with lysozyme (20 mg/mL) and mutanolysin (5  
223 U/µL), then Rnase (25 µg/mL; Qiagen, Germany) and lyticase (0.5 U/µL; Sigma-Aldrich,  
224 Germany) were added. Samples were incubated at 37°C for 2.5 h followed by mechanical  
225 lysis with 300 µL of 0.1mm zirconium beads per tube. Samples were homogenized for 2x 40 s  
226 with a 30 s break and this was repeated twice with a 1 min break on ice using a Precellys  
227 Evolution homogeneizer (Bertin technologies, Germany). A proteinase K (20 mg/mL)  
228 treatment was finally applied for 1 h at 56°C. The remaining extraction and purification steps  
229 were performed according to the manufacturer's instructions. Extracted DNA quality and  
230 quantity were verified using a Nanodrop spectrophotometer and samples were stored at -20°C  
231 until further analysis.

#### 232 **2.4.3.** Quantitative PCR of *Lactiplantibacillus paracasei*

233 The number of *Lactiplantibacillus paracasei* (ex-*Lactobacillus paracasei*) in total DNA  
234 extracts from day 14, 62 and 90 were determined using quantitative PCR targeting the  
235 monocopy *tuf* gene, as described by Achilleos and Berthier (2013). To build the standard  
236 curve, DNA was extracted from pure cultures from 1 mL of an overnight culture of *L.*  
237 *paracasei* CIRM-BIA 1517. Serial dilutions of standard DNA were prepared and submitted to  
238 the same amplification conditions. The curve was then created by plotting qPCR Ct values

239 against the gene copy numbers previously determined and the copy number of *L. paracasei*  
240 was calculated for each cheese sample by comparing the Ct of the sample with that of the  
241 standard curve.

#### 242 **2.4.4.** Amplification and sequencing parameters

243 To study bacterial and fungal diversity in both cheese and environmental samples, PCR  
244 were performed as follows. For bacteria, V3-V4 region of the 16S rRNA gene was targeted  
245 using S-D-Bact-0341-b-S-17 and S-D-Bact-0785-a-A-21 primers (Klindworth et al., 2013).  
246 For fungi, ITS3f/ITS4\_Kyo1 primers (Toju et al., 2012) were used, targeting the ITS2 region.  
247 ITS2 and V3-V4 amplifications and sequencing steps were performed in the same run at  
248 Genome Quebec sequencing platform (MacGill University, Canada) using Illumina Miseq  
249 PE300 technology generating 2x300 bp reads and a total of 8.2 Gb of data for both amplicon  
250 types derived from the DNA extracts.

#### 251 **2.4.5.** Bioinformatic and phylogenetic analysis

252 Sequences were pre-processed for quality and length using the following parameters:  
253 amplicon size between 370 and 490 bp for V3-V4 contigs and 90-500 bp for ITS2 contigs,  
254 mismatch rate was set at 0.1 for both data types. Sequencing data were then analyzed using  
255 the FROGS pipelines as developed by Escudié et al. (2018) under Galaxy (Afgan et al., 2018).  
256 Briefly, raw paired-end reads were assembled and sequences were clustered using the Swarm  
257 algorithm (Mahé et al., 2015) with an aggregation distance of 3 into Operational Taxonomic  
258 Units (OTUs). Sequences underwent some filtering steps: chimeras were detected using  
259 Uparse (Edgar, 2013) and 'de novo parameter' and removed, then sequences with a relative  
260 abundance below  $5 \times 10^{-5}$  or present in only one sample were excluded. Finally, the affiliation  
261 step was performed using SILVA database (v132 pintail 100) and UNITE (v7.1) for 16S and  
262 ITS2 data, respectively. When 16S sequences were multi-affiliated by FROGS because the



263 targeted V3-V4 region was unable to discriminate species, the resulting possible species level  
264 assignments were implemented into the final OTU table.

#### 265 **2.4.6. Biodiversity and statistical analyses**

266 Processing and statistical analyses of microbial communities were performed using  
267 Phyloseq package (McMurdie and Holmes, 2013) under R software. Data were normalized  
268 based on the sample that had the lowest number of sequences. Alpha-diversity indexes  
269 taxonomic composition and abundance distribution were then determined for each sample and  
270 compared.

### 271 **2.5. Biochemical analyses of cheese samples**

#### 272 **2.5.1. pH measurement**

273 pH of the whey and day 2 cheeses was measured using a WTW 3100 pH meter  
274 (Weilheim, Germany) equipped with a puncture electrode (LoT406-M6-DXK GmbH, Mettler  
275 Toledo, Urdorf, Switzerland) and temperature probe (WTW 325/HC) by direct insertion. For  
276 cheese samples from day 8, 14, 62 and 90, pH was measured in core and rind separately.. All  
277 measurements were performed on cheese triplicates.

#### 278 **2.5.2. Quantification of sugars and organic acids by HPLC**

279 Two sugars and four organic acids were quantified in samples at six sampling points by  
280 high-performance liquid chromatography (HPLC). Compound extraction was performed as  
281 follows: all samples except whey were first blended in sterile distilled water (1:5) using a  
282 Stomacher (Merck Eurolab, Strasbourg, France) and incubated at 40°C for 1 h. The obtained  
283 dilutions were then centrifugated (3000 g, 30 min, 4°C) and filtered on a Whatmann 40  
284 membrane. The filtrates were then diluted (3:4) with H<sub>2</sub>SO<sub>4</sub> to reach a final concentration of  
285 0.05 M. Samples were then frozen overnight at -20°C and, after thawing, centrifuged (8000 g,

286 20 min, 4°C). For each sample, supernatant was finally recovered and filtered using 0.45 µm  
287 PTFE membrane into a 2 mL-vial and stocked at -20°C until analysis.  
288 HPLC analyses were performed using the conditions described by Leyva Salas et al. (2019).  
289 Compounds were identified and quantified by comparing retention times and peak area of  
290 standard solutions injected at different concentrations (ranging from 0.05 to 1 mg/mL). All  
291 samples were analyzed in the same run. When necessary, supernatants were diluted with 0.05  
292 M H<sub>2</sub>SO<sub>4</sub> and re-injected.

### 293 **2.5.3. Volatile compounds profile analyses by headspace-GC-MS**

294 Volatile profiles were obtained for six sampling points including whey and cheeses (from  
295 days 2, 8, 14, 62 and 90) using headspace (HS) gas chromatography-mass spectrometry (GC-  
296 MS). Compound extraction was performed using a Perkin Elmer Turbomatrix HS-40 trap  
297 automatic headspace sampler with trap enrichment on 2.5 g of mixed cheese placed in 22 mL  
298 vials. Analyses were performed according to methods previously described by Harlé et al.,  
299 (2020). Prior to compounds identification, data were processed using PerkinElmer Turbomass  
300 software, version 5.4.2.1617 and by converting the raw data to time- and mass-aligned  
301 chromatographic peaks areas using the open source XCMS package implemented with the R  
302 statistical language (Smith et al., 2006). Parameters were set as follows: width at half  
303 maximum=5, group bandwidth=3. Volatile compound identification was achieved by  
304 comparing the retention index and mass spectral values (1) from the NIST 2008 Mass  
305 Spectral Library (Scientific Instrument Services, Ringoes, NJ, USA) and (2) when possible  
306 with those of authentic standards (Sigma Aldrich, France) analysed in the same system and  
307 (3) those reported in the literature. Finally, volatile profile changes between samples were  
308 investigated by comparing relative abundance of the identified compounds using ANOVA  
309 analyses with R software. Data were then normalized by centering and scaling the abundances.

310 Hierarchical clustering and correlation of the data was then performed using Ward's  
311 minimum variance linkage and Euclidean distance method under R software.

## 312 **2.6. Statistical analyses of microbial and biochemical data**

313 Data from both culture-dependent and -independent analyses were implemented into a  
314 Principal Component analysis (PCA). Prior to the analysis, culture-dependent identified  
315 variables were reduced to one medium per species as many species were identified on more  
316 than one medium. The medium with the highest specificity and abundance of each species  
317 was kept and the abundance values multiplied by the corresponding microbial population  
318 counts. Pearson correlation coefficients between microbial species (culture-dependent  
319 approach) and biochemical compounds were then calculated and represented on a heatmap.  
320 Significance levels of correlations between microbial genera and biochemical compounds  
321 were then determined. All analyses were performed with R software using FactomineR,  
322 Hmisc, and ggplot2 packages (Lê et al., 2008; Wickham, 2016).

## 323 **3. Results**

324 **3.1. Culture-dependent analyses to determine the succession of microbial species during**  
325 **cheese-making and ripening**

### 326 **3.1.1. Microbial population monitoring**

327 Eight populations of interest were monitored during cheese-making and ripening.  
328 Microbial counts obtained for raw materials and cheese core and rind samples are presented in  
329 Fig. 2.A, 2.B and 2.C, respectively. Raw milk harbored low mesophilic bacterial populations  
330 ( $3.4 \log_{10}$  CFU/mL) and fungal populations ( $2 \log_{10}$  CFU/mL). Higher bacterial counts by up  
331 to  $5 \log_{10}$  CFU/mL were observed in the whey used for backslopping than in raw milk except  
332 for enterococci and presumptive aerobic lactobacilli. Indeed, presumptive aerobic lactobacilli  
333 were encountered at low levels in raw milk (around  $2 \log_{10}$  CFU/mL) and undetected in whey,

334 while enterococci counts were higher in raw milk ( $2.3 \log_{10}$  CFU/mL) compared to whey (2  
335  $\log_{10}$  CFU/mL). In day 2 curds, presumptive lactococci and anaerobic lactobacilli dominated  
336 (up to  $9.2 \log_{10}$  CFU/g), while enterococci and presumptive aerobic lactobacilli remained  
337 undetected. Fungal growth occurred during milk acidification and renneting step with  
338 populations reaching  $4.6 \log_{10}$  CFU/g at day 2.

339 From the post-drying stage (day 8) onwards, differences were observed between core and rind  
340 populations. Aerobic bacterial populations remained relatively stable in rinds while a decrease  
341 by  $1.2 \log_{10}$  was observed in cores and up to  $3 \log_{10}$  for halotolerant bacteria. A similar trend  
342 was observed for presumptive lactococci and anaerobic lactobacilli populations in cores  
343 which decreased by  $\sim 1.8 \log_{10}$  during drying (day 8) and, then remained stable ( $\sim 6 \log_{10}$   
344 CFU/g) up to 90 days. Inversely, both presumptive aerobic lactobacilli and enterococci  
345 populations significantly increased during the cheese-making and ripening in both core and  
346 rinds to reach  $6 \log_{10}$  CFU/g for lactobacilli and  $5 \log_{10}$  CFU/g for enterococci over the course  
347 of ripening. Fungal populations were consistently  $2 \log_{10}$  higher on rinds and also increased  
348 during the drying step then remained at high levels until the end of ripening ( $7.91 \log_{10}$   
349 CFU/g). In contrast, they remained fairly constant in core samples, around  $5.5 \log_{10}$  CFU/g.

350 An increase in halotolerant fungal populations was also observed in rinds between day 8 and  
351 day 14 while no such effect was observed in the core.

### 352 **3.1.2.** Identification of viable microbial communities by MALDI-TOF and sequencing

353 A total of 2001 bacterial and 563 fungal isolates were collected from the raw materials  
354 and cheese-making samples and 143 bacterial and 160 fungal isolates from environmental  
355 samples.

356 Isolates were first subjected to a dereplication step using MALDI-TOF analysis to select  
357 representative isolates for further molecular identifications. Overall, good-quality spectra  
358 were acquired for 88 % bacterial isolates (n=1876) and 99 % fungal (n= 727) isolates. These

359 spectra were thus used to build similarity dendrograms and grouped into clusters based on  
360 their similarity and presumptive identifications. In total, 146 representative bacterial and 38  
361 fungal isolates were identified by sequencing. Based on MALDI-TOF spectra clustering,  
362 presumptive identifications and sequencing data, 1769 bacterial and 683 fungal isolates,  
363 respectively representing 82 % and 93 % of the initial collection, were successfully identified  
364 and included in the final analysis. Noteworthy, the presumptive identifications done by  
365 MALDI-TOF analyses were confirmed by species-level molecular identification for 78 % and  
366 86 % of the sequenced isolates for bacteria and fungi, respectively. Results based on all these  
367 data are presented in Fig. 3.

368 Regarding raw materials, **only few lactic acid bacteria species** (LAB) were identified in raw  
369 milk whereas high species diversity was observed for staphylococci among the isolated  
370 aerobic bacteria **with three species only identified in these samples namely *Staphylococcus***  
371 ***arlettae*, *Staphylococcus caprae* and *Staphylococcus epidermidis***. In contrast, in whey used  
372 for backslopping and in curd (day 2), *Lactococcus lactis* was the dominant bacterial species  
373 followed by *Leuconostoc mesenteroides*. **Among aerobic bacteria and presumptive lactococci,**  
374 *L. lactis* abundance was highest at day 2 and progressively decreased throughout ripening  
375 **(aerobic bacteria)** and was no longer detected neither in core nor rind at three months of  
376 ripening. A similar trend was observed for *L. mesenteroides* as its relative abundance was  
377 highest at day 8 in cores **(aerobic bacteria and presumptive lactococci)** and progressively  
378 decreased until day 90, and was no longer detectable in rinds after day 14. These observations  
379 were concomitant with the progressive growth of *Lactiplantibacillus paracasei/casei* from  
380 day 8 to the end of ripening and it was the most abundant bacterial species in cores at 62 and  
381 90 days of ripening **(aerobic bacteria and presumptive anaerobic lactobacilli)**. During  
382 ripening, the growth of several **actinobacteria** was observed such as *Glutamicibacter* spp.,  
383 *Arthrobacter* spp. and *Brevibacterium* spp.; they represented up to half of the isolates among

384 aerobic bacteria in rinds. Although their populations levels were lower than the most  
385 dominant bacterial groups, interesting dynamics were observed among enterococci and  
386 staphylococci. Regarding enterococci, their diversity increased in rinds during ripening, with  
387 only *Enterococcus durans* identified during the post-drying step (day 8) and three other  
388 species namely *E. faecalis*, *E. faecium* and *E. hirae* identified at days 62 and 90.  
389 Staphylococci species diversity also increased during ripening and numerous species were  
390 identified such as *Staphylococcus xylosus*, *Staphylococcus equorum*, *Staphylococcus succinus*  
391 and *Staphylococcus fleurettii*, which were not previously isolated in raw material or curds  
392 (day 2).  
393 Regarding fungal communities, the isolates collected from raw milk could not be identified  
394 while whey was dominated by *Geotrichum candidum*. This yeast-like fungus remained  
395 dominant until day 14. After curdling (day 2), despite different population levels, no striking  
396 differences between core and rind composition were observed in cheese samples. Fungal  
397 diversity then increased from day 14 in the rinds and from day 62 in cores. In particular,  
398 *Penicillium commune* and *Scopulariopsis brevicaulis* relative abundances increased and *S.*  
399 *brevicaulis* even became dominant in both core and rind at days 62 and 90. Other yeast  
400 species such as *Debaryomyces hansenii* and *Yarrowia lipolytica* were also encountered in the  
401 same samples but at lower abundances.

402 **3.2.** Culture-independent analyses to determine the succession of microbial species during  
403 cheese-making and ripening

404 **3.2.1.** Microbial dynamics determined using 16S rRNA gene and ITS2 metabarcoding  
405 analyses

406 V3-V4 16S rRNA gene sequencing analysis resulted after quality filtering in a total of 3  
407 537 077 contigs with an average length of 467 nucleotides. Sequences were subsequently  
408 clustered into 43 OTUs after chimera and singleton removal. Concerning metabarcoding

409 analysis of the ITS2 region, a total of 6 945 569 quality-filtered contigs (average length of 245  
410 nucleotides) were obtained. After chimera and singleton removal, sequences were clustered  
411 into 15 OTUs. After normalization, 67 540 and 54 622 sequences per sample were kept for  
412 V3-V4 and ITS2 data analysis, respectively. Rarefaction curves are available in  
413 Supplementary Fig. S2.

414 For both bacterial and fungal communities, alpha-diversity indexes were calculated from  
415 OTU data to investigate the impact of three factors: cheese-making and ripening stage, sample  
416 type and the two factors associated (see Supplementary Fig. S3). For bacterial communities,  
417 the richness and Chao1 indexes (defined as OTU number in samples and estimated richness  
418 respectively) showed respectively significant differences between sample type and stage  
419 ( $p < 0.001$ ) as well as Shannon evenness index ( $p < 0.01$ ). Bacterial diversity significantly  
420 increased during cheese-making and ripening, especially in rinds. Regarding fungal  
421 communities, diversity increased throughout ripening. Significant differences regarding  
422 richness indexes were however observed between cores and rinds and overall values were  
423 lower ( $p < 0.001$ ) in cores. For both communities, low Shannon and Simpson indexes ( $p > 0.05$ )  
424 were observed compared to Chao1 index. As Shannon and Simpson indexes consider relative  
425 abundances, results suggested that bacterial and fungal populations were dominated by a few  
426 abundant taxa.

427 The succession of bacterial communities during Pélardon cheese-making and ripening are  
428 illustrated in Fig. 4. All samples considered, a total of three phyla (Firmicutes 94 %,  
429 Actinobacteria 5.6 % and Proteobacteria 0.4 % of total reads), five orders, 10 families, 12  
430 genera and 21 species were identified. Overall, regardless of sample type or stage, the most  
431 abundant genera were *Lactococcus* (88.9 % of all sequences) followed *Glutamicibacter* (3.8  
432 %) *Leuconostoc* and *Brevibacterium* (1% each). Other genera were encountered at an  
433 abundance below 1 %. Raw milk samples were characterized by a low number of reads

434 (below 200 reads per samples) and thus not formally included in this analysis. In the whey  
435 used for backslopping, the most abundant genus was *Lactococcus*, with two identified species  
436 (*L. lactis*, *L. raffinolactis*) and one OTU that could not be identified at the species-level. These  
437 three OTUs represented up to 99.5 % of sequences. At day 2, after acidification, curd samples  
438 were dominated by *L. lactis* and it remained dominant during the following stages in both  
439 core and rind samples although its abundance progressively decreased in rinds concomitantly  
440 with the appearance of other genera and species, especially belonging to staphylococci (i.e. *S.*  
441 *succinus* and *S. saprophyticus*) and actinobacteria. For these groups, high variations in relative  
442 abundances were observed among replicates in the late stages of ripening (days 62 and 90).  
443 Indeed, *Brevibacterium* spp. abundances ranged from 2.8 to 7.6 %, *Glutamicibacter* spp. from  
444 4.4 to 47 % and those of *Brachybacterium* spp. from 0.8 to 11 %.

445 Regarding fungal communities, 15 species belonging to 11 genera, nine families and two  
446 phyla were identified in raw materials and cheeses (see Supplementary Fig. S4). *G. candidum*  
447 was the most abundant species with a relative abundance of 99.5 % in all cases, regardless of  
448 the sample type or stage. However, fungal diversity increased at the beginning of the ripening  
449 period (day 14) as shown by the presence of *Penicillium* species, and in late ripening stages  
450 (days 62 and 90) with the presence of *S. brevicaulis* in rinds. However, no differences in the  
451 relative abundances between core and rind were observed for these species at the end of  
452 ripening. Several other fungal species were punctually identified e.g. *Cladosporium* was only  
453 found in whey and raw milk.

### 454 3.2.2. *Lactiplantibacillus paracasei* quantification during ripening

455 Because of some discrepancies between culture-dependent and independent analyses  
456 regarding the relative abundance of *L. paracasei* at days 14, 62 and 90, its presence was  
457 further investigated by qPCR, using *TufLcpara* specific primers. In these conditions, qPCR  
458 efficiency was 94 %. Based on the standard curve, *tuf* gene copy number of *L. paracasei* per



459 gram of cheese was determined and used as a proxy of *L. paracasei* cells per gram of cheese.  
460 We observed that *tuf* gene copy numbers per gram of cheese were 33-fold (cheese core) and  
461 3-fold (cheese rind) higher at day 62 than at day 14, indicating that *L. paracasei* population  
462 increased over ripening time (Table 1). It then remained stable until day 90. Noteworthy, at  
463 day 62 and 90, the *L. paracasei* population was 10-fold higher in the cheese core than in the  
464 rind.

### 465 **3.3. Study of potential environmental microbial sources using culture-dependent and -** 466 **independent approaches**

467 To investigate the potential sources of microorganisms in shaping Pélardon cheese  
468 microbial communities, sampling was done inside the farmhouse. Detailed sample  
469 information and viable microbial counts are shown in Table 2. All samples were analyzed  
470 using culture-dependent (Fig. 5) and -independent approaches (Supplementary Fig. S5),  
471 except air samples which were only assessed by the culture-dependent approach. Low  
472 microbial counts for mesophilic aerobic bacteria, fungi and LAB were encountered especially  
473 for surface samples ( $< 4 \log_{10} \text{CFU/cm}^2$  or  $\text{m}^3$ ) and these results were correlated with low  
474 amplification by culture-independent analysis. Concerning the main bacterial species  
475 identified in cheese, *L. mesenteroides* was identified on cheese trays used in the drying room  
476 (PS5) by culture-dependent analyses and surfaces from the cheese-making room (PS2, PS4  
477 and PS9) by the culture-independent approaches. *L. lactis* was only isolated from a tray in the  
478 drying room (PS5), while it was also identified at low relative abundance (8 %) on a clean  
479 tray used for curdling (PS9) by the culture-independent approach. *Staphylococcus* species  
480 were among the most frequently isolated species on trays used for ripening (PS6) and in the  
481 cheese-making room (PS7) by both approaches (see Supplementary Fig. S6 for a detailed  
482 comparison). The same species were also found in the air of the drying and ripening rooms.  
483 Bacterial genera identified in Pélardon cheese rinds such as *Glutamicibacter* and

484 *Brevibacterium* were also isolated from surface samples collected from the cheese-making,  
485 drying and ripening rooms at high abundances. Noteworthy, enterococci and lactobacilli were  
486 not isolated nor detected in environmental samples. For fungi, air sample analysis showed that  
487 *G. candidum* was present at high abundances in the air of the cheese-making, drying and  
488 ripening rooms. *P. commune* and *S. brevicaulis* were also present but at lower abundances.  
489 These three species were also identified on the surface of the trays in the different rooms  
490 using culture-independent analysis.

#### 491 **3.4. pH, sugar and non-volatile organic acid changes**

492 pH measurements were similar in the whey used for backslopping and in day 2 curd (~4.3  
493 units) and continuously increased ( $p < 0.001$ ) throughout ripening until day 62 to reach 6.07  
494 and 6.73 pH units in cores and in rinds, respectively. pH values were systematically and  
495 significantly higher in rinds as compared to cores except on day 90 (Table 3).

496 The main sugars (lactose, galactose) and organic acids (acetic, citric, lactic acids) were  
497 quantified in whey used for backslopping and during the different stages of Pélardon cheese-  
498 making and ripening. Lactose, galactose as well as acetic and lactic acids showed a similar  
499 trend throughout the process. Their highest concentrations were in curd after acidification  
500 (day 2) and progressively decreased: lactose and acetate were detected until day 62 whereas  
501 galactose and lactic acid were no longer detected as early as day 14 (early ripening). In  
502 contrast, citric and 3-phenyllactic acid concentrations were below the detection level during  
503 acidification (day 2) and increased during drying (for 3-phenyllactic acid) and ripening (both  
504 acids).

#### 505 **3.5. Changes in volatile profiles during cheese-making and ripening**

506 Using headspace GC-MS, 54 volatile compounds belonging to six main families were  
507 detected and identified in Pélardon cheese samples during cheese-making and ripening.

508 Compound names, identification criteria and their respective abundances are listed in the  
509 Supplementary Table S2.

510 To better understand these changes and the main compounds involved, a one-way ANOVA  
511 analysis followed by Tukey's highest significant difference test were performed on abundance  
512 data. Out of the 54 volatile compounds, 47 were detected at a significantly different  
513 abundance at, at least, one stage ( $p < 0.05$ ). To confirm these trends, a correlation analysis was  
514 then performed on the volatile abundances and the results are presented using a heatmap with  
515 sample clustering. Overall, the identified compounds could be linked to specific cheese-  
516 making or ripening stages based on their high abundances and four main groups of  
517 **compounds** could be differentiated (Fig. 6). A first group (group I) gathered six compounds  
518 present in high abundances in curd (day 2) and whey samples, i.e. acetic acid, 3-  
519 hydroxybutan-2-one, 2-methylthiolan-3-one, pentane-2,3-dione, butane-2,3-dione and  
520 hexanal. The post-drying (day 8) and early ripening (day 14) stages were associated with  
521 compounds of group II corresponding to branched-chain and aromatic compounds (i.e. 3-  
522 methylbutanoic acid, 2-phenylacetaldehyde). Finally, in the late ripening stages (day 62 and  
523 90), 33 compounds were quantified at their highest abundances, mainly belonging to  
524 methylketones, secondary alcohols and free fatty acids and could be separated in two groups  
525 (III and IV). Group III gathered compounds identified in particularly high abundances at day  
526 62 and day 90 among which were seven free fatty acids (butanoic, hexanoic, pentanoic,  
527 heptanoic octanoic, nonanoic and decanoic acids), five ketones (non-8-en-2-one, octan-3-one,  
528 pentan-2-one, heptan-2-one, hexan-2-one, and nonan-2-one) and one alcohol (heptan-2-ol).

529 Group IV mostly gathered compounds with high abundances in day 90 cheeses and  
530 noteworthy, it included the four esters as well as ketones and secondary alcohols.

531 **3.6. Correlation between microbial communities and changes in biochemical profiles**  
532 **during cheese-making and ripening**

533 A principal component analysis was conducted with all microbial and biochemical data to  
534 correlate temporal changes in microbial communities with biochemical profiles  
535 (Supplementary Fig. S7). Dimensions 1 and 3, which explained 34.2 % and 11.6 % of the  
536 variance respectively, provided the best separation of samples according to the cheese-making  
537 and ripening times. Day 2 and whey samples were grouped together and associated with the  
538 *Lactococcus* genus, identified by both culture-dependent and -independent approaches (three  
539 OTUs), and two sugars (lactose and galactose) as well as six other compounds including  
540 acetic acid, lactic acid and 3-hydroxybutanone. Rind samples from day 8 and day 14 were  
541 linked to 2-phenylacetaldehyde. During ripening (day 62), rinds were associated with a high  
542 pH and six volatile compounds as well as three species, *S. xylosus*, *L. casei/paracasei*, and *P.*  
543 *commune*, identified by culture-dependent approach. At the end of ripening (day 90), multiple  
544 species identified by both culture-dependent analyses (*E. faecalis*, *S. equorum*, *S. brevicaulis*,  
545 *Brevibacterium* spp.) and metabarcoding (*L. lactis*\_62, *Lactococcus* sp.\_78, *Brachybacterium*  
546 sp\_56 and *Brachybacterium* sp.\_8) were correlated to 13 compounds mainly free fatty acids,  
547 methyl ketones and esters.

548 To get a better insight into the potential links between aroma compounds and  
549 microorganisms, a Pearson correlation analysis was performed based on culture-dependent  
550 data, organic acid and volatile data. Pair-wise correlation coefficients and corresponding p-  
551 value were thus calculated (Fig. 7). Focusing on the strongest correlations ( $|r| > 0.8$  and  $p <$   
552  $0.001$ ), four microbial groups stood out. *L. lactis* and *G. candidum* were strongly and  
553 positively correlated with lactose, acetic acid, lactic acid, and 3-hydroxybutan-2-one. *L.*  
554 *mesenteroides*, whose correlation profile was relatively similar to that of *L. lactis*, was  
555 strongly and positively correlated to 3-methylpropanoic, 3-methylbutanoic acids and 2-  
556 phenylacetaldehyde; highest abundances of these compounds being at day 8. *S. xylosus*, *S.*  
557 *equorum* and *E. hirae* were also positively correlated with 11 compounds, mainly ketones,

558 carboxylic acids and alcohols. *P. commune*, *L. casei/paracasei*, *E. faecalis* and *S. brevicaulis*  
559 were positively correlated with the same 11 compounds but also showed high correlations to  
560 11 others compounds, namely octan-3-one, oct-1-en-3-ol, 2-ethylhexanoic acid, octanoic acid,  
561 2-hydroxypentan-3-one, 3-methylheptan-2-one, nonanal, decan-2-one, nonanoic acid,  
562 methylsulfonylmethane and ethyl butanoate. Among these species, *S. brevicaulis* showed a  
563 slightly different correlation profile as strong and positive correlations were also observed for  
564 ethyl hexanoate, ethyl acetate and 1-hydroxypropan-2-one. High correlations for those  
565 compounds were otherwise observed only with *E. faecium*. Finally, strong positive  
566 correlations were also punctually observed.

#### 567 **4. Discussion**

568 In this study, microbial community dynamics during PDO Pélardon cheese-making and  
569 ripening were unraveled for the first time by culture-dependent and -independent approaches  
570 and linked to the cheese aroma profile.

571 The experimental set-up aimed at combining extensive culture-dependent and culture-  
572 independent approaches to precisely and efficiently study the microbial communities  
573 associated to Pélardon cheese, raw materials and the cheese-making environment. Cultural  
574 methods and isolate identifications rely on tedious work and time-consuming techniques that  
575 often limit the analysis scale in terms of sample number and collected isolates. In the present  
576 study, we used MALDI-TOF MS as a high-throughput technique combined with molecular  
577 sequencing to describe cheese microbial communities at a species-level, based on a recently  
578 described microbial culturomics strategy (Lagier et al., 2015; Sarhan et al., 2019). Although  
579 MALDI-TOF identifications have been recently used in food-related microbial diversity  
580 studies (Nacef et al., 2017; Peruzzy et al., 2019), this is the first time, to our best knowledge,  
581 that this method is used to study both bacterial and fungal dynamics in a fermented product. It  
582 allowed us to analyze a large collection of autochthonous isolates (n = 2877) from different

583 media targeting dominant and sub-dominant populations. Taking into account both culture-  
584 dependent and independent data, 29 genera were identified during Pélardon cheese-making,  
585 12 of which were identified by both approaches (Supplementary Fig. S8). Among those,  
586 *Lactococcus* and *Leuconostoc* for bacteria and *Geotrichum* for fungi were identified by both  
587 approaches as the most dominant during curdling while during ripening and in rinds,  
588 *Brevibacterium*, *Arthrobacter*, *Staphylococcus*, *Penicillium* and *Scopulariopsis* were found  
589 but in different proportion according to the method. Interestingly, enterococci were only  
590 identified by the culture-dependent approach with increasing species diversity during  
591 ripening; a result that was completely overlooked by the culture-independent approach.  
592 Culture-independent approach however, further identified five genera, mainly highlighting the  
593 subdominant halophilic bacteria (i.e. *Brachybacterium*, *Salinococcus*, *Marinhabitans*) on the  
594 Pélardon cheese surface.

595 Our dynamic study of Pélardon microbial diversity revealed inconsistencies regarding  
596 microbial dominances during ripening between approaches. Indeed, culture-dependent results  
597 revealed that *L. lactis*, after dominating during acidification, decreased during ripening. In  
598 contrast, metabarcoding results suggested that *L. lactis* remained dominant in core during the  
599 entire ripening. Moreover, most surprisingly, no *Lactiplantibacillus* OTUs were detected in  
600 any samples by the culture-independent approach, while it was shown to increase and  
601 dominate during ripening by the culture-dependent approach. To eliminate any potential  
602 discrepancies due to technical errors qPCR tests targeting *L. paracasei* were performed on  
603 ripening samples to confirm culture-dependent data. *L. paracasei* was detected at high levels  
604 during ripening in both cores and rinds. These results have thus led us to assume that the  
605 dominance of *L. lactis* DNA from both live and dead cells following curdling masked any  
606 lactobacilli DNA in the studied samples using the culture-independent approach. Indeed,  
607 autolysis of starter bacteria, such as *L. lactis* in our study, during cheese ripening is a well-

608 known phenomenon although the involved mechanisms are not well established (Lazzi et al.,  
609 2016; Treimo et al., 2006). It is also possible that *L. lactis* cells were in a Viable But Not  
610 Cultivable state (VBNC). Bacteria in VBNC state has been discussed in the general context of  
611 cheese production (Edalatian et al., 2012) and to a lesser extent for cheese starter cultures  
612 (Falentin et al. 2012; Ruggirello, Dolci and Cocolin 2014), however their active metabolic  
613 contribution have yet to be determined. Similar hypotheses could be done to explain the  
614 differences in the proportions of *G. candidum*, *Penicillium* spp. and *S. brevicaulis* at each  
615 ripening stage analyzed by the two approaches. Discrepancies between culture-dependent and  
616 culture-independent results have been repeatedly reported regardless of the culture-  
617 independent method used (Coton et al., 2017; Delbès et al., 2007) and our study led to a  
618 different “picture” of the microbial communities present during ripening. Based on the  
619 literature findings reported above and on our observations, the 3-step culture-dependent  
620 approach we implemented in this study provided the most thorough overview of the microbial  
621 community dynamics during Pélardon cheese-making although metabarcoding highlighted the  
622 presence of some specific micro-organisms. Moreover, similar microbial analyses were  
623 performed on another batch six months before and the same dominant species were found and  
624 discrepancies between the two approaches also observed (data not shown). We thus decided  
625 to mainly consider the culture-dependent results for correlation analysis.

626 As previously mentioned, Pélardon cheese is produced without any starter. Thus, the  
627 plausible origin of the microorganisms identified during cheese-making and ripening was  
628 investigated in the raw materials and cheese-making environment. Raw milk microbiota was  
629 dominated by coagulase negative Staphylococci as previously reported by Tormo et al. (2011)  
630 although most of the identified species did not persisted after acidification. However, results  
631 highlighted the contribution of whey as a major source of starter LAB, mainly *L. lactis* and in  
632 a lesser extent *L. mesenteroides*. This finding is consistent with other cheese technologies

633 using backslopping such as Rocamadour or Parmigiano Regiano, in which the main acidifying  
634 bacteria are also brought by the whey (Coloretti et al., 2016; Demarigny et al., 2006).  
635 However, in the case of Pélardon cheese, the whey also acted as a source of fungal  
636 populations involved in ripening as it contained high populations of *G. candidum*. Besides, *G.*  
637 *candidum* and the other most dominant fungal species were also found all over the cheese-  
638 factory. In addition, some species identified on Pélardon cheeses, especially *Glutamicibacter*  
639 and *Staphylococcus* species, were also found on several surfaces but not in raw materials. Our  
640 results thus strongly suggested that the working environment likely contributes to the  
641 enrichment of the surface microbiota of these cheeses during ripening. Overall, it would be of  
642 interest in the future to investigate to what extent the microbiota revealed here is shared  
643 between other Pélardon producers and if the specific microorganisms we identified as part of  
644 the core microbiota can be considered as a “house microbiota” as previously described by  
645 Bokulich and Mills (2013) for two artisanal washed-rinds cheeses.

646 Cheese aroma compound composition is particularly relevant for artisanal cheeses as it  
647 strongly contributes to their typicity and influences the consumer’s preference. In this study, a  
648 correlation between the identified bacteria and fungi, the main fermentation products and  
649 aroma compounds was established. Our results revealed dynamics importance of both  
650 communities early-on in the cheese-making process. Indeed, whey used as an inoculum was  
651 dominated by *L. lactis*, the main acidifying microorganism, along with *L. mesenteroides* and  
652 none of the other species identified in raw milk persisted in curd. These two species were well  
653 correlated to the major lactose fermentation products such as lactic acid and acetic acid but  
654 also to hexanal and 3-hydroxybutan-2-one (acetoin) compounds associated with buttery and  
655 fresh aromas. Lactic and acetic acids rapidly decreased during drying suggesting their  
656 utilization, which is consistent with the significant pH increase. *L. mesenteroides* was also  
657 correlated to high abundances of leucine and valine degradation products such as 3-



658 methylbutanoic and 2-methylpropanoic acid (Delgado et al., 2011), which abundances were  
659 highest at day 8. This would suggest that *L. mesenteroides* is also actively involved during the  
660 post-acidification stage which is often related to proteolysis activities (McSweeney and  
661 Sousa, 2000). *G. candidum* was highly associated with the drying and early ripening stages  
662 and is most likely responsible for the observed deacidification phenomenon. Moreover, lactic  
663 acid catabolism by *G. candidum* is well-established in cheeses such as Camembert and Brie  
664 (Boutrou and Guéguen, 2005). Its deamination activity has also been reported and linked to  
665 the production of 2-phenylacetaldehyde and 2-phenylethanol from phenylalanine, compounds  
666 that we found to considerably increase in Pélardon cheese during drying (day 8) and early  
667 ripening (day 14) stages. Interestingly, both compounds, found at high abundances at the  
668 same stages, are known to bring floral and sweet aromas to the product (Majcher et al., 2014).  
669 Prolonged ripening for up to three months seems to be linked to the dominance of two non-  
670 starter lactic acid bacteria, *L. paracasei* and *E. faecalis*, and two filamentous fungi, *P.*  
671 *commune* and *S. brevicaulis*, and associated with most of the identified volatile compounds.  
672 The strongest positive correlations were observed with some ketones, which originate from  
673 free fatty acid catabolism by different microorganisms in particular fungi and have low odor  
674 thresholds with strong “earthy” and “cheesy” notes. Interestingly, *S. brevicaulis* was  
675 previously identified in cow milk ripened cheeses (Schornsteiner et al., 2014) and natural  
676 ripened cheeses as a core microorganism (Wolfe et al., 2014). This species has also been  
677 associated with high proteolytic activity (Bothast et al., 1975) and considered as a cheese  
678 spoiler when it predominates. Pélardon late ripening stages were also characterized by high  
679 abundances of free fatty acids, which can be released by either microbial or milk lipases  
680 (Collins et al., 2003). Their concentrations have similarly been shown to increase with aging  
681 in other goat cheeses (Delgado et al., 2011) and hexanoic and decanoic acids, also designated  
682 as caprylic and caproic acids, respectively, have previously been reported as aroma markers of

683 goat cheese. In contrast, day 14 cheeses, which corresponds to the minimal ripening time  
684 required for the PDO Pélardon denomination, were not correlated with high abundances of  
685 specific compounds. Thus, the typical aroma of Pélardon cheese probably results from a  
686 balance between compounds, associated with fresh buttery and floral aromas, and as the  
687 ripening progresses by compounds that bring increased strong, earthy and goaty aromas,  
688 potentially attenuated by esters associated with fruity aromas.

689 Overall, these results highlight the dynamics of Pélardon **cheese microbial community**.  
690 Through correlation analysis we identified the key-players of Pélardon cheese microbial  
691 communities. However, more in-depth studies are necessary to first validate and secondly  
692 further explore their functional role. By implementing a culture-dependent approach, we were  
693 able to constitute a large collection of autochthonous isolates, hence working on individual  
694 species or reconstructed consortia could help to better characterize their metabolic potential,  
695 decipher interactions between the key-players and further demonstrate their contribution to  
696 Pélardon cheese aroma typicality.

697

## 698 **Funding**

699 This work, which was conducted in the framework of the ProMedFoods project ‘Promotion of  
700 local Mediterranean fermented foods through a better knowledge and management of  
701 microbial resources’, was funded through the ARIMNet2 2016 Call by the following funding  
702 agencies: ANR (France), MERS (Algeria), ELGO-DEMETER (Greece), MIPAF (Italy), INIA  
703 (Spain) and MHESR (Tunisia). ARIMNet2 (ERA-NET) has received funding from the  
704 European Union’s Seventh Framework Programme for research, technological development  
705 and demonstration under grant agreement [618127].

## 706 **Declaration of competing interest**

707 The authors declare that they have no known competing financial interests or personal  
708 relationships that could have appeared to influence the work reported in this paper.

## 709 **Acknowledgements**

710 We would like to thank our cheesemaker partner for accepting to be part of our project, for  
711 welcoming us at their cheese-making facilities and providing the samples. We would like to  
712 thank Geraldine Pascal and Olivier Rué for their technical support and advice for the FROGS  
713 bioinformatic pipelines. Finally, we are grateful to the INRAE MIGALE bioinformatics  
714 facility (MIGALE, INRAE, 2020. Migale bioinformatics Facility, doi:  
715 10.15454/1.5572390655343293E12) for providing computing and storage resources.

## 716 **Data and microbial resources availability**

717 Sequence data generated in this study are publicly available on the European Bioinformatics  
718 Institute database (www.ebi.ac.uk) under accession numbers PRJEB37923 (bacterial 16S  
719 rRNA sequencing run data) and PRJEB37922 (fungal ITS2 sequencing run  
720 data). Representative isolates will be deposited in culture collections: UBOCC for fungi and  
721 CIRM-BIA for bacteria and accessible on demand.

## 722 **References**

- 723 Achilleos, C., Berthier, F., 2013. Quantitative PCR for the specific quantification of  
724 *Lactococcus lactis* and *Lactobacillus paracasei* and its interest for *Lactococcus lactis* in  
725 cheese samples. *Food microbiology* 36, 286–95.  
726 <https://doi.org/10.1016/j.fm.2013.06.024>
- 727 Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J.,  
728 Clements, D., Coraor, N., Grüning, B.A., Guerler, A., Hillman-Jackson, J., Hiltemann,  
729 S., Jalili, V., Rasche, H., Soranzo, N., Goecks, J., Taylor, J., Nekrutenko, A.,  
730 Blankenberg, D., 2018. The Galaxy platform for accessible, reproducible and  
731 collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* 46, W537–W544.  
732 <https://doi.org/10.1093/nar/gky379>
- 733 Bertuzzi, A.S., Walsh, A.M., Sheehan, J.J., Cotter, P.D., Crispie, F., McSweeney, P.L.H.,  
734 Kilcawley, K.N., Rea, M.C., 2018. Omics-Based Insights into Flavor Development  
735 and Microbial Succession within Surface-Ripened Cheese. *mSystems* 3.  
736 <https://doi.org/10.1128/mSystems.00211-17>
- 737 Bokulich, N.A., Mills, D., 2013. Facility-Specific “House” Microbiome Drives Microbial

738 Landscapes of Artisan Cheesemaking Plants. Applied and environmental  
739 microbiology 79. <https://doi.org/10.1128/AEM.00934-13>

740 Bothast, R.J., Lancaster, E.B., Hesseltine, C.W., 1975. Scopulariopsis brevicaulis: Effect of  
741 pH and substrate on growth. European J. Appl Microbiol. 1, 55–66.  
742 <https://doi.org/10.1007/BF01880620>

743 Boutrou, R., Guéguen, M., 2005. Interests in Geotrichum candidum for cheese technology.  
744 Int. J. Food Microbiol. 102, 1–20. <https://doi.org/10.1016/j.ijfoodmicro.2004.12.028>

745 Calasso, M., Ercolini, D., Mancini, L., Stellato, G., Minervini, F., Di Cagno, R., De Angelis,  
746 M., Gobbetti, M., 2016. Relationships among house, rind and core microbiotas during  
747 manufacture of traditional Italian cheeses at the same dairy plant. Food Microbiology  
748 54, 115–126. <https://doi.org/10.1016/j.fm.2015.10.008>

749 Carpino, S., Randazzo, C., Pino, A., Russo, N., Rapisarda, T., Belvedere, G., Caggia, C.,  
750 2016. Influence of PDO Ragusano cheese biofilm microbiota on flavour compounds  
751 formation. Food Microbiology 61. <https://doi.org/10.1016/j.fm.2016.09.006>

752 Collins, Y.F., McSweeney, P.L.H., Wilkinson, M.G., 2003. Lipolysis and free fatty acid  
753 catabolism in cheese: a review of current knowledge. International Dairy Journal 13,  
754 841–866. [https://doi.org/10.1016/S0958-6946\(03\)00109-2](https://doi.org/10.1016/S0958-6946(03)00109-2)

755 Coloretti, F., Chiavari, C., Nocetti, M., Reverberi, P., Bortolazzo, E., Musi, V., Grazia, L.,  
756 2016. Whey starter addition during maturation of evening milk: effects on some  
757 characteristics of cheese milk and Parmigiano–Reggiano cheese. Dairy Sci. &  
758 Technol. 96, 185–197. <https://doi.org/10.1007/s13594-015-0257-y>

759 Coton, M., Pawtowski, A., Taminiau, B., Burgaud, G., Deniel, F., Coulloume-Labarthe, L.,  
760 Fall, A., Daube, G., Coton, E., 2017. Unraveling microbial ecology of industrial-scale  
761 Kombucha fermentations by metabarcoding and culture-based methods. FEMS  
762 Microbiol Ecol 93. <https://doi.org/10.1093/femsec/fix048>

763 De Filippis, F., Parente, E., Ercolini, D., 2018. Recent Past, Present, and Future of the Food  
764 Microbiome. Annu Rev Food Sci Technol 9, 589–608.  
765 <https://doi.org/10.1146/annurev-food-030117-012312>

766 Delbès, C., Ali-Mandjee, L., Montel, M.-C., 2007. Monitoring bacterial communities in raw  
767 milk and cheese by culture-dependent and -independent 16S rRNA gene-based  
768 analyses. Appl. Environ. Microbiol. 73, 1882–1891.  
769 <https://doi.org/10.1128/AEM.01716-06>

770 Delcenserie, V., Taminiau, B., Delhalle, L., Nezer, C., Doyen, P., Crevecoeur, S., Roussey,  
771 D., Korsak, N., Daube, G., 2014. Microbiota characterization of a Belgian protected  
772 designation of origin cheese, Herve cheese, using metagenomic analysis. Journal of  
773 Dairy Science 97, 6046–6056. <https://doi.org/10.3168/jds.2014-8225>

774 Delgado, F.J., González-Crespo, J., Cava, R., Ramírez, R., 2011. Formation of the aroma of a  
775 raw goat milk cheese during maturation analysed by SPME-GC-MS. Food Chem 129,  
776 1156–1163. <https://doi.org/10.1016/j.foodchem.2011.05.096>

777 Demarigny, Y., Sabatier, C., Laurent, N., Prestoz, S., Rigobello, V., Blachier, M.-J., 2006.  
778 Microbial diversity in natural whey starters used to make traditional Rocamadour goat  
779 cheese and possible relationships with its bitterness. Italian Journal of Food Science  
780 18, 261–276.

781 Dolci, P., Alessandria, V., Zeppa, G., Rantsiou, K., Cocolin, L., 2008. Microbiological  
782 characterization of artisanal Raschera PDO cheese: Analysis of its indigenous lactic  
783 acid bacteria. Food Microbiology 25, 392–399.  
784 <https://doi.org/10.1016/j.fm.2007.09.006>

785 Dugat-Bony, E., Straub, C., Teissandier, A., Onésime, D., Loux, V., Monnet, C., Irlinger, F.,  
786 Landaud, S., Leclercq-Perlat, M.-N., Bento, P., Fraud, S., Gibrat, J.-F., Aubert, J., Fer,  
787 F., Guédon, E., Pons, N., Kennedy, S., Beckerich, J.-M., Swennen, D., Bonnarme, P.,

788 2015. Overview of a Surface-Ripened Cheese Community Functioning by Meta-  
789 Omics Analyses. *PLOS ONE* 10, e0124360.  
790 <https://doi.org/10.1371/journal.pone.0124360>

791 Edalatian, M.R., Najafi, M.B.H., Mortazavi, S.A., Alegría, Á., Nassiri, M.R., Bassami, M.R.,  
792 Mayo, B., 2012. Microbial diversity of the traditional Iranian cheeses Lighvan and  
793 Koozeh, as revealed by polyphasic culturing and culture-independent approaches.  
794 *Dairy Science & Technol.* 92, 75–90. <https://doi.org/10.1007/s13594-011-0045-2>

795 Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads.  
796 *Nat. Methods* 10, 996–998. <https://doi.org/10.1038/nmeth.2604>

797 Ercolini, D., Frisso, G., Mauriello, G., Salvatore, F., Coppola, S., 2008. Microbial diversity in  
798 Natural Whey Cultures used for the production of Caciocavallo Silano PDO cheese.  
799 *International Journal of Food Microbiology* 124, 164–170.  
800 <https://doi.org/10.1016/j.ijfoodmicro.2008.03.007>

801 Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S.,  
802 Hernandez-Raquet, G., Combes, S., Pascal, G., 2018. FROGS: Find, Rapidly, OTUs  
803 with Galaxy Solution. *Bioinformatics* 34, 1287–1294.  
804 <https://doi.org/10.1093/bioinformatics/btx791>

805 Falentin, H., Henaff, N., Le Bivic, P., Deutsch, S.-M., Parayre, S., Richoux, R., Sohier, D.,  
806 Thierry, A., Lortal, S., Postollec, F., 2012. Reverse transcription quantitative PCR  
807 revealed persistency of thermophilic lactic acid bacteria metabolic activity until the  
808 end of the ripening of Emmental cheese. *Food Microbiol.* 29, 132–140.  
809 <https://doi.org/10.1016/j.fm.2011.09.009>

810 Glass, N.L., Donaldson, G.C., 1995. Development of primer sets designed for use with the  
811 PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ*  
812 *Microbiol* 61, 1323–1330.

813 Harlé, O., Falentin, H., Niay, J., Valence, F., Courselaud, C., Chuat, V., Maillard, M.-B.,  
814 Guédon, É., Deutsch, S.-M., Thierry, A., 2020. Diversity of the metabolic profiles of a  
815 broad range of lactic acid bacteria in soy juice fermentation. *Food Microbiology* 89,  
816 103410. <https://doi.org/10.1016/j.fm.2019.103410>

817 Kears, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S.,  
818 Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P.,  
819 Drummond, A., 2012. Geneious Basic: An integrated and extendable desktop software  
820 platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–  
821 1649. <https://doi.org/10.1093/bioinformatics/bts199>

822 Kergourlay, G., Taminiau, B., Daube, G., Champomier Vergès, M.-C., 2015. Metagenomic  
823 insights into the dynamics of microbial communities in food. *Int. J. Food Microbiol.*  
824 213, 31–39. <https://doi.org/10.1016/j.ijfoodmicro.2015.09.010>

825 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O.,  
826 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and  
827 next-generation sequencing-based diversity studies. *Nucleic Acids Res* 41, e1.  
828 <https://doi.org/10.1093/nar/gks808>

829 Kurtzman, C.P., Robnett, C.J., 1998. Identification and phylogeny of ascomycetous yeasts  
830 from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences.  
831 *Antonie Van Leeuwenhoek* 73, 331–371. <https://doi.org/10.1023/A:1001761008817>

832 Lagier, J.C., Hugon, P., Khelaifia, S., Fournier, P.E., La Scola, B., Raoult, D., 2015. The  
833 rebirth of culture in microbiology through the example of culturomics to study human  
834 gut microbiota. *Clinical Microbiology Reviews* 28, 237–264.  
835 <https://doi.org/10.1128/cmr.00014-14>

836 Lazzi, C., Povolo, M., Locci, F., Bernini, V., Neviani, E., Gatti, M., 2016. Can the  
837 development and autolysis of lactic acid bacteria influence the cheese volatile

838 fraction? The case of Grana Padano. *International Journal of Food Microbiology* 233,  
839 20–28. <https://doi.org/10.1016/j.ijfoodmicro.2016.06.009>

840 Lê, S., Josse, J., Husson, F., 2008. FactoMineR: An R Package for Multivariate Analysis.  
841 *Journal of Statistical Software* 25, 1–18. <https://doi.org/10.18637/jss.v025.i01>

842 Leyva Salas, M., Mounier, J., Maillard, M.-B., Valence, F., Coton, E., Thierry, A., 2019.  
843 Identification and quantification of natural compounds produced by antifungal  
844 bioprotective cultures in dairy products. *Food Chemistry* 301, 125260.  
845 <https://doi.org/10.1016/j.foodchem.2019.125260>

846 Lindgren, Å., Karami, N., Karlsson, R., Åhrén, C., Welker, M., Moore, E.R.B., Stadler, L.S.,  
847 2018. Development of a rapid MALDI-TOF MS based epidemiological screening  
848 method using MRSA as a model organism. *Eur. J. Clin. Microbiol. Infect. Dis.* 37, 57–  
849 68. <https://doi.org/10.1007/s10096-017-3101-x>

850 Mahé, F., Rognes, T., Quince, C., de Vargas, C., Dunthorn, M., 2015. Swarm v2: highly-  
851 scalable and high-resolution amplicon clustering. *PeerJ* 3, e1420.  
852 <https://doi.org/10.7717/peerj.1420>

853 Majcher, M.A., Myszka, K., Kubiak, J., Jeleń, H.H., 2014. Identification of key odorants of  
854 fried cottage cheese and contribution of *Galactomyces geotrichum* MK017 to the  
855 formation of 2-phenylethanol and related rose-like aroma compounds. *International*  
856 *Dairy Journal* 39, 324–329. <https://doi.org/10.1016/j.idairyj.2014.08.008>

857 McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive  
858 Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 8, e61217.  
859 <https://doi.org/10.1371/journal.pone.0061217>

860 McSweeney, P.L.H., Sousa, M.J., 2000. Biochemical pathways for the production of flavour  
861 compounds in cheeses during ripening: A review. *Lait* 80, 293–324.  
862 <https://doi.org/10.1051/lait:2000127>

863 Montel, M.-C., Buchin, S., Mallet, A., Delbes-Paus, C., Vuitton, D.A., Desmasures, N.,  
864 Berthier, F., 2014. Traditional cheeses: rich and diverse microbiota with associated  
865 benefits. *Int. J. Food Microbiol.* 177, 136–154.  
866 <https://doi.org/10.1016/j.ijfoodmicro.2014.02.019>

867 Nacef, M., Chevalier, M., Chollet, S., Drider, D., Flahaut, C., 2017. MALDI-TOF mass  
868 spectrometry for the identification of lactic acid bacteria isolated from a French  
869 cheese: The Maroilles. *International Journal of Food Microbiology, Special Issue:*  
870 *CBL 20th edition: New challenges for research and industry* 247, 2–8.  
871 <https://doi.org/10.1016/j.ijfoodmicro.2016.07.005>

872 Peruzzy, M.F., Murru, N., Yu, Z., Kerkhof, P.-J., Neola, B., Joossens, M., Proroga, Y.T.R.,  
873 Houf, K., 2019. Assessment of microbial communities on freshly killed wild boar  
874 meat by MALDI-TOF MS and 16S rRNA amplicon sequencing. *International Journal*  
875 *of Food Microbiology* 301, 51–60. <https://doi.org/10.1016/j.ijfoodmicro.2019.05.005>

876 Quéro, L., Girard, V., Pawtowski, A., Tréguer, S., Weill, A., Arend, S., Cellière, B.,  
877 Polsinelli, S., Monnin, V., van Belkum, A., Vasseur, V., Nodet, P., Mounier, J., 2019.  
878 Development and application of MALDI-TOF MS for identification of food spoilage  
879 fungi. *Food Microbiol.* 81, 76–88. <https://doi.org/10.1016/j.fm.2018.05.001>

880 Quigley, L., O’Sullivan, O., Beresford, T.P., Ross, R.P., Fitzgerald, G.F., Cotter, P.D., 2012.  
881 High-Throughput Sequencing for Detection of Subpopulations of Bacteria Not  
882 Previously Associated with Artisanal Cheeses. *Appl. Environ. Microbiol.* 78, 5717–  
883 5723. <https://doi.org/10.1128/AEM.00918-12>

884 Ruggirello, M., Cocolin, L., Dolci, P., 2016. Fate of *Lactococcus lactis* starter cultures during  
885 late ripening in cheese models. *Food Microbiol.* 59, 112–118.  
886 <https://doi.org/10.1016/j.fm.2016.05.001>

887 Ruggirello, M., Dolci, P., Cocolin, L., 2014. Detection and viability of *Lactococcus lactis*

888 throughout cheese ripening. PLoS ONE 9, e114280.  
889 <https://doi.org/10.1371/journal.pone.0114280>  
890 Sarhan, M.S., Hamza, M.A., Youssef, H.H., Patz, S., Becker, M., ElSawey, H., Nemr, R.,  
891 Daanaa, H.-S.A., Mourad, E.F., Morsi, A.T., Abdelfadeel, M.R., Abbas, M.T., Fayez,  
892 M., Ruppel, S., Hegazi, N.A., 2019. Culturomics of the plant prokaryotic microbiome  
893 and the dawn of plant-based culture media – A review. *Journal of Advanced Research*,  
894 *Special Issue on Plant Microbiome* 19, 15–27.  
895 <https://doi.org/10.1016/j.jare.2019.04.002>  
896 Schornsteiner, E., Mann, E., Bereuter, O., Wagner, M., Schmitz-Esser, S., 2014. Cultivation-  
897 independent analysis of microbial communities on Austrian raw milk hard cheese  
898 rinds. *International Journal of Food Microbiology* 180, 88–97.  
899 <https://doi.org/10.1016/j.ijfoodmicro.2014.04.010>  
900 Smith, C.A., Want, E.J., O’Maille, G., Abagyan, R., Siuzdak, G., 2006. XCMS: processing  
901 mass spectrometry data for metabolite profiling using nonlinear peak alignment,  
902 matching, and identification. *Anal. Chem.* 78, 779–787.  
903 <https://doi.org/10.1021/ac051437y>  
904 Toju, H., Tanabe, A.S., Yamamoto, S., Sato, H., 2012. High-Coverage ITS Primers for the  
905 DNA-Based Identification of Ascomycetes and Basidiomycetes in Environmental  
906 Samples. *PLOS ONE* 7, e40863. <https://doi.org/10.1371/journal.pone.0040863>  
907 Tormo, H., Agabriel, C., Lopez, C., Lekhal, D. a. H., Roques, C., 2011. Relationship between  
908 the production conditions of goat’s milk and the microbial profiles of milk.  
909 *International Journal of Dairy Science* 6, 13–28.  
910 Treimo, J., Vegarud, G., Langsrud, T., Rudi, K., 2006. Use of DNA quantification to measure  
911 growth and autolysis of *Lactococcus* and *Propionibacterium* spp. in mixed  
912 populations. *Appl. Environ. Microbiol.* 72, 6174–6182.  
913 <https://doi.org/10.1128/AEM.00515-06>  
914 Van Hoorde, K., Vandamme, P., Huys, G., 2008. Molecular identification and typing of lactic  
915 acid bacteria associated with the production of two artisanal raw milk cheeses. *Dairy*  
916 *Science & Technology* 88, 445–455.  
917 Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA  
918 amplification for phylogenetic study. *J. Bacteriol.* 173, 697–703.  
919 <https://doi.org/10.1128/jb.173.2.697-703.1991>  
920 Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*, 2nd ed, Use R! Springer  
921 International Publishing. <https://doi.org/10.1007/978-3-319-24277-4>  
922 Wolfe, B.E., Button, J.E., Santarelli, M., Dutton, R.J., 2014. Cheese Rind Communities  
923 Provide Tractable Systems for In Situ and In Vitro Studies of Microbial Diversity.  
924 *Cell* 158, 422–433. <https://doi.org/10.1016/j.cell.2014.05.041>  
925

926 **Table 1. *Lactiplantibacillus paracasei* quantification by qPCR during cheese ripening.** Results are expressed as number of copies per gram of  
 927 cheese based on mean values and standard deviation of triplicate analyses. Means with different letters in the last column are significantly  
 928 different (Tukey's test p<0.05).  
 929  
 930

Sampling time	Sample type	Ct value	<i>tuf</i> gene copy number/g of cheese
day 14	core	27.63 ± 0.71	1.81x10 <sup>6</sup> ± 8.07x10 <sup>5</sup> <sup>a</sup>
day 62	core	22.35 ± 0.95	6.08x10 <sup>7</sup> ± 3.80x10 <sup>7</sup> <sup>b</sup>
	rind	25.66 ± 0.77	5.46x10 <sup>6</sup> ± 2.07x10 <sup>6</sup> <sup>a</sup>
day 90	core	21.81 ± 0.90	7.63x10 <sup>7</sup> ± 3.06x10 <sup>7</sup> <sup>b</sup>
	rind	25.05 ± 0.54	9.09x10 <sup>6</sup> ± 3.04x10 <sup>6</sup> <sup>a</sup>

---

Ct Cyle threshold



931 **Table 2. Microbial counts in environmental samples of the cheese factory**

932

933

934

Sample code	Sample type	Microbial counts		
		Mesophilic aerobic bacteria	Fungi	Presumptive LAB <sup>a</sup>
<b>Surface</b> (log <sub>10</sub> CFU/cm <sup>2</sup> )				
PS1	-Cleaned plastic trays surface used for renneting A	2.95	3.62	-
PS2	Cleaned inox bench used for cheese draining	3.26	3.04	-
PS3	Cleaned plastic mould A	-	-	-
PS4	Cleaned tray stored in the cheese-making room	3.15	3.46	-
PS5	Cheese tray in the drying room	5.96	6.22	6.00
PS6	Cheese tray in the ripening room	6.48	6.23	5.26
PS7	Wall tiles in the making room	4.84	2.00	-
PS8	Cleaned plastic mould B	-	-	-
PS9	Cleaned plastic trays surface used for renneting B	3.38	3.80	2.30
PS10	Milk arrival pipe in the cheesemaking room	-	-	-
<b>Intrants</b> (log <sub>10</sub> CFU/mL)				
PL1	Tap water arriving in the cheesemaking room	-	-	-
PL2	Tap water arriving in the cleaning area	2.77	-	-
PL3	Salt used for cheese dry salting	-	-	-
<b>Air</b> (log <sub>10</sub> CFU/m <sup>3</sup> )				
PAF	Main cheese-making room	>3	2.38	NP
PAS	Drying room	>3	2.34	NP
PAA	Ripening room	>3	2.45	NP

a LAB : Lactic acid bacteria ; - : Population below delection level ; NP : enumeration Not Performed

935 **Table 3. Changes in pH values, sugar and organic acid concentrations (mg/g of cheese) during cheesemaking and ripening.** Mean values  
 936 and standard deviations are given based on triplicate cheeses. Means with different letters within pH values or within a row for acids and sugars  
 937 are significantly different (Tukey's test  $p < 0.05$ ).

	<b>pvalue</b>	<b>Whey</b>	<b>Day 2</b>	<b>Day 8</b>	<b>Day 14</b>	<b>Day 62</b>	<b>Day 90</b>
<b><i>pH (Units)</i></b>	<b><math>3.42E^{-20}</math></b>						
pH in core		$4.33 \pm 0.05^a$	$4.31 \pm 0.07^a$	$4.62 \pm 0.08^b$	$5.02 \pm 0.15^c$	$6.07 \pm 0.03^e$	$6.56 \pm 0.02^f$
pH in rind		/	/	$4.82 \pm 0.06^{bc}$	$5.62 \pm 0.05^d$	$6.73 \pm 0.09^f$	$6.69 \pm 0.05^f$
<b><i>Acids and sugars (mg/g of cheese)</i></b>							
Lactose	<b><math>7.3E^{-17}</math></b>	$23.61 \pm 0.01^a$	$19.96 \pm 0.59^b$	$10.11 \pm 0.32^c$	$0.87 \pm 0.73^d$	ND	ND
Galactose	<b><math>2.36E^{-10}</math></b>	$0.12 \pm 0.03^a$	$0.14 \pm 0^b$	$0.01 \pm 0.01^c$	ND	ND	ND
Lactic acid	<b><math>6.14E^{-13}</math></b>	$5.77 \pm 0^a$	$7.3 \pm 0.32^b$	$4.55 \pm 0.37^c$	ND	ND	ND
Acetic acid	<b><math>2.86E^{-10}</math></b>	$0.29 \pm 0^a$	$0.47 \pm 0.04^b$	$0.08 \pm 0.03^c$	$0.1 \pm 0.02^c$	ND	ND
3-Phenyllactic acid	<b><math>4.91E^{-10}</math></b>	$0.01 \pm 0^a$	$0.01 \pm 0^b$	$0.05 \pm 0^{bc}$	$0.14 \pm 0.01^c$	$0.07 \pm 0.01^d$	$0.06 \pm 0^d$
Citric acid	<b><math>2.13E^{-04}</math></b>	ND	ND	ND	ND	$0.61 \pm 0.33^a$	$0.42 \pm 0.03^a$

ND below quantification and detection limit

938 **Figure captions**

939 **Fig. 1. Sampling strategy and experimental design used in this study for the analysis of**  
940 **an artisanal goat's milk cheese: Pélardon**

941  
942  
943 **Fig. 2. Microbial population counts estimated by culture-dependent method during the**  
944 **cheese-making and ripening of Pélardon goat cheese: (A) in raw materials and curd, (B)**  
945 **in cheese core and (C) in cheese rind.** Mean values and standard deviations are given based  
946 on triplicates. Cultivation conditions for each population are indicated between brackets. In  
947 order to facilitate the comprehension, the curd (day 2) is represented on A, B and C.

948  
949 **Fig. 3. Microbial communities diversity revealed by culture-dependent method during**  
950 **Pélardon cheese-making and ripening in the core (A) and in the rind (B).** Dot plots show  
951 microbial population counts as detailed on Fig. 1 (average values of 3 replicates) whereas bar  
952 plots represent species composition based on isolates identification obtained via MALDI-TOF  
953 analysis followed by sequencing of 16S rRNA gene (bacteria) or ITS/beta-tubulin region  
954 (fungi). The isolate proportion represents the number of isolates identified as a species  
955 divided by the number of isolates all three replicates. In order to facilitate the comprehension,  
956 raw milk, whey used for backslopping and day 2 (curd) are represented on A and B.


957  
958 **Fig. 4. Bacterial community composition during Pélardon cheese-making and ripening**  
959 **determined by a culture-independent approach.** Data were obtained by metabarcoding  
960 analysis targeting the V3-V4 region of the 16S rRNA gene. Results are expressed as  
961 normalized counts.

962  
963 **Fig. 5. Microbial diversity encountered in the environment of the cheese-factory**  
964 **determined using a culture-dependent approach.** Microbial composition based on isolates  
965 identification obtained via MALDI-TOF MS analysis followed by sequencing of 16S rRNA  
966 gene (bacteria) or ITS/beta-tubulin region (fungi). PS refers to surface samples, PL to intrants  
967 samples while PAF, PAS, PAA refer to air samples from cheese-making, drying and ripening  
968 rooms respectively. For detailed information of the samples see Table 2.

969  
970 **Fig. 6. Normalized heatmap showing volatile compounds changes determined by HS-**  
971 **GC-MS in Pélardon cheese samples during cheese-making and ripening.** Clustering based  
972 on Ward's linkage using Euclidean distance. Samples names are on the right, R1, R2 and R3  
973 refer to the cheese replicates used for analysis. Identified compounds are at the bottom part of  
974 the plot. Abundance data were centered and scaled prior to analysis. Color ranging from blue  
975 to red correspond to normalized abundance mean levels. The different frames highlight  
976 compounds with similar profile during cheese-making and ripening.

977  
978 **Fig. 7. Heat-map representing Pearson correlation analysis between species identified**  
979 **using culture-dependent methods and biochemical compounds.** Species are shown at the  
980 bottom, compounds are displayed on the right and both were clustered using on Ward's  
981 linkage using Euclidean distance. Color ranging from red to blue correspond to negative and  
982 positive correlation respectively. Red square highlight the strongest positive correlations  
983 ( $p < 0.001$ ) - \*, compounds quantified using HPLC.


Fig. 1





**Fig 3**


**A**



**Bacteria**

- Acidovorax temperans*
- Arthrobacter russicus*
- Brevibacterium* spp.
- Enterococcus casseliflavus*
- Enterococcus faecalis*
- Enterococcus hirae*
- Enterococcus malodoratus*
- Glutamicibacter* spp.
- Lactobacillus brevis*
- Lactobacillus casei/paracasei*
- Lactococcus lactis*
- Microbacterium* spp.
- Micrococcus luteus*
- Serratia liquefaciens*
- Staphylococcus arlettae*
- Staphylococcus epidermidis*
- Staphylococcus fleurettii*
- Staphylococcus saprophyticus*
- Staphylococcus succinus*
- Staphylococcus warneri*
- Aerococcus viridans*
- Bacillus cereus/thuringiensis*
- Curtobacterium flaccumfaciens*
- Enterococcus durans*
- Enterococcus faecium*
- Enterococcus italicus*
- Enterococcus mundtii*
- Kocuria* spp.
- Lactobacillus casei/paracasei*
- Lactococcus lactis*
- Staphylococcus caprae*
- Staphylococcus equorum*
- Staphylococcus haemolyticus*
- Staphylococcus sciuri*
- Staphylococcus vitulinus/fleurettii*
- Staphylococcus xylosus*


**B**



**Fungi**

- Candida parapsilosis*
- Geotrichum candidum*
- Meyerozyma guilliermondii*
- Penicillium commune*
- Scopulariopsis alboflavescens*
- Trichosporon ovoides*
- Unidentified*
- Debaryomyces hansenii*
- Kluyveromyces lactis*
- Penicillium christenseniae*
- Penicillium glabrum*
- Scopulariopsis brevicaulis*
- Yarrowia lipolytica*


**Fig. 4**



**Species**

- Acetobacter cibinongensis
- Arthrobacter sp.
- Brachybacterium massiliense
- Brachybacterium sp.
- Brevibacterium aureum/linens/oceani
- Brevibacterium sp.
- Glutamicibacter arilaitensis/bergerei/nicotianae
- Glutamicibacter bergerei
- Klebsiella pneumoniae
- Lactococcus lactis
- Lactococcus raffinolactis
- Lactococcus sp.
- Leuconostoc mesenteroides
- Marihabitans sp.
- Psychrobacter alimentarius
- Salinicoccus kunmingensis
- Staphylococcus equorum
- Staphylococcus fleurettii
- Staphylococcus saprophyticus/xylosus
- Other

Fig 5



**Bacteria**

- Acidovorax temperans
- Lactococcus lactis
- Spingomonas paucimobilis
- Staphylococcus sciuri
- Bacillus cereus/thuringiensis
- Leuconostoc mesenteroides
- Staphylococcus epidermidis
- Staphylococcus succinus
- Brevibacterium spp.
- Microbacterium spp.
- Staphylococcus equorum
- Staphylococcus vitulinus/fleurettii
- Curtobacterium flacumfaciens
- Micrococcus luteus
- Staphylococcus fleurettii
- Staphylococcus warneri
- Glutamicibacter spp.
- Rhodococcus spp.
- Staphylococcus saprophyticus
- Staphylococcus xylosus

**Fungi**

- Geotrichum candidum
- Kluyveromyces lactis
- Candida parapsilosis
- Debaryomyces hansenii
- Penicillium commune
- Penicillium glabrum
- Meyerozyma guilliermondii
- Penicillium christenseniae
- Trichosporon ovoides
- Yarrowia lipolytica
- Scopulariopsis alboflavescens
- Scopulariopsis brevicaulis



Fig. 6





Fig. 7

Fig. 7



## Supplementary material

### Supplementary Fig .S1.

**Example of a dendrogram of presumptive *Staphylococcus* species spectra obtained by MALDI-ToF analysis.** Representative isolates chosen for identification by sequencing are indicated with “\*” symbol. Choice was based on heterogeneity within the same cluster and discrepancies between presumptive identification by commercial database and position within the species clusters.

### Supplementary Fig. S2.

**Rarefaction curves of Illumina sequencing data after normalization grouped by cheese-making/ripening stage and sample type.** (A) V3-V4 region of the 16S rRNA gene data (B) ITS2 sequencing data

### Supplementary Fig. S3.

**Alpha-diversity indexes of bacterial (A) and fungal (B) communities during cheese-making and ripening of Pélardon cheese.** Samples are grouped according to the process stage and type.

### Supplementary Fig. S4.

**Composition of fungal communities during Pélardon cheese-making and ripening by culture-independent approach.** Data were obtained by high-throughput metabarcoding analysis targeting the ITS2 region.

### Supplementary Fig. S5.

**Microbial community composition encountered in Pélardon cheese-making environment samples using a culture-independent approach.** (A) bacterial communities (B) fungal communities using metabarcoding analyses targeting the V3-V4 region of 16S rRNA gene for bacteria and the ITS2 region for fungi. See Table 2 for legend of environmental samples

### Supplementary Fig. S6.

**Venn diagrams representing diversity and shared species in cheese and environmental samples using (A) culture-dependent and (B) culture-independent approaches.**

### Supplementary Fig. S7.

**Principal component analysis biplot of microbial counts, culture-dependent (red), culture-independent (green) and biochemical data (acids and volatile compounds in blue) at the different cheese-making and ripening stages.** Data are projected on dimensions 1 and 3 and only variables with a  $\cos^2$  superior to 0.6 are displayed.

### Supplementary Fig.S8.

**Venn diagrams comparing the number of genera and species identified using culture-dependent (red circles) and culture-independent (green circles) approaches for both bacteria (A) and fungi (B).** Dominant genera identified by either of culture-dependent or independent approaches throughout the cheese-making or ripening are indicated in bold.

### Supplementary Table S1.

**Primers used in this study**


### Supplementary Table S2.

**Volatile compounds identified and quantified (AU) during Pélardon cheese-making and ripening using HS-GSMS.**





## Supplementary Fig. S1.

Example of a dendrogram of presumptive *Staphylococcus* species spectra obtained by MALDI-TOF analysis. Representative isolates chosen for identification by sequencing are indicated with “\*” symbol. Choice was based on heterogeneity within the same cluster and discrepancies between presumptive identification by commercial MALDI-TOF database and position within the species clusters.



**Supplementary Fig. S2**


**Rarefaction curves of Illumina sequencing data after normalization grouped by cheese-making/ripening stage and sample type. (A) V3-V4 region of the 16S rRNA gene data (B) ITS2 sequencing data**




**Supplementary Fig. S3.**

**Alpha-diversity indexes of bacterial (A) and fungal (B) communities during cheese-making and ripening of Pélardon cheese. Samples are grouped according to the process stage and type.**

**A**  $\alpha$ -diversity indexes of bacterial communities as a function of sample type and process stage



**B**  $\alpha$ -diversity indexes of fungal communities as a function of sample type and process stage










**Supplementary Fig.S5.**

**Microbial community composition encountered in Pélardon cheese-making environment samples using a culture-independent approach.** (A) bacterial communities (B) fungal communities using metabarcoding analyses targeting the V3-V4 region of 16S rRNA gene for bacteria and the ITS2 region for fungi. See Table 2 for legend of environmental samples.




Supplementary Fig. S6.

Venn diagrams representing diversity and shared species in cheese and environmental samples using (A) culture-dependent and (B) culture-independent approaches.



## Supplementary Fig. S7


Principal component analysis biplot of microbial counts, culture-dependent (red), culture-independent (green) and biochemical data (acids and volatile compounds in blue) at the different cheese-making and ripening stages. Data are projected on dimensions 1 and 3 and only variables with a  $\cos^2$  superior to 0.6 are displayed.



**Supplementary Fig. S8.**

**Venn diagrams comparing the number of genera and species identified using culture-dependent (red circles) and culture-independent (green circles) approaches for both bacteria (A) and fungi (B).**

Dominant genera identified by either of culture-dependent or independent approaches throughout the cheese-making or ripening are indicated in bold.



**Supplementary Table S1. Primers used in this study**

<b>Primer</b>	<b>Sequence</b>	<b>Target</b>	<b>Reference</b>
ITS4	5'-TCCTCCGCTTATTGATATGC-3'	ITS region	
ITS5	5'-GGAAGTAAAAGTCGTAACAAGG-3'		
Bt2a	5'-GGTAACCAAATCGGTGCTGCTTTC-3'	$\beta$ -tubulin gene	Glass and Donaldson (1995)
Bt2b	5'-ACCCTCAGTGTAGTGACCCTTGGC-3'		
NL1	5'-GCATATCAATAAGCGGAGGAAAAG-3'	D1-D2 domain	Kurtzman and Robnett (1998)
NL4	5'-GGTCCGTGTTTCAAGACGG-3'		
fD1	5'-CCGAATTCGTCGACAACAGAGTTTGATCCTGGCTCAG-3'	16S rRNA gene	Weisburg et al. (1991)
rP2	5'-CCCGGGATCCAAGCTTACGGCTACCTTGTTACGACTT-3'		
TufIparacF	5'-TCCGGGAAGTCTCAGC-3'	tuf gene	modified from Achilleos and Berthier (2013)
TufIparacR	5'-TGTTTCACGAACAGGTG-3'		
S-D-Bact-0341-b-S-17	5'-CCTACGGGNGGCWGCAG-3'	V3-V4 region of 16S rDNA gene	Klindworth et al. (2013)
S-D-Bact-0785-a-A-21	5'-GACTACHVGGGTATCTAATCC-3'		
ITS3f	5'-GCATCGATGAAGAACGCAGC-3'	ITS2 region	Toju et al. (2012)
ITS4_Kyo1	5'-TCCTCCGCTTWTGWTWTGC-3'		

**Supplementary Table S2. Volatile compounds identified and quantified (AU) during Pélardon cheese-making and ripening using HS-GSMS.**

Compound	Pubchem #	LRI	EIC (m/z)	Identification <sup>a</sup>	Pvalue <sup>b</sup>	Abundance (AU) <sup>b</sup>						Odor descriptor <sup>c</sup>
						Whey	Curd (day 2)	Post-drying (day 8)	Ripening -2weeks (day 14)	Ripening-2months (day 62)	Ripening-3months (day 90)	
<i>Alcohols</i>												
Ethanol	702	950	57	RI, DB	1.98E-01	6.8E+08 ± 4.8E+07 <sup>a</sup>	6.0E+07 ± 4.2E+06 <sup>a</sup>	3.2E+08 ± 2.2E+08 <sup>a</sup>	4.0E+08 ± 5.1E+08 <sup>a</sup>	5.1E+08 ± 2.9E+08 <sup>a</sup>	4.7E+08 ± 2.0E+08 <sup>a</sup>	Alcoholic
Heptan-2-ol	10976	1355	83	RI, DB	<b>2.06E-04</b>	5.1E+05 ± 4.9E+04 <sup>c</sup>	1.2E+05 ± 1.9E+04 <sup>c</sup>	1.0E+05 ± 4.0E+04 <sup>c</sup>	5.7E+07 ± 1.1E+07 <sup>bc</sup>	1.9E+08 ± 8.6E+07 <sup>ba</sup>	2.4E+08 ± 9.7E+07 <sup>a</sup>	Citrus, earthy, bitter
Nonan-2-ol	12367	1568	69	RI, DB	<b>5.83E-04</b>	4.1E+06 ± 1.4E+06 <sup>bc</sup>	1.2E+06 ± 4.3E+05 <sup>c</sup>	7.7E+06 ± 1.1E+07 <sup>bc</sup>	3.7E+07 ± 1.5E+07 <sup>bc</sup>	9.6E+08 ± 3.6E+08 <sup>ba</sup>	1.5E+09 ± 7.8E+08 <sup>a</sup>	Waxy, fruity, dairy, musty
Oct-1-en-3-ol	18827	1482	57	RI, DB	<b>2.44E-07</b>	2.5E+07 ± 1.3E+06 <sup>c</sup>	6.8E+06 ± 1.3E+06 <sup>c</sup>	1.2E+07 ± 3.1E+06 <sup>c</sup>	8.7E+07 ± 3.7E+07 <sup>bc</sup>	1.9E+09 ± 4.2E+08 <sup>a</sup>	5.9E+08 ± 2.2E+08 <sup>b</sup>	Fungal, earthy
2-Phenylethanol	6054	NA	91	DB	<b>3.68E-05</b>	2.7E+07 ± 1.4E+06 <sup>c</sup>	4.5E+06 ± 3.0E+06 <sup>c</sup>	3.5E+08 ± 1.4E+08 <sup>ab</sup>	5.1E+08 ± 8.8E+07 <sup>bc</sup>	1.4E+08 ± 8.9E+07 <sup>bc</sup>		Floral, sweet
<i>Aldehydes</i>												
3-Methylbut-2-enal	61020	1193	84	RI, DB	<b>1.60E-09</b>	2.9E+07 ± 7.7E+05 <sup>a</sup>	1.6E+07 ± 8.1E+05 <sup>b</sup>	1.3E+07 ± 2.2E+06 <sup>c</sup>	9.7E+06 ± 2.6E+05 <sup>c</sup>	1.3E+07 ± 1.3E+06 <sup>c</sup>	1.3E+07 ± 1.3E+05 <sup>c</sup>	Fruity, nutty, almond
3-Methylbutanal	11552	908	44	RI, DB, S	<b>6.30E-06</b>	5.2E+08 ± 6.1E+07 <sup>b</sup>	2.3E+08 ± 1.4E+07 <sup>b</sup>	8.0E+09 ± 2.6E+09 <sup>a</sup>	1.1E+09 ± 1.2E+08 <sup>b</sup>	6.8E+08 ± 2.0E+08 <sup>b</sup>	6.7E+08 ± 4.1E+07 <sup>b</sup>	Fruity, green
Hexanal	6184	1084	56	RI, DB	<b>8.26E-10</b>	1.6E+08 ± 2.5E+07 <sup>a</sup>	1.2E+08 ± 1.1E+07 <sup>b</sup>	3.3E+06 ± 5.4E+05 <sup>c</sup>	2.0E+06 ± 3.1E+05 <sup>c</sup>	4.0E+06 ± 5.9E+05 <sup>c</sup>	4.9E+06 ± 3.8E+06 <sup>c</sup>	Green, fresh
Heptanal	8130	1183	70	RI, DB, S	<b>1.09E-06</b>	4.2E+07 ± 2.7E+06 <sup>c</sup>	1.8E+07 ± 1.8E+06 <sup>c</sup>	9.9E+05 ± 4.6E+05 <sup>c</sup>	2.5E+07 ± 1.3E+07 <sup>a</sup>	1.9E+08 ± 3.7E+06 <sup>b</sup>	2.8E+08 ± 8.2E+07 <sup>a</sup>	Green, fatty, fresh
Nonanal	31289	1390	70	RI, DB	<b>2.13E-03</b>	2.6E+07 ± 8.3E+05 <sup>b</sup>	3.8E+06 ± 2.5E+06 <sup>b</sup>	4.3E+06 ± 6.8E+04 <sup>b</sup>	1.3E+07 ± 9.7E+06 <sup>b</sup>	2.7E+08 ± 7.2E+07 <sup>ba</sup>	5.3E+08 ± 3.3E+08 <sup>a</sup>	Fatty, rose
2-Phenylacetaldehyde	998	1641	120	RI, DB, S	<b>7.43E-10</b>	1.9E+06 ± 6.2E+05 <sup>c</sup>	1.3E+06 ± 7.7E+04 <sup>c</sup>	5.7E+07 ± 8.4E+06 <sup>a</sup>	3.8E+06 ± 3.6E+06 <sup>b</sup>	3.5E+06 ± 1.5E+06 <sup>c</sup>	4.4E+05 ± 1.2E+05 <sup>c</sup>	Green, floral, honey
<i>Fatty acids</i>												
Acetic acid	176	1451	45	RI, DB, S	<b>1.62E-06</b>	2.2E+10 ± 3.3E+09 <sup>a</sup>	2.4E+10 ± 3.3E+09 <sup>a</sup>	1.1E+10 ± 3.4E+09 <sup>b</sup>	1.1E+10 ± 2.3E+09 <sup>b</sup>	3.7E+09 ± 3.3E+09 <sup>bc</sup>	4.1E+08 ± 7.0E+08 <sup>c</sup>	Acidic, sour
Butanoic acid	264	1630	73	RI, DB, S	<b>9.86E-07</b>	9.7E+06 ± 5.7E+06 <sup>c</sup>	5.3E+06 ± 3.9E+06 <sup>c</sup>	1.1E+07 ± 1.1E+07 <sup>c</sup>	2.7E+09 ± 4.6E+08 <sup>bc</sup>	3.6E+09 ± 2.4E+09 <sup>a</sup>	9.0E+09 ± 2.4E+09 <sup>a</sup>	Cheesy, sour, dairy
Propanoic acid	1032	1544	74	RI, DB, S	8.50E-01	3.9E+08 ± 2.4E+08 <sup>a</sup>	4.0E+08 ± 1.9E+08 <sup>a</sup>	5.5E+08 ± 4.2E+08 <sup>a</sup>	5.7E+08 ± 2.7E+08 <sup>a</sup>	3.7E+08 ± 2.6E+08 <sup>a</sup>	5.6E+08 ± 1.8E+08 <sup>a</sup>	Acidic, dairy, fruity
Pentanoic acid	7991	1747	73	DB, S	<b>9.04E-07</b>	8.2E+06 ± 1.5E+06 <sup>b</sup>	9.0E+06 ± 1.5E+06 <sup>b</sup>	8.2E+06 ± 1.5E+06 <sup>b</sup>	2.6E+07 ± 3.5E+06 <sup>a</sup>	5.8E+07 ± 1.1E+07 <sup>a</sup>	7.5E+07 ± 1.7E+07 <sup>a</sup>	Cheesy, acidic
Hexanoic acid	8892	NA	73	DB, S	<b>6.95E-09</b>	3.6E+08 ± 7.7E+07 <sup>d</sup>	5.5E+08 ± 1.1E+08 <sup>d</sup>	1.5E+08 ± 5.3E+07 <sup>d</sup>	2.0E+09 ± 3.0E+08 <sup>c</sup>	3.7E+09 ± 2.2E+08 <sup>a</sup>	6.2E+09 ± 2.2E+08 <sup>a</sup>	Fatty, cheesy, goaty
Heptanoic acid	8094	NA	87	DB, S	<b>1.46E-06</b>	5.8E+06 ± 9.8E+05 <sup>c</sup>	6.1E+06 ± 2.4E+06 <sup>c</sup>	5.7E+06 ± 1.6E+06 <sup>c</sup>	1.5E+07 ± 3.1E+05 <sup>b</sup>	2.5E+07 ± 5.1E+06 <sup>a</sup>	2.5E+07 ± 5.1E+06 <sup>a</sup>	Waxy, cheesy
Octanoic acid	379	NA	101	DB, S	<b>1.06E-02</b>	7.3E+07 ± 2.7E+07 <sup>b</sup>	1.3E+08 ± 3.4E+07 <sup>ab</sup>	5.4E+07 ± 2.7E+07 <sup>b</sup>	1.7E+08 ± 3.3E+07 <sup>ab</sup>	2.5E+08 ± 7.4E+07 <sup>ab</sup>	4.1E+08 ± 2.3E+08 <sup>a</sup>	Fatty, rancid, cheesy
Nonanoic acid	8158	NA	115	DB	<b>2.77E-02</b>	6.3E+06 ± 2.1E+06 <sup>ab</sup>	6.1E+06 ± 2.7E+06 <sup>ab</sup>	5.4E+06 ± 2.0E+06 <sup>b</sup>	8.8E+06 ± 7.0E+05 <sup>ab</sup>	8.1E+06 ± 6.6E+05 <sup>ab</sup>	1.1E+07 ± 8.0E+06 <sup>a</sup>	Waxy, cheesy mild sweet
Decanoic acid	2969	NA	73	DB, S	6.22E-01	8.1E+07 ± 2.4E+07 <sup>a</sup>	8.8E+07 ± 3.2E+07 <sup>a</sup>	5.7E+07 ± 3.5E+07 <sup>a</sup>	8.1E+07 ± 1.8E+07 <sup>a</sup>	9.2E+07 ± 3.0E+07 <sup>a</sup>	1.0E+08 ± 4.1E+07 <sup>a</sup>	Rancid (-), sour, fruity
<i>Branched-chain fatty acids</i>												
3-Methylbutanoic acid	10430	1674	60	RI, DB, S	<b>1.20E-04</b>	1.1E+08 ± 1.3E+07 <sup>b</sup>	2.2E+07 ± 5.9E+06 <sup>b</sup>	2.8E+09 ± 1.1E+09 <sup>a</sup>	2.0E+09 ± 6.9E+08 <sup>b</sup>	3.0E+08 ± 2.1E+08 <sup>b</sup>	3.2E+08 ± 2.1E+08 <sup>b</sup>	Cheesy, sour, ripe
2-Methylpropanoic acid	6590	1573	73	RI, DB, S	<b>2.62E-03</b>	9.4E+07 ± 2.9E+06 <sup>bc</sup>	9.7E+06 ± 2.2E+06 <sup>c</sup>	6.2E+08 ± 3.2E+08 <sup>a</sup>	4.3E+08 ± 1.5E+08 <sup>ab</sup>	1.3E+08 ± 5.2E+07 <sup>bc</sup>	1.4E+08 ± 1.1E+08 <sup>bc</sup>	Acidic, cheesy, sour
2-Ethylhexanoic acid	8897	NA	88	DB	<b>7.64E-03</b>	5.2E+06 ± 1.1E+06 <sup>b</sup>	4.7E+06 ± 3.9E+06 <sup>ab</sup>	4.7E+06 ± 3.9E+06 <sup>ab</sup>	1.4E+07 ± 4.9E+06 <sup>a</sup>	2.4E+07 ± 4.8E+06 <sup>ab</sup>		-
<i>Esters</i>												
Ethyl acetate	8857	874	61	RI, DB, S	<b>1.14E-04</b>	4.1E+07 ± 6.9E+05 <sup>b</sup>	4.2E+07 ± 2.1E+06 <sup>b</sup>	1.9E+07 ± 9.5E+06 <sup>b</sup>	6.9E+07 ± 5.1E+06 <sup>b</sup>	2.5E+07 ± 1.5E+07 <sup>b</sup>	2.0E+08 ± 7.6E+07 <sup>a</sup>	Ethereal
Ethyl butanoate	7762	1032	71	RI, DB, S	<b>1.28E-04</b>	8.2E+07 ± 1.1E+07 <sup>b</sup>	5.0E+06 ± 5.5E+05 <sup>b</sup>	2.6E+07 ± 3.8E+06 <sup>b</sup>	1.1E+08 ± 1.6E+07 <sup>b</sup>	3.8E+08 ± 2.2E+08 <sup>b</sup>	2.6E+09 ± 1.2E+09 <sup>a</sup>	Fruity, fsweet, fresh
Ethyl hexanoate	31265	1233	88	RI, DB, S	<b>1.74E-02</b>	7.0E+06 ± 2.1E+06 <sup>b</sup>	1.3E+04 ± 1.5E+04 <sup>b</sup>	1.2E+08 ± 2.1E+07 <sup>b</sup>	5.9E+07 ± 7.9E+06 <sup>b</sup>	7.9E+07 ± 3.9E+07 <sup>b</sup>	7.7E+08 ± 6.0E+08 <sup>a</sup>	Fruity, etery, pineapple
Ethyl octanoate	7799	1436	88	RI, DB, S	1.37E-01	9.1E+05 ± 3.4E+05 <sup>a</sup>	2.6E+05 ± 9.1E+04 <sup>a</sup>	3.1E+06 ± 1.2E+06 <sup>a</sup>	8.3E+06 ± 4.7E+06 <sup>a</sup>	4.1E+06 ± 3.5E+06 <sup>a</sup>	4.1E+06 ± 3.5E+06 <sup>a</sup>	Waxy, musty, mushroom
Ethyl decanoate	8048	1639	101	RI, DB, S	6.44E-02	5.6E+06 ± 4.3E+05 <sup>a</sup>	1.6E+05 ± 1.2E+05 <sup>a</sup>	1.2E+06 ± 3.8E+05 <sup>a</sup>	3.7E+06 ± 3.7E+05 <sup>a</sup>	1.0E+07 ± 5.9E+06 <sup>a</sup>	8.8E+07 ± 8.6E+07 <sup>a</sup>	Waxy, fruity
<i>Ketones</i>												
Butan-2-one	6569	893	43	RI, DB, S	<b>1.10E-07</b>	2.2E+09 ± 7.7E+07 <sup>a</sup>	7.8E+08 ± 5.5E+07 <sup>cd</sup>	3.1E+08 ± 3.2E+07 <sup>e</sup>	3.2E+08 ± 5.5E+07 <sup>de</sup>	9.9E+08 ± 1.7E+08 <sup>bc</sup>	1.4E+09 ± 3.7E+08 <sup>b</sup>	Fruity, green
Butane-2,3-dione	650	975	43	RI, DB, S	<b>2.68E-02</b>	7.7E+09 ± 1.6E+08 <sup>ab</sup>	1.2E+10 ± 2.8E+09 <sup>a</sup>	8.2E+09 ± 1.7E+09 <sup>ab</sup>	8.5E+09 ± 3.2E+09 <sup>ab</sup>	4.0E+09 ± 5.4E+08 <sup>b</sup>	7.4E+09 ± 2.7E+09 <sup>ab</sup>	Buttery, creamy
Pentan-2-one	7895	967	86	RI, DB	<b>2.40E-04</b>	3.4E+07 ± 6.3E+06 <sup>b</sup>	1.4E+08 ± 1.6E+08 <sup>b</sup>	1.7E+07 ± 3.7E+09 <sup>a</sup>	3.8E+09 ± 3.7E+09 <sup>a</sup>	3.8E+09 ± 7.3E+08 <sup>ab</sup>	7.4E+09 ± 3.1E+09 <sup>a</sup>	Fruity, fermented, sweet
Pentane-2,3-dione	11747	1066	57	RI, DB	<b>2.22E-15</b>	4.4E+07 ± 1.1E+06 <sup>b</sup>	1.7E+08 ± 1.3E+06 <sup>a</sup>	1.3E+07 ± 1.9E+06 <sup>d</sup>	1.4E+07 ± 2.5E+06 <sup>cd</sup>	2.0E+07 ± 3.1E+06 <sup>cd</sup>	2.2E+07 ± 6.6E+06 <sup>c</sup>	Buttery, toasted
Hexan-2-one	11583	1080	100	RI, DB, S	<b>3.29E-04</b>	3.8E+05 ± 1.8E+05 <sup>b</sup>	6.0E+05 ± 2.6E+05 <sup>b</sup>	1.9E+04 ± 1.4E+05 <sup>b</sup>	2.8E+07 ± 1.3E+07 <sup>b</sup>	5.4E+07 ± 6.2E+06 <sup>a</sup>	5.8E+07 ± 3.1E+07 <sup>a</sup>	Fruity, fungal, buttery
Heptan-2-one	8051	1181	43	RI, DB, S	<b>1.43E-07</b>	1.1E+08 ± 1.0E+07 <sup>b</sup>	6.6E+07 ± 1.2E+07 <sup>b</sup>	4.7E+07 ± 9.5E+06 <sup>b</sup>	8.7E+09 ± 2.8E+09 <sup>a</sup>	9.7E+09 ± 6.1E+08 <sup>a</sup>	9.7E+09 ± 1.2E+09 <sup>a</sup>	Cheesy, green, waxy
3-Methylheptan-2-one	92927	1195	72	RI, DB	<b>2.29E-05</b>	9.2E+05 ± 5.3E+04 <sup>b</sup>	4.4E+04 ± 3.9E+04 <sup>b</sup>	1.1E+05 ± 3.4E+04 <sup>b</sup>	2.0E+06 ± 7.2E+05 <sup>b</sup>	9.2E+07 ± 7.2E+06 <sup>a</sup>	2.8E+08 ± 1.1E+08 <sup>a</sup>	-
(E)-Hept-5-en-2-one	5363108	1219	112	RI, DB	<b>1.07E-04</b>	1.6E+05 ± 1.4E+05 <sup>b</sup>	1.0E+04 ± 7.3E+03 <sup>b</sup>	9.9E+03 ± 1.0E+04 <sup>b</sup>	9.1E+04 ± 6.9E+04 <sup>b</sup>	2.3E+06 ± 4.1E+05 <sup>ba</sup>		-
Octan-2-one	8093	1280	58	RI, DB, S	<b>7.86E-05</b>	5.6E+06 ± 1.5E+06 <sup>c</sup>	4.1E+06 ± 3.7E+05 <sup>c</sup>	9.0E+06 ± 5.8E+05 <sup>c</sup>	5.0E+07 ± 2.1E+07 <sup>bc</sup>	4.6E+08 ± 1.2E+08 <sup>ab</sup>	8.6E+08 ± 3.7E+08 <sup>a</sup>	Musty, cheesy, parmesan
Octan-3-one	246728	1250	72	RI, DB	<b>3.96E-09</b>	3.4E+06 ± 4.1E+05 <sup>c</sup>	4.9E+04 ± 3.1E+04 <sup>c</sup>	7.0E+06 ± 9.9E+05 <sup>c</sup>	3.4E+07 ± 1.3E+07 <sup>c</sup>	4.6E+08 ± 7.8E+07 <sup>a</sup>	1.4E+08 ± 6.2E+06 <sup>b</sup>	Mushroom, cheesy, herbal
Nonan-2-one	13187	1387	43	RI, DB	<b>6.04E-06</b>	7.4E+07 ± 5.6E+06 <sup>b</sup>	3.9E+07 ± 1.6E+06 <sup>b</sup>	4.4E+07 ± 2.4E+07 <sup>b</sup>	2.2E+09 ± 1.3E+09 <sup>b</sup>	1.0E+10 ± 2.5E+09 <sup>ab</sup>	1.2E+10 ± 3.7E+09 <sup>a</sup>	Green, cheesy
Non-8-en-2-one	21108	1443	58	RI, DB	<b>3.66E-02</b>	1.3E+06 ± 4.2E+05 <sup>a</sup>	2.8E+05 ± 4.8E+04 <sup>a</sup>	4.4E+05 ± 2.6E+05 <sup>a</sup>	2.9E+07 ± 2.2E+07 <sup>a</sup>	3.3E+08 ± 2.9E+08 <sup>a</sup>	2.7E+07 ± 7.5E+06 <sup>a</sup>	Fruity, baked
Decan-2-one	12741	1492	58	RI, DB	<b>1.46E-02</b>	3.4E+06 ± 6.1E+05 <sup>b</sup>	6.8E+05 ± 1.5E+05 <sup>b</sup>	8.0E+05 ± 2.7E+05 <sup>b</sup>	6.5E+06 ± 3.7E+06 <sup>b</sup>	1.0E+08 ± 3.8E+07 <sup>ab</sup>	2.1E+08 ± 1.7E+08 <sup>a</sup>	Floral, fermented, fatty
Undecan-2-one	8163	1596	58	RI, DB	<b>3.07E-02</b>	8.4E+06 ± 9.0E+05 <sup>ab</sup>	3.4E+06 ± 7.5E+05 <sup>b</sup>	3.0E+06 ± 9.8E+05 <sup>b</sup>	3.2E+07 ± 2.8E+07 <sup>ab</sup>	7.1E+08 ± 3.8E+08 <sup>ab</sup>	2.1E+09 ± 1.9E+09 <sup>a</sup>	Fruity, floral, creamy, cheesy
3-Hydroxybutan-2-one	179	1282	43	RI, DB, S	<b>1.12E-09</b>	2.0E+10 ± 6.2E+08 <sup>a</sup>	1.7E+10 ± 5.7E+08 <sup>a</sup>	1.7E+10 ± 3.3E+09 <sup>a</sup>	6.1E+09 ± 4.5E+08 <sup>b</sup>	6.1E+08 ± 8.2E+07 <sup>c</sup>	9.6E+08 ± 2.8E+08 <sup>c</sup>	Buttery, creamy, milky
1-Hydroxypropan-2-one	8299	1301	74	RI, DB	<b>2.64E-02</b>	6.4E+06 ± 4.8E+05 <sup>b</sup>	4.2E+06 ± 4.8E+05 <sup>b</sup>	6.8E+06 ± 1.7E+06 <sup>b</sup>	2.1E+07 ± 1.7E+07 <sup>ab</sup>	4.3E+07 ± 3.6E+07 <sup>ab</sup>	2.8E+08 ± 2.3E+08 <sup>a</sup>	caramellic, sweet
2-Hydroxypentan-3-one	521790	1363	57	RI, DB	<b>3.94E-05</b>	7.7E+07 ± 4.3E+05	9.5E+07 ± 1.4E+06	1.1E+07 ± 3.7E+06	3.8E+07 ± 1.0E+07	1.2E+08 ± 4.0E+07	2.0E+08 ± 5.5E+07	Truffle, earthy, nutty
2-Methylthiolan-3-one	61664	1519	116	RI, DB	<b>4.63E-18</b>	7.4E+07 ± 1.0E+06 <sup>a</sup>	7.2E+07 ± 1.4E+06 <sup>a</sup>	1.0E+07 ± 6.0E+05 <sup>b</sup>	7.4E+06 ± 1.6E+06 <sup>b</sup>	8.6E+05 ± 1.2E+06 <sup>c</sup>	5.2E+05 ± 3.7E+05 <sup>c</sup>	Sulfurous, fruity
4-Methyl-2,3-dihydropyran-6-one	557445	NA	82	DB	<b>6.28E-07</b>	1.1E+06 ± 1.0E+06 <sup>c</sup>	2.7E+05 ± 1.4E+05 <sup>c</sup>	6.5E+05 ± 4.0E+05 <sup>c</sup>	2.3E+06 ± 3.0E+05 <sup>c</sup>	2.6E+07 ± 1.1E+07 <sup>b</sup>	5.0E+07 ± 9.0E+06 <sup>a</sup>	-
<i>Others</i>												
(Methylsulfonyl)methane	12232	1067	94	RI, DB, S	<b>6.45E-12</b>	3.0E+07 ± 1.2E+06 <sup>c</sup>	8.5E+06 ± 1.7E+06 <sup>c</sup>	1.1E+08 ± 1.2E+07 <sup>c</sup>	8.1E+08 ± 5.3E+07 <sup>b</sup>	9.8E+08 ± 4.4E+07 <sup>a</sup>	8.9E+08 ± 9.7E+07 <sup>b</sup>	Sulfurous, malty
Methylsulfonylmethane	6213	NA	79	DB	<b>6.13E-05</b>	4.8E+07 ± 2.3E+07						

**Manuscript : Linking Pélardon artisanal goat cheese microbial communities to aroma compounds during cheese-making and ripening**

**Authors :**

Marine Penland

Hélène Falentin

Sandrine Parayre

Audrey Pawtowski

Marie-Bernadette Maillard

Anne Thierry

Jérôme Mounier

Monika Coton

Stéphanie-Marie Deutsch

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: