Detection and complete genome sequence of a divergent isolate of cucumber fruit mottle mosaic virus on *Coccinia grandis* in Sudan

Cecile Desbiez, Pauline Millot, Catherine Wipf-Scheibel, Gazim Dafalla, Hervé Lecoq

► To cite this version:

Cecile Desbiez, Pauline Millot, Catherine Wipf-Scheibel, Gazim Dafalla, Hervé Lecoq. Detection and complete genome sequence of a divergent isolate of cucumber fruit mottle mosaic virus on *Coccinia grandis* in Sudan. *Archives of Virology*, 2021, 166, pp.1779-1782. 10.1007/s00705-021-05038-6 . hal-03190083

HAL Id: hal-03190083

<https://hal.inrae.fr/hal-03190083>

Submitted on 9 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

Detection and complete sequence of a divergent isolate of cucumber fruit mottle mosaic virus on
Coccinia grandis in Sudan

C. Desbiez¹, P. Millot¹, C. Wipf-Scheibel¹, G. Dafalla², H. Lecoq¹

¹INRAE, UR0407 Pathologie Végétale, F-84140 Montfavet, France

²University of Gezira, Plant Pathology Center, Wad medani, Sudan

Corresponding author: C. Desbiez

Email: cecile.desbiez@inrae.fr

Tel: 33 (0)4 32 72 28 07

Fax: 33 (0)4 32 72 28 42

ORCID number 0000-0001-8691-7423

Abstract

Cucurbit-infecting tobamoviruses known so far belong to six acknowledged or tentative species. Except cucumber green mottle mosaic virus (CGMMV) present worldwide, they are geographically restricted, mostly to Asia, and have not been observed in Africa so far. A tobamovirus isolate infecting a wild *coccinia grandis* plant was collected in central Sudan in 2012. Its host range appeared mostly limited to cucurbits. A full-length sequence was established and it presented 85% nt identity with isolates of cucumber fruit mottle mosaic virus (CFMMV) described in Israel and Korea, whereas aa identity with CFMMV isolates was 92 to 95% depending on the protein. Based on its biological and molecular properties, we suggest to consider the Sudanese isolate as a divergent isolate of CFMMV. This constitutes the first description of CFMMV in Africa. Its high divergence with isolates from Israel and Korea suggests a lack of recent exchanges between CFMMV from Sudan and the other known populations.

A sample of the wild cucurbit *Coccinia grandis* was collected during a survey performed in the Singa region in central Sudan in December 2012. After mechanical inoculation on *Cucurbita pepo*, severe mosaic symptoms were observed, and electron microscope observations revealed the presence of rod-shaped particles suggestive of the presence of a tobamovirus, as well as flexuous particles and pinwheels that were found to correspond to the ipomovirus coccinia mottle virus (CocMoV) [4]. CocMoV was eliminated after mechanical inoculation of *Nicotiana benthamiana*, as confirmed by RT-PCR with specific CocMoV primers [4], but the tobamovirus-like particles were still present and the inoculated *N. benthamiana* plants displayed systemic mosaic symptoms. Mosaic and deformations were observed after mechanical inoculation from *N. benthamiana* to melon and cucumber. Two primer pairs were designed through alignments of full-length sequences of known cucurbit-infecting tobamoviruses: cucumber green mottle mosaic virus (CGMMV) present worldwide [6], cucumber fruit mottle mosaic virus (CFMMV) detected so far only in Israel and Korea [2, 7], kyuri green mottle mosaic virus (KGMMV) found in Korea, Japan, Indonesia and China, cucumber mottle virus (CMoV) described in Japan and zucchini green mottle mosaic virus (ZGMMV) described in Korea and China. Primer pair 1 (TobCuc-1150-5': ACGTTYTCTGTGTGGTTCC and TobCuc-2700-3': CCACADCCVGGMACNCCATC targeted a 1550nt fragment in the replicase region, and pair 2 (TobCuc-4980-5': TMGAYGATGCWATCCACGA and TobCuc-6240-3': GCRGCNGTAGACGCRTCGTC targeted a 1300nt fragment encompassing the complete movement protein and part of the coat protein regions. RT-PCR fragments of the expected size were obtained for the coccinia isolate as well as an isolate from *Luffa acutangula* collected in the same area. Blast analyses on partial sequences revealed 82% and 88% nt identity in the first and second regions respectively between the Sudan isolates and CFMMV isolates from Israel and Korea (GenBank accessions AF321057 and JN226146 respectively), and more than 99% nt identity between the Sudanese coccinia and loofah isolates. Specific primers (Supporting information S1) were designed using the coccinia partial sequences and complete CFMMV sequences of Israeli and Korean isolates (AF321057 and JN226146, 98.2% identity between the two sequences). After RT-PCR amplification, Sanger sequencing and manual assembly, a full-length sequence of 6574

nt was obtained (GenBank accession MT989352). The sequenced virus had a typical tobamovirus genetic organization [1] with a 131 kDa nonstructural protein (1165 amino-acids) terminated by an amber (UAG) stop codon, a 188.6 kDa (1669 aa) protein expressed by readthrough of the amber stop codon, a putative 27.7 kDa (258 aa) movement protein and a 17.4 kDa (163 aa) coat protein. It showed circa 85% overall nt identity (92 to 95% aa identity depending on the protein) with CFMMV isolates from Korea and Israel (6571 and 6562 nt respectively) (Table 1). Nt identity was 72-75% with KGMMV and ZGMMV, 62-63% with CGMMV, CMoV and the tentative tobamovirus watermelon green mottle mosaic virus (WGMMV) described recently in Taiwan and the USA [3], and less than 55% with tobamoviruses outside the “cucurbit” group (Table 1). Multiple sequence alignments were performed with ClustalW included in MEGA6 [8]. The best nt and aa substitution models were determined with MEGA6 and used for building maximum likelihood trees (Figure 1) with 500 bootstrap replicates, confirming the clustering of the Sudanese isolate with CFMMV within the “cucurbit tobamovirus” group, and its closer relation with KGMMV and ZGMMV than with the other cucurbit-infecting tobamoviruses. The species demarcation threshold for tobamoviruses is 90% overall nt identity, but numerous pairs of tobamovirus isolates showing 85 to 90% identity are considered as belonging to the same species (<https://www.ncbi.nlm.nih.gov/sutils/pasc/>). Also, for all proteins, the Sudanese isolate presented more than 92% aa identity with Israeli and Korean CFMMV isolates (Table 1). After mechanical inoculation, *Chenopodium amaranticolor*, *C. quinoa* and tobacco cv. Xanthi showed no symptoms and the virus was not detected by RT-PCR at the apex of the plants (data not shown), whereas CFMMV isolates from Israel and Korea did not infect tobacco but were reported to induce local lesions on *C. amaranticolor* and symptomless systemic infection of *C. quinoa*. Indeed, the biological and molecular differences between the Sudanese isolate and isolates of CFMMV are similar to those between the Y and C isolates of KGMMV [9]. Too few isolates are characterized to determine if there is a molecular continuum between CFMMV-Su and the other isolates, and if differences in biological properties are frequent for this virus. Based on the current knowledge on its biological and

molecular properties, we propose to consider the Sudanese tobamovirus as a divergent isolate of CFMMV rather than a member of a new species

To our knowledge, this is the first observation of CFMMV in Africa and its first detection on *Coccinia grandis* and *Luffa acutangula*. These plants could constitute reservoirs and contribute to virus maintenance and dissemination, whereas CFMMV had been observed previously only on cultivated cucurbits in natural conditions [2, 7]. As observed for the ipomovirus cucumber vein yellowing virus [5], the high sequence divergence between the Sudanese CFMMV isolate and the other isolates indicates that the presence of CFMMV in distant parts of the world is not related to recent exchanges of contaminated material between Sudan and the other places.

Figure legends

Fig. 1: Maximum-likelihood tree obtained for full-length nucleotide sequences of tobamoviruses. Bootstrap values above 70% (500 bootstraps) are indicated for each node. The scale bar represents a genetic distance of 0.2. Subgroups of tobamoviruses infecting members of different plant families are indicated. The Sudanese isolate analyzed here is indicated in bold

Acknowledgements:

We thank the Experimental Infrastructure team of INRAE in Montfavet for their help in the greenhouse experiments, and the Microscopy Platform at INRAE in Montfavet for the electron microscope observations.

Declarations

Funding: this study did not receive any specific funding

Ethical statement: This research did not involve any human participants and/or animals.

Conflict of interest: The authors declare that they have no conflict of interest.

References:

1. Adams MJ, Heinze C, Jackson AO, Kreuze JF, Macfarlane SA, Torrance L (2012) Tobamoviruses. In: King AM, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus Taxonomy*. Elsevier/Academic Press, London, pp 1153-1156
2. Antignus Y, Wang Y, Pearlsman M, Lachman O, Lavi N, Gal-On A (2001) Biological and molecular characterization of a new cucurbit-infecting tobamovirus. *Phytopathology* 91: 565-571
3. Cheng YH, Huang CH, Chang CJ, Jan FJ (2019) Identification and characterisation of watermelon green mottle mosaic virus as a new cucurbit-infecting tobamovirus. *Ann Appl Biol* 174: 31-39
4. Desbiez C, Verdin E, Tepfer M, Wipf-Scheibel C, Millot P, Dafalla G, Lecoq H (2016) Characterization of a new cucurbit-infecting ipomovirus from Sudan. *Arch Virol* 161: 2913-2915
5. Desbiez C, Caciagli P, Wipf-Scheibel C, Millot P, Ruiz Garcia L, Marian D, Dafalla G, Lecoq H (2019) Evidence for long-term prevalence of cucumber vein yellowing virus in Sudan and genetic variation of the virus in Sudan and the Mediterranean Basin. *Plant Pathol* 68: 1268-1275
6. Dombrovsky A, Tran-Nguyen LTT, Jones RAC (2017) Cucumber green mottle mosaic virus: Rapidly Increasing Global Distribution, Etiology, Epidemiology, and Management. *Ann Rev Phytopathol* 55: 231-256
7. Rhee SJ, Hong JS, Lee GP (2014) Infectivity and complete nucleotide sequence of cucumber fruit mottle mosaic virus isolate Cm cDNA. *Arch Virol* 159: 1807-1811
8. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* 30: 2725-2729

9. Yoon JY, Min BE, Choi SH, Ryu KH (2001) Completion of nucleotide sequence and generation of highly infectious transcripts to cucurbits from full-length cDNA clone of Kyuri green mottle mosaic virus. Arch Virol 146: 2085-2096

Table 1 : sequence identity (%) between the Sudanese coccinia tobamovirus isolate and other tobamoviruses, for the full-length nucleotide sequence and for deduced amino-acid sequences of the polymerase (RdRp), movement protein (MP) and coat protein (CP)

Accession	Tobamovirus	Genome			
		(nt)	RdRp (aa)	MP (aa)	CP (aa)
NC_002633.1	Cucumber fruit mottle mosaic virus (CFMMV)	85.5	92.7	95.3	93.8
JN226146	CFMMV-Cm	85.9	94.9	95.3	93.8
NC_003610.1	Kyuri green mottle mosaic virus (KGMMV)	74.6	84.2	78.7	83.2
AB015145	KGMMV-Yodo	73.9	83.3	79.1	82.0
AB162006	KGMMV-YM	72.4	78.7	78.3	80.7
NC_003878.1	Zucchini green mottle mosaic virus (ZGMMV)	73.5	83.5	77.9	77.6
NC_001801.1	Cucumber green mottle mosaic virus (CGMMV)	63.3	66.1	62.8	45.3
NC_008614.1	Cucumber mottle virus (CMoV)	62.1	64.1	59.3	41.4
MH837097	Watermelon green mottle mosaic virus (WGMMV)	61.7	64.0	58.4	42.6
NC_016519.1	Clitoria yellow mottle virus (CIYMV)	49.4	44.8	24.6	41.7
NC_025381.1	Hibiscus latent Fort Pierce virus (HLFPV)	53.5	51.5	27.2	39.9
NC_008310.2	Hibiscus latent Singapore virus (HLSV)	53.3	51.0	28.4	38.7
NC_015552.1	Passion fruit mosaic virus (PafMV)	50.2	47.7	22.2	33.3
NC_008716.1	Maracuja mosaic virus (MarMV)	50.7	47.7	24.9	33.3
NC_014546.1	Frangipani mosaic virus (FrMV)	51.2	45.6	36.4	36.2
NC_026816.1	Plumeria mosaic virus (PluMV)	50.6	45.4	33.9	36.8
NC_001873.1	Turnip vein-clearing virus (TVCV)	50.8	48.1	31.0	34.4
NC_002792.2	Ribgrass mosaic virus (RMV)	51.0	47.8	31.5	33.8
NC_003355.1	Wasabi mottle virus (WMoV)	50.7	47.9	31.9	31.2
NC_004422.1	Youcai mosaic virus (YoMV)	50.7	47.9	32.8	31.8
NC_008365.1	Streptocarpus flower break virus (SFBV)	51.0	46.2	31.9	34.3
NC_010944.1	Brugmansia mild mottle virus (BrMMV)	51.5	47.4	32.9	33.3
NC_003852.1	Obuda pepper virus (ObPV)	50.8	46.0	32.5	32.7
NC_004106.1	Paprika mild mottle virus (PaMMV)	51.7	46.6	31.5	34.0
NC_001556.1	Tobacco mild green mosaic virus (TMGMV)	50.0	46.0	31.9	33.3
NC_030229.1	Tropical soda apple mosaic virus (TSAMV)	51.1	46.4	34.5	36.9
NC_003630.1	Pepper mild mottle virus (PMMoV)	50.8	46.3	33.9	36.3
NC_001728.1	Odontoglossum ringspot virus(ORSV)	50.8	45.4	32.6	35.3
NC_009642.1	Bell pepper mottle virus (BPeMV)	51.1	47.1	31.3	33.8
NC_002692.1	Tomato mosaic virus (ToMV)	51.2	47.5	30.5	32.1
NC_022230.1	Tomato mottle mosaic virus (ToMMV)	50.7	47.8	29.7	30.8

NC_028478.1	Tomato brown rugose fruit virus (ToBRFV)	51.0	47.7	31.1	32.7
NC_022801.1	Yellow tailflower mild mottle virus (YTMMV)	51.4	46.8	34.9	34.8
NC_001367.1	Tobacco mosaic virus (TMV)	51.0	47.3	31.8	35.2
NC_009041.1	Rehmannia mosaic virus (ReMV)	51.2	47.8	30.5	34.0
NC_016442.1	Rattail cactus necrosis associated virus (RCNaV)	49.1	44.3	22.9	27.0
NC_011803.1	Cactus mild mottle virus (CMMoV)	48.7	44.4	20.5	25.2

Figure 1