



HAL
open science

Oak decaying wood harbors taxonomically and functionally different bacterial communities in sapwood and heartwood

S. Mieszkin, P. Richet, C. Bach, C. Lambrot, Laurent Augusto, Marc Buée, S. Uroz

► To cite this version:

S. Mieszkin, P. Richet, C. Bach, C. Lambrot, Laurent Augusto, et al.. Oak decaying wood harbors taxonomically and functionally different bacterial communities in sapwood and heartwood. *Soil Biology and Biochemistry*, 2021, 155, pp.1-12. 10.1016/j.soilbio.2021.108160 . hal-03195736

HAL Id: hal-03195736

<https://hal.inrae.fr/hal-03195736>

Submitted on 14 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Title**

2 Oak decaying wood harbors taxonomically and functionally different bacterial
3 communities in sapwood and heartwood

4

5 **Authors**

6 S. Mieszkin^{a,*}, P. Richet^a, C. Bach^a, C. Lambrot^b, L. Augusto^b, M. Buée^a, S. Uroz^{a,c}

7

8 **Authors' affiliation**

9 ^a Université de Lorraine - INRAE, UMR1136 Interactions Arbres-Microorganismes,
10 Centre INRAE grand Est de Nancy, F-54280 Champenoux, France

11 ^b INRAE, Bordeaux Sciences Agro, UMR 1391 Interactions Sol Plante Atmosphère
12 (ISPA), 33140 Villenave d'Ornon, France

13 ^c INRAE UR1138 Biogéochimie des Ecosystèmes Forestiers, Centre INRA grand Est
14 de Nancy, F-54280 Champenoux, France

15 Corresponding author: Sophie Mieszkin; sophie.mieszkin@univ-brest.fr

16

17 **Running title:** Sapwood and heartwood select different bacterial communities

18

19 **Keywords**

20 Oak decaying wood; bacteria; sapwood; heartwood; functional and metabolic
21 potential; taxonomic diversity

22

23 **Current adress: Laboratoire de Microbiologie des Environnements Extrêmes LM2E, Univ*
24 *Brest, CNRS, Ifremer, UMR 6197, IUEM, Rue Dumont d'Urville, F-29280 Plouzané, France*

25 **Abstract**

26 Wood decay is an important process in forest ecosystems, which relies on wood
27 chemical properties and the action of a complex community of decomposers. While
28 the important role of fungi in this process is recognized, our knowledge concerning
29 the colonization of decaying wood by bacteria, their relative distribution as well as
30 their potential functional roles remain under-investigated. In this context, our aim was
31 to characterize how the bacterial communities are structured at both taxonomic and
32 functional levels along the soil-wood continuum, using oak discs positioned on the
33 forest floor for nine months. Using a combination of culture-dependent and -
34 independent methods associated to a physical measure of wood decay, we
35 evidenced that at the middle stage of decay investigated, the heartwood- and
36 sapwood-inhabiting bacterial communities significantly differed from one another in
37 term of richness and taxonomic composition, but also from those of the bulk soil. The
38 functional screening revealed low metabolic potentials and a higher frequency of
39 cellulose decomposing bacteria in wood than in the bulk soil, suggesting an
40 adaptation of these communities to this habitat and to the physical-chemical
41 conditions occurring in decaying wood. Together, our data evidence that the
42 colonization of decaying wood by bacteria is based on a deterministic process linking
43 extrinsic and intrinsic factors.

44 **1. Introduction**

45 Forest ecosystems, due to the large area they cover on earth (up to 31% of land
46 area in 2015), provide major ecosystem services. One of the most important is their
47 capacity to stock carbon (C) and to act as a C sink (Pan et al., 2011; Seidl et al.,
48 2014). In addition, trees accumulate inorganic nutrients taken from the soil in their
49 biomass. Even if a significant C fraction is directly transferred to the surrounding
50 environment through belowground allocation and root exudation (Litton et al., 2004;
51 Hobbie, 2006), a major transfer comes from the seasonal litter deposition and after
52 the death of trees or harvesting, during the wood decay. Consequently, decaying
53 wood strongly impacts nutrient cycling and plays an important role in term of organic
54 and inorganic nutrients recycling (Augusto et al., 2015; Ricker et al., 2016). In
55 temperate regions, woody debris can be represented by logs, stumps or fine
56 branches and can reach 2-65 m³ ha⁻¹ in managed forests where wood is harvested,
57 while it can reach up to 1200 m³ ha⁻¹ in non-managed forests (Hahn and
58 Christensen, 2005; Stokland et al., 2012). This material is a specific habitat for
59 various organisms, such as insects and fungi, providing them an important source of
60 cellulose, lignin and hemicellulose (Stokland et al., 2012; Seibold et al., 2015;
61 Baldrian, 2017). However, the nutrients entrapped in the wood structure are not
62 directly accessible to the wood biota, due to the recalcitrance of wood to
63 decomposition. Indeed, the high content of lignin and the low nutrient content (mainly
64 nitrogen (N) and phosphorus (P)) are not favorable to microbial development
65 (Meerts, 2002). In addition, presence of toxic wood extractives and low pH conditions
66 prevent wood microbial decomposition. All these parameters vary according to the
67 wood species and the stage of decay, with some wood easily degradable (e.g. *Acer*
68 sp. or *Tilia* sp.) and other more recalcitrant (i.e., *Quercus* sp. or *Acacia* sp.) (Weedon

69 et al., 2009; Rajala et al., 2012; Augusto et al., 2015).

70 A large body of literature suggests that fungi but also insects are the main
71 decomposers of decaying wood. Insects, due to their capability of altering wood
72 structure by tunneling and fragmentation and of decomposing wood through
73 enzymatic digestion, are important engineers of early stage decay (Baldrian et al.,
74 2016; Ulyshen, 2016). Fungi are well known for their ability to produce a large panel
75 of extracellular lignocellulolytic enzymes to decompose the lignin and then the
76 cellulose (Schneider et al., 2012; Mathieu et al., 2013; Baldrian et al., 2016; Noll et
77 al., 2016). In comparison, less attention has been paid to wood-inhabiting bacteria,
78 while they are known to colonize wood since the early steps of decomposition
79 (Clausen, 1996; Blanchette, 2000; Kielak et al., 2016). Based on the analysis of the
80 bacterial communities occurring at different stages of decomposition of pine-wood,
81 Kielak et al. (2016) proposed that bacterial community assembly may be the result of
82 a stochastic process at the initial stages of dead wood decomposition, and on the
83 contrary determined by the wood properties at advanced stages. It was also reported
84 that decaying bacteria communities could be diverse, potentially active at
85 decomposing dead wood and capable of interacting with fungi (Valášková et al.,
86 2009; Hervé et al., 2016; Johnston et al., 2016; Baldrian, 2017). Among the different
87 studies performed on these communities, a dominance of *Proteobacteria*,
88 *Actinobacteria*, *Bacteroidetes* and *Acidobacteria* was reported, suggesting an
89 adaptation to the wood environment (Valášková et al., 2009; Hervé et al., 2014; Sun
90 et al., 2014; Kielak et al., 2016; Rinta-Kanto et al., 2016). Noticeably, some bacterial
91 classes or even genera were shown to differ according to the tree species (Prewitt et
92 al., 2014; Moll et al., 2018) and/or the stage of decay (Hoppe et al., 2015). The
93 potential structuring effect of the wood compartments (heartwood (HW), the dead

94 inner part of wood; sapwood (SW), the living outermost part of wood) on the bacterial
95 communities was also investigated, but such studies remain scarce (Zhang et al.,
96 2008; Moll et al., 2018). However, most of these studies described the community
97 composition, while the relative role of the bacterial communities in the decomposition
98 of wood is still largely underexplored.

99 Most of the current knowledge on the ability of bacteria to decompose wood
100 comes from the analysis of soil bacteria and on their capacities to hydrolyse cellulose
101 (Nicolitch et al., 2017), hemicelluloses (Lee et al., 2010), lignin (Větrovský et al.,
102 2014) or even pectin (Clausen, 1996). For instance, these functional abilities were
103 found in culturable representatives of *Acidobacteria* (*Silvibacterium bohemicum*)
104 (Lladó et al., 2016), of *Actinobacteria* (e.g. *Streptomyces*) (Nicolitch et al., 2016) or of
105 *Proteobacteria* (e.g., *Burkholderia*) (Hervé et al., 2016). Together, these results
106 suggest that the soil may act as a reservoir of bacteria capable of adapting to woody
107 debris and of participating to wood decomposition. Comparatively, few functional
108 analyses have been done on bacteria directly collected *in situ* from decaying wood
109 (Murray and Woodward, 2007; Valášková et al., 2009; Vorob'ev et al., 2009).
110 Consequently, determining whether bacteria are functionally structured according to
111 the stage of wood decay, the wood compartment or according to the tree species are
112 questions of importance in microbial ecology. As bacteria are not considered as the
113 main actors of wood decay in the early stages of decomposition, we investigated the
114 taxonomic and functional distribution of the bacterial communities at middle stage of
115 wood decay and across a soil-wood continuum. At this stage of decay, we
116 hypothesized that i) the direct role and the structure of the bacterial communities vary
117 according to the compartments (*i.e.*, bulk soil (BS) *versus* HW *versus* SW), and ii) the
118 two main wood compartments (HW and SW) select different bacterial communities

119 according to their own properties (i.e., nutritive content, wood biopolymers). To test
120 these hypotheses, we combined metabolic and functional bioassays, with a culture-
121 independent approach on oak wood samples incubated nine months on a forest floor.

122

123 **2. Materials and Methods**

124 **2.1. Experimental site and sampling description**

125 The present study was performed in the forest experimental site of Champenoux
126 (north-eastern of France; Lat: 48.718°; Long: 6.347°; Alt: 248 m; 0.5 ha of surface).
127 The site is characterized as an oak forest aged of 110 years old where *Quercus*
128 *petraea* (*Q. petraea*) and *Carpinus betulus* represent the dominant tree species,
129 followed by some *Fagus sylvatica* and *Quercus robur* trees. The soil is classified as
130 an acidic luvic cambisol (pH-H₂O = 4.6 in the A1 horizon, 0-5 cm) and is
131 characterized by a mull-type humus and a loamy texture. The site is flat with
132 understory vegetation (i.e. oak and hornbeam seedlings, *Convallaria majalis* L.,
133 *Deschampsia cespitosa* L.).

134 At this site, four living oak trees of similar age (ca. 110 years old; 42 cm
135 diameter), spatially distant from 5 to 20 m, were selected. Oak was selected, as it
136 represents a major tree species in Europe and its coverage is expected to increase
137 with changes in temperatures and precipitations due to climate change (Hanewinkel
138 et al., 2013). An area of 25 m² (~5 m x ~5 m) was delimited around each oak tree,
139 and the litter was removed. Then, twenty oak discs were placed around each oak
140 tree directly on the surface of the soil and spatially separated from one another by a
141 distance of approximately 50 cm, giving a total of 80 wood discs. The wood discs
142 came from a sessile oak tree (*Q. petraea*) aged of approximately 48 years and cut off
143 in January 2015 from a neighboring forest plot and then stored and air-dried in the

144 laboratory protected from light and moisture for six months. Their average diameter
145 without the bark was 14.5 cm and their average thickness 2 cm. They were placed
146 directly in contact with the soil and incubated nine months in natural conditions to
147 enable wood decomposition (Fig. SI-1).

148 After a nine months incubation period, sampling was performed (April 2016) to
149 focus on the middle stage of wood decay and on the associated bacterial
150 communities. For each tree area and each oak disc, SW and HW as well as the soil
151 (BS) in contact with the wood discs were sampled. SW and HW samples were
152 recovered by drilling using a sterilized 10-mm drill bit. For each tree area, the
153 sawdust of the twenty oak discs was recovered from the two wood compartments
154 (HW and SW). The samples were then pooled to obtain four composite HW sawdust
155 samples ($n=20$ per composite samples) and four composite SW sawdust samples
156 ($n=20$ per composite samples). To compare the wood-inhabiting bacterial community
157 structures and functions, with those of the adjacent soil, 5 g of BS samples
158 underlying each wood-decaying disc (5 cm depth corresponding to the A1 horizon)
159 were sampled. The BS samples were then pooled for each tree area, leading to four
160 composite BS samples ($n=20$ per composite samples). Each sample was stored in
161 sterile containers and once at the laboratory directly analyzed.

162 **2.2. Soil chemical analyses**

163 For each replicate tree area, the chemical properties of the four BS samples have
164 been determined. Total N and C contents, both obtained after combustion at 1000°C,
165 and P content, were determined according to methods published by Olsen (1954),
166 Duchaufour and Bonneau, (1959) and Duval, (1963). The cation exchange capacity
167 (CEC) was determined according to the cobalti-hexamine method, which is based on
168 the titration of the cobalti-hexamine chloride soil extract. Exchangeable cations (Ca,

169 Mg, Na, K, Fe, Mn and Al) and protons were extracted using cobaltihexamine and
170 determined by inductively coupled plasma spectrometry-atomic emission
171 spectrometry (ICP-AES) for cations and by potentiometric measurement using 0.05
172 M KOH for protons. The pH was determined by the water method using a soil/water
173 ratio of 1:5 (w/v).

174 **2.3. Wood decaying stage and chemical analyses**

175 The variation of density of the SW and HW was assessed during the wood decay
176 directly on specific oak discs incubated on the experimental site. For each replicate
177 tree area, one oak disk at T0 and five oak disks at T3, T6 and T9 months were
178 considered. A Pilodyn penetrometer with a 3 mm pin was used to measure
179 penetration depth (i.e., the density) (Mäkipää and Linkosalo, 2011; Fundova et al.,
180 2018). For each sampling time and oak disc, three measures were performed for
181 both the SW and HW samples. Results were expressed as a mean percentage of
182 penetration depth for each wood compartment at each time point.

183 For each tree area and each wood compartment, the chemical properties of the
184 samples have been determined (four SW and four HW sawdust samples). All the
185 samples were dried at 65°C during 72h, homogenized and then sub-sampled for the
186 different analyses described in the supplementary section. Briefly, C and N were
187 analyzed through total combustion. Nutrients (i.e., P, K, Ca, and Mg) were analyzed
188 after wet mineralisation. Quantification of the cellulose was obtained by successive
189 separations with different solvents using a Soxhlet extractor. Quantification of the
190 lignin content in the wood samples was performed with the Klason method. Finally,
191 the pH was determined for each sample.

192 **2.4. Collection of bacterial strains**

193 For each sample (four BS, four SW and four HW), 1 g of sawdust (SW or HW) or
194 BS was added to 20 mL of sterile distilled water and vortexed three times during 90
195 s. After a serial dilution in sterile distilled water, a volume of 0.1 mL from the dilutions
196 10^{-2} to 10^{-5} , was then spread in triplicate onto 1/10 diluted Tryptic Soy Agar (TSA)
197 medium (Tryptic Soy Broth (TSB) from Difco, 3 g.L⁻¹ and agar 15 g.L⁻¹) containing
198 cycloheximide (100 µg.L⁻¹, final concentration). The pH of the medium was adjusted
199 at 5 to fit with the pH found in oak sawdust and soil. All plates were then incubated at
200 25°C for 5 days. After incubation, the number of colony-forming units (CFU).g⁻¹ from
201 sawdust or soil samples was determined using the appropriate dilutions (10^{-3} and 10^{-4}
202 according to the sample considered) permitting to quantify the densities of
203 culturable bacteria in each compartment. For each compartment (BS, HW and SW)
204 and replicate, and considering the dilution giving a total of 30 colonies per Petri dish,
205 all the bacterial colonies were recovered. Bacterial isolates were then purified by
206 three successive platings on 1/10 diluted TSA at pH 5 to obtain pure cultures. All the
207 bacterial isolates were cryopreserved at -80°C in 35% glycerol. The bacterial
208 collection was composed of 308 bacterial isolates that were distributed as follow: 117
209 BS isolates, 107 SW isolates and 84 HW isolates.

210 **2.5. Metabolic and functional bioassays performed on bacterial isolates**

211 The metabolic potential of each bacterial isolate was determined using Biolog
212 microplates, which contain ninety-five different C sources. The GN2 microplates were
213 selected in our study because they contain relevant substrates related to wood
214 derivatives (ie, cellulose) or fungal metabolites (trehalose). The screening was
215 performed according to the manufacturer's instruction. After 48 h incubation at 25°C,
216 the absorbance was measured at 595 nm using an iMark microplate reader (Bio-Rad,
217 USA) (see supplementary section for detailed information).

218 The capacity of each bacterial isolate to degrade organic matter and specific
219 compounds found in wood (cellulose, lignin, xylan and chitin) as well as their ability to
220 recover iron through the production of siderophores was evaluated. Five different
221 bioassays allowing to measure: i) their cellulolytic ability (carboxy-methyl-cellulose
222 (CMC) assay), ii) their lignolytic ability (Remazol brilliant blue R (RBBR) assay), iii)
223 their xylanolytic ability (xylan assay); iv) their chitinolytic ability (chitin assay) and v)
224 their ability to mobilise iron (the CAS (chromoazurol-S assay), were used. Each
225 bioassay is based on a gelosed medium allowing the measure of a discoloration
226 halo. When compatible with the bioassay, the pH of the medium was adjusted to 5 to
227 stay close to the pH of the wood and the soil. Composition of each medium and the
228 experimental procedure are fully described in the supplementary section.

229 ***2.6. Molecular identification of bacterial strains and phylogenetic*** 230 ***analyses***

231 A fragment of the 16S rRNA gene was amplified using the universal set of primers
232 pA and 907r (Edwards et al., 1989; Lane, 1991). The polymerase chain reaction
233 (PCR) was performed in a total reaction volume of 50 μ L containing 20 μ L of master
234 mix (PCR Taq PCR Master Mix, Qiagen), 2 μ L of each primer (10 μ M) and 2 μ L of
235 bacterial cells inoculum. PCR conditions were the following: 94°C for 4 min, then 30
236 cycles of 30 s denaturation at 94°C, 60 s annealing at 53°C, 90 s extension at 72°C,
237 and a final extension step of 10 min at 72°C. PCR products were then purified using
238 the QIAquick PCR purification kit (Qiagen) and concentration determined using a
239 Nanodrop-1000 spectrometer (NanoDrop Technologies, Wilmington, DE, USA). The
240 Sanger sequencing reactions were performed by MWG Eurofin Operon (Ebersberg,
241 Germany) using the primer 907r. All the sequences were identified using the
242 nucleotide BLAST program (Altschul, 1997). Sequence alignment and distance tree

243 were constructed using SeaView (version 4.7) (Gouy et al., 2010) and by using
244 neighbor-joining algorithms with the Kimura two-parameter correction. Distance trees
245 were constructed and bootstrap values were obtained using 1,000 replicates. Trees
246 were generated using the Dendroscope 3 software (version 3.6.3) (Huson and
247 Scornavacca, 2012).

248 The 16S rRNA Sanger sequences generated for this study were submitted to the
249 Genbank database of NCBI under the accession numbers MN193104 - MN193411.

250 ***2.7. Illumina 16S high-throughput sequencing and data processing***

251 Total DNA was extracted from 0.2 g of soil or oak sawdust (SW and HW) using
252 the Fast DNA SPIN kit for soil extraction kit (MPbiomedical, Illkirch, France) following
253 manufacturer's instructions with slight modifications including a step to grind the
254 samples using a Retsch MM301 mixer (Retsch GbmH, Haan, Germany) and three
255 additional washings with guanidine thiocyanate (5.5 M, pH 7) before to add the
256 binding matrix. This step allows to efficiently denature endogenous nucleases and
257 would facilitate the separation of proteins from the related sugars or polyols (Cox,
258 1968; Mason et al., 2003). To prepare the amplicon libraries for the paired-end
259 Illumina sequencing, the primers 515f and 806r targeting the V4 region have been
260 used (Caporaso et al., 2011). The sample multiplexing, the paired-end Illumina
261 sequencing using 2x250 bp Miseq v3 and the de-multiplexing of the sequences were
262 performed at GeT-PlaGe (Genotoul, Toulouse, France) according to the
263 manufacturer's instructions.

264 Raw sequences processing was performed using FROGS (Find Rapidly OTU with
265 Galaxy Solution) pipeline (Escudié et al., 2018). Paired sequences were merged into
266 contigs with flash (Magoc and Salzberg, 2011), cleaned with cutadapt (Martin, 2011)
267 and chimeras were removed with VSEARCH (Rognes et al., 2016). The clustering to

268 generate Operational Taxonomic Units (OTU) was then performed with SWARM
269 algorithm (Mahé et al., 2014) using a local clustering threshold (97% similarity) and
270 the taxonomic affiliation of each OTU was based on the reference databases NCBI
271 Blast+ on SILVA123 16S (Camacho et al., 2009, Quast et al., 2013). Finally, a table
272 presenting the number of 16S rRNA sequences assigned to each OTU as well as
273 their taxonomic affiliation was generated. OTUs belonging to *Eukaryota* or *Archaea*,
274 chloroplasts or mitochondria were discarded as well as contaminant OTUs generated
275 by the extraction and amplification steps. All the singletons were removed for the
276 OTUs analyses. Finally, the number of sequences per sample ranged from 17,806 to
277 39,132 giving a total of 334,380 sequences and 3,079 OTUs. OTU analyses were
278 then performed using packages in R with a random subsampling to normalize each
279 sample with the smallest number of sequences. The bacterial richness and alpha-
280 diversity (Chao1, ACE, Shannon and inverse Simpson indexes) were calculated
281 based on OTU relative abundance. The Illumina 16S rRNA sequences generated for
282 this study were submitted to the Sequence Read Archive (SRA) of NCBI under the
283 accession number PRJNA55031.

284 To determine both sequence homology and the relative environmental
285 representativeness of the bacterial isolates recovered in our study, the Sanger 16S
286 rRNA gene sequences were compared to the Illumina 16S rRNA gene sequences
287 using the Basic Local Alignment Search Tool (BLAST) algorithm (Altschul et al.,
288 1990). The Illumina 16S rRNA gene sequences of each compartment were used as
289 database and an alignment of the Sanger 16S rRNA gene sequences was performed
290 with NCBI-Blast+ using an e-value cutoff of 1×10^{-5} . This analysis was done on the 253
291 bp overlapping between the Sanger and Illumina 16S rRNA gene sequences.

292 **2.8. Statistical analyses**

293 All statistical analyses were performed using R 3.2.3 software (R Core Team,
294 2015). The effect of the compartment (i.e., BS vs HW vs SW) was tested on the
295 chemistry data and the bioassay data using an analysis of variance (one-way
296 ANOVA, $P < 0.05$), followed by a Tukey test. Wood density differences between the
297 two oak wood compartments (SW and HW) was determined by a t-test. For the
298 culture-dependent approaches, a Chi^2 test ($P < 0.05$) analysis was performed to
299 determine whether the frequency of distribution of the bacterial isolates according to
300 their efficacy depended on their ecological origin.

301 For the Illumina 16S rRNA gene sequences and the related OTUs, the analyses
302 were performed using R packages: phyloseq (McMurdie and Holmes, 2013), ape
303 (Paradis et al., 2004) and vegan (Oksanen et al., 2007). The data were visualized
304 using ggplot 2 (Wickham, 2009). The distribution of the bacterial communities
305 according to their compartments was analyzed by performing multidimensional
306 scaling (MDS) ordination using Bray Curtis matrix distance. Permutational
307 multivariate analysis of variance (PERMANOVA) based on Bray-Curtis distances with
308 999 permutations were performed on the OTU table using the vegan package and
309 the Adonis command. Analysis of variance and subsequent post-hoc Tukey's tests to
310 compare taxonomic and functional differences between the compartments were also
311 performed. Relative abundance values at all taxonomic levels (phylum, class, order,
312 family, genus, OTU) were compared for taxa or OTUs presenting a minimum of 100
313 sequences. These values were transformed using the arcsine square root to achieve
314 a normal distribution. When required, a p adjustment was done using the false-
315 Discovery-Procedure (FDR) of Benjamini and Hochberg (1995) to determine the p-
316 value threshold. Correlations between the major wood phyla and wood chemistry
317 was assessed using Pearson correlations ($P < 0.05$).

318

319 **3. Results**

320 **3.1. Chemical analyses and wood decay stage assessment**

321 The chemical analyses were done after nine months of incubation on the BS and
322 wood samples (SW and HW sawdust) (Table SI-1A, B). The analysis of the pH
323 revealed that BS harbored similar pH to the SW. The C/N ratio was 10 to 26 times
324 lower in BS than in the wood compartments (Tab. SI-1A, B). The comparison of the
325 wood compartments revealed that SW was characterized by significantly higher pH
326 (pH-SW=4.63 vs pH-HW=3.75), higher amounts of N, P and Mg (ANOVA, $P<0.05$)
327 and lower (approximately two times) C/N ratio (ANOVA, $P<0.05$). The content of
328 cellulose and lignin (expressed on percentage relative to the global composition)
329 appeared similar in HW and SW (ANOVA, $P>0.05$) (Tab. SI-1A).

330 After nine months of *in situ* decay, the visual inspection showed a strong fungal
331 colonization of the SW compartment, while the HW compartment appeared poorly
332 colonized (Fig. 1A, B). Significant variations of the wood density were observed
333 based on the pilodyn measurements. Indeed, the penetration increased from T0
334 (25.8% \pm 7.4 depth penetration in SW vs 16.4% \pm 3.2 in HW) to T9 months (55.8% \pm
335 12.6 in SW vs 30.4% \pm 4.5 in HW). A significantly higher decay stage was measured
336 for SW from T3 to T9 months compared to HW (t-Test; $P<0.05$) (Fig. 1C). According
337 to these data, the decay stage of the oak discs after nine months of decay was
338 considered as a middle stage.

339 **3.2. OTUs distribution and diversity indices in soil and decaying wood**

340 From the 213,672 16SrRNA sequences retained after bioinformatics treatments
341 (Tab. SI-2), a total of 2,901 OTUs were generated. Comparison between the
342 compartments revealed that the BS harbors significantly more OTUs (1,141 \pm 189)

343 than the wood compartments (SW: 548 ± 127 and HW: 351 ± 123) (ANOVA,
344 $P < 0.05$). No difference was observed between the SW and HW (ANOVA, $P > 0.05$)
345 (Tab. SI-3). Among the total number of OTUs obtained, 9.8% were shared between
346 the three compartments, while some were compartment specific (BS: 50.2%; SW:
347 14.3% and HW: 7.7% OTUs). Furthermore, 7.4% of the OTUs were shared between
348 the two wood compartments and ca 5% of the OTUs were shared between BS and
349 the wood compartments (Fig. SI-2). The observed (i.e., number of OTU observed)
350 and the estimated richness (i.e., chao1, ACE) as well as the alpha-diversity (i.e.,
351 Shannon, Inv. Simpson) were significantly higher in BS compared to the wood
352 compartments (ANOVA, $P < 0.05$). These indices tended to be higher in the SW than
353 in the HW, but only significantly for the Shannon values (Tab. SI-3). Most of the
354 OTUs obtained were assigned at the phylum (99.8%-23 different phyla) to family
355 (88.1%) level, but only 66.4% of these sequences were assigned at the genus level.
356 A global description of the taxonomic affiliation of these OTUs is presented in the
357 table SI-4.

358 **3.3. Structure and taxonomic composition of the bacterial communities in** 359 **soil and decaying wood.**

360 The properties of the different compartments considered (BS, SW and HW)
361 strongly determined the bacterial community structure (i.e., BS vs HW vs SW;
362 $F=10.6$; $R^2=0.70$; $P=0.01$; PERMANOVA and Fig. 2A). Noticeably, the bacterial
363 community structure significantly differed between SW and HW (i.e., $F=3.17$; $R^2=$
364 0.35 ; $P=0.03$; PERMANOVA). A MDS analysis done on the OTUs confirmed this
365 compartment effect (Fig. 2A, B). A detailed analysis of the taxonomic composition
366 revealed that the wood-inhabiting bacterial community was significantly dominated by
367 the *Proteobacteria* (59 to 73.4% of the 16S rRNA sequences in SW and HW,

368 respectively) compared to the BS compartment (ANOVA, $P < 0.05$) (Fig. 2A, Tab. SI-
369 5). *Alphaproteobacteria* was the most abundant class in wood with 28.8% in SW and
370 41.7% in HW. A similar trend was observed for *Betaproteobacteria* (wood > BS;
371 ANOVA, $P < 0.05$). Notably, our analyses revealed that the SW was significantly
372 enriched in *Bacteroidetes* (ANOVA, $P < 0.05$). In contrast, the BS bacterial community
373 was mainly represented by *Acidobacteria*, followed by *Proteobacteria* (with
374 *Alphaproteobacteria* being the most represented class) and *Verrucomicrobia*
375 (12.8%). The relative abundance of *Acidobacteria* and *Verrucomicrobia* was
376 significantly higher in BS than in HW and SW (ANOVA, $P < 0.05$). Noticeably, the
377 relative abundance of the *Bacteroidetes* (mainly present in SW) was positively
378 correlated with wood pH ($P = 0.01$) and P concentration ($P = 0.02$), while it was
379 negatively correlated with the C/N ratio ($P = 0.02$). At the opposite, the relative
380 abundance of the *Alphaproteobacteria* (dominant in HW) was positively correlated
381 with the C/N ratio ($P = 0.001$) but negatively correlated with P concentrations ($P = 0.02$)
382 and the total amount of N ($P = 0.02$) (Fig. SI-3).

383 Differences, at the OTUs level, between wood compartments and between wood
384 and BS were also observed. Dominant and specific OTUs of each compartment are
385 presented in Fig. 3, Table SI-6 and 7.

386 **3.4. Taxonomic affiliation of the bacterial isolates and relationship with** 387 **their ecological origin**

388 The quantification of the culturable bacteria by dilution-plating revealed that BS
389 harbored significantly higher densities ($9.03 \pm 0.14 \text{ Log}_{10} \text{ CFU g}^{-1}$ of soil) than the
390 SW (8.09 ± 0.08 ; ANOVA, $P < 0.05$), but was not different from HW (8.80 ± 0.30 ;
391 ANOVA, $P > 0.05$). HW harbored slightly higher densities than the SW (ANOVA,
392 $P < 0.05$).

393 The taxonomic affiliation of the bacterial isolates revealed that most of them were
394 affiliated to *Proteobacteria* (77.7%), *Actinobacteria* (15.5%), *Acidobacteria* (4.5%)
395 and *Bacteroidetes* (2.3%). The relative distribution of the 30 genera detected is
396 presented in Figure 4 and Table SI-8. Notably, ca. 10% of the bacterial isolates had
397 less than 97% similarity with the 16S rRNA gene sequence of a known culturable
398 bacterial strain deposited in NCBI. Most of these particular isolates came from SW.

399 When the taxonomic affiliation of the bacterial isolates was linked to their
400 ecological niche (BS vs HW vs SW), our analysis showed that isolates belonging to
401 the *Beta*- and *Gammaproteobacteria* were common to the three compartments (Fig.
402 4; Tab. SI-8). However, we noticed that the SW compartment was significantly
403 enriched in *Gammaproteobacteria* isolates (SW, 57.8% of the isolates; BS, 24.4%
404 and HW, 17.8%; Chi^2 , $P < 0.002$), whereas, the BS compartment was significantly
405 enriched in *Betaproteobacteria* isolates (BS, 49.6% of the isolates; SW, 30.6% and
406 HW, 19.8%; Chi^2 , $P < 0.003$). Interestingly, all the isolates belonging to *Acidobacteria*
407 and *Bacteroidetes* (*Alphaproteobacteria*) were exclusively found in wood (SW and/or
408 HW). Several genera were strongly represented and only detected in wood. This is
409 the case of *Sphingomonas*, *Mucilaginibacter* and *Luteibacter* in SW, and *Acidocella*,
410 *Acidipila* and *Novosphingobium* in HW. At the opposite, *Actinobacteria* strains
411 assigned to *Streptomyces*, *Kitasatospora*, *Arthrobacter* and *Streptacidiphilus* were
412 dominant in BS. Strains assigned to the *Paraburkholderia* and *Dyella* genus were
413 recovered in the three compartments considered, with *Paraburkholderia* representing
414 34.7% of the total of the isolates.

415 **3.5. Metabolic and functional potentials of the culturable bacterial** 416 **communities**

417 The metabolic bioassays based on Biolog plates revealed that the BS bacterial
418 isolates harbored significantly higher metabolic potentials than those of the decaying-
419 wood compartments (BS>SW=HW) (ANOVA; $P<0.05$) (Fig. SI-4). Such difference
420 was observed at both substrate and substrate category levels (Fig. SI-4 and Fig 5).
421 Comparisons between wood bacterial isolates showed that HW isolates had a higher
422 metabolic potentials for substrates belonging to (i) the carboxylic acid group (ca. 2
423 fold higher than SW for quinic acid, sebacic acid, succinamic acid, D-glucosaminic
424 acid, glycyl-L-glutamic acid) (ii) the amino acid group (ca. 2 fold higher than SW for
425 L-histidine); (iii) the miscellaneous group (15 fold higher than SW for thymidine); and
426 finally, (iv) the amine-amide group (aminoethanol, 3 fold higher). Beta-methyl-
427 D-glucoside was almost the unique substrate more effectively metabolized by the SW
428 bacterial isolates (4 fold higher than HW isolates) (Fig. 5).

429 Among the functional bioassays performed, the cellulose decomposition assay
430 gave the highest number of positive bacterial isolates. Indeed, ca. 12.6% of the
431 bacterial isolates tested appeared capable of degrading cellulose (Fig. SI-5).
432 Significantly more bacteria isolated from HW (ca. 28.6%) were capable of degrading
433 cellulose, than those coming from SW (13.4%) and BS (10.3%) (χ^2 ; $P<0.0001$). In
434 term of effectiveness, the BS and HW bacterial isolates tended to be more effective
435 at degrading cellulose (halo diameters: $0.21\text{cm}\pm 0.04$ for BS isolates and
436 $0.20\text{cm}\pm 0.02$ for HW isolates) than the SW isolates ($0.12\text{cm}\pm 0.02$; ANOVA
437 $P=0.07$). In the wood compartments, the ability to degrade cellulose was mainly
438 encountered for bacterial isolates assigned to the genus *Mucilaginibacter* (relative
439 efficacy to degrade cellulose (halo diameter): $0.81\text{cm}\pm 0.51$), *Novosphingobium*
440 ($0.50\text{cm}\pm 0.51$) and *Acidisoma* ($0.48\text{cm}\pm 0.41$). Noticeably, 71.4% of the
441 *Mucilaginibacter* isolates were capable of hydrolyzing cellulose, while only 54.5% for

442 *Acidisoma* and 38.5% for *Novosphingobium* (Fig. 6). In contrast, in the BS
443 compartment this function was mainly encountered for bacterial isolates assigned to
444 the genus *Kitasatospora* ($2.05\text{cm}\pm 1.61$) and *Streptomyces* ($0.73\text{cm}\pm 1.21$) (Fig. 6).

445 Concerning the ability to degrade chitin, only 8.5% of the bacterial isolates were
446 capable of degrading this compound and were mainly isolated from BS (45.5% of the
447 isolates capable to hydrolyze chitin were originated from BS) (Fig. SI-5). The BS and
448 HW bacterial isolates tended to be more effective at degrading chitin. The bacterial
449 genera presenting the highest potential were assigned to the genus *Collimonas* (halo
450 diameter: $1.83\text{cm}\pm 0.49$; only recovered from the BS compartment), *Luteibacter*
451 (0.91 ± 0.74 ; only recovered from the HW compartment) and *Streptomyces*
452 ($0.62\text{cm}\pm 0.72$; only recovered from the BS compartment). While all of the *Collimonas*
453 isolates were effective at hydrolyzing chitin, only 57.1% and 25% of the *Luteibacter*
454 and *Streptomyces* isolates did it (Fig. 6).

455 For the other functions related to organic matter decomposition, the ability to
456 degrade xylan was only detected for seven bacterial isolates (2.8% of the bacterial
457 collection) (Fig. 6) and none of the bacterial isolates was capable of degrading lignin.
458 Concerning the ability to mobilize iron, ca. 24% of the bacterial isolates tested were
459 positives (Fig. SI-5). Significantly higher proportions were observed in BS (72.9% of
460 the isolates mobilizing iron) than in HW (18.6%) or SW (8.5%) (Chi^2 ; $P < 0.001$). The
461 bacterial genera presenting the highest potentials to mobilize iron were assigned to
462 the genus *Pseudomonas* (relative efficacy to mobilize iron (halo diameter):
463 $2.08\text{cm}\pm 0.98$) and in a lesser extend to *Collimonas* isolates ($1.48\text{cm}\pm 0.69$) with
464 respectively 85.7% and 100% of the isolates of these genera being effective for iron
465 mobilization. The two other genera: *Paraburkholderia* ($0.7\text{cm}\pm 0.93$) and

466 *Streptomyces* (0.2cm±0.49) were significantly less efficient than *Pseudomonas*
467 isolates (ANOVA, $P<0.05$) (Fig.6).

468 **3.6. Comparison of the Sanger and Illumina 16S RNA gene sequences**

469 This analysis highlighted that all the 308 bacterial isolates of our collection had a
470 minimum of 97% of sequence homology with at least one OTU of the Illumina
471 sequence dataset. When the highest homology was considered, this analysis
472 revealed that the bacterial isolates corresponded to 65 OTUs identified in the Illumina
473 dataset. These OTUs represented 55.4% of the total number of Illumina 16S rRNA
474 sequences. Interestingly, the most represented bacterial isolates in our collection (at
475 the genus level) were also among the dominant in the OTUs (Rank in the collection /
476 rank in the OTU table; *Paraburkholderia* (1/1), *Acidisoma* (2/5), *Dyella* (3/12),
477 *Acidocella* (6/4) and more distantly *Pseudomonas* (7/114) (data not shown).

478

479 **4. Discussion**

480 To date, a few studies have investigated the taxonomic diversity and especially
481 the metabolic and functional role of decaying wood-inhabiting bacteria (Murray and
482 Woodward 2007; Folman et al., 2008; Hervé et al., 2016). However, this aspect is of
483 importance to better understand the decaying wood process and how bacteria
484 contribute to the recycling of the nutritive elements accumulated in the tree biomass.
485 Therefore, to investigate the structure and the potential role of the bacterial
486 communities, we used oak discs installed on the floor of an oak forest site, mimicking
487 partially in this way the conditions when trees fall on the soil and initiate a
488 decomposition process. A combination of culture-dependent and -independent
489 approaches combined with chemical analyses were used and comparisons with bulk
490 soil bacteria was made to highlight the soil-decaying wood continuum.

491 **4.1. Edaphic origin of decaying oak wood-inhabiting bacteria.**

492 How decaying-wood is colonized by microorganisms is rather difficult to
493 determine. Indeed, part of the microorganisms found in decaying wood may originate
494 from the tree tissues (i.e., endophytes), atmospheric deposits or from the surrounding
495 BS (Baldrian et al., 2016). Previous analyses provided clear evidences that whether
496 or not an initial microbiota exists inside trees during their life, its biomass and
497 diversity changed importantly and quickly between the early and later stages of wood
498 decay (Tláškal et al., 2017). The Venn diagram analyses done in our study suggest
499 that a part of the wood-inhabiting bacterial communities is also part of the BS
500 bacterial communities (10.6%). The other OTUs only detected in wood may
501 represent minor groups occurring in BS but not detected by sequencing or may be
502 coming from other locations (i.e., atmosphere, endophytes of the oak discs, woody
503 debris fallen on the oak discs). Interestingly, Probst et al. (2018) also observed that a
504 significant part of the OTUs identified in decaying wood were shared with the
505 surrounding BS, supporting an edaphic origin of wood-inhabiting bacteria. The
506 detailed analysis of the culture-dependent approach also indicated that several
507 bacterial genera were detected in both the BS and wood compartments including
508 bacterial isolates assigned to *Paraburkholderia*, *Dyella* and *Pseudomonas*. These
509 bacterial genera have frequently been described in forest soils, underlying their ability
510 to adapt to various habitats such as decaying wood (Folman et al., 2008; Lladó et al.,
511 2016; Nicolitch et al., 2017). Supporting this hypothesis, Hervé et al. (2014)
512 demonstrated that *Burkholderia* and *Dyella* strains isolated from BS and inoculated in
513 wood sawdust microcosms became dominant after a succession of enrichment
514 steps. Those results support the idea that the BS in forest ecosystem represents a
515 seedbank with a high diversity of functional groups capable of decomposing wood.

516 This is not surprising as the topsoil is continuously in contact with litter and wood
517 debris. Consequently, the abundance of wood-inhabiting bacteria depends on the
518 turnover of tree biomass (wood, litter) with bacteria returning to the surrounding soil
519 at the end of the wood decomposition process (Probst et al., 2018).

520 **4.2. Metabolic and functional potential of decaying oak wood-inhabiting** 521 **bacterial communities differ from soil bacterial communities.**

522 Decaying oak wood-inhabiting bacterial communities may directly or indirectly
523 contribute to wood decomposition. In this study, we tested the hypothesis of a direct
524 role of bacteria in wood decay. To do it, a functional screening targeting the main
525 wood constituents (cellulose, xylan and lignin) was performed. We also focused on
526 chitin because it is an essential constituent of fungal cell walls, and the
527 decomposition of the wood is characterized also by a dynamic succession of different
528 fungi during the whole process (Rajala et al., 2015), generating an important fungal
529 necromass potentially recyclable by bacteria. Mobilization of iron was also studied
530 because iron plays a role in the oxidative process of wood decomposition (Goodell et
531 al., 1997). Our results showed that cellulose degrading bacteria were more frequent
532 in wood, especially in HW, than in the surrounding BS. However, BS was
533 characterized by bacterial isolates presenting a higher effectiveness at degrading
534 cellulose, xylan or chitin or at mobilizing iron than HW and SW. These results differ
535 from those obtained by Hervé et al. (2016) where there was no functional difference
536 observed between bacteria isolated from BS and those adapted to beech decaying
537 wood. However, our experiment was done *in situ* and over a longer period, which
538 may explain the differences we observed. Looking at the taxonomic identity of the
539 strains, we found that representatives of the *Kitasatospora* genus, isolated from BS,
540 were among the most effective at degrading cellulose. Nonetheless, representatives

541 of the *Mucilaginibacter*, *Acidisoma* and *Novosphingobium* genera, all isolated from
542 wood (SW and HW), were also effective. Except for *Novosphingobium*, these genera
543 were previously identified as important cellulose decomposers (Štursová et al., 2012;
544 Lladó et al., 2016; Lopez-Mondejar et al., 2016; Nicolitch et al., 2016). The absence
545 of bacterial strains able to degrade lignin confirms the results obtained by Murray and
546 Woodward (2007) and Hervé et al. (2016). Those results suggest that the ligninolytic
547 activity is probably mainly attributable to fungi (Janusz et al. 2017). However,
548 bacterial strains affiliated to the *Actinomycetes*, *Alphaproteobacteria* and
549 *Gammaproteobacteria* have been reported for their ability to break down lignin (Bugg
550 et al., 2011; Janusz et al., 2017). The metabolic assays (strain by strain) revealed
551 significant differences between wood and soil bacteria. The decaying wood-inhabiting
552 bacteria presented weaker metabolic potentials than the BS isolates, but a conserved
553 ability to consume cellulose and hemicellulose derived monomers such as alpha D-
554 glucose, D-mannitol, D-mannose, L-arabinose, N-acetyl-D-glucosamine, L-fucose, D-
555 sorbitol, D-fructose, L-rhamnose and beta-methyl-D-glucoside. The preferential use
556 of these wood monomers by bacterial strains isolated from a wood environment was
557 also demonstrated by Hervé et al. (2016). Another hypothesis explaining the
558 enrichment of particular bacteria in decaying wood is their indirect role in the wood
559 decay process (Clausen 1996; Johnston et al., 2016). Indeed, it was proposed that
560 wood-colonizing fungi enrich their habitat with bacteria capable of fixing nitrogen or
561 providing them other services, promoting fungal development and wood
562 decomposition. Although this hypothesis was not tested in our study, the chitin
563 degradation assay performed evidenced that some HW bacteria may have a
564 negative impact on fungi. Altogether, the results obtained through the functional
565 bioassays suggest that the decaying wood-inhabiting bacteria are adapted to their

566 environment where substrates are less diverse than in the soil and potentially more
567 recalcitrant to decomposition.

568 **4.3 Wood properties determine the structure of the bacterial communities.**

569 Intrinsic (chemistry) and extrinsic (environment) factors may impact the
570 structuration of the wood-inhabiting bacterial communities. Indeed, wood is not a
571 homogenous material and it is established that HW and SW present different
572 properties. In our study, the chemical analyses done on the SW and HW
573 compartments clearly evidenced that SW presented significantly higher content of N,
574 P and Mg, and lower C/N ratio than HW. In addition, HW presented a significantly
575 more acidic pH than SW. Interestingly, C/N ratio and pH have been proposed as
576 driving factors governing the composition of the wood-inhabiting bacterial
577 communities (Folman et al., 2008; Hoppe et al., 2015). In this sense, the correlation
578 analyses done between the Illumina 16S rRNA sequences and the wood properties
579 revealed that the C/N ratio was negatively correlated with the abundance of
580 *Bacteroidetes*, while it was positively correlated with *Alphaproteobacteria*. In addition,
581 pH appeared positively correlated with the abundance of *Bacteroidetes*. These two
582 wood chemical properties were also reported as main drivers of bacterial community
583 structures in decaying wood (Tláškal et al., 2017; Moll et al., 2018). The HW
584 compartment is also known to present a higher phenolic extractives content (up to
585 10%), which may have an antimicrobial activity (Kebbi-Benkeder et al., 2015;
586 Miranda et al., 2017; Moll et al., 2018). The colonization of the wood compartments
587 by other types of decaying wood organisms (i.e., fungi and insects; Kubartova' et al.,
588 2012; Ulyshen, 2016; Leonhardt et al., 2019) may also be important extrinsic factors
589 driving the structure of the bacterial communities. Indeed, fungi and insects are
590 known to be important wood decomposers (Baldrian et al., 2016; Ulyshen, 2016) and

591 their interactions with bacteria are known to have an important ecosystemic role
592 (Folman et al., 2008; Frey-Klett et al., 2011; Hervé et al., 2016; Johnston et al.,
593 2016). Notably, bacterial–fungal consortia were shown to degrade wood more
594 effectively than fungi alone (Murray and Woodward, 2003; Hervé et al., 2016). In our
595 study, a visual inspection of the oak discs at the end of the incubation revealed an
596 important fungal development (i.e., hyphae) on the surface of SW, which was not
597 visible on HW. Consequently, it is strongly plausible that differences between HW
598 and SW in term of bacterial community structure are also due to the presence of
599 fungal communities. Whether the density of culturable bacteria was similar between
600 the HW and BS, the taxonomic structuration differed. Although it was only significant
601 for the richness index (i.e., Shannon), all the richness and diversity indices tended to
602 have lower values for the HW compartment compared to SW. A similar trend was
603 obtained by Moll et al. (2018) considering 30 tree species, including oak tree. In our
604 study, the decrease of diversity and richness corresponded to higher abundances of
605 *Alpha-* and *Betaproteobacteria* and lower abundance of *Bacteroidetes* in HW than in
606 SW. These results differ from the study of Moll et al. (2018) where the major
607 difference between HW and SW of decaying oak wood relied on an increase of
608 *Acidobacteria* in SW. The culturable approach also permitted to differentiate the SW
609 and HW compartments. Notably, several genera were only isolated in HW (i.e.,
610 *Acidocella*) or only in SW (i.e., *Mucilaginibacter*, *Luteibacter*, *Yersinia*), evidencing a
611 relatively good overlap between the conclusions obtained by the culture-dependent
612 and -independent approaches. Together, these results highlight that specific bacterial
613 communities are enriched in each wood compartment, potentially according to the
614 wood properties as well as due to the interactions with the fungal communities
615 occurring in decaying wood.

616

617 **5. Conclusion**

618 Our study highlighted that at middle stage of decay, a significant part of the
619 microbiota enriched in wood is shared with the surrounding bulk soil (BS), evidencing
620 that decaying wood is not disconnected from its environment. At the same time, ca
621 30% of the OTUs identified in our study were only detected in the wood
622 compartments, highlighting the enrichment of particular taxa and/or functional group
623 in decaying wood. Interestingly, the bacterial communities occurring in the SW and
624 HW compartments significantly differed in their taxonomic and functional
625 compositions. The distribution of some taxa appeared significantly correlated to the
626 wood chemical properties. This differentiation clearly evidences that the colonization
627 of decaying wood by bacteria is based on a deterministic process linking extrinsic
628 and intrinsic factors. Although the bacteria isolated from the wood compartments
629 presented lower metabolic potentials compared to those of the surrounding BS, they
630 were characterized by a conserved ability to decompose cellulose/hemicellulose and
631 derivatives. In contrast, no bacteria capable of decomposing lignin was identified.
632 Those results suggest that wood-inhabiting bacteria are capable of degrading the
633 easily degradable part of wood, but that the decomposition of the recalcitrant part
634 (i.e., lignin) is probably done by fungi (van der Wal et al., 2015; Hoppe et al., 2015;
635 Leonhardt et al., 2019; Tláskal et al., 2017). Another hypothesis may be that these
636 bacterial communities indirectly contribute to wood decay through the promotion of
637 fungal growth, highlighting a possible functional complementation between bacteria
638 and fungi (Frey-Klett et al., 2011). Future work will need to focus i) on the functional
639 structuration of the bacterial communities at different stage of decay, ii) on bacterial-
640 fungal interactions and/or co-occurrence studies during wood decay and iii) on the

641 indirect role of bacteria such as nitrogen fixation during wood decay. Combination of
642 field experiments and controlled microcosms will be necessary to better understand
643 the role of microbial interactions and community successions in wood decay.

644

645 **6. Acknowledgments**

646 This work was supported by the Laboratory of Excellence Arbre (ANR-11- LABX-
647 0002-01; DivBacto project) and by the EFABA project (SF 4242; AI-16-3). The
648 UMR1136 is supported by the French Agency through the Laboratory of Excellence
649 Arbre (ANR-11-LABX-0002-01). We would like also to thank A. Deveau for her help
650 concerning the analysis of the Illumina Miseq 16S rRNA sequences.

651

652 **7. Author contributions**

653 S.M., S.U. and M.B., conceived and designed the study; S.M., C.B., P.R. and
654 C.L., performed field and laboratory work; S.M. and S.U., analyzed the data and
655 wrote the paper. All authors reviewed and edited the manuscript.

656

657 **Figure Legends:**

658

659 **Figure 1. Evidence of the oak decay stage.** **A.** Picture of a freshly cut oak disc at
660 the beginning of the experiment (T0 month). **B.** Picture of an oak disc after 9 months
661 of *in situ* decay (T9 months) showing the side in direct contact with the soil. **C.** Mean
662 percentage of the penetration depth for the sapwood (SW) and heartwood (HW)
663 compartments assessed using a Pilodyn penetrometer at T0 ($n=3$) and T3, T6 and
664 T9 months ($n=20$ per time point). Significant differences between the two wood
665 compartments for each time point are presented using a star ($P<0.05$ according to a
666 t-test).

667

668 **Figure 2. A. Relative abundance of the Illumina Miseq 16S rRNA sequences**
669 **from bacterial taxonomic phyla for each compartment (oak decaying sapwood**
670 **(SW) and heartwood (HW) and bulk soil (BS)).** The ten most abundant phyla were
671 considered. **B. Multidimensional scaling (MDS) ordination using Bray Curtis**
672 **matrix distance representing the distribution of the bacterial communities**
673 **according to each compartment (oak decaying sapwood (SW) and heartwood**
674 **(HW) and bulk soil (BS)).**

675

676 **Figure 3. Heatmap showing the relative abundance of the 10 most important**
677 **OTUs per compartment (oak decaying sapwood (SW) or heartwood (HW) or**
678 **bulk soil (BS)) expressed in percentage.**

679

680 **Figure 4. Neighbor-joining tree showing the phylogenetic relationships of**
681 **bacterial isolates depending on their compartment (oak decaying sapwood**
682 **(SW) and heartwood (HW), Bulk soil (BS)), based on PCR sequencing of a**
683 **portion of the 16S rRNA gene.** A bootstrap analysis was performed with 1000
684 repetitions. Due to the relatively large number of sequences analysed, we have
685 chosen one representative sequence in each genus for the tree representation. Pie
686 chart on the right side of each genus represents the distribution of bacterial isolates
687 assigned to this genus according to their origin (blue: BS, red: SW, green: HW).
688 Number accessions for reference strains from NCBI are indicated in brackets.

689

690 **Figure 5. Heatmap analysis of the metabolic potentials obtained using Biolog**
691 **GN2 microplates.** A total of 304 bacterial isolates was tested. Detailed analysis of
692 the substrates for which metabolic potentials varied significantly among the bacterial
693 isolates depending on the compartment (oak decaying sapwood (SW) and heartwood
694 (HW) and bulk soil (BS)) was done. Statistics on the right of the heatmap were
695 obtained according to a one-way ANOVA followed by a Tukey test (NS: not
696 significant, $*P<0.05$, $**P<0.01$, $***P<0.001$).

697

698 **Figure 6. Relationship between functional efficacy and taxonomic affiliation.**
699 For each bacterial genus represented by a minimum of three effective isolates, their

700 relative efficacy to hydrolyze cellulose, xylan and chitin and to mobilize iron is
701 presented, whatever their compartment (oak decaying sapwood (SW) and heartwood
702 (HW) and bulk soil (BS)) of the bacterial isolates. Lowercase letters (a, b, c, d or e)
703 indicate significant differences between genus efficacies. Statistics were obtained
704 according to a one-way ANOVA followed by a Tukey test ($P < 0.05$). The number of
705 isolates tested per genus as well as their repartition according to their compartment
706 (BS, SW or HW) is presented under the bars. The error bars indicate the standard
707 error.

708 **References**

709

710 Altschul, S.F., 1997. Evaluating the statistical significance of multiple distinct local
711 alignments. In *Theoretical and Computational Methods in Genome Research*,
712 Springer, Boston, MA. 1-14.

713

714 Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman,
715 D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database
716 search programs. *Nucleic Acids Research* 25, 3389-3402.

717

718 Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local
719 alignment search tool. *Journal of Molecular Biology* 215, 403-410.

720

721 Augusto, L., De Schrijver, A., Vesterdal, L., Smolander, A., Prescott, C., Ranger, J.,
722 2015. Influences of evergreen gymnosperm and deciduous angiosperm tree species
723 on the functioning of temperate and boreal forests. *Biological Reviews* 90, 444-466.

724

725 Baldrian, P., 2017. Forest microbiome: diversity, complexity and dynamics. *FEMS*
726 *Microbiology Reviews* 41, 109-130.

727

728 Baldrian P., Zrůstová, P., Tláškal, V., Davidová, A., Merhautová, V., Vrška. T., 2016.
729 Fungi associated with decomposing deadwood in a natural beech-dominated forest.
730 *Fungal Ecology* 23, 109-122.

731

732 Benjamini, Y., Hochberg, Y. 1995. Controlling the false discovery rate: a practical and
733 powerful approach to multiple testing. *Journal of the Royal statistical society: series B*
734 *(Methodological)*, 57, 289-300.

735

736 Blanchette R.A., 2000. A review of microbial deterioration found in archaeological
737 wood from different environments. *International Biodeterioration and Biodegradation*
738 46, 189-204.

739

740 Bugg, T.D., Ahmad, M., Hardiman, E.M., Singh, R., 2011. The emerging role for
741 bacteria in lignin degradation and bio-product formation. *Current Opinion in Chemical*
742 *Biology* 22, 394-400

743

744 Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K.,
745 Madden, T.L., 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10,
746 421.

747

748 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A.,
749 Turnbaugh, P.J., *et al.*, 2011. Global patterns of 16S rRNA diversity at a depth of
750 millions of sequences per sample. *Proceedings of the National Academy of Sciences*
751 108, 4516-4522.

752

753 Clausen, C.A., 1996. Bacterial associations with decaying wood: a review.
754 *International Biodeterioration and Biodegradation* 37, 101-107.

755

756 Cox, R.A., 1968. The use of guanidinium chloride in the isolation of nucleic acids. In

757 Methods in enzymology, Academic Press 12, 120-129.
758
759 Duchaufour, P., Bonneau, M., 1959. Une nouvelle méthode de dosage du phosphore
760 assimilable dans les sols forestiers. Bulletin AFES 4, 193-198.
761
762 Duval, L., 1963. Etude des conditions de validité du dosage céruléomolybdique de
763 l'acide phosphorique. Conséquences Pratiques. Chimie Analytique 45, 237-240.
764
765 Edwards, U., Rogall, T., Blöcker, H., Emde, M., Böttger, E.C., 1989. Isolation and
766 direct complete nucleotide determination of entire genes. Characterization of a gene
767 coding for 16S ribosomal RNA. Nucleic Acids Research 17, 7843-7853.
768
769 Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman,
770 S., Hernandez-Raquet, Guillerma, Combes, Sylvie., Pascal, G. 2018. FROGS: find,
771 rapidly, OTUs with galaxy solution. Bioinformatics 34, 1287-1294.
772
773 Folman, L.B., Klein Gunnewiek, P.J., Boddy, L., De Boer, W., 2008. Impact of white-
774 rot fungi on numbers and community composition of bacteria colonizing beech wood
775 from forest soil. FEMS Microbiology Ecology 63, 181-191.
776
777 Frey-Klett, P., Burlinson, P., Deveau, A., Barret, M., Tarkka, M., Sarniguet, A., 2011.
778 Bacterial-fungal interactions: hyphens between agricultural, clinical, environmental,
779 and food microbiologists. Microbiology and Molecular Biology Reviews 75, 583-609.
780
781 Fundova, I., Funda, T. Wu, H.X., 2018. Non-destructive wood density assessment of
782 Scots pine (*Pinus sylvestris* L.) using Resistograph and Pilodyn. PLoS one 13,
783 e0204518.
784
785 Goodell, B., Jellison, J., Liu, J., Daniel, G., Paszczynski, A., Fekete, F.,
786 Krishnamurthy, S., Jun, L., Xu, G., 1997. Low molecular weight chelators and
787 phenolic compounds isolated from wood decay fungi and their role in the fungal
788 biodegradation of wood. Journal of Biotechnology, 53, 133-162.
789
790 Gouy, M., Guindon, S., Gascuel, O., 2010. SeaView version 4: a multiplatform
791 graphical user interface for sequence alignment and phylogenetic tree building.
792 Molecular Biology and Evolution 27, 221-224.
793
794 Hahn, K., Christensen, M., 2005. Dead wood in European forest reserves - a
795 reference for forest management. In: Marchetti M (ed.). Monitoring and Indicators of
796 Forest Biodiversity in Europe - From Ideas to Operationality. European Forest
797 Institute 181-91.
798
799 Hanewinkel, M., Cullmann, D.A., Schelhaas, M.J., Nabuurs, G.J., Zimmermann, N.E.,
800 2013. Climate change may cause severe loss in the economic value of European
801 forest land. Nature Climate Change 3, 203-207.
802
803 Hervé, V., Le Roux, X., Uroz, S., Gelhaye, E., Frey-Klett, P., 2014. Diversity and
804 structure of bacterial communities associated with *Phanerochaete chrysosporium*
805 during wood decay. Environmental Microbiology 16, 2238-2252.
806

807 Hervé, V., Ketter, E., Pierrat, J.C., Gelhaye, E., Frey-Klett, P., 2016. Impact of
808 *Phanerochaete chrysosporium* on the functional diversity of bacterial communities
809 associated with decaying wood. PLoS one 11, e0147100.
810
811 Hobbie, E.A., 2006. Carbon allocation to ectomycorrhizal fungi correlates with
812 belowground allocation in culture studies. Ecology 87, 563-569.
813
814 Hoppe, B., Krüger, D., Kahl, T., Arnstadt, T., Buscot, F., Bauhus, J., Wubet, T., 2015.
815 A pyrosequencing insight into sprawling bacterial diversity and community dynamics
816 in decaying deadwood logs of *Fagus sylvatica* and *Picea abies*. Scientific Reports 5,
817 9456.
818
819 Huson, D.H., Scornavacca, C., 2012. Dendroscope 3: an interactive tool for rooted
820 phylogenetic trees and networks. Systematic Biology 61, 1061-1067.
821
822 Janusz, G., Pawlik, A., Sulej, J., Świdorska-Burek, U., Jarosz-Wilkolazka, A.,
823 Paszczyński, A., 2017. Lignin degradation: microorganisms, enzymes involved,
824 genomes analysis and evolution. FEMS microbiology reviews 41, 941-962.
825
826 Johnston, S.R., Boddy, L., Weightman, A.J., 2016. Bacteria in decomposing wood
827 and their interactions with wood-decay fungi. FEMS Microbiology Ecology 92, fiw179.
828
829 Kebbi-Benkeder, Z., Colin, F., Dumarçay, S., Gérardin, P., 2015. Quantification and
830 characterization of knotwood extractives of 12 European softwood and hardwood
831 species. Annals of Forest Sciences 72, 277-284.
832
833 Kielak, A.M., Scheublin, T.R., Mendes, L.W., Van Veen, J.A., Kuramae, E.E., 2016.
834 Bacterial community succession in pine-wood decomposition. Frontiers in
835 Microbiology 7, 231.
836
837 Kubartová, A., Ottosson, E., Dahlberg, A., Stenlid, J., 2012. Patterns of fungal
838 communities among and within decaying logs revealed by 454 sequencing.
839 Molecular Ecology 21, 4514-4532.
840
841 Lane, D.J., 1991. 16S/23S rRNA sequencing. Nucleic acid techniques in bacterial
842 systematics 115-175.
843
844 Lee, C.S., Jung, Y.T., Park, S., Oh, T.K., Yoon, J.H., 2010. *Lysinibacillus xylanilyticus*
845 sp. nov., a xylan-degrading bacterium isolated from forest humus. International
846 Journal of Systematic and Evolutionary Microbiology 60, 281-286.
847
848 Leonhardt, S., Hoppe, B., Stengel, E., Noll, L., Moll, J., Bässler, C., Dahl, A., Buscot,
849 F., Hofrichter, M., Kellner H., 2019. Molecular fungal community and its
850 decomposition activity in sapwood and heartwood of 13 temperate European tree
851 species. PLoS one 14, e0212120.
852
853 Litton, C.M., Ryan, M.G., Knight, D.H., 2004. Effects of tree density and stand age on
854 carbon allocation patterns in postfire lodgepole pine. Ecological Applications 14, 460-
855 475.
856

857 Lladó, S., Žifčáková, L., Větrovský, T., Eichlerová, I., Baldrian, P., 2016. Functional
858 screening of abundant bacteria from acidic forest soil indicates the metabolic
859 potential of Acidobacteria subdivision 1 for polysaccharide decomposition. *Biology
860 and Fertility of Soils* 52, 251-260.
861

862 Lopez-Mondejar, R., Zuhlke, D., Becher, D., Riedel, K., Baldrian, P., 2016. Cellulose
863 and hemicellulose decomposition by forest soil bacteria proceeds by the action of
864 structurally variable enzymatic systems. *Scientific Reports* 6, 25279.
865

866 Mahé, F., Rognes, T., Quince, C., de Vargas, C., Dunthorn, M., 2014. Swarm: robust
867 and fast clustering method for amplicon-based studies. *PeerJ* 2, e593.
868

869 Magoc, T., Salzberg, S.L., 2011. FLASH: fast length adjustment of short reads to
870 improve genome assemblies. *Bioinformatics* 27, 2957-2963.
871

872 Mäkipää, R., Linkosalo, T., 2011. A non-destructive field method for measuring wood
873 density of decaying logs. *Silva Fennica* 45, 1135-1142.
874

875 Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput
876 sequencing reads. *European Molecular Biology network Journal* 17, 10-12.
877

878 Mason, P.E., Neilson, G.W., Dempsey, C.E., Barnes, A.C., Cruickshank, J.M., 2003.
879 The hydration structure of guanidinium and thiocyanate ions: implications for protein
880 stability in aqueous solution. *Proceedings of the National Academy of Sciences* 100,
881 4557-4561.
882

883 Mathieu, Y., Gelhaye, E., Dumarçay, S., Gérardin, P., Harvengt, L., Buée, M., 2013.
884 Selection and validation of enzymatic activities as functional markers in wood
885 biotechnology and fungal ecology. *Journal of Microbiological Methods* 92, 157-163.
886

887 McMurdie, P.J., Holmes, S., 2013. phyloseq: An R package for reproducible
888 interactive analysis and graphics of microbiome census data. *PLoS ONE* 8, e61217.
889

890 Meerts, P. 2002. Mineral nutrient concentrations in sapwood and heartwood: a
891 literature review. *Annals of Forest Science* 59, 713-722.
892

893 Miranda, I., Sousa, V., Ferreira, J., Pereira, H., 2017. Chemical characterization and
894 extractives composition of heartwood and sapwood from *Quercus faginea*. *PloS one*
895 12, e0179268.
896

897 Moll, J., Kellner, H., Leonhardt, S., Stengel, E., Dahl, A., Bässler, C., Buscot, F.,
898 Hofrichter, M., Hoppe, B., 2018. Bacteria inhabiting deadwood of 13 tree species are
899 heterogeneously distributed between sapwood and heartwood. *Environmental
900 Microbiology* 20, 3744-3756.
901

902 Murray, A., Woodward, S., 2003. In vitro interactions between bacteria isolated from
903 Sitka spruce stumps and *Heterobasidion annosum*. *Forest Pathology* 33, 53-67.
904

905 Murray, A., Woodward, S., 2007. Temporal changes in functional diversity of
906 culturable populations in Sitka spruce stumps. *Forest Pathology* 37, 217-235.

907
908 Nicolitch, O., Colin, Y., Turpault, M.P., Uroz, S., 2016. Soil type determines the
909 distribution of nutrient mobilizing bacterial communities in the rhizosphere of beech
910 trees. *Soil Biology and Biochemistry* 103, 429-445.
911
912 Nicolitch, O., Colin, Y., Turpault, M.P., Fauchery, L., Uroz, S., 2017. Tree roots select
913 specific bacterial communities in the subsurface critical zone. *Soil Biology and*
914 *Biochemistry* 115, 109-123.
915
916 Noll L., Leonhardt, S., Arnstadt, T., Hoppe, B., Poll, C., Matzner, E., Hofrichter, M.,
917 Kellner, H., 2016. Fungal biomass and extracellular enzyme activities in coarse
918 woody debris of 13 tree species in the early phase of decomposition. *Forest Ecology*
919 *Management* 378, 181-192.
920
921 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J.,
922 Suggests M.A.S.S., 2007. The vegan package. *Community ecology package* 10,
923 631-637.
924
925 Olsen, S.R., 1954. Estimation of available phosphorus. In soils by extraction with
926 sodium bicarbonate. US Department of Agriculture 939, 1-19.
927
928 Pan, Y., Birdsey, R.A., Fang, J., Houghton, R., Kauppi, P.E. Kurz, W.A. *et al.*, 2011.
929 A large and persistent carbon sink in the world's forests. *Science* 333, 988–93.
930
931 Paradis, E., Claude, J., Strimmer, K., 2004 APE: analyses of phylogenetics and
932 evolution in R language. *Bioinformatics* 20, 289-290.
933
934 Prewitt, L., Kang, Y., Kakumanu, M.L., Williams, M., 2014. Fungal and bacterial
935 community succession differs for three wood types during decay in a forest soil.
936 *Microbial Ecology* 68, 212-221.
937
938 Probst, M., Gómez-Brandón, M., Bardelli, T., Egli, M., Insam, H., Ascher-Jenull, J.,
939 2018. Bacterial communities of decaying Norway spruce follow distinct slope
940 exposure and time-dependent trajectories. *Environmental Microbiology* 20, 3657-
941 3670.
942
943 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J.,
944 Glöckner, FO., 2012. The SILVA ribosomal RNA gene database project: improved
945 data processing and web-based tools. *Nucleic Acids Research* 41, D590-D596.
946
947 R Core Team., 2015. R: a Language and Environment for Statistical Computing.
948
949 Rajala, T., Peltoniemi, M., Pennanen, T., Mäkipää, R., 2012. Fungal community
950 dynamics in relation to substrate quality of decaying Norway spruce (*Picea abies* L.
951 Karst.) logs in boreal forests. *FEMS Microbiology Ecology* 81, 494-505.
952
953 Rajala, T., Tuomivirta, T., Pennanen, T., Mäkipää, R., 2015. Habitat models of wood-
954 inhabiting fungi along a decay gradient of Norway spruce logs. *Fungal Ecology* 18,
955 48-55.
956

957 Ricker, M.C., Lockaby, B.G., Blosser, G.D., Conner, W.H., 2016. Rapid wood decay
958 and nutrient mineralization in an old-growth bottomland hardwood forest.
959 Biogeochemistry 127, 323-338.
960

961 Rinta-Kanto, J.M., Sinkko, H., Rajala, T., Al-Soud, W.A., Sørensen, S.J., Tamminen,
962 M.V., Timonen, S., 2016. Natural decay process affects the abundance and
963 community structure of Bacteria and Archaea in *Picea abies* logs. FEMS
964 Microbiology Ecology 92, fiw087.
965

966 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a
967 versatile open source tool for metagenomics. PeerJ. 4, e2584.
968

969 Schneider, T., Keiblinger, K. M., Schmid, E., Sterflinger-Gleixner, K., Ellersdorfer, G.,
970 Roschitzki, B., Richter, A., Eberl, L., Zechmeister-Boltenstern, S., Riedel, K. (2012).
971 Who is who in litter decomposition? Metaproteomics reveals major microbial players
972 and their biogeochemical functions. The ISME journal 6, 1749-1762.
973

974 Seibold, S., Bäessler, C., Brandl, R., Gossner, M.M., Thorn, S., Ulyshen, M.D., Müller,
975 J., 2015. Experimental studies of dead-wood biodiversity—a review identifying global
976 gaps in knowledge. Biological Conservation 191, 139-149.
977

978 Seidl, R, Schelhaas, M-J, Rammer, W, Verkerk, P.J., 2014. Increasing forest
979 disturbances in Europe and their impact on carbon storage. Nature Climate Change
980 4, 806-810.
981

982 Stokland J.N., Siitonen, J., Jonsson, B.G., 2012. Biodiversity in Dead Wood.
983 Cambridge: Cambridge University Press.
984

985 Štursová, M., Žifčáková, L., Leigh, M.B., Burgess, R. Baldrian, P., 2012. Cellulose
986 utilization in forest litter and soil: identification of bacterial and fungal decomposers.
987 FEMS Microbiology Ecology 80, 735-746.
988

989 Sun, H., Terhonen, E., Kasanen, R., Asiegbu, F.O., 2014. Diversity and community
990 structure of primary wood-inhabiting bacteria in boreal forest. Geomicrobiology
991 Journal 31, 315-324.
992

993 Tláškal, V., Zrůstová, P., Vrška, T., Baldrian, P., 2017. Bacteria associated with
994 decomposing dead wood in a natural temperate forest. FEMS Microbiology Ecology
995 93, fix157.
996

997 Ulyshen, M.D., 2016. Wood decomposition as influenced by invertebrates. Biological
998 Reviews 91, 70-85.
999

1000 Valášková, V., De Boer, W., Gunnewiek, P.J.K., Pospisek, M., Baldrian, P., 2009.
1001 Phylogenetic composition and properties of bacteria coexisting with the fungus
1002 *Hypholoma fasciculare* in decaying wood. The ISME Journal 3, 1218.
1003


1004 Větrovský, T., Steffen, K.T., Baldrian, P., 2014. Potential of cometabolic
1005 transformation of polysaccharides and lignin in lignocellulose by soil Actinobacteria.
1006 PLoS one 9, e89108.

1007
1008 van der Wal, A., Ottosson, E., De Boer, W., 2015. Neglected role of fungal
1009 community composition in explaining variation in wood decay rates. *Ecology* 96, 124-
1010 133.
1011
1012 Vorob'ev, A.V., de Boer, W., Folman, L.B., Bodelier, P.L., Doronina, N.V., Suzina,
1013 N.E., Trotsenko Y.A., Dedysh, S. N., 2009. *Methylovirgula ligni* gen. nov., sp. nov.,
1014 an obligately acidophilic, facultatively methylotrophic bacterium with a highly
1015 divergent mxaF gene. *International Journal of Systematic and Evolutionary*
1016 *Microbiology* 59, 2538-2545.
1017
1018 Weedon, J.T., Cornwell, W.K., Cornelissen, J.H.C., Zanne, A.E., Wirth, C., Coomes.
1019 D.A., 2009. Global meta-analysis of wood decomposition rates: a role for trait
1020 variation among tree species? *Ecological Letters* 12, 45-56.
1021
1022 Wickham, H., 2009. *ggplot2: elegant graphics for data analysis*. New York: Springer.
1023
1024 Zhang, H.B., Yang, M.X., Tu, R., 2008. Unexpectedly high bacterial diversity in
1025 decaying wood of a conifer as revealed by a molecular method. *International*
1026 *Biodeterioration and Biodegradation* 62, 471-474.


A.




B.




C.




A. Composition within bacteria (top 10 phyla)



B. MDS+BC





Bacteroidetes
Actinobacteria
Acidobacteria
Alphaproteobacteria
Gammaproteobacteria
Betaproteobacteria

