



HAL
open science

Bacterial Conjugation Protocol for Ruminant Mycoplasmas

Eveline Sagné, Christine Citti, Emilie Dordet-Frisoni

► **To cite this version:**

Eveline Sagné, Christine Citti, Emilie Dordet-Frisoni. Bacterial Conjugation Protocol for Ruminant Mycoplasmas. Bio-protocol , 2021, 11 (2), pp.e3893. 10.21769/BioProtoc.3893 . hal-03196778

HAL Id: hal-03196778

<https://hal.inrae.fr/hal-03196778>

Submitted on 13 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Bacterial Conjugation Protocol for Ruminant Mycoplasmas

Eveline Sagné[#], Christine Citti and Emilie Dordet-Frisoni^{#, *}

IHAP, Université de Toulouse, INRAE, ENVT, Toulouse, France

*For correspondence: emilie.dordet-frisoni@envt.fr

[#]Contributed equally to this work

[Abstract] In *Mycoplasma agalactiae*, two simultaneous processes of DNA transfer have been described that require direct cell-to-cell contact and are similar to conjugation. One involves the self-transmission of an integrative conjugative element (ICE) while the second concerns the horizontal transfer of large and small fragments of chromosomal DNA. Here, we describe an optimized conjugation protocol for the horizontal transfer of ICE or chromosomal DNA carrying antibiotic resistance markers (*i.e.*, tetracycline, gentamicin, puromycin) from donor to recipient mycoplasma cells. Calculation of the conjugation frequencies, selection and characterization of transconjugants are detailed. This protocol has been developed with *M. agalactiae* but has been successfully used for *M. bovis* and can be adapted to other related mycoplasma species.

Keywords: Mycoplasma, Horizontal gene transfer, Conjugation, Mating, Integrative and conjugative element

[Background] Conjugative, horizontal DNA transfer is a key player of microbial diversification. By promoting an active transfer of DNA through close cell-to-cell contact (Lederberg and Tatum, 1946), this phenomenon facilitates the rapid acquisition of new traits from external resources. Mycoplasmas (class Mollicutes) are wall-less bacteria with reduced genomes whose evolution has been considered to be only driven by gene loss and in which horizontal gene transfer (HGT) was long thought to be marginal. During this past decade, comparative genomic analyses have revisited this paradigm and have shown (i) the occurrence of HGTs events in between mycoplasma species sharing same hosts (Sirand-Pugnet *et al.*, 2007) and (ii) the presence of integrative and conjugative elements (ICEs) in a large number of sequenced mycoplasma genomes (Calcutt *et al.*, 2002; Marenda *et al.*, 2006; Dordet-Frisoni *et al.*, 2013; Tardy *et al.*, 2015; Meygret *et al.*, 2019), raising the prospect that these minimal bacteria might be able to conjugate.

In vitro conjugation-like process has been demonstrated in *Mycoplasma agalactiae* (Dordet-Frisoni *et al.*, 2013, 2014 and 2019), a mycoplasma species pathogenic for ruminants. Mating assays were based on mixing in liquid culture two strains tagged with antibiotic resistant genes and resulted in a bidirectional DNA transfer process (Dordet-Frisoni *et al.*, 2013 and 2014). The first is the conventional horizontal dissemination of a mycoplasma ICE (MICE), from ICE-positive to ICE-negative cells (Dordet-Frisoni *et al.*, 2013). The second concerns the transfer of large and small blocks of chromosomal DNA from ICE-negative to ICE-positive cells and was further referred to as MCT for mycoplasma chromosomal transfer (Dordet-Frisoni *et al.*, 2014 and 2019; Faucher *et al.*, 2019). *M. agalactiae*

conjugation has been shown to rely on the presence of an ICE in at least one parent cell, but both DNA transfers are physically independent (taking place in opposite direction) and occur at the same frequency (around 10^{-8} transconjugants/total CFU) (Dordet-Frisoni *et al.*, 2014).

The detailed protocol presented here describes an optimized mating procedure developed for the conjugation of *M. agalactiae* and related species. For instance, as described in Dordet-Frisoni *et al.* (2014), this protocol has also been successfully applied to *M. bovis* and can most likely be adapted to other mycoplasma species with some adjustments.

Materials and Reagents

1. Sterile tips for micropipettes (STARLAB, catalog numbers: S1122-1830 [1,000 µl], S1120-3810 [10-20 µl])
2. Sterile 1.5 ml and 2 ml microcentrifuge tubes (SORENSEN, catalog number: 017040A; TREFF, catalog number: 96.09329.9.01)
3. Sterile plastic Petri dish plates (90 mm diameter, Clearline, catalog number: 076084BS)
4. 0.22 µm pore filter (PVDF 33 mm, ClearLine, Dutscher, catalog number: 051733)
5. Donor and recipient cells, harbouring different selective markers for antibiotic selection #

#Note: M. agalactiae strains were tagged with selectable markers previously introduced by transposition using a pMT85-derived plasmid (Figure 1) (Zimmerman and Herrmann, 2005). pMT85 and derivatives bear a mini-Tn with either the aacA-aphD gentamicin resistance cassette, the tetM determinant for resistance to tetracycline or the pac gene conferring puromycin resistance. Marker genes are flanked by two Tn4001 inverted repeats (IRs). Since the mini-Tn backbone contains no transposase gene (it is located elsewhere in the pMT85), its insertion into the cell chromosome is stable (Baranowski *et al.*, 2010).*

*Several Mycoplasma agalactiae strains such as 5632, PG2, 4867, 14628 and 4055, have been successfully employed in mating assays (Dordet-Frisoni *et al.*, 2019), but conjugation experiment has been also successful with other ruminant mycoplasmas species such as M. bovis PG45 strain (Dordet-Frisoni *et al.*, 2014) and M. mycoides subsp. capri GM12 strain (personal data). For conjugation to occur, at least one of the two partners should carry an ICE. "Concerning chromosomal transfers (MCT), only two M. agalactiae strains have been documented as recipient cells, namely 5632 and 4867, while all strains tested so far can act as donor. As for ICEA, all strains carrying an active ICE are potentially donor cell. All parental M. agalactiae strains cited above can be acquired at the Vigimyc collection (Chazel *et al.*, 2010).*




Figure 1. Schematic representing *M. agalactiae* antibiotic tagging procedure. Selectable markers are introduced by transposition using the mini transposon pMT85, which bear antibiotic resistance genes as selection markers (Antibiotic^R). The Antibiotic^R marker is flanked by two inverted (IR) Tn4001 repeats. During insertion, the transposase (Tnp) which is located outside the IRs is eliminated and the selection marker is integrated into the chromosome in a random (in coding (CDS) or non-coding sequences) and a stable manner. This experiment results in mycoplasmas transformants tagged with resistance markers (ATB^R) at different locations on the chromosome.

6. Appropriate selectable antibiotics:
 - Gentamicin (Gm^R) (Gibco, catalog number: 15750037)
 - Tetracycline (Tet^R) (Sigma-Aldrich, catalog number: T3383)
 - Puromycin (Pur^R) (Fisher, catalog number: 10054207)
 - Ampicillin (Sigma-Aldrich, catalog number: A9518)
7. PPLO broth (Difco, catalog number: 255420)
8. Tryptone peptone (Difco, catalog number: 211705)
9. Bacto peptone (Difco, catalog number: 211677)
10. FBS (Fetal Bovine Serum) (Gibco, catalog number: 10270-106)
11. CMRL 1066 (10x) (Gibco, catalog number: 21540-026)
12. TC yeastolate (Difco, catalog number: 255772)
13. Yeast extract 15% (Gibco, catalog number: 18180-059)
14. D-(+)-Glucose (Sigma-Aldrich, catalog number: G6152)
15. Phenol red 0.5% (Sigma-Aldrich, catalog number: P0290)
16. Na Pyruvate (Sigma-Aldrich, catalog number: P2256)
17. Select agar (Invitrogen, catalog number: 30391-023)
18. NaOH (Sigma-Aldrich, catalog number: S8045)
19. D-PBS (Gibco, catalog number: 14190169)
20. Tween-20 (Sigma-Aldrich, catalog number: P9416)
21. Ultra pure Tris (Invitrogen, catalog number: 15504020)
22. HCl (Sigma-Aldrich, catalog number: 320331)
23. Proteinase K 20 mg/ml (Qiagen catalog number: 19131)
24. Taq DNA Polymerase (New England Biolabs catalog number: M0267L)
25. dNTP mix 10 mM (New England Biolabs catalog number: N0447S)
26. Taq Buffer 10x (New England Biolabs catalog number: M0267L)

27. Sterile SP4 liquid medium (see Recipes)
28. Sterile SP4 agar plates (see Recipes)
29. *M. agalactiae* Lysis buffer (see Recipes)

Equipment

1. Air-flow cabinet (Faster, model: SF Classic 212)
2. 37 °C incubator (Firlabo, model: FA Type B 407L-FA407B)
3. Microcentrifuge with rotor for 1.5-2 ml tubes (Eppendorf, model: 5424)
4. Vortex (Scientific Industries, model: GENIE 2)
5. Binocular magnifier (Olympus, model: SZ 60)
6. Multiblock heater (Lab Line Instruments, model: 2052-1CCE)
7. Thermocycler (Mastercycler EP Eppendorf Gradient, model: 5341)
8. Micropipettes (Gilson Pipetman Classic 10 µl to 1 ml range, Dutscher, catalog numbers: 066002, 066003, 066004, 066005 and 066006)
9. Graduated pipettes (Falcon 5 to 25 ml range pipettes, Dutscher, catalog numbers: 357543, 357551 and 357535)
10. Pipettor (Pipetboy Acu 2, INTEGRA, DUTSCHER, catalog number: 060201)

Procedure

A schematic of the procedure is also provided bellow (Figure 2).

A. Culture preparation

1. Inoculate 1 ml of SP4 liquid medium with 10 µl of each saturated liquid growth culture of *M. agalactiae* (standard bacterial stock at $\sim 10^9$ CFU/ml) previously stored at -80 °C in the absence of glycerol or particular storage conditions and defrosted at room temperature.
 2. Incubate the culture at 37 °C for 24 h without shaking in an incubator #.
- #Note: Growth rate depends on the M. agalactiae strain and can vary from 16 h to 48 h of incubation in SP4 medium. Change from red to orange of the phenol-red present in the SP4 medium indicates optimal growth.*

B. Mating procedure

1. Mix 1 ml# of each saturated mycoplasma culture (between 10^8 and 10^9 CFU/ml) in a 2 ml sterile microcentrifuge tubes.
- # Note: The ratio of donor:recipient mycoplasma cells should be close to 1:1. Depending on the assays or on the concentration of the two mycoplasma cultures (e.g., due to differences in growth rates), the ratio can be changed to 1:2 (vol:vol) or 1:10 (vol:vol) in the same final volume of 2 ml.*
2. Centrifuge the 2 ml mixed cultures for 5 min at 8,000 x g at room temperature (20-25 °C).

3. Carefully discard the supernatant using a 1 ml micropipette.
4. Resuspend the pellets with 1 ml of SP4 liquid medium at room temperature, proceed to 2 to 3 drawing/discharging cycles with a 1 ml micropipette.
5. Incubate over night# at 37 °C without shaking.

#Note: Conjugation events can be observed after 3 h of incubation, but the highest mating frequency is observed after at least 16 h.

C. Plating

1. A part of the mating mix (10 to 50 µl) is used to determine the total CFU#. For this purpose, prepare serial dilutions in DPBS 1x + 5% decomplexed FBS (from 10⁻¹ to 10⁻⁷) and spot 10 µl of each dilution onto a SP4 agar plate containing no selective pressure.

#Note: If the conjugation efficiency is expressed by the ratio of transconjugants/donor cells or transconjugant/recipient cells, spot 10 µl of each dilution onto a SP4 agar plate containing the antibiotic marker corresponding to the donor/recipient strain, at the appropriate concentration.

2. Inoculate 300 µl of the mating mix onto selective SP4 agar plates. Spread the liquid onto the surface of the Petri dish (90 mm) by gentle circular movements so that the entire dish (except the edges) is covered#.

#Note: All the mating mix can be plated in 3 Petri dishes. For highly efficient matings, dilutions should be made before plating (1/10x, 1/100x maximum). These dilutions are made with sterile DPBS 1x complemented with 5% decomplexed FBS in microcentrifuge tubes. The mating mixture can be stored as such directly at -80 °C if necessary.

3. Incubate the SP4 agar plates at 37 °C for 3 to 5 days#.

#Note: To prevent desiccation during incubation the Petri dishes should be stored inside closed plastic bags, in the 37 °C incubator.

D. Calculation of conjugation frequencies

1. Using the binocular magnifier (10x-63x), count the Colony Forming Unit (CFU) on the 10µl dilution spots or the 300 µl spreads (Figure 2) of each selective and non-selective SP4 agar plates.
2. Calculate the mating frequency# as the CFUs grown in both selective markers (*i.e.*, Tet^R and Gm^R) divided by the total CFU obtained in dilution spots spread on SP4 agar without selective pressure.

#Note: The conjugation frequency can also be expressed as the number of transconjugants per donor or recipient cell by dividing the CFU cultivated in the two selective markers, by the CFU cultivated with only one of the two selective marker (that of the donor, or that of the recipient, see note above).

3. At least three independent experiments should be performed for reliable and reproducible data.

E. Picking, cloning and storage of transconjugants

1. Pick colonies with a sterile 200 μ l micropipette sterile tip and transfer it into 1 ml of SP4 liquid medium supplemented with the 2 appropriate antibiotics.
2. Incubate at 37 $^{\circ}$ C for 24 to 96 h[#].
#Note: Transconjugants can be directly stored at -80 $^{\circ}$ C at this stage .
3. To ensure the clonality of transconjugants, three successive subcloning steps can be performed as previously described (Rosengarten *et al.*, 1994), with the last broth culture being filtered through a 0.22 μ m pore filter before storage at -80 $^{\circ}$ C.




Figure 2. Schematic representing *M. agalactiae* mating procedure. Here, as an example, the *M. agalactiae* strain 5632 (ICE containing strain, ICE+) was mated with strain PG2 (ICE-). Prior to mating, strains were independently tagged by stably inserting the gentamicin resistance gene (Gm^R, in purple) or the tetracycline resistance gene (Tet^R, in green) in their genome, using the pMT85 plamid (Baranowski *et al.*, 2010). Saturated cultures of each of these strains ($\approx 10^9$ CFU/ml) were mixed in equal proportion (1 ml each), gently centrifuged, resuspended in fresh SP4 medium and incubated together. The mix culture was then directly plated on appropriate selective medium and dilutions were inoculated onto selective and non-selective plates to determine the conjugation frequency.

F. Verification of putative transconjugants

1. Centrifuge a 1 ml culture of cloned transconjugants at 12,000 \times g for 25 min at 4 $^{\circ}$ C.
2. Remove the supernatant and add 50 μ l of lysis buffer (see Recipes section) to the cell pellet
3. Incubate at 37 $^{\circ}$ C for 1 h.

4. Inactivate the proteinase K contained in the lysis buffer by heating at 95 °C for 10 min.
5. Perform PCR amplifications using specific primers targeting the resistance genes present in the parental strains used for mating (Table 1) (Dordet-Frisoni *et al.*, 2014). To do this, take 2 µl of the lysed transconjugants culture and perform the PCR using the standard Taq DNA polymerase procedure (here, Taq DNA polymerase from NEB).
6. Identify transconjugant genomic backbone by PCR using a set of 7 primer pairs (Table 2) that covers clockwise the *M. agalactiae* genome and produces PCR fragments specific of the *M. agalactiae* genomes (Dordet-Frisoni *et al.*, 2014) or by whole genome sequencing (Illumina) (Dordet-Frisoni *et al.*, 2019).

Table 1. Oligonucleotide used for the detection of antibiotic markers: annealing temperature and expected PCR product size (bp)

	Sequence	Tm	Size (bp)
Gm1 (Gentamicin)	ACATGAATTACACGAGGGC	55 °C	401
Gm2 (Gentamicin)	GTTCTTCTTCTGACATAGTAG	55 °C	
IntMtet1 (Tetracycline)	TGGCGTACAAGCACAAACTC	55 °C	444
IntMtet2 (Tetracycline)	GCAAAGTTCAGACGGACCTC	55 °C	
PuroF (Puromycin)	GTTGCTGTTTGGACTACTCCTG	52 °C	359
PuroR (Puromycin)	CACCAAGTTCCTAGGACCTTCAGG	52 °C	

Table 2. Oligonucleotides used for transconjugant typing: sequences, annealing temperature and expected PCR products (bp) according to the strains

Sequence	Tm	<i>M. agalactiae</i> strains				
		5632	PG2	4055	4867	14628
MAGa0220F CGCCATTGAAGGAACTTGCTCC	52	946	0	946	946	946
MAGa0220R TATGCCTGCAGATCAGTCACCA						
MAGa0900F ATGGCAATTGATTTACTCCC	55	1500	900	1500	1500	1500
MAGa0900R ATGATGTTGGCCGTATTTGG						
MAG1560F GCCGGTGCCCGGGTTAAT	55	0	1311	0	0	0
MAG1560R TCGACACCGCATTTAGCAGCA						
MAG2450F AGGACGCTGTAGAGTTTGCAAAGAA	55	0	1500	0	0	0
MAG2450R TACCACTAGCTGCCGCACCA						
MAGa3840F GAAGCACTAAAATCAGGCAA	55	1223	1067	1223	1223	1067
MAGa3840R GCGGCAGCAGAGTTATCAAG						
MAGa4400F CTCTGGCTACAGATTCATCA	55	490	607	490	490	0
MAGa4400R GGTGAACGAGAACAATCGCC						
MAG5860F ACCATTGAACTACATTTGCA	52	494	1318	0	nd	nd
MAGa6510R AAAACCGCAGACCCAAATGGT						
MAG5890R ACATTGCGATGTAGCGGGAACAG						

Recipes

1. SP4 liquid medium (for 1 L)
 - 3.5 g PPLO broth
 - 10 g Tryptone peptone
 - 5.325 g Bacto peptone
 - 690 ml distilled water
 - pH 7.8 (pH may be adjusted by the addition of NaOH 5 N solution)
 - This mixture should be autoclaved prior to adding:*
 - 170 ml decomplexed FBS (Fetal Bovine Serum, sterile solution)
 - 50 ml CMRL 1066 (10x, sterile solution)
 - 50 ml TC yeastolate (sol 4% in H₂O, filtered 0.22 µm)
 - 25 ml Yeast extract (15%, sterile solution)
 - 10 ml D-(+)-Glucose (sol 50% in H₂O, filtered 0.22 µm)
 - 1 g Ampicillin
 - 4 ml Phenol red, 0.5%
 - 10 ml Sodium pyruvate (sol 0.5 M in H₂O, filtered 0.22 µm)
2. SP4 agar medium
 - Proceed as for SP4 liquid medium but with some modifications
 - Before autoclaving add:*
 - 9 g select agar
 - After autoclaving*
 - Do not add phenol red and sodium pyruvate
3. *M. agalactiae* Lysis buffer
 - 1 ml Tris-HCl (1 M, pH 8.5)
 - 5 µl Tween-20
 - 125 µl Proteinase K 20 mg/ml
 - Add sterile ultrapure water up to a final volume of 10 ml

Acknowledgments

This work was supported by Grant ANR09MIE016 (MycXgene) from the French national funding research agency (ANR), by INRA and ENVT. The protocol is derived from works published in Dordet-Frisoni *et al.* (2013 and 2019).

Competing interests

The authors state that there are no competing interests

References

1. Baranowski, E., Guiral, S., Sagne, E., Skapski, A. and Citti, C. (2010). [Critical role of dispensable genes in *Mycoplasma agalactiae* interaction with mammalian cells.](#) *Infect Immun* 78(4): 1542-1551.
2. Calcutt, M. J., Lewis, M. S. and Wise, K. S. (2002). [Molecular genetic analysis of ICEF, an integrative conjugal element that is present as a repetitive sequence in the chromosome of *Mycoplasma fermentans* PG18.](#) *J Bacteriol* 184(24): 6929-6941.
3. Chazel, M., Tardy, F., Le Grand, D., Calavas, D. and Poumarat, F. (2010). [Mycoplasmoses of ruminants in France: recent data from the national surveillance network.](#) *BMC Vet Res* 6: 32.
4. Dordet-Frisoni, E., Marena, M. S., Sagne, E., Nouvel, L. X., Guerillot, R., Glaser, P., Blanchard, A., Tardy, F., Sirand-Pugnet, P., Baranowski, E. and Citti, C. (2013). [ICEA of *Mycoplasma agalactiae*: a new family of self-transmissible integrative elements that confers conjugative properties to the recipient strain.](#) *Mol Microbiol* 89(6): 1226-1239.
5. Dordet-Frisoni, E., Faucher, M., Sagne, E., Baranowski, E., Tardy, F., Nouvel, L. X. and Citti, C. (2019). [Mycoplasma Chromosomal Transfer: A Distributive, Conjugative Process Creating an Infinite Variety of Mosaic Genomes.](#) *Front Microbiol* 10: 2441.
6. Dordet-Frisoni, E., Sagne, E., Baranowski, E., Breton, M., Nouvel, L. X., Blanchard, A., Marena, M. S., Tardy, F., Sirand-Pugnet, P. and Citti, C. (2014). [Chromosomal transfers in mycoplasmas: when minimal genomes go mobile.](#) *mBio* 5(6): e01958.
7. Faucher, M., Nouvel, L. X., Dordet-Frisoni, E., Sagne, E., Baranowski, E., Hygonenq, M. C., Marena, M. S., Tardy, F. and Citti, C. (2019). [Mycoplasmas under experimental antimicrobial selection: The unpredicted contribution of horizontal chromosomal transfer.](#) *PLoS Genet* 15(1): e1007910.
8. Lederberg, J. and Tatum, E. L. (1946). [Gene recombination in *Escherichia coli*.](#) *Nature* 158(4016): 558.
9. Marena, M., Barbe, V., Gourgues, G., Mangenot, S., Sagne, E. and Citti, C. (2006). [A new integrative conjugative element occurs in *Mycoplasma agalactiae* as chromosomal and free circular forms.](#) *J Bacteriol* 188(11): 4137-4141.
10. Meygret, A., Peuchant, O., Dordet-Frisoni, E., Sirand-Pugnet, P., Citti, C., Bebear, C., Beven, L. and Pereyre, S. (2019). [High Prevalence of Integrative and Conjugative Elements Encoding Transcription Activator-Like Effector Repeats in *Mycoplasma hominis*.](#) *Front Microbiol* 10: 2385.
11. Rosengarten, R., Behrens, A., Stetefeld, A., Heller, M., Ahrens, M., Sachse, K., Yogev, D. and Kirchhoff, H. (1994). [Antigen heterogeneity among isolates of *Mycoplasma bovis* is generated by high-frequency variation of diverse membrane surface proteins.](#) *Infect Immun* 62(11): 5066-5074.
12. Sirand-Pugnet, P., Citti, C., Barre, A. and Blanchard, A. (2007). [Evolution of mollicutes: down a bumpy road with twists and turns.](#) *Res Microbiol* 158(10): 754-766.

13. Tardy, F., Mick, V., Dordet-Frisoni, E., Marenda, M. S., Sirand-Pugnet, P., Blanchard, A. and Citti, C. (2015). [Integrative conjugative elements are widespread in field isolates of *Mycoplasma* species pathogenic for ruminants](#). *Appl Environ Microbiol* 81(5): 1634-1643.
14. Zimmerman, C. U. and Herrmann, R. (2005). [Synthesis of a small, cysteine-rich, 29 amino acids long peptide in *Mycoplasma pneumoniae*](#). *FEMS Microbiol Lett* 253: 315-321.