



HAL
open science

Whole-Genome Sequence of a *Brucella pinnipedialis* Sequence Type 54 Strain Isolated from a Hooded Seal (*Cystophora cristata*) from the North Atlantic Ocean, Norway

Michel S Zygmunt, Gilles Vergnaud, Axel Cloeckaert

► **To cite this version:**

Michel S Zygmunt, Gilles Vergnaud, Axel Cloeckaert. Whole-Genome Sequence of a *Brucella pinnipedialis* Sequence Type 54 Strain Isolated from a Hooded Seal (*Cystophora cristata*) from the North Atlantic Ocean, Norway. *Microbiology Resource Announcements*, 2021, 10 (18), 2 p. 10.1128/MRA.00271-21 . hal-03222575

HAL Id: hal-03222575

<https://hal.inrae.fr/hal-03222575>

Submitted on 7 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Whole-Genome Sequence of a *Brucella pinnipedialis* Sequence Type 54 Strain Isolated from a Hooded Seal (*Cystophora cristata*) from the North Atlantic Ocean, Norway

Michel S. Zygmunt,^a Gilles Vergnaud,^b Axel Cloeckaert^a

^aINRAE, UMR ISP, Université de Tours, Nouzilly, France

^bUniversité Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell, Gif-sur-Yvette, France

ABSTRACT Since the 1990s, *Brucella* strains have been isolated from a wide variety of marine mammal species. We report the first complete genome sequence of a *Brucella* strain isolated from a hooded seal (*Cystophora cristata*), *Brucella pinnipedialis* strain 23a-1 of sequence type 54, found in the North Atlantic Ocean surrounding Norway.

B *Brucella* constitutes a genus comprising pathogenic species affecting livestock and causing disease in humans, of which the main representative, in terms of global health impact, is *Brucella melitensis* (1). Since the 1990s, a number of new species, mostly from wildlife, have been identified. Among them are the species *Brucella ceti* and *Brucella pinnipedialis* isolated from marine mammal species, with cetaceans and pinnipeds, respectively, as preferred hosts (2). These species could be further subdivided into a large number of subtypes by genotyping. In particular, multilocus sequence typing (MLST) and multiple-locus variable-number tandem repeats (VNTR) analysis (MLVA) proved to be highly discriminatory (3–5). The latest MLST21 scheme (5) identified over 100 *Brucella* sequence types (STs), including 16 STs for marine mammal brucellae. Similarly, MLVA16 identified seven major clusters of strains within *B. ceti* and *B. pinnipedialis*, with distinct MLVA16 genotypes in each cluster (3). Some clusters appeared to be restricted to specific marine mammal hosts. For example, MLVA16 cluster C3 genotypes are restricted to a single seal species host, i.e., hooded seal (*Cystophora cristata*). In MLST21, hooded seal isolates currently constitute two STs, ST53 and ST54 (5). The hooded seals form two populations, the Northeast and Northwest Atlantic populations, and their specific biology, physiology, habitat, and behavior, relative to those of other seal species, have been reviewed by Nymo et al. (6).

Among hooded seal strains previously typed by MLVA16 (3), we selected for whole-genome sequencing (WGS) *B. pinnipedialis* strain 23a-1, constituting a single specific genotype in the MLVA16 C3 cluster (<https://microbesgenotyping.i2bc.paris-saclay.fr/databases/public>) (3). This strain was isolated from a hooded seal found in the Northeast Atlantic Ocean in Norway (3). The genomic DNA was extracted from the trypticase soy agar with 0.6% yeast extract (TSAYE)-cultured strain with the DNeasy blood and tissue kit (Qiagen, Les Ulis, France). WGS and library construction were performed by GenoScreen (Lille, France), using the PacBio Sequel platform. The sequencing libraries were generated with a SMRTbell template preparation kit v3.0. SMRT Link (v5.1.0.26412) was used for quality control and filtering, which resulted in 127,654 subreads with an average length of 6,868 bp and an N_{50} value of 8,281 bp. The assembly was circularized by Circlator v1.5.5 using the raw reads corrected by Canu v1.6. Default parameters were applied for all software packages. The genome sequence was annotated by the NCBI Prokaryotic Genome Annotation Pipeline v4.13 (7).

The whole genome of strain 23a-1 consisted of two circular chromosomes, with

Citation Zygmunt MS, Vergnaud G, Cloeckaert A. 2021. Whole-genome sequence of a *Brucella pinnipedialis* sequence type 54 strain isolated from a hooded seal (*Cystophora cristata*) from the North Atlantic Ocean, Norway. Microbiol Resour Announc 10:e00271-21. <https://doi.org/10.1128/MRA.00271-21>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2021 Zygmunt et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Michel S. Zygmunt, michel.zygmunt@inrae.fr.

Received 11 March 2021

Accepted 15 April 2021

Published 6 May 2021

sizes of 2,120,314 bp and 1,192,109 bp and with a deduced G+C content of 57.3%, as seen for other *Brucella* species. WGS confirmed the specific MLVA16 genotype of strain 23a-1, and the deduced ST in MLST21 was ST54. The PacBio technology also allowed identification of the distribution of the genus-specific insertion sequence IS711 (8). A total number of 32 IS711 elements were identified, dispersed on both chromosomes. In addition, the lack of a 67-kb DNA fragment flanked by IS711 elements located on chromosome 2, relative to *B. pinnipedialis* strains from other seal species (9), was confirmed.

Data availability. The complete genome sequence of *B. pinnipedialis* strain 23a-1 can be retrieved from the European Nucleotide Archive within BioProject [PRJNA664313](https://www.ebi.ac.uk/bioproject/131313), with assembly accession number [GCA_015624465](https://www.ncbi.nlm.nih.gov/assembly/GCA_015624465) (accession number [CP061816.1](https://www.ncbi.nlm.nih.gov/assembly/CP061816.1) for chromosome 1 and accession number [CP061815.1](https://www.ncbi.nlm.nih.gov/assembly/CP061815.1) for chromosome 2). The raw sequencing results were deposited in the NCBI SRA database under accession number [SRR14150359](https://www.ncbi.nlm.nih.gov/sra/SRR14150359).

ACKNOWLEDGMENTS

We thank Jacques Godfroid for supplying *B. pinnipedialis* strain 23a-1. We also thank Sébastien O. Leclercq for his help in the submission of the genome sequence to GenBank.

This work was funded by Agence Nationale de la Recherche grant ASTRID-Maturation ANR-14-ASMA-0002-02.

REFERENCES

- Moreno E. 2021. The one hundred year journey of the genus *Brucella* (Meyer and Shaw 1920). *FEMS Microbiol Rev* 45:fuaa045. <https://doi.org/10.1093/femsre/fuua045>.
- Foster G, Osterman BS, Godfroid J, Jacques I, Cloeckaert A. 2007. *Brucella ceti* sp. nov. and *Brucella pinnipedialis* sp. nov. for *Brucella* strains with cetaceans and seals as their preferred hosts. *Int J Syst Evol Microbiol* 57:2688–2693. <https://doi.org/10.1099/ijs.0.65269-0>.
- Maquart M, Le Flèche P, Foster G, Tryland M, Ramisse F, Djønne B, Al Dahouk S, Jacques I, Neubauer H, Walravens K, Godfroid J, Cloeckaert A, Vergnaud G. 2009. MLVA-16 typing of 295 marine mammal *Brucella* isolates from different animal and geographic origins identifies 7 major groups within *Brucella ceti* and *Brucella pinnipedialis*. *BMC Microbiol* 9:145. <https://doi.org/10.1186/1471-2180-9-145>.
- Vergnaud G, Hauck Y, Christy D, Daoud B, Pourcel C, Jacques I, Cloeckaert A, Zygmunt MS. 2018. Genotypic expansion within the population structure of classical *Brucella* species revealed by MLVA16 typing of 1404 *Brucella* isolates from different animal and geographic origins, 1974–2006. *Front Microbiol* 9:1545. <https://doi.org/10.3389/fmicb.2018.01545>.
- Whatmore AM, Koylass MS, Muchowski J, Edwards-Smallbone J, Gopaul KK, Perrett LL. 2016. Extended multilocus sequence analysis to describe the global population structure of the genus *Brucella*: phylogeography and relationship to biovars. *Front Microbiol* 7:2049. <https://doi.org/10.3389/fmicb.2016.02049>.
- Nymo IH, Tryland M, Godfroid J. 2011. A review of *Brucella* infection in marine mammals, with special emphasis on *Brucella pinnipedialis* in the hooded seal (*Cystophora cristata*). *Vet Res* 42:93. <https://doi.org/10.1186/1297-9716-42-93>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Suárez-Esquivel M, Chaves-Olarte E, Moreno E, Guzmán-Verri C. 2020. *Brucella* genomics: macro and micro evolution. *Int J Mol Sci* 21:7749. <https://doi.org/10.3390/ijms21207749>.
- Audic S, Lescot M, Claverie JM, Cloeckaert A, Zygmunt MS. 2011. The genome sequence of *Brucella pinnipedialis* B2/94 sheds light on the evolutionary history of the genus *Brucella*. *BMC Evol Biol* 11:200. <https://doi.org/10.1186/1471-2148-11-200>.