



HAL
open science

The pig as a medical model for acquired respiratory diseases and dysfunctions: An immunological perspective

Nicolas Bertho, François Meurens

► To cite this version:

Nicolas Bertho, François Meurens. The pig as a medical model for acquired respiratory diseases and dysfunctions: An immunological perspective. *Molecular Immunology*, 2021, 135, pp.254-267. 10.1016/j.molimm.2021.03.014 . hal-03229038

HAL Id: hal-03229038

<https://hal.inrae.fr/hal-03229038>

Submitted on 9 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

1 **The pig as a medical model for acquired respiratory diseases and dysfunction: an immunological**
2 **perspective**

3 Nicolas Bertho* and François Meurens
4 INRAE, Oniris, BIOEPAR, 44300 Nantes, France
5 *Correspondence: nicolas.bertho@inrae.fr
6

7 **Abstract**

8 By definition no model is perfect, and this also holds for biology and health sciences. In medicine,
9 murine models are, and will be indispensable for long, thanks to their reasonable cost and huge
10 choice of transgenic strains and molecular tools. On the other side, non-human primates remain the
11 best animal models although their use is limited because of financial and obvious ethical reasons. In
12 the field of respiratory diseases, specific clinical models such as sheep and cotton rat for
13 bronchiolitis, or ferret and Syrian hamster for influenza and Covid-19, have been successfully
14 developed, however, in these species, the toolbox for biological analysis remains scarce. In this view
15 the porcine medical model is appearing as the third, intermediate, choice, between murine and
16 primate. Herein we would like to present the pros and cons of pig as a model for acquired respiratory
17 conditions, through an immunological point of view. Indeed, important progresses have been made
18 in pig immunology during the last decade that allowed the precise description of immune molecules
19 and cell phenotypes and functions. These progresses might allow the use of pig as clinical model of
20 human respiratory diseases but also as a species of interest to perform basic research explorations.
21

22 **Generalities**

23 In its ‘Global Impact of Respiratory Disease’, the World Health Organization (WHO) identifies
24 ‘the big five’ respiratory diseases as chronic obstructive pulmonary disease (COPD), asthma, acute
25 lower respiratory tract infections, tuberculosis and lung cancer. Moreover, according to Eurostat
26 report, diseases of the respiratory system accounted for 7.5 % of all deaths in the European Union in
27 2016 (Eurostat, 2020), and these statistics do not take into account the current SARS-CoV-2
28 pandemic which will probably become the infectious disease associated to the highest number of
29 fatalities in 2020. In the pathophysiology of all these diseases, the immune system has definitely a
30 central role and we need to better understand how it operates in order to prevent and treat the
31 respiratory diseases.

32 Current Food and Drug Administration (FDA) guidelines require testing of new
33 pharmaceutical agents in both small and large animal models when proving a therapeutic concept
34 (FDA, 2015). We advocate herein to consider pig as a model of choice when investigating the
35 immunological side of respiratory diseases.

36 The *Suidae* family (swine, wild boar, red river hog, and warthog to cite the best known) is
37 evolutionary more distant from the *Hominidae* family (including human being and all the great apes)
38 than the *Muridae* family (mouse, rat, hamster and naked mole rat). Indeed, men and mice belong to
39 the same *Euarchontoglires* superorder whereas pigs belong to the *Laurasiatheria* superorder. Despite
40 this, and probably as a consequence of the faster genetic evolutionary rate of mouse, the pig genome
41 has remained more similar to the human one both in terms of DNA sequences than in term of genes’
42 synteny (Humphray et al., 2007; Wernersson et al., 2005). Because of this genetic proximity and for
43 other reasons listed below, the pig has been largely used as a biomedical model and notably as a
44 model for the study of human infectious diseases and for vaccine development (Gerdtts et al., 2015;
45 Käser et al., 2018; Lunney, 2007; Meurens et al., 2012). In this review we focused especially on the
46 pig as a model to study acquired respiratory diseases except cancer, at the light of the accumulating
47 knowledge concerning the swine respiratory immune system.

48

49 **Porcine Respiratory System Structure**

50 The size and the weight of the pig is age-, sex- and breed-dependent. Some minipig breeds
51 like the Yucatan or the Göttingen present adult pigs with size and weight similar to human adults
52 making the breeds attractive in biomedical research (Meurens et al., 2012; Swindle et al., 2012).
53 Overall, the porcine anatomy and physiology are quite similar to human anatomy and physiology,
54 both species are all-eating for instance (Käser et al., 2018; Meurens et al., 2012; Swindle et al., 2012).
55 Moreover, pigs are widely available with usually manageable size allowing for smooth handling and
56 easy experimental interventions.

57 Regarding specifically the porcine respiratory tract, the pig has been extensively
58 characterized and compared with the human one (Judge et al., 2014; Krejci et al., 2013) as well as in
59 exhaustive reviews that were assessing its potential as a model for cystic fibrosis (Rogers et al., 2008)
60 and asthma (Kirschvink and Reinhold, 2008). The respiratory tract can be divided into the upper and
61 lower respiratory tracts. The upper respiratory tract is including the nasal cavities, the sinuses, the
62 nasopharynx, the larynx and the trachea while the lower respiratory tract includes overall the lung
63 with the bronchi, the bronchioles, the terminal and respiratory bronchioles, and alveoli. Air enters
64 the porcine respiratory tract through round nostrils and a pair of nasal cavities situated in the snout
65 (*rostrum*). In the nasal cavities, the epithelium is typically pseudostratified with ciliated columnar
66 cells. The epithelium contains loose lymphocytes, lympho-reticular tissue aggregates and tubo-
67 alveolar glands. Regarding submucosal glands in nasal turbinates and trachea, some differences have
68 been shown between pigs and humans using the porcine cystic fibrosis model (Cho et al., 2011).
69 Similarly, to humans, pigs have ethmoid and maxillary sinuses at birth and develop frontal and
70 sphenoid sinuses (Sisson et al., 1975). Then, after its circulation in the nasal cavities, the air moves
71 through the nasopharyngeal tube into the pharynx with the possibility to gain access to the
72 esophagus. The respiratory part of the pharynx, called nasopharynx, also presents a pseudostratified
73 ciliated columnar epithelium with goblet cells. In the upper respiratory tract, its nasal-associated
74 lymphoid tissue (NALT) resembles the human's Waldeyer ring and pigs, like humans, possess several
75 tonsils (*veli palatine*, *pharyngea*, *tubaria*, *paraepiglottica* and *lingualis*) (Horter et al., 2003; Liebler-
76 Tenorio and Pabst, 2006; Swindle et al., 2012). This lymphoid tissue is comprised of the *Lamina*
77 *propria mucosae* as well as fine glandules. Then, the larynx links the nasopharynx and the lower
78 respiratory tract, including epiglottis and vocal cords. In the larynx the mucosa showed again a
79 pseudostratified columnar epithelium with isolated goblet cells. In the submucosa of epiglottis, *plicae*
80 *aryepiglotticae* and *vestibulum larynges*, an accumulation of lymphoid tissue is observed. The trachea
81 consists of 32-36 C-shaped rings of hyaline cartilage with strong fibro-elastic membranes observed
82 between the rings. The trachea shows a ciliated pseudostratified columnar epithelium and isolated
83 goblet cells. Below, in the mucosa, small islets of lympho-reticular tissue and combined tubo-alveolar
84 glands are observed. Regarding the gross anatomy of the porcine lung, two lobes on the left sides
85 and four on the right have been described (Rogers et al., 2008; Swindle et al., 2012). On the left side,
86 the lobes are designated as left cranial and caudal and on the right as right cranial, right middle, right
87 caudal and right accessory. Interestingly, the right cranial lobe is directly linked to the trachea.
88 Looking at subgross anatomy, pig like human lung, and in contrast to murine lung shows extensive
89 intralobular and interlobular connective tissue. This tissue joins the bronchi and the main blood
90 vessels to the pleural surface (Tyler, 1983). However, the interlobular collagenous network is
91 incomplete in humans whereas in the pig the interlobular septa are "complete" blocking the
92 collateral ventilation (Rogers et al., 2008; Woolcock and Macklem, 1971). The porcine pleura like its
93 human counterpart is relatively thick and collagenous and has a vascular supply originating from the
94 bronchial arteries (McLaughlin et al., 1961; Tyler, 1983). Regarding the lymph nodes (LN) in the
95 thoracic cavity, they are concentrated into four lymphoid centres, two parietal (*thoracicum dorsal*

96 and *ventral*) and two visceral (*mediastinal* and *bronchale*). Moreover, there are the LN adjacent to
97 the bronchi (*tracheobronchales dextri, sinistri* and *medii*). While the vascular circulation of the pig
98 follows a common pattern of organization and development with human vascular circulation there is
99 an inverted structure of swine LN compared to their human and murine counterparts. Indeed, in
100 most of the mammals, including mouse and man, afferent lymph percolates in a centripetal
101 manner from the periphery to the center (hilum) of the node. In swine, the afferent lymph diffuses
102 from the center to the periphery in a centrifugal manner (Mcfarlin et al., 1973). In addition, in pig,
103 lymphocytes exit the LN through the high-endothelial venules whereas in mouse and man they exit
104 through the efferent lymphatics (Sasaki et al., 1994). The functional implications of these oddities
105 have not been explored yet. Overall, even if similar to human airways, the porcine airways show
106 small differences (Haworth and Hislop, 1981; McLaughlin et al., 1961; Rendas et al., 1978): i) the
107 cartilage around porcine airways is more developed than in humans; ii) a lower number of bronchi
108 generations, after the last cartilage plate, has been reported in pigs compared to what is reported in
109 humans; and iii) longer terminal bronchioles and overall less-defined respiratory bronchioles are
110 described. At the microscopic level, porcine lungs show similar cell lineages to human lungs (Haworth
111 and Hislop, 1981; Jones et al., 1975; Mills et al., 1986; Rogers et al., 2008; Winkler and Cheville,
112 1984).

113

114 **General systemic immunology**

115 Before addressing specifically the porcine lung immunology, a brief overview of the systemic
116 porcine immune system is needed. Comparably to the whole genome comparisons, the immunome
117 analysis demonstrated a greater similarity between human and pig than between human and mouse.
118 However, a peculiarity of swine immune system that can be highlighted is the expansion of the type I
119 IFN gene families, especially the IFN δ and the IFN ω (Dawson et al., 2013). IFN δ clearly distinguishes
120 pig from human and mouse since pig has 11 functional IFN δ genes and mouse and man have none.
121 IFN δ bind to the same type 1 receptor as IFN α , it has high antiviral and anti-proliferative activities on
122 porcine cells, but not on human cells (Lefèvre et al., 1998). Conversely IFN ω separates mouse from
123 the two other species since mouse has no functional IFN ω gene, man 1, and pig 7. IFN ω presented
124 highly cross-species antiviral (but little anti-proliferative) activity (Jennings and Sang, 2019; Shields et
125 al., 2019). A recent review summarized the different porcine cytokines, chemokines and growth
126 factors, and described the tools available to study them (Dawson et al., 2020).

127 Looking more precisely at the different immune cell types, three relatively recent reviews
128 have been published on the porcine innate immune response (Mair et al., 2014), mononuclear
129 phagocytes (Fairbairn et al., 2011) and the B and T lymphocytes development (Sinkora and Butler,
130 2016). However, we would like here to recall some important swine specific features:

131 **Myeloid cells: Neutrophil** serine proteases (NSP) have critical roles in neutrophil-associated tissue-
132 destructive diseases. Human and mouse NSP present different peptide substrate specificities
133 (Kalupov et al., 2009) whereas porcine NSP, present on the surface of triggered neutrophils and
134 neutrophil extracellular traps (NETs) are efficiently inhibited by human NSP inhibitors (Bréa et al.,
135 2012). In addition to neutrophils, porcine basophils and eosinophils can be identified by flow
136 cytometry (Blanc et al., 2020; Haverson et al., 1994).

137 **Peripheral blood monocytes** present in pig a similar division in CCR2+/CX3CR1-/CD14+/CD163-
138 /MHC-II- and CX3CR1+ CCR2-/CD14-/CD163+/MHC-II+ monocytes (Moreno et al., 2010). When
139 human, mouse and porcine monocytes were compared by a transcriptomic study, porcine CD14+ and
140 CD14- blood monocytes resembled more to their human counterparts than mouse monocytes
141 (Fairbairn et al., 2013).

142 **Dendritic cells** (DC) subpopulations corresponding to cDC1, cDC2, pDC and moDC (Guilliams et al.,
143 2014) can be readily identified in blood (Auray et al., 2016), skin (Marquet et al., 2014), spleen
144 (Soldevila, personal communication), tonsils (Soldevila et al., 2018), tracheal epithelium and lung
145 (Maisonasse et al., 2016), with minimal species-specific features. The main swine DC peculiarity is
146 the differential expression of TLR3, which is almost exclusively expressed on pDC in swine (Auray et
147 al., 2016; Soldevila et al., 2018) but on cDC1 in human (Poulin et al., 2010) and mouse (Desch et al.,
148 2011). Despite this difference, the calculation of a similarity score between swine, human and mouse
149 cDC1, cDC2 and pDC transcriptomic signatures highlighted that the three human's blood DC
150 populations resembled more to their porcine than to their murine counterparts, especially cDC2,
151 which appears to be the most similar between pig and man (Auray et al., 2016).

152 **Lymphoid cells:** It has long been thought that porcine B cells were generated in ileal Peyer's patches,
153 however it is now clear that porcine B lymphopoiesis is located in the bone marrow as for other
154 mammals (Sinkora and Sinkorova, 2014). To note, the porcine B cell development studies are still
155 impaired because of the absence of antibodies recognizing pan-B cells markers such as CD19 and
156 CD20 (Dawson and Lunney, 2018) and the absence of antibodies discriminating porcine IgG isotypes.
157 Despite these difficulties, the porcine peripheral B lymphocyte differentiation steps have been
158 described: IgM+CD2+CD21+ B cells are composed of naive B cells, IgM+CD2-CD21+ are activated or
159 primed B cells and CD79 α +CD2+CD21- or CD2-CD21- represent antibody producing B cells (Sinkora et
160 al., 2013), memory and effector antibody-forming B cells remaining CD79 α +CD2+CD21- while
161 CD79 α +CD2-CD21- B cells are probably composed of resting plasma cells. In a recent paper (Bordet
162 et al., 2019), we investigated the structure of the trachea-bronchial inverted porcine LN and, thanks
163 to the conservation of the B cell affinity maturation master-genes expressions (BCL6, Pax5, IRF4,
164 XBP1 and Blimp1), we were able to distinguish the centroblasts, centrocytes, plasmablasts and
165 plasma cells in their respective follicular and extrafollicular LN sublocalisations, as observed in mouse
166 and human (for review see (Klein and Dalla-Favera, 2008)). Interestingly, we also identified a specific
167 feature of swine inverted LN which is a centroblasts-like B cell type decorated with surface CD169
168 proteins (Bordet et al., 2019), that interact probably with perifollicular macrophages in order to
169 capture antigens and transport them to follicular DC (fDC). This feature might be a consequence of
170 LN inversion in pig. In the regular murine LN, naive B cells have been shown to play this antigen-
171 transporting role (Phan et al., 2009). Follicular DC have not been described so far in pig, however
172 they have been identified in an evolutionary close species, the sheep, and their phenotype appeared
173 similar to murine and human fDC (Melzi et al., 2016).

174 Development of thymic $\alpha\beta$ T cells follows mice and humans T-cell maturation model (Sinkora
175 et al., 2013). However, it is important to note that swine possess extrathymic CD4+/CD8 α + T cells,
176 that appear to be regular memory/activated peripheral CD4 T cells (Gerner et al., 2009). The
177 peripheral blood re-expression of CD8 α has been endowed with no specific role. Indeed, so far the
178 same CD4 T cell subtypes have been described in pig, namely Th1, Th2, Th17, Treg expressing the
179 same transcription factors (respectively T-bet, GATA-3, ROR γ T and FoxP3) than human and murine
180 CD4 T cells (for review see (Gerner et al., 2015)). To note, in swine some clues are in agreement with
181 a B cell activation/Th2 pathways relying more on IL13 than on IL4 (Murtaugh et al., 2009). Finally
182 peripheral CD8 T cells expressing Eomesodermin (Eomes), a transcription factor involved in the
183 differentiation and long-term survival of CD8 memory T cells have been described in pig like in mouse
184 and human blood, although these cells are less numerous in swine than in human (Rodríguez-Gómez
185 et al., 2016). Finally, by using CD27 staining, Talker *et al* were able to distinguish *bona fide* effector
186 memory cells (CD27^{neg}) from naive and early activated lymphocytes (Reutner et al., 2013).

187 As in other *Laurasiatheria*, pigs present a high proportion of $\gamma\delta$ T cells (Binns et al., 1992),
188 paralleled with more D-segments in the TCR delta gene than in human and mouse (Dawson et al.,
189 2013). Thus porcine $\gamma\delta$ TCR have a strong diversity potential, with no tissue-specificity (Holtmeier et

190 al., 2004) or subset restrictions (Stepanova and Sinkora, 2013) unlike in humans and mice. Moreover,
191 conversely to murine and human ones, a consistent proportion of porcine $\gamma\delta$ T cells retained a strong
192 expression of GATA3 (Rodríguez-Gómez et al., 2019).

193 **Invariant NK T cells** expressed a semi-invariant TCR which recognizes glycolipids presented on the
194 non-polymorphic CD1d molecules and corresponding either to stress-related self-ligand or to
195 microorganism surface-antigens. Interestingly, iNKT cells frequency (Artiaga et al., 2014) as well as
196 specific CD1d/TCR molecular interactions (Yang et al., 2019) are closer between swine and human
197 than with mouse.

198 Conversely to the majority of mammals, swine possess an **NK cell subset** that does not
199 express Nkp46 (Mair et al., 2012). This might be relevant here since Nkp46 recognizes
200 hemagglutinins (HA) of influenza, parainfluenza, and Sendai virus and that its ligation leads to lysis of
201 infected cells (Arnon et al., 2004). In swine, NKP46 expression has also been observed on CD3+ cells
202 expressing $\alpha\beta$ or $\gamma\delta$ TCR but presenting the main functions of NK cells (Mair et al., 2016). Cells with a
203 similar phenotype have been described in mice (Arnon et al., 2004) humans (Arnon et al., 2004) and
204 cattle (Connelley et al., 2014).

205

206 **Specific Lung immunology**

207 Macrophages

208 In all mammalian species, lung macrophages are prominently composed of alveolar
209 macrophages (AM). AM main roles are the phagocytosis of inhaled particles and the clearance of
210 surfactant (for review see (Hussell and Bell, 2014)). AM have been shown in mouse to originate from
211 embryonic monocytes that settled before birth in the alveoli and self-renew independently of
212 peripheral blood adult monocytes (Guilliams et al., 2013). This scheme can be modified upon events
213 that trigger both strong inflammation and partial or complete AM depletion. In this case, as
214 documented in mice, recruited inflammatory monocytes will differentiate in pro-inflammatory
215 monocyte-derived macrophages (moM θ) before entering AM 'niche' (for review see (Guilliams and
216 Scott, 2017)) and differentiating finally in 'true' AM, undistinguishable from the original AM
217 population (Aegerter et al., 2020). Although not yet formally demonstrated, the belonging of human
218 AM to the local-self-renewable macrophage type is also postulated. By analyzing AM from lung
219 grafted patients, several teams tried to clarify the importance of AM self-renewal *versus* blood
220 monocytes replacement in human, leading to contradictory conclusions (Bittmann et al., 2001;
221 Eguíluz-Gracia et al., 2016; Nayak et al., 2016). However a recent work using up to date single cell
222 technology (Byrne et al., 2020) arbitrated toward a strong contribution of blood monocytes for long
223 term AM maintenance, the authors arguing that human beings, conversely to lab's mice, are
224 constantly challenged by inflammatory stimuli, in a much longer time frame, and that in human
225 everyday life, the AM pool might be replaced by moM θ within a year. In porcine lung, we observed
226 that AM did not express blood-related genes such as cKit, CSF1R CCR2 or CX3CR1 but did expressed
227 HDAC10, PU.1 (Bordet et al., 2018; Maisonnasse et al., 2016) whose expressions are related to
228 embryonically-derived macrophage precursors in mouse (Guilliams et al., 2013; Schulz et al., 2012),
229 in agreement with a similar local-self-renewable capacity of porcine and mouse AM. However,
230 according to the 'niche theory', these local-self-renewable AM could well be former moM θ that
231 transdifferentiated in true AM upon AM-niche occupancy as postulated in human.

232 To note, similarly to mouse and man (Keller et al., 2015), porcine AM expressed the
233 immunoproteasome subunits LMP2, LMP7, and MECL-1 upon respiratory viral infection (Liu et al.,
234 2018, 2017).

235 One intriguing feature of pulmonary macrophage network in swine, and the majority of the
236 *Laurasiatheria*, is the presence of pulmonary intravascular macrophages (PIM, for review
237 (Schneberger et al., 2012)) that are not present, at least at steady state, in mouse, rat and human

238 (Brain et al., 1999). PIM are resident lung intravascular macrophages intimately tight with endothelial
239 cells, what differentiates them from marginated blood monocytes (for review see (Kuebler and
240 Goetz, 2002)). We recently isolated porcine PIM and demonstrated that they were very similar to AM
241 (Bordet et al., 2018; Crisci et al., 2020), indeed PIM and AM overexpressed genes such as PU.1 and
242 HDAC10, leading us to suggest than PIM were, like AM, self-renewable embryonically derived
243 macrophages, and/or that PIM may directly originate from AM (Bordet et al., 2018). PIM have been
244 involved in acute lung inflammation in *Laurasiatheria* species (Cantu et al., 2007; Singh et al., 2004).
245 Interestingly PIM can be induced in rat model of systemic induced inflammation triggered by biliary
246 duct ligation (Gill et al., 2008) and some clues of PIM induction in humans suffering of liver
247 dysfunctions (Klingensmith et al., 1978, 1976) have been reported. In a recent report using mice
248 grafted with human bone-marrow ('humanized' mice), the authors detected PIM which were, in this
249 model, monocyte-derived human macrophages in tight contact with murine endothelial cells (Evren
250 et al., 2021).

251

252 Interstitial macrophages

253 Interstitial lung moM θ identities and functions have been recently revisited in mouse and
254 man by the identification of two distinct populations (Chakarov et al., 2019), one Lyve1^{low}/MHC-
255 II^{high}/CD163^{low}, and the other Lyve1^{high}/MHC-II^{low}/CD163^{high} respectively residing adjacent to nerve
256 bundles and to blood vessels. The later Lyve1^{high}/MHC-II^{low} moM θ secreted a high basal level of IL10,
257 supported blood vessel integrity, and decreased inflammatory cells infiltrations upon pulmonary
258 inflammation in mice. This degree of precision has not been reached in pig, although we identified an
259 MHC-II^{high}/CD163^{low}/CD169^{high} moM θ population (Maisonasse et al., 2016) as well as an MHC-
260 II^{low}/CD163^{high}/CD169^{low} (unpublished data), whose differential roles and locations remain to be
261 explored.

262

263 Dendritic cells

264 Swine DC have been identified in tonsils (Soldevila et al., 2018), tracheal epithelium and
265 pulmonary parenchyma (Maisonasse et al., 2016). They can be divided in cDC1, cDC2 and moDC
266 whose RNAseq signatures cluster with their mouse spleen and/or lung counterparts (Crisci et al.,
267 2020). Plasmacytoid DC have also been identified in swine tonsil, they are E2.2 and IRF7 positive as
268 observed in mouse and man. The porcine lung cDC and moDC populations present the functional
269 characteristics of *bona fide* DC, migrating toward LN chemokine upon maturation, whereas only cDC,
270 but not moDC significantly trigger naïve CD8 and CD4 T cells proliferation. Conventional DC2 are
271 better in activating CD4 T cells than cDC1, and this in lung as well as in tonsil (Maisonasse et al.,
272 2016; Soldevila et al., 2018), as previously described in mouse and human (Schlitzer and Ginhoux,
273 2014). Again consistent in tonsil and in lung, both porcine cDC1 and cDC2 activate CD8 T cells,
274 conversely to what is observed in mouse (Ng et al., 2018). Interestingly, in humanized mice, human
275 cDC1 and cDC2 expend similarly influenza specific CD8 T cells, however only cDC2 induced mucosal
276 effector CD8 T cells (Yu et al., 2013). This property of cDC2 has not been explored yet in swine lung.

277 Another intriguing transpecies feature of DC is the expression of Langerin on cDC2 in pig
278 (Maisonasse et al., 2016) and human (Bigley et al., 2015) but on cDC1 in mouse (Sung et al., 2006).
279 It has been shown in human that Langerin is rapidly induced in blood cDC2 upon TGF β exposure
280 (Bigley et al., 2015). Interestingly, this might be interpreted in conjunction with the expression of
281 CD103 (α E β 7 integrin) on lung porcine cDC2, as well as with the sub-epithelial location of cDC2 in
282 man (Yu et al., 2013) and pig (Maisonasse et al., 2016). Indeed, TGF β is largely produced by
283 respiratory epithelial cells and Langerin, like CD103, is induced upon TGF β exposure (Parker et al.,
284 1992; Picarda et al., 2016). The sub-epithelial location of cDC2 in humans and pigs might explain both
285 the Langerin and CD103 expression phenotype and the higher involvement of cDC2 in the activation

286 of CD8 response compared with their mouse counterparts (Yu et al., 2013). FcεRIα is expressed in pig
287 (Maisonnasse et al., 2016) and human (Greer et al., 2014) but not in mouse cDC2. Indeed in mouse
288 inflammatory moDC expressed the FcεRIα (for review see (Lambrecht and Hammad, 2012)). This
289 difference might be of great importance in the feedback of allergic response since IgE/FcεRIα
290 signaling in human cDC2 trigger an inhibition of the inflammatory response (Platzer et al., 2015)
291 whereas it is thought that FcεRIα expression on mouse moDC would allow an IgE-mediated allergen
292 presentation in a positive amplification loop. Interestingly, a recent work in mice complicated the
293 cDC1/cDC2/moDC picture by demonstrating that upon inflammation lung cDC2 can arbor cDC1 and
294 moDC features such as IRF8, CD64 and FcεRIα expressions (Bosteels et al., 2020). It would be
295 interesting to investigate if these modifications stand true for human and porcine cDC2. Finally, in
296 pig, monocyte-derived DC have been identified upon influenza infection and their phenotype and
297 functions are compatible with inflammatory DC, harboring low migratory capacities and strong
298 production of IL1β and IL8 upon stimulation (Maisonnasse et al., 2016).

299

300 Innate lymphocytes

301 NK cells were shown to decrease in peripheral blood and to preferentially localize close to
302 infected area in porcine influenza infected lung (Forberg et al., 2014), in agreement with an
303 important role in the control of influenza infection as observed in mice (Gazit et al., 2006). The
304 CD3+NKp46+ lymphocyte population described by Mair et al. (Mair et al., 2016) was minimally
305 detected in spleen, blood and mediastinal LN (< 2%) but presented a higher frequency, associated to
306 a large variability according to the animal, in lung (1% to 10%), what may suggest a role of this
307 population in lung immune surveillance in swine. Invariant NKT (iNKT) cells binding human CD1d
308 tetramers loaded with an α-GalCer analog have been described in the porcine lung. They are CD3 and
309 CD44 positive but do not express Nkp46 and CD11b, and can be subdivided in three main populations
310 expressing no, low or high levels of CD8α (Yang et al., 2017a). Interestingly, it has been shown
311 recently that porcine iNKT cells were able to respond to influenza-exposed myeloid cells *in vitro*
312 (Schäfer et al., 2019).

313 To our knowledge, an interesting observation has been done only in pig lung, which is the
314 demonstration that γδ T cells were the main alveolar population able to migrate to bronchial LN
315 (Pabst and Binns, 1995).

316

317 T Lymphocytes

318 As expected from mouse and man data, it has been observed in pig that parasitic *Ascaris*
319 *suum* infection bias the lung immune response toward a Th2 response (GATA3, IL4, IL5 upregulation
320 in the whole lung tissue) (Steenhard et al., 2009) and that *Actinobacillus pleuropneumoniae* bacterial
321 infection triggered a Th17 immune response, as validated by the presence of antigen-specific IL17A
322 secreting CD4 T cells (Sassu et al., 2017). Intranasal influenza vaccination triggered resident memory
323 T cells (Holzer et al., 2018). Moreover, proliferating poly-functional Th1 (IFNγ, TNFα and IL2-
324 secreting) and CD8 (IFNγ, TNFα-secreting) lymphocytes can be detected in the trachea-bronchial LN
325 and the lung of influenza infected animals, from 6 to 44 days post-infection (Talker et al., 2016).
326 Interestingly flu-specific IFNγ-secreting CD8 T cells expressed perforine and were 30 times
327 overrepresented in the lung than in the peripheral blood. Cytokine production was dominated by T
328 cells with an early effector phenotype or central memory phenotype (CD27^{pos}) as observed in mice
329 (Ballesteros-Tato et al., 2010) in agreement with the existence in porcine lung of tissue-resident
330 memory T cells.

331

332 B lymphocytes

333 To our knowledge the B cell response in the porcine respiratory tract has only been
334 addressed through nasal swabs and broncho-alveolar lavages antibody monitoring (among others
335 (Bernelin-Cottet et al., 2019; Heinen et al., 2000)).

336

337 In conclusion, considering the respiratory immune system, mice and swine present both similarities
338 and differences compared with human beings (Table 1), which must be consciously considered
339 before choosing one species as an animal model for the pathogen of interest.

340

341 **Model**

342 We will consider here the porcine models developed to tackle the main respiratory diseases
343 identified by the WHO, at the exception of lung cancer. *Id est*, respiratory infections and vaccination,
344 allergy and asthma, acute and chronic inflammations, to which we will add two short chapters on
345 microbiota and lung graft (Table 2).

346 Respiratory infections

347 Swine has been frequently used to study human respiratory pathogens naturally infecting the
348 pigs, including, amongst others, *Mycobacterium* sp (Bolin et al., 1997; Ramos et al., 2017),
349 *Pseudomonas aeruginosa* (Dehring et al., 1983; Luna et al., 2009), *Staphylococcus aureus* (Martínez-
350 Olondris et al., 2010) and *Alphainfluenzavirus* (Rajao and Vincent, 2015). Alternatively, pigs can be
351 susceptible to experimental infections of otherwise strictly human pathogens such as *Bordetella*
352 *pertussis* (Foreman-Wykert and Miller, 2005). One of the main gap of naturally occurring respiratory
353 infections in swine is the absence of a porcine equivalent of the human respiratory syncytial virus
354 (RSV), a very significant pathogen in human, although an orthopneumovirus close to RSV has been
355 recently detected in pigs (Hause et al., 2016) and might be present in France too (Richard et al.,
356 2018). In this chapter we will expose the interest of the pig model for *P. aeruginosa*, influenza A and
357 coronavirus infections.

358 -The bacterium *P. aeruginosa* rarely infects human lungs unless the host immune system has been
359 impaired because of cystic fibrosis (CF), chronic obstructive disease (COPD) or ventilator-associated
360 pneumonia. An infectious model has been developed in pig that recapitulates all the main features of
361 this human infection, including bronchial contraction, transient increase in pro-inflammatory
362 cytokines (IL8, IL6 and TNF α), neutrophilia, neutrophil extracellular trap (NET)osis, and the secretion
363 of massive amounts of neutrophil serine proteases leading to lung damages (Chevaleyre et al., 2016).

364 -*Orthomyxoviridae* and more specifically *Alphainfluenzavirus* (IAV) are pathogens of major
365 importance in animal and human medicines. As recently reviewed, pig is more and more considered
366 as an alternative model to consider with ferret and mouse, for the study of human influenza infection
367 (Starbæk et al., 2018). Pigs are natural hosts of different strains of IAV (Kuntz-Simon and Madec,
368 2009), presenting frequent interspecies transmissions from pig to man and the reverse (Chastagner
369 et al., 2019a, 2019b), what underlines the proximity of swine and human IAV as well as the similarity
370 of the respiratory systems in both species. The most frequently encountered subtypes in pigs, H1N1,
371 H1N2 and H3N2, are the same than in humans. Besides that, the interest for the pig model is
372 strengthened by the fact that pig is considered as a mixing vessel of human, porcine and avian
373 influenza virus for the generation of new influenza virus reassortants (Ma et al., 2009). The
374 susceptibility of pigs to infections with both avian and mammalian influenza virus was recently
375 explained to some extent in a study showing that the porcine host factor ANP32A, contrary to
376 porcine ANP32B and other mammalian ANP32, had stronger supporting activity to the avian viral
377 RNA polymerase (Zhang et al., 2020). In pigs, IAV are responsible of mild diseases (for review (Crisci
378 et al., 2013)) with low fever (40.5°C versus 39.5°C for normal body temperature), low inflammatory
379 signs of the upper respiratory tract (nasal/ ocular discharge and conjunctivitis), dyspnea and

380 coughing (Talker et al., 2016) very similar to human common, mild influenza infections. In mice,
381 which are not the natural hosts for IAV, adapted strains trigger whole lung infection and
382 inflammation, leading to death, more similar to highly pathogenic avian influenza infections in
383 human (Gao et al., 2013). Ferrets are recognized among the best model for testing pathogenicity and
384 transmission of human respiratory viruses including IAV (Enkirch and von Messling, 2015), although
385 the evolutionary reasons for this convergence are still a mystery. Interestingly, a recent comparison
386 of influenza heterosubtypic vaccination in pigs and ferrets, showed different outcome between
387 species and underlined the interest of using different preclinical models to assess new vaccines
388 against flu (Holzer et al., 2018). Which animal model, the pig or the ferret, reflects best the situation
389 in human remains however to be determined.

390 By keeping the immune response under control, AM are recognized as the 'peace-keepers' of
391 the lung (Schneider et al., 2014). In mouse (Schneider et al., 2014) like in pig (Kim et al., 2008), AM
392 depletion aggravate the IAV-mediated disease. In pigs, similarly to humans, disease severity has been
393 associated with increased local pro-inflammatory cytokines (Barbé et al., 2011; Khatri et al., 2010).
394 This cytokine storm has been associated in mice to the arrival of inflammatory moDC in the lung,
395 indeed, pharmacological or genetic downregulation of moDC trafficking moderates the early
396 inflammation, reduces morbidity and mortality without impacting the mounting of the adaptive
397 immune response (Aldridge et al., 2009). This model might also stand for IAV infection in swine, since
398 we observed the arrival of similar pro-inflammatory moDC in the porcine lung as soon as 2 days post-
399 IAV inoculation (Maisonasse et al., 2016). As in mouse (Westerhof et al., 2019) and human
400 (Bonduelle et al., 2014), influenza virus infections in pigs trigger multifunctional blood (Talker et al.,
401 2015) and lung (Talker et al., 2016) antigen-specific CD8 and CD4 T cells. Interestingly a strain of
402 inbred animals, the Babrahama pigs, allowed the use of swine major histocompatibility class I (MHC-
403 I) tetramers to analyze the anti-NP CD8 T cells raised upon influenza intranasal vaccination (Tungatt
404 et al., 2018), and influenza infection (Edmans et al., 2021). This last study also detected a strong
405 influx of $\gamma\delta$ T cells in the alveolar space during influenza infection. Similarly, and in agreement with a
406 role in respiratory immunity, the CD3+NKp46+ lymphocyte population described in swine by Mair et
407 al. (Mair et al., 2016) presented a strong increase in the lung of influenza infected animal, most
408 probably due to local proliferation. SCID human beings and pigs presenting an autologous deficiency
409 in Artemis gene (DCLRE1C) (Moshous et al., 2001) do not develop adaptive immune T and B
410 lymphocytes but harbor functional NK cells (Powell et al., 2016). Human affected by this mutation
411 present recurrent respiratory infections but not recorded influenza infections (Volk et al., 2015), and
412 survived early childhood thanks to intravenous Ig therapy. Interestingly SCID/Artemis pigs presented
413 higher virus excretion and delay in virus clearance, but milder lung lesions and clinical signs
414 compared with their heterologous counterparts (Rajao et al., 2017), in agreement with a role of the
415 adaptive immune response both in the control of the viremia and in the inflammatory pathology. In
416 short, pigs can be an interesting model of human influenza infection using endemic porcine or
417 human influenza viruses. The height of the influenza porcine model being to use pig-originated
418 pdmH1N1 (Mena et al., 2016) human pandemic virus in porcine infectious assay as a model of the
419 human infection (Schwaiger et al., 2019).

420 **-Coronavirus:** With the recent COVID-19 crisis there is new need for the development of relevant
421 animal models to study coronavirus infections. In pig, the only respiratory coronavirus (Porcine
422 Respiratory Coronavirus, PrCoV) is an *Alphacoronavirus* which is a variant of the Transmissible
423 Gastroenteritis Virus (TGEV) (Wang et al., 2019). Indeed, a large 5' deletion in the Spike gene of TGEV
424 altered the tropism and the virulence of PrCoV. PrCoV receptor is the aminopeptidase N (APN also
425 named CD13), mainly expressed on respiratory and intestinal epithelial cells, whereas human SARS-
426 CoV and SARS-CoV-2 bind to ACE2, a protein similarly expressed on respiratory and intestinal

427 epithelial cells but also on endothelial cells, which may have strong implication in virus pathogeny
428 (Saponaro et al., 2020). Indeed, engineering transgenic mice expressing the human ACE2 suffice to
429 trigger the full clinical picture of SARS-CoV-2 in mice (Bao et al., 2020; Israelow et al., 2020; Sun et al.,
430 2020). Regarding the first SARS-CoV, the pig has been demonstrated to be susceptible (Chen et al.,
431 2005). With the SARS-CoV-2, the situation is less clear and contradictory results have been reported
432 (Meekins et al., 2020; Pickering et al., 2020; ScloTTau et al., 2020; Shi et al., 2020). In 3 out of 4 of
433 these reports (Meekins et al., 2020; ScloTTau et al., 2020; Shi et al., 2020), pigs inoculated using nasal
434 or oral routes did not develop any lesions nor clinical signs. Moreover, authors did not detect any
435 viral excretion. However in one of these 3 studies (Meekins et al., 2020) SARS-CoV-2 replicated in
436 porcine epithelial cell lines. In the fourth study (Pickering et al., 2020), mild clinical signs were
437 observed in some inoculated pigs and viral RNA was detected in nasal lavage and oral fluid.
438 Furthermore, infectious virus was isolated from submandibular LN in one pig 13 days post-
439 inoculation and a serological response was evidenced in two animals 11 days post-inoculation
440 (Meekins et al., 2020). Thus, at this early stage, it is very difficult to definitely conclude about the
441 relevance of the pig model for the study of SARS-CoV-2.

442 The SARS-Cov-2 emergence revealed the lack of knowledge on seasonal human coronavirus.
443 The prototype viruses from the two main HCoV lineages are 229E (*Alphacoronavirus*, like the porcine
444 PrCOV) and OC43 (*Betacoronavirus*). They cause 15–29% of all common colds, and are the best
445 characterized HCoV, although 2 other human coronavirus are also considered as endemic (NL63 and
446 HKU1) (Su et al., 2016). In addition, HCOV-229E uses the same receptor as PrCOV. Thus swine might
447 be a good model of seasonal coronavirus infections allowing, in addition, the investigation of
448 coronavirus and IAV co-infections.

449 Regarding respiratory infections, it is worth to report here that pig lungs are suitable for the
450 generation of precision cut lung slices (PCLS) permitting the analysis, using multiple *in vitro*
451 conditions, of monoinfections (Delgado-Ortega et al., 2014) as well as of coinfecting viruses' complex
452 interactions on primary tissue-samples (Dobrescu et al., 2014; Saade et al., 2020).

453

454 Vaccination

455 Reviews arguing for the use of large animal models, including swine, have been published in
456 2015 that inventoried pig usage for vaccination and challenge tests against *Bordetella pertussis*,
457 *Chlamydia trachomatis*, human norovirus, rotavirus and IAV (Gerdtts et al., 2015; Rajao and Vincent,
458 2015). In addition to allow the evaluation of vaccines directly through protection against an
459 infectious challenge, animal models can permit to evaluate vaccines indirectly through immune
460 responses measurement without the final step of the infectious challenge. In this case, the animal
461 model is not anymore needed to be susceptible to the pathogen. For instance, swine are commonly
462 used to assess the potency of delivery devices and adjuvants.

463 **-Delivery:** The place and mode of delivery determine the localization of memory and effector cells
464 upon challenge. For instance it has been shown in mice that epicutaneous (EP) vaccination allows
465 mucosal immune response in mice (Belyakov et al., 2004). Pig skin is recognized as a good model of
466 human skin because of structure (hairiness, epidermis and dermis thickness, subcutaneous fat) and
467 lipid composition (Hammond et al., 2000) similarities, which makes pig skin permeability similar to
468 the human one, as well as because of dendritic cells and macrophages resemblance (Marquet et al.,
469 2014; Summerfield et al., 2015). Thus pig is a model of choice to test EP, transcutaneous (TC) and
470 intradermic (ID) vaccination protocols for the induction of respiratory immune protection. Direct
471 intramuscular (IM) DNA vaccination presented early encouraging success in mouse (Ulmer et al.,
472 1993) that were not reproduced in large animals including humans and pigs. Although not further
473 investigated, this discrepancy might rely on tissue stiffness differences according to the animal size.
474 Vaccination against respiratory diseases can also favor a pulmonary localized immune response by

475 directly exposing the target tissue to the antigens using aerosol (AS), intranasal (IN) or intratracheal
476 (IT) deliveries. For that purpose, the size of swine upper respiratory tract, more similar to the human
477 ones than small animals such as mouse or ferret, might guarantee a better compliance with clinical
478 situations (Kirschvink and Reinhold, 2008; Rogers et al., 2008). As an illustration, herein some
479 technics used in vaccination (mainly against influenza) in swine: prime-boost DNA/whole virus
480 vaccine delivered ID/IM (Hewitt et al., 2019), single ID Aujeszky's Disease Virus glycoproteins vaccine
481 (Le Luduec et al., 2016), influenza-proteins, DC-targeted ID or IM vaccination (Bernelin-Cottet et al.,
482 2016), protein, epicutaneous anti-RSV vaccination (Hervé et al., 2016), single-cycle influenza virus
483 delivered AS (Holzer et al., 2018), influenza virus-derived-replicon delivered IM (Ricklin et al., 2017),
484 nanoparticules whole influenza virus encapsulation delivered IN (Dhakal et al., 2017) and aerosol
485 intranasal delivery (Martini et al., 2020).

486 - **Adjuvant:** In addition to antigen and delivery, an essential part of a vaccine is the adjuvant used to
487 increase the immunogenicity of dead vaccines. One of the main developing arm of adjuvant is the
488 use of TLR-ligands, that trigger inflammation and antigen presenting cells activation. In that instance,
489 swine present a restrictive expression of TLR3 on pDC (Auray et al., 2016; Soldevila et al., 2018)
490 compared to the TLR3 expression on cDC1 in human (Poulin et al., 2010) and mouse (Desch et al.,
491 2011). Conversely, pig and human, but not mouse, expressed an active TLR8 receptor that can detect
492 RNA specifically from live bacteria (Ugolini et al., 2018), leading to a better activation of follicular DC
493 and the differentiation of high affinity antibodies. Finally, it has been proposed to use α GalSer as an
494 adjuvant in human vaccination that would harness iNKT cells (Speir et al., 2017) and help for CD8 T
495 cells activation. This strategy has been prospected in pig (for review (Yang et al., 2017b)). A first
496 assay using α GalSer as immunostimulant before IAV infection did not allow protection (Gu et al.,
497 2021), although this result does not preclude the use of α GalSer as true vaccine adjuvant.

498

499 Microbiota

500 In the last years the importance of the microbiota and especially the gut microbiota in the
501 preservation of homeostasis and health has been extensively studied. The gut microbiota can
502 influence the lung health and the susceptibility of porcine host to respiratory infections (Bassaganya-
503 Riera et al., 2003; Niederwerder, 2017). It is also known that all the mucosa in the body have some
504 connections forming the common mucosal immune system in man (Dang and Marsland, 2019) as in
505 pig (Wilson and Obradovic, 2020). Because of the use of pigs in biomedical research, some
506 comparisons between human and pig gut microbiota have been carried out. By deep metagenome
507 sequencing of faecal DNA from 287 pigs, Xiao and collaborators showed that among functional
508 pathways found in humans, 96% were present in pigs (Xiao et al., 2016). However, in other studies
509 comparing gut microbiota (Xiao et al., 2016) and fecal microbiota (Kobayashi et al., 2020) in different
510 species, it was shown that pig was less close to humans than marmosets and three shrews. Moreover
511 the pig, even if omnivorous like humans, had fecal microbiota showing some common features with
512 herbivores as the presence of *Fibrobacter*, a cellulolytic bacterium (Kobayashi et al., 2020). Recently a
513 further demonstration of the gut lung axis has been demonstrated in swine with an impact of a
514 porcine herpesvirus (Aujeszky's Disease Virus) on microbial community and immune status in the
515 ileum and colon of piglets (Zhang et al., 2019). Finally, the last decade firmly established the
516 existence and precisely described a symbiotic, stable respiratory microbiota in mouse and man (for
517 review see (Man et al., 2017). These investigations remain to be done in pig.

518

519 Respiratory Allergy and Asthma

520 Respiratory allergy corresponds to the improper activation of a Th2 response, leading to the
521 establishment of an IgE mediated anamnestic immune response that will trigger eosinophils

522 degranulation and asthma. Although food allergy is well-established in swine (for review see (Rupa et
523 al., 2009)), allergic asthma does not occur spontaneously in the animal world, albeit guinea pigs,
524 ferrets and pigs can be artificially sensitized. The establishment of a stable chronic porcine asthma
525 model appears to be difficult because the sensitivity to the antigen declines after repeated allergen
526 exposure (Szelenyi, 2000). However, upon *Ascaris suum* antigen sensitization consisting of an *A.*
527 *suum* extract in Al(OH)₃ followed by two booster doses and a challenge with nebulized allergen, pigs
528 developed airway inflammation associated with eosinophils and neutrophils infiltration (Fornhem et
529 al., 1996). On the induction side, cDC2 have been demonstrated to be the main trigger of the Th2
530 bias involved in the allergic response. As specified in the dendritic cells chapter above, porcine cDC2
531 resemble more to human than to mice cDC2 on several aspects in blood (Auray et al., 2016) as well
532 as in the respiratory tract (Crisci et al., 2020). For instance, the two main features important to recall
533 here are the expression by cDC2 of the FcεR1α, a receptor of IgE, and the intra-epithelial and sub-
534 epithelial cDC2 location in the trachea and the bronchia respectively (Maisonnasse et al., 2016; Yu et
535 al., 2013). Invariant NKT cells have been strongly involved in the Th2 bias of allergic responses (for
536 review see (Matangkasombut et al., 2009)), interestingly Renukaradhya *et al.* demonstrated that
537 intra-tracheal instillation of α-GalCer analog in pigs triggered iNKT cells activation and acute airway
538 hyperactivity associated with a Th2 cytokine profile (Renukaradhya et al., 2011). On the effector side,
539 dendrograms indicate that porcine IgE is more similar to IgE of carnivores, horses and humans than
540 to other artiodactyls IgE (Butler et al., 2009). In swine as in human, mast cell tryptase inhibitor
541 administrated prior to allergen challenge prevented acute bronchoconstrictive response and
542 decreased histamine release (Ploeg et al., 2002), while corticoid administration decreased airways
543 inflammatory cells infiltration (Fornhem et al., 1996). Still on the clinical side, porcine lung smooth
544 muscles reactions to inflammatory stimuli are a current model of asthma-muscle contraction (Ram-
545 Mohan et al., 2020; Sieck et al., 2019). Interestingly, airway contractions upon *in vivo* acid treatment
546 in piglets (Reznikov et al., 2019) recapitulated the airway hyper-responsiveness (AHR) gender bias
547 observed in infants (Van Merode et al., 2007).

548 Several intravenously administered compound such as nanomedicines, radiologic contrast agents
549 and other pharmaceutical products may trigger in human rare events of pseudoallergic infusion
550 reactions which are hypersensitivity reactions. Thanks to their high sensitivity, pigs have been used
551 for decades to detect such reactions, using the “complement activation-related pseudoallergy”
552 (CARPA) model (for review (Szebeni and Bawa, 2020)). Interestingly, pig high sensitivity seems
553 strongly related to the presence of constitutive PIM (Csukás et al., 2015), what might be a clue of the
554 presence of induced PIM in humans susceptible to such reactions.

555

556 Acute and Chronic inflammations

557 Acute lung injury (ALI) and its severe form acute respiratory distress syndrome (ARDS) are
558 responsible for more than 10% of intensive care unit admissions. ARDS conditions can be artificially
559 reproduced in swine by using lung repeated lavages, and oleic acid or endotoxin instillations (for
560 review see (Ballard-Croft et al., 2012)). Porcine models more related to human medical conditions
561 have been developed recently such as hemorrhagic shock (Morris et al., 2020), hyperoxia (Katalan et
562 al., 2017) or ricin induced-ARDS (Katalan et al., 2017). To our knowledge, no porcine model of ARDS
563 triggered upon infection-induced pneumonia has been reported. The early ARDS stage is
564 characterized by an acute inflammatory response that includes release of IL1, TNF, IL8 and
565 subsequent neutrophil recruitment (for review see (Ware, 2006)), that can be mimicked in pig
566 (Morris et al., 2020).

567 In developed countries, chronic obstructive disease (COPD) is mostly provoked by smoking
568 and result in the remodeling and narrowing of small airway associated with pulmonary emphysema,

569 likely due to chronic inflammation leading to increased numbers of neutrophils, macrophages, DC, T
570 and B lymphocytes. Chronic inflammation is maintained by macrophages and neutrophils infiltration
571 although autoimmunity related to autoantibodies and activated CD8 is suspected (for review see
572 (Caramori et al., 2016)). The emphysema appears to be linked to metalloproteinases (MMP) catalytic
573 activity. A mutated-pig phenotype has been described presenting emphysema associated with MMP9
574 and 12 hyper-expression which might be the first step toward the development of a COPD porcine
575 model (Bruun et al., 2013). To note, pigs are already used as model for cigarette smoke short (Gilman
576 et al., 1981) and middle (Gilman et al., 1981) term effects. The development of pig as a COPD model
577 would allow to explore an interesting hypothesis linking COPD and non-alcoholic fatty liver diseases
578 (Lonardo et al., 2017) that we would like to bring together with the potential pro-inflammatory PIM
579 induction in humans suffering of liver dysfunctions (Klingensmith et al., 1978, 1976). Whether the
580 constitutional presence of PIM in swine might favor or impaired this model remains to be
581 established.

582

583 Lung graft

584 Herein we will not tackle the field of xenotransplantation, that is beyond the scope of this
585 review and has been reviewed recently elsewhere (Burdorf et al., 2018). Regular lung transplantation
586 is the final treatment option for patients presenting end-stage lung diseases. The 1-year survival has
587 greatly improved in the last 40 years, reaching now 84% (Chambers et al., 2017). Pig is used as a
588 model of lung transplantation (Mariscal et al., 2018) to study primary graft dysfunctions and test new
589 preventive interventions to reduce or avoid these conditions (Iskender et al., 2016; Martins et al.,
590 2004). Interestingly, pig lungs are currently used to improve and further develop the promising *ex*
591 *vivo* lung perfusion (EVLV) method (for review (Tane et al., 2017)) leading to the reconditioning of
592 lung graft rejected in first instances because of not reaching the graft quality criteria. Porcine model
593 allowed to validate methods that decreased the expression of acute lung injury related genes
594 (Dromparis et al., 2019) or that increased the extracorporeal surviving time (Hozain et al., 2020).
595 Moreover, EVLP development give accessibility and time for intervention on the lung graft before
596 grafting. The pig model allows to test extracorporeal intervention such as IL10 gene-therapy in order
597 to act specifically on the immune tolerability of the transplant (MacHuca et al., 2017).

598

599 **Conclusions and perspectives:**

600 In conclusion, we think that the development of pig as a respiratory model of medical
601 conditions might be encouraged following three main research fields:

602 Because of the resemblances between porcine and human cutaneous, respiratory and
603 intestinal systems, associated to the immunological similarities of these species, pig appears to be a
604 model of choice to study the interactions between the different mucosae. In this perspective, a
605 better knowledge of porcine skin, lung and intestinal microbiota will be needed.

606 The second important point would be to develop pig as a model of respiratory allergies
607 according to the high similarity of cDC2 in pig and man, as well as to the good knowledge of porcine
608 neutrophils and iNKT cells. The main immunological hurdle for this development remains the lack of
609 reagents to follow the B cell response (B cells markers, such as CD19 and CD20 and tools to
610 discriminate the different IgG isotypes).

611 Finally, according to the large knowledge accumulated on the pig blood monocytes and
612 respiratory macrophages, as well as to the relative long life span of pigs, the long term consequences
613 of immune system alterations triggered by immunostimulants or primary infections, leading to
614 trained immunity (Angulo et al., 2020; Guillon et al., 2020; Stylianou et al., 2019; Yao et al., 2018) or
615 tolerance (Bouras et al., 2018; Didierlaurent et al., 2008) might be advantageously studied in this
616 model.

617 To summarize, thanks to the knowledge and tools developed during the last years, pig
618 appears now as the third medical model after mice and non-human primates, and might be seriously
619 considered when looking for a respiratory experimental model, especially in the fields of mucosal
620 interactions, allergies and trained immunity.

621

622 **Bibliography**

623 Aegerter, H., Kulikauskaite, J., Crotta, S., Patel, H., Kelly, G., Hessel, E.M., Mack, M., Beinke, S., Wack,
624 A., 2020. Influenza-induced monocyte-derived alveolar macrophages confer prolonged
625 antibacterial protection. *Nat. Immunol.* <https://doi.org/10.1038/s41590-019-0568-x>

626 Aldridge, J.R., Moseley, C.E., Boltz, D.A., Negovetich, N.J., Reynolds, C., Franks, J., Brown, S.A.,
627 Doherty, P.C., Webster, R.G., Thomas, P.G., 2009. TNF/iNOS-producing dendritic cells are the
628 necessary evil of lethal influenza virus infection. *Proc. Natl. Acad. Sci.* 106, 5306–5311.
629 <https://doi.org/10.1073/pnas.0900655106>

630 Angulo, M., Reyes-Becerril, M., Cepeda-Palacios, R., Angulo, C., 2020. Oral administration of
631 *Debaryomyces hansenii* CBS8339- β -glucan induces trained immunity in newborn goats. *Dev.*
632 *Comp. Immunol.* 105, 103597. <https://doi.org/10.1016/J.DCI.2019.103597>

633 Arnon, T.I., Achdout, H., Lieberman, N., Gazit, R., Gonen-Gross, T., Katz, G., Bar-Ilan, A., Bloushtain,
634 N., Lev, M., Joseph, A., Kedar, E., Porgador, A., Mandelboim, O., 2004. The mechanisms
635 controlling the recognition of tumor- and virus-infected cells by NKp46. *Blood* 103, 664–672.
636 <https://doi.org/10.1182/blood-2003-05-1716>

637 Artiaga, B.L., Whitener, R.L., Staples, C.R., Driver, J.P., 2014. Adjuvant effects of therapeutic
638 glycolipids administered to a cohort of NKT cell-diverse pigs. *Vet. Immunol. Immunopathol.* 162,
639 1–13. <https://doi.org/10.1016/j.vetimm.2014.09.006>

640 Auray, G., Keller, I., Python, S., Gerber, M., Bruggmann, R., Ruggli, N., Summerfield, A., 2016.
641 Characterization and Transcriptomic Analysis of Porcine Blood Conventional and Plasmacytoid
642 Dendritic Cells Reveals Striking Species-Specific Differences. *J. Immunol.* 197, 4791–4806.
643 <https://doi.org/10.4049/jimmunol.1600672>

644 Ballard-Croft, C., Wang, D., Sumpter, L.R., Zhou, X., Zwischenberger, J.B., 2012. Large-animal models
645 of acute respiratory distress syndrome. *Ann. Thorac. Surg.* 93, 1331–1339.
646 <https://doi.org/10.1016/j.athoracsur.2011.06.107>

647 Ballesteros-Tato, A., León, B., Lund, F.E., Randall, T.D., 2010. Temporal changes in dendritic cell
648 subsets, cross-priming and costimulation via CD70 control CD8+T cell responses to influenza.
649 *Nat. Immunol.* 11, 216–224. <https://doi.org/10.1038/ni.1838>

650 Bao, L., Deng, W., Huang, B., Gao, H., Liu, J., Ren, L., Wei, Q., Yu, P., Xu, Y., Qi, F., Qu, Y., Li, F., Lv, Q.,
651 Wang, W., Xue, J., Gong, S., Liu, M., Wang, G., Wang, S., Song, Z., Zhao, Linna, Liu, P., Zhao, Li,
652 Ye, F., Wang, H., Zhou, W., Zhu, N., Zhen, W., Yu, H., Zhang, X., Guo, L., Chen, L., Wang, C.,
653 Wang, Y., Wang, X., Xiao, Y., Sun, Q., Liu, H., Zhu, F., Ma, C., Yan, L., Yang, M., Han, J., Xu, W.,
654 Tan, W., Peng, X., Jin, Q., Wu, G., Qin, C., 2020. The pathogenicity of SARS-CoV-2 in hACE2
655 transgenic mice. *Nature* 583, 830–833. <https://doi.org/10.1038/s41586-020-2312-y>

656 Barbé, F., Atanasova, K., Van Reeth, K., 2011. Cytokines and acute phase proteins associated with
657 acute swine influenza infection in pigs. *Vet. J.* 187, 48–53.
658 <https://doi.org/10.1016/j.tvjl.2009.12.012>

659 Bassaganya-Riera, J., Pogramichniy, R.M., Jobgen, S.C., Halbur, P.G., Yoon, K.J., O’Shea, M., Mohede,
660 I., Hontecillas, R., 2003. Conjugated linoleic acid ameliorates viral infectivity in a pig model of
661 virally induced immunosuppression. *J. Nutr.* 133, 3204–3214.

- 662 <https://doi.org/10.1093/jn/133.10.3204>
- 663 Belyakov, I.M., Hammond, S.A., Ahlers, J.D., Glenn, G.M., Berzofsky, J.A., 2004. Transcutaneous
664 immunization induces mucosal CTLs and protective immunity by migration of primed skin
665 dendritic cells. *J. Clin. Invest.* 113, 998–1007. <https://doi.org/10.1172/JCI20261>
- 666 Bernelin-Cottet, C., Deloizy, C., Stanek, O., Barc, C., Bouguyon, E., Urien, C., Boulesteix, O., Pezant, J.,
667 Richard, C.-A., Moudjou, M., Da Costa, B., Jouneau, L., Chevalier, C., Leclerc, C., Sebo, P., Bertho,
668 N., Schwartz-Cornil, I., 2016. A universal influenza vaccine can lead to disease exacerbation or
669 viral control depending on delivery strategies. *Front. Immunol.* 7.
670 <https://doi.org/10.3389/fimmu.2016.00641>
- 671 Bernelin-Cottet, C., Urien, C., McCaffrey, J., Collins, D., Donadei, A., McDaid, D., Jakob, V., Barnier-
672 Quer, C., Collin, N., Bouguyon, E., Bordet, E., Barc, C., Boulesteix, O., Leplat, J.-J., Blanc, F.,
673 Contreras, V., Bertho, N., Moore, A., Schwartz-Cornil, I., 2019. Electroporation of a nanoparticle-
674 associated DNA vaccine induces higher inflammation and immunity compared to its delivery
675 with microneedle patches in pigs. *J. Control. Release.*
676 <https://doi.org/10.1016/J.JCONREL.2019.06.041>
- 677 Bigley, V., McGovern, N., Milne, P., Dickinson, R., Pagan, S., Cookson, S., Haniffa, M., Collin, M., 2015.
678 Langerin-expressing dendritic cells in human tissues are related to CD1c+ dendritic cells and
679 distinct from Langerhans cells and CD141high XCR1+ dendritic cells. *J. Leukoc. Biol.* 97, 627–
680 634. <https://doi.org/10.1189/jlb.1HI0714-351R>
- 681 Binns, R.M., Duncan, I.A., Powis, S.J., Hutchings, A., Butcher, G.W., 1992. Subsets of null and gamma
682 delta T-cell receptor+ T lymphocytes in the blood of young pigs identified by specific
683 monoclonal antibodies. *Immunology* 77, 219–27.
- 684 Bittmann, I., Dose, T., Baretton, G.B., Müller, C., Schwaiblmair, M., Kur, F., Löhres, U., 2001. Cellular
685 chimerism of the lung after transplantation: An interphase cytogenetic study. *Am. J. Clin.*
686 *Pathol.* 115, 525–533. <https://doi.org/10.1309/GAFN-5MPA-LY8E-DTPQ>
- 687 Blanc, F., Prévost-Blondel, A., Piton, G., Bouguyon, E., Leplat, J.-J., Andréoletti, F., Egidy, G., Bourneuf,
688 E., Bertho, N., Vincent-Naulleau, S., 2020. The Composition of Circulating Leukocytes Varies
689 With Age and Melanoma Onset in the MeLiM Pig Biomedical Model. *Front. Immunol.* 11, 291.
690 <https://doi.org/10.3389/fimmu.2020.00291>
- 691 Bolin, C.A., Whipple, D.L., Khanna, K. V., Risdahl, J.M., Peterson, P.K., Molitor, T.W., 1997. Infection of
692 swine with *Mycobacterium bovis* as a model of human tuberculosis. *J. Infect. Dis.* 176, 1559–
693 1566. <https://doi.org/10.1086/514155>
- 694 Bonduelle, O., Carrat, F., Luyt, C.E., Lepoint, C., Mosnier, A., Benhabiles, N., Krivine, A., Rozenberg, F.,
695 Yahia, N., Samri, A., Rousset, D., Van Der Werf, S., Autran, B., Combadiere, B., 2014.
696 Characterization of pandemic influenza immune memory signature after vaccination or
697 infection. *J. Clin. Invest.* 124, 3129–3136. <https://doi.org/10.1172/JCI74565>
- 698 Bordet, E., Frétaud, M., Crisci, E., Bouguyon, E., Rault, S., Pezant, J., Pleau, A., Renson, P., Giuffra, E.,
699 Larcher, T., Bourge, M., Bourry, O., Boulesteix, O., Langevin, C., Schwartz-Cornil, I., Bertho, N.,
700 2019. Macrophage-B Cell Interactions in the Inverted Porcine Lymph Node and Their Response
701 to Porcine Reproductive and Respiratory Syndrome Virus. *Front. Immunol.* 10, 953.
702 <https://doi.org/10.3389/fimmu.2019.00953>
- 703 Bordet, E., Maisonnasse, P., Renson, P., Bouguyon, E., Crisci, E., Tiret, M., Descamps, D., Bernelin-
704 Cottet, C., Urien, C., Lefèvre, F., Jouneau, L., Bourry, O., Leplat, J.-J., Schwartz-Cornil, I., Bertho,
705 N., 2018. Porcine alveolar macrophage-like cells are pro-inflammatory pulmonary intravascular
706 macrophages that produce large titers of porcine reproductive and respiratory syndrome virus.

707 Sci. Rep. 8. <https://doi.org/10.1038/s41598-018-28234-y>

708 Bosteels, C., Neyt, K., Vanheerswynghels, M., van Helden, M.J., Sichien, D., Debeuf, N., De Prijck, S.,
709 Bosteels, V., Vandamme, N., Martens, L., Saeys, Y., Louagie, E., Lesage, M., Williams, D.L., Tang,
710 S.C., Mayer, J.U., Ronchese, F., Scott, C.L., Hammad, H., Guilliams, M., Lambrecht, B.N., 2020.
711 Inflammatory Type 2 cDCs Acquire Features of cDC1s and Macrophages to Orchestrate
712 Immunity to Respiratory Virus Infection. *Immunity* 52, 1039-1056.e9.
713 <https://doi.org/10.1016/j.immuni.2020.04.005>

714 Bouras, M., Asehounne, K., Roquilly, A., 2018. Contribution of Dendritic Cell Responses to Sepsis-
715 Induced Immunosuppression and to Susceptibility to Secondary Pneumonia. *Front. Immunol.* 9,
716 2590. <https://doi.org/10.3389/fimmu.2018.02590>

717 Brain, J.D., Molina, R.M., DeCamp, M.M., Warner, A.E., 1999. Pulmonary intravascular macrophages:
718 their contribution to the mononuclear phagocyte system in 13 species. *Am. J. Physiol. Lung Cell.*
719 *Mol. Physiol.* 276, L146-154.

720 Bréa, D., Meurens, F., Dubois, A.V., Gaillard, J., Chevaleyre, C., Jourdan, M.-L., Winter, N., Arbeille, B.,
721 Si-Tahar, M., Gauthier, F., Attucci, S., 2012. The pig as a model for investigating the role of
722 neutrophil serine proteases in human inflammatory lung diseases. *Biochem. J.* 447, 363–370.
723 <https://doi.org/10.1042/BJ20120818>

724 Bruun, C.S., Jensen, L.K., Leifsson, P.S., Nielsen, J., Cirera, S., Jørgensen, C.B., Jensen, H.E., Fredholm,
725 M., 2013. Functional characterization of a porcine emphysema model. *Lung* 191, 669–675.
726 <https://doi.org/10.1007/s00408-013-9504-2>

727 Burdorf, L., Azimzadeh, A.M., Pierson, R.N., 2018. Progress and challenges in lung
728 xenotransplantation: an update. *Curr. Opin. Organ Transplant.* 23, 621–627.
729 <https://doi.org/10.1097/MOT.0000000000000582>

730 Butler, J.E., Zhao, Y., Sinkora, M., Wertz, N., Kacskovics, I., 2009. Immunoglobulins , antibody
731 repertoire and B cell development 33, 321–333. <https://doi.org/10.1016/j.dci.2008.06.015>

732 Byrne, A.J., Powell, J.E., O’Sullivan, B.J., Ogger, P.P., Hoffland, A., Cook, J., Bonner, K.L., Hewitt, R.J.,
733 Wolf, S., Ghai, P., Walker, S.A., Lukowski, S.W., Molyneaux, P.L., Saglani, S., Chambers, D.C.,
734 Maher, T.M., Lloyd, C.M., 2020. Dynamics of human monocytes and airway macrophages during
735 healthy aging and after transplant. *J. Exp. Med.* 217. <https://doi.org/10.1084/jem.20191236>

736 Cantu, E., Balsara, K.R., Li, B., Lau, C., Gibson, S., Wyse, A., Baig, K., Gaca, J., Gonzalez-Stawinski, G. V.,
737 Nichols, T., Parker, W., Davis, R.D., 2007. Prolonged function of macrophage, von Willebrand
738 factor-deficient porcine pulmonary xenografts. *Am. J. Transplant.* 7, 66–75.
739 <https://doi.org/10.1111/j.1600-6143.2006.01603.x>

740 Caramori, G., Casolari, P., Barczyk, A., Durham, A.L., Di Stefano, A., Adcock, I., 2016. COPD
741 immunopathology. *Semin. Immunopathol.* 38, 497–515. [https://doi.org/10.1007/s00281-016-](https://doi.org/10.1007/s00281-016-0561-5)
742 0561-5

743 Chakarov, S., Lim, H.Y., Tan, L., Lim, S.Y., See, P., Lum, J., Zhang, X.-M., Foo, S., Nakamizo, S., Duan, K.,
744 Kong, W.T., Gentek, R., Balachander, A., Carbajo, D., Bleriot, C., Malleret, B., Tam, J.K.C., Baig, S.,
745 Shabeer, M., Toh, S.-A.E.S., Schlitzer, A., Larbi, A., Marichal, T., Malissen, B., Chen, J., Poidinger,
746 M., Kabashima, K., Bajenoff, M., Ng, L.G., Angeli, V., Ginhoux, F., 2019. Two distinct interstitial
747 macrophage populations coexist across tissues in specific subtissular niches. *Science (80-)*. 363,
748 eaau0964. <https://doi.org/10.1126/SCIENCE.AAU0964>

749 Chambers, D.C., Yusen, R.D., Cherikh, W.S., Goldfarb, S.B., Kucheryavaya, A.Y., Khusch, K., Levvey,
750 B.J., Lund, L.H., Meiser, B., Rossano, J.W., Stehlik, J., 2017. The Registry of the International

- 751 Society for Heart and Lung Transplantation: Thirty-fourth Adult Lung And Heart-Lung
752 Transplantation Report—2017; Focus Theme: Allograft ischemic time. *J. Hear. Lung Transplant.*
753 36, 1047–1059. <https://doi.org/10.1016/j.healun.2017.07.016>
- 754 Chastagner, A., Bonin, E., Fablet, C., Quéguiner, S., Hirchaud, E., Lucas, P., Gorin, S., Barbier, N.,
755 Béven, V., Garin, E., Blanchard, Y., Rose, N., Hervé, S., Simon, G., 2019a. Virus persistence in pig
756 herds led to successive reassortment events between swine and human influenza A viruses,
757 resulting in the emergence of a novel triple-reassortant swine influenza virus. *Vet. Res.* 50, 1–9.
758 <https://doi.org/10.1186/s13567-019-0699-y>
- 759 Chastagner, A., Enouf, V., Peroz, D., Hervé, S., Lucas, P., Leneveu, P., Garin, E., Simon, G., 2019b.
760 Bidirectional Human–Swine Transmission of Seasonal Influenza A(H1N1)pdm09 Virus in Pig
761 Herd, France, 2018. *Emerg. Infect. Dis.* 25, 1940–1943. <https://doi.org/10.3201/eid2510.190068>
- 762 Chen, W., Yan, M., Yang, L., Ding, B., He, B., Wang, Y., Liu, X., Liu, C., Zhu, H., You, B., Huang, S.,
763 Zhang, J., Mu, F., Xiang, Z., Feng, X., Wen, J., Fang, J., Yu, J., Yang, H., Wang, J., 2005. SARS-
764 associated coronavirus transmitted from human to pig. *Emerg. Infect. Dis.* 11, 446–448.
765 <https://doi.org/10.3201/eid1103.040824>
- 766 Chevalayre, C., Riou, M., Bréa, D., Vandebrouck, C., Barc, C., Pezant, J., Melo, S., Olivier, M.,
767 Delaunay, R., Boulesteix, O., Berthon, P., Rossignol, C., Gaillard, J.B., Becq, F., Gauthier, F., Si-
768 Tahar, M., Meurens, F., Berri, M., Caballero-Posadas, I., Attucci, S., 2016. The pig: A relevant
769 model for evaluating the neutrophil serine protease activities during acute pseudomonas
770 aeruginosa lung infection. *PLoS One* 11, 1–17. <https://doi.org/10.1371/journal.pone.0168577>
- 771 Cho, H.J., Joo, N.S., Wine, J.J., 2011. Defective fluid secretion from submucosal glands of nasal
772 turbinates from CFTR -/- and CFTRΔ ΔF508/ΔF508 pigs. *PLoS One* 6.
773 <https://doi.org/10.1371/journal.pone.0024424>
- 774 Connelley, T.K., Longhi, C., Burrells, A., Degnan, K., Hope, J., Allan, A.J., Hammond, J.A., Storset, A.K.,
775 Morrison, W.I., 2014. NKp46 + CD3 + Cells: A Novel Nonconventional T Cell Subset in Cattle
776 Exhibiting Both NK Cell and T Cell Features . *J. Immunol.* 192, 3868–3880.
777 <https://doi.org/10.4049/jimmunol.1302464>
- 778 Crisci, E., Morolodo, M., Vu Manh, T.-P., Mohammad, A., Jourden, L., Urien, C., Bouguyon, E.,
779 Bordet, E., Bevilacqua, C., Bourge, M., Pezant, J., Pléau, A., Boulesteix, O., Schwartz, I., Bertho,
780 N., Giuffra, E., 2020. Distinctive Cellular and Metabolic Reprogramming in Porcine Lung
781 Mononuclear Phagocytes Infected With Type 1 PRRSV Strains. *Front. Immunol.* 11, 1–17.
782 <https://doi.org/10.3389/fimmu.2020.588411>
- 783 Crisci, E., Mussá, T., Fraile, L., Montoya, M., 2013. Review: Influenza virus in pigs. *Mol. Immunol.* 55,
784 200–211. <https://doi.org/10.1016/j.molimm.2013.02.008>
- 785 Csukás, D., Urbanics, R., Wéber, G., Rosivall, L., Szebeni, J., 2015. Pulmonary intravascular
786 macrophages: Prime suspects as cellular mediators of porcine CARPA. *Eur. J. Nanomedicine* 7,
787 27–36. <https://doi.org/10.1515/ejnm-2015-0008>
- 788 Dang, A.T., Marsland, B.J., 2019. Microbes, metabolites, and the gut–lung axis. *Mucosal Immunol.* 12,
789 843–850. <https://doi.org/10.1038/s41385-019-0160-6>
- 790 Dawson, H.D., Loveland, J.E., Pascal, G., Gilbert, J.G.R., Uenishi, H., Mann, K.M., Sang, Y., Zhang, J.,
791 Carvalho-Silva, D., Hunt, T., Hardy, M., Hu, Z., Zhao, S.H., Anselmo, A., Shinkai, H., Chen, C.,
792 Badaoui, B., Berman, D., Amid, C., Kay, M., Lloyd, D., Snow, C., Morozumi, T., Cheng, R.P.Y.,
793 Bystrom, M., Kapetanovic, R., Schwartz, J.C., Kataria, R., Astley, M., Fritz, E., Steward, C.,
794 Thomas, M., Wilming, L., Toki, D., Archibald, A.L., Bed’Hom, B., Beraldi, D., Huang, T.H., Ait-Ali,
795 T., Blecha, F., Botti, S., Freeman, T.C., Giuffra, E., Hume, D.A., Lunney, J.K., Murtaugh, M.P.,

796 Reecy, J.M., Harrow, J.L., Rogel-Gaillard, C., Tuggle, C.K., 2013. Structural and functional
797 annotation of the porcine immunome. *BMC Genomics* 14. [https://doi.org/10.1186/1471-2164-](https://doi.org/10.1186/1471-2164-14-332)
798 14-332

799 Dawson, H.D., Lunney, J.K., 2018. Porcine cluster of differentiation (CD) markers 2018 update. *Res.*
800 *Vet. Sci.* 118, 199–246. <https://doi.org/10.1016/J.RVSC.2018.02.007>

801 Dawson, H.D., Sang, Y., Lunney, J.K., 2020. Porcine cytokines, chemokines and growth factors: 2019
802 update. *Res. Vet. Sci.* 131, 266–300. <https://doi.org/10.1016/j.rvsc.2020.04.022>

803 Dehring, D.J., Crocker, S.H., Wismar, B.L., Steinberg, S.M., Lowery, B.D., Cloutier, C.T., 1983.
804 Comparison of live bacteria infusions in a porcine model of acute respiratory failure. *J. Surg.*
805 *Res.* 34, 151–158. [https://doi.org/10.1016/0022-4804\(83\)90054-9](https://doi.org/10.1016/0022-4804(83)90054-9)

806 Delgado-Ortega, M., Melo, S., Punyadarsaniya, D., Ramé, C., Olivier, M., Soubieux, D., Marc, D.,
807 Simon, G., Herrler, G., Berri, M., Dupont, J., Meurens, F., 2014. Innate immune response to a
808 H3N2 subtype swine influenza virus in newborn porcine trachea cells, alveolar macrophages,
809 and precision-cut lung slices. *Vet. Res.* 45, 42. <https://doi.org/10.1186/1297-9716-45-42>

810 Desch, A.N., Randolph, G.J., Murphy, K., Gautier, E.L., Kedl, R.M., Lahoud, M.H., Caminschi, I.,
811 Shortman, K., Henson, P.M., Jakubzick, C. V., 2011. CD103⁺ pulmonary dendritic cells
812 preferentially acquire and present apoptotic cell-associated antigen. *J. Exp. Med.* 208, 1789–
813 1797. <https://doi.org/10.1084/jem.20110538>

814 Dhakal, S., Hiremath, J., Bondra, K., Lakshmanappa, Y.S., Shyu, D.L., Ouyang, K., Kang, K. il,
815 Binjawadagi, B., Goodman, J., Tabynov, K., Krakowka, S., Narasimhan, B., Lee, C.W.,
816 Renukaradhya, G.J., 2017. Biodegradable nanoparticle delivery of inactivated swine influenza
817 virus vaccine provides heterologous cell-mediated immune response in pigs. *J. Control. Release*
818 247, 194–205. <https://doi.org/10.1016/j.jconrel.2016.12.039>

819 Didierlaurent, A., Goulding, J., Patel, S., Snelgrove, R., Low, L., Bebien, M., Lawrence, T., Van Rijjt, L.S.,
820 Lambrecht, B.N., Sirard, J.C., Hussell, T., 2008. Sustained desensitization to bacterial Toll-like
821 receptor ligands after resolution of respiratory influenza infection. *J. Exp. Med.* 205, 323–329.
822 <https://doi.org/10.1084/jem.20070891>

823 Dobrescu, I., Levast, B., Lai, K., Delgado-Ortega, M., Walker, S., Banman, S., Townsend, H., Simon, G.,
824 Zhou, Y., Gerds, V., Meurens, F., 2014. In vitro and ex vivo analyses of co-infections with swine
825 influenza and porcine reproductive and respiratory syndrome viruses. *Vet. Microbiol.* 169, 18–
826 32. <https://doi.org/10.1016/j.vetmic.2013.11.037>

827 Dromparis, P., Aboelnazar, N.S., Wagner, S., Himmat, S., White, C.W., Hatami, S., Luc, J.G.Y., Rotich,
828 S., Freed, D.H., Nagendran, J., Mengel, M., Adam, B.A., 2019. Ex vivo perfusion induces a time-
829 and perfusate-dependent molecular repair response in explanted porcine lungs. *Am. J.*
830 *Transplant.* 19, 1024–1036. <https://doi.org/10.1111/ajt.15123>

831 Edmans, M., McNee, A., Porter, E., Vatzia, E., Paudyal, B., Martini, V., Gubbins, S., Francis, O., Harley,
832 R., Thomas, A., Burt, R., Morgan, S., Fuller, A., Sewell, A., Charleston, B., Bailey, M., Tchilian, E.,
833 2021. Magnitude and Kinetics of T Cell and Antibody Responses During H1N1pdm09 Infection in
834 Inbred Babraham Pigs and Outbred Pigs. *Front. Immunol.* 11, 1–17.
835 <https://doi.org/10.3389/fimmu.2020.604913>

836 Eguíluz-Gracia, I., Schultz, H.H.L., Sikkeland, L.I.B., Danilova, E., Holm, A.M., Pronk, C.J.H., Agace,
837 W.W., Iversen, M., Andersen, C., Jahnsen, F.L., Baekkevold, E.S., 2016. Long-Term persistence of
838 human donor alveolar macrophages in lung transplant recipients. *Thorax* 71, 1006–1011.
839 <https://doi.org/10.1136/thoraxjnl-2016-208292>

- 840 Enkirch, T., von Messling, V., 2015. Ferret models of viral pathogenesis. *Virology* 479–480, 259–270.
841 <https://doi.org/10.1016/j.virol.2015.03.017>
- 842 Eurostat, 2020. Respiratory diseases statistics [WWW Document].
- 843 Evren, E., Ringqvist, E., Tripathi, K.P., Sleiers, N., Rives, I.C., Alisjahbana, A., Gao, Y., Sarhan, D., Halle,
844 T., Sorini, C., Lepzien, R., Marquardt, N., Michaëlsson, J., Smed-Sörensen, A., Botling, J.,
845 Karlsson, M.C.I., Villablanca, E.J., Willinger, T., 2021. Distinct developmental pathways from
846 blood monocytes generate human lung macrophage diversity. *Immunity* 54, 259-275.e7.
847 <https://doi.org/10.1016/j.immuni.2020.12.003>
- 848 Fairbairn, L., Kapetanovic, R., Beraldi, D., Sester, D.P., Tuggle, C.K., Archibald, A.L., Hume, D.A., 2013.
849 Comparative Analysis of Monocyte Subsets in the Pig. *J. Immunol.* 190, 6389–6396.
850 <https://doi.org/10.4049/jimmunol.1300365>
- 851 Fairbairn, L., Kapetanovic, R., Sester, D.P., Hume, D.A., 2011. The mononuclear phagocyte system of
852 the pig as a model for understanding human innate immunity and disease. *J. Leukoc. Biol.* 89,
853 855–871. <https://doi.org/10.1189/jlb.1110607>
- 854 FDA, 2015. Product Development Under the Animal Rule - Guidance for Industry, Food and Drug
855 Administration.
- 856 Forberg, H., Hauge, A.G., Valheim, M., Garcon, F., Nunez, A., Gerner, W., Mair, K.H., Graham, S.P.,
857 Brookes, S.M., Storset, A.K., 2014. Early responses of natural killer cells in pigs experimentally
858 infected with 2009 pandemic H1N1 influenza a virus. *PLoS One* 9.
859 <https://doi.org/10.1371/journal.pone.0100619>
- 860 Foreman-Wykert, A.K., Miller, J.F., 2005. A new animal model of Bordetella pertussis infection and
861 immunity. *Trends Microbiol.* 13, 559–560. <https://doi.org/10.1016/j.tim.2005.09.007>
- 862 Fornhem, C., Peterson, C.G.B., Dahlbackf, M., Scheyniusj, A., Alving, K., 1996. Granulocyte function in
863 the airways of allergen-challenged pigs : effects of inhaled and systemic budesonide 26, 1436–
864 1448.
- 865 Gao, H.-N., Lu, H.-Z., Cao, B., Du, B., Shang, H., Gan, J.-H., Lu, S.-H., Yang, Y.-D., Fang, Q., Shen, Y.-Z.,
866 Xi, X.-M., Gu, Q., Zhou, X.-M., Qu, H.-P., Yan, Z., Li, F.-M., Zhao, W., Gao, Z.-C., Wang, G.-F.,
867 Ruan, L.-X., Wang, W.-H., Ye, J., Cao, H.-F., Li, X.-W., Zhang, W.-H., Fang, X.-C., He, J., Liang, W.-
868 F., Xie, J., Zeng, M., Wu, X.-Z., Li, J., Xia, Q., Jin, Z.-C., Chen, Q., Tang, C., Zhang, Z.-Y., Hou, B.-M.,
869 Feng, Z.-X., Sheng, J.-F., Zhong, N.-S., Li, L.-J., 2013. Clinical Findings in 111 Cases of Influenza A
870 (H7N9) Virus Infection. *N. Engl. J. Med.* 368, 2277–2285.
871 <https://doi.org/10.1056/nejmoa1305584>
- 872 Gazit, R., Gruda, R., Elboim, M., Arnon, T.I., Katz, G., Achdout, H., Hanna, J., Qimron, U., Landau, G.,
873 Greenbaum, E., Zakay-Rones, Z., Porgador, A., Mandelboim, O., 2006. Lethal influenza infection
874 in the absence of the natural killer cell receptor gene Ncr1. *Nat. Immunol.* 7, 517–523.
875 <https://doi.org/10.1038/ni1322>
- 876 Gerdts, V., Wilson, H.L., Meurens, F., Van den Hurk, S. van D.L., Wilson, D., Walker, S., Wheler, C.,
877 Townsend, H., Potter, A.A., 2015. Large animal models for vaccine development and testing.
878 *ILAR J.* 56, 53–62. <https://doi.org/10.1093/ilar/ilv009>
- 879 Gerner, W., Käser, T., Saalmüller, A., 2009. Porcine T lymphocytes and NK cells - An update. *Dev.*
880 *Comp. Immunol.* 33, 310–320. <https://doi.org/10.1016/j.dci.2008.06.003>
- 881 Gerner, W., Talker, S.C., Koinig, H.C., Sedlak, C., Mair, K.H., Saalmüller, A., 2015. Phenotypic and
882 functional differentiation of porcine $\alpha\beta$ T cells: Current knowledge and available tools. *Mol.*
883 *Immunol.* 66, 3–13. <https://doi.org/10.1016/j.molimm.2014.10.025>

- 884 Gill, S.S., Suri, S.S., Janardhan, K.S., Caldwell, S., Duke, T., Singh, B., 2008. Role of pulmonary
885 intravascular macrophages in endotoxin-induced lung inflammation and mortality in a rat
886 model. *Respir. Res.* 9, 69. <https://doi.org/10.1186/1465-9921-9-69>
- 887 Gilman, M., Sylvester, J., Kennedy, T., Menkes, H., Traystman, R.J., 1981. Vascular effects of cigarette
888 smoke in isolated pig lungs. *Am Rev Respir Dis* 124, 549–553.
889 <https://doi.org/10.1164/arrd.1981.124.5.549>
- 890 Greer, A.M., Matthay, M.A., Kukreja, J., Bhakta, N.R., Nguyen, C.P., Wolters, P.J., Woodruff, P.G.,
891 Fahy, J. V., Shin, J.S., 2014. Accumulation of BDCA1+ dendritic cells in interstitial fibrotic lung
892 diseases and Th2-high asthma. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0099084>
- 893 Gu, W., Madrid, D.M.D., Yang, G., Artiaga, B.L., Loeb, J.C., Castleman, W.L., Richt, J.A., Lednicky, J.A.,
894 Driver, J.P., 2021. Unaltered influenza disease outcomes in swine prophylactically treated with
895 α -galactosylceramide. *Dev. Comp. Immunol.* 114, 103843.
896 <https://doi.org/10.1016/j.dci.2020.103843>
- 897 Guilliams, M., De Kleer, I., Henri, S., Post, S., Vanhoutte, L., De Prijck, S., Deswarte, K., Malissen, B.,
898 Hammad, H., Lambrecht, B.N., 2013. Alveolar macrophages develop from fetal monocytes that
899 differentiate into long-lived cells in the first week of life via GM-CSF. *J. Exp. Med.* 210, 1977–
900 1992. <https://doi.org/10.1084/jem.20131199>
- 901 Guilliams, M., Ginhoux, F., Jakubzick, C., Naik, S.H., Onai, N., Schraml, B.U., Segura, E., Tussiwand, R.,
902 Yona, S., 2014. Dendritic cells, monocytes and macrophages: A unified nomenclature based on
903 ontogeny. *Nat. Rev. Immunol.* 14, 571–578. <https://doi.org/10.1038/nri3712>
- 904 Guilliams, M., Scott, C.L., 2017. Does niche competition determine the origin of tissue-resident
905 macrophages? *Nat. Rev. Immunol.* 17, 451–460. <https://doi.org/10.1038/nri.2017.42>
- 906 Guillon, A., Arafa, E.I., Barker, K.A., Belkina, A.C., Martin, I., Shenoy, A.T., Wooten, A.K., De Ana, C.L.,
907 Dai, A., Labadorf, A., Escalante, J.H., Doms, H., Blasco, H., Traber, K.E., Jones, M.R., Quinton,
908 L.J., Mizgerd, J.P., 2020. Pneumonia recovery reprograms the alveolar macrophage pool. *JCI*
909 *Insight* 5. <https://doi.org/10.1172/jci.insight.133042>
- 910 Hammond, S.A., Tsonis, C., Sellins, K., Rushlow, K., Scharton-Kersten, T., Colditz, I., Glenn, G.M., 2000.
911 Transcutaneous immunization of domestic animals: Opportunities and challenges. *Adv. Drug*
912 *Deliv. Rev.* 43, 45–55. [https://doi.org/10.1016/S0169-409X\(00\)00076-4](https://doi.org/10.1016/S0169-409X(00)00076-4)
- 913 Hause, B.M., Padmanabhan, A., Pedersen, K., Gidlewski, T., 2016. Feral swine virome is dominated by
914 single-stranded DNA viruses and contains a novel Orthopneumovirus which circulates both in
915 feral and domestic swine. *J. Gen. Virol.* 97, 2090–2095. <https://doi.org/10.1099/jgv.0.000554>
- 916 Haverson, K., Bailey, M., Higgins, V.R., Bland, P.W., Stokes, C.R., 1994. Characterization of monoclonal
917 antibodies specific for monocytes, macrophages and granulocytes from porcine peripheral
918 blood and mucosal tissues. *J. Immunol. Methods* 170, 233–245. [https://doi.org/10.1016/0022-1759\(94\)90398-0](https://doi.org/10.1016/0022-1759(94)90398-0)
- 920 Haworth, S.G., Hislop, A.A., 1981. Adaptation of the pulmonary circulation to extra-uterine life in the
921 pig and its relevance to the human infant. *Cardiovasc. Res.* 15, 108–119.
922 <https://doi.org/10.1093/cvr/15.2.108>
- 923 Heinen, P., van Nieuwstadt, A., Pol, J., Boer-Luijze, E., van Oirschot, J., Bianchi, A., 2000. Systemic
924 and Mucosal Isotype-Specific Antibody. *Viral Immunol.* 13, 237–247.
- 925 Hervé, P.-L., Descamps, D., Deloizy, C., Dhelft, V., Laubreton, D., Bouguyon, E., Boukadiri, A.,
926 Dubuquoy, C., Larcher, T., Benhamou, P.-H., Eléouët, J.-F., Bertho, N., Mondoulet, L., Riffault, S.,
927 2016. Non-invasive epicutaneous vaccine against Respiratory Syncytial Virus: Preclinical proof of

928 concept. *J. Control. Release* 243. <https://doi.org/10.1016/j.jconrel.2016.10.003>

929 Hewitt, J.S., Karuppanan, A.K., Tan, S., Gauger, P., Halbur, P.G., Gerber, P.F., De Groot, A.S., Moise,
930 L., Opriessnig, T., 2019. A prime-boost concept using a T-cell epitope-driven DNA vaccine
931 followed by a whole virus vaccine effectively protected pigs in the pandemic H1N1 pig challenge
932 model. *Vaccine* 37, 4302–4309. <https://doi.org/10.1016/j.vaccine.2019.06.044>

933 Holtmeier, W., Geisel, W., Bernert, K., Butler, J.E., Šinkora, M., Řehákova, Z., Šinkora, J., Caspary,
934 W.F., 2004. Prenatal development of the porcine TCR δ repertoire: Dominant expression of an
935 invariant T cell receptor V δ 3-J δ 3 chain. *Eur. J. Immunol.* 34, 1941–1949.
936 <https://doi.org/10.1002/eji.200425055>

937 Holzer, B., Morgan, S.B., Matsuoka, Y., Edmans, M., Salguero, F.J., Everett, H., Brookes, S.M., Porter,
938 E., MacLoughlin, R., Charleston, B., Subbarao, K., Townsend, A., Tchilian, E., 2018. Comparison
939 of Heterosubtypic Protection in Ferrets and Pigs Induced by a Single-Cycle Influenza Vaccine. *J.*
940 *Immunol.* 200, 4068–4077. <https://doi.org/10.4049/jimmunol.1800142>

941 Horter, D.C., Yoon, K.-J., Zimmerman, J.J., 2003. A review of porcine tonsils in immunity and disease.
942 *Anim. Heal. Res. Rev.* 4, 143–155. <https://doi.org/10.1079/ahrr200358>

943 Hozain, A.E., Tipograf, Y., Pinezich, M.R., Cunningham, K.M., Donocoff, R., Queen, D., Fung, K., 2020.
944 Multiday maintenance of extracorporeal lungs using cross-circulation with conscious swine. *J.*
945 *Thorac. Cardiovasc. Surg.* 159, 1640–1653.

946 Humphray, S.J., Scott, C.E., Clark, R., Marron, B., Bender, C., Camm, N., Davis, J., Jenks, A., Noon, A.,
947 Patel, M., Sehra, H., Yang, F., Rogatcheva, M.B., Milan, D., Chardon, P., Rohrer, G., Nonneman,
948 D., de Jong, P., Meyers, S.N., Archibald, A., Beever, J.E., Schook, L.B., Rogers, J., 2007. A high
949 utility integrated map of the pig genome. *Genome Biol.* 8. [https://doi.org/10.1186/gb-2007-8-7-](https://doi.org/10.1186/gb-2007-8-7-r139)
950 [r139](https://doi.org/10.1186/gb-2007-8-7-r139)

951 Hussell, T., Bell, T.J., 2014. Alveolar macrophages: Plasticity in a tissue-specific context. *Nat. Rev.*
952 *Immunol.* 14, 81–93. <https://doi.org/10.1038/nri3600>

953 Iskender, I., Sakamoto, J., Nakajima, D., Lin, H., Chen, M., Kim, H., Guan, Z., Del Sorbo, L., Hwang, D.,
954 Waddell, T.K., Cypel, M., Keshavjee, S., Liu, M., 2016. Human α 1-antitrypsin improves early
955 post-transplant lung function: Pre-clinical studies in a pig lung transplant model. *J. Hear. Lung*
956 *Transplant.* 35, 913–921. <https://doi.org/10.1016/j.healun.2016.03.006>

957 Israelow, B., Song, E., Mao, T., Lu, P., Meir, A., Liu, F., Alfajaro, M.M., Wei, J., Dong, H., Homer, R.J.,
958 Ring, A., Wilen, C.B., Iwasaki, A., 2020. Mouse model of SARS-CoV-2 reveals inflammatory role
959 of type i interferon signaling. *J. Exp. Med.* 217. <https://doi.org/10.1084/JEM.20201241>

960 Jennings, J., Sang, Y., 2019. Porcine interferon complex and co-evolution with increasing viral
961 pressure after domestication. *Viruses* 11. <https://doi.org/10.3390/v11060555>

962 Jones, R., Baskerville, A., Reid, L., 1975. Histochemical Identification Of Glycoproteins In Pig Bronchial
963 Epithelium: (a) Normal And (b) Hypertrophied From Enzootic Pneumonia 116, 1–11.

964 Judge, E.P., Hughes, J.M.L., Egan, J.J., Maguire, M., Molloy, E.L., O’Dea, S., 2014. Anatomy and
965 bronchoscopy of the porcine lung: A model for translational respiratory medicine. *Am. J. Respir.*
966 *Cell Mol. Biol.* 51, 334–343. <https://doi.org/10.1165/rcmb.2013-0453TR>

967 Kalupov, T., Brillard-Bourdet, M., Dadé, S., Serrano, H., Wartelle, J., Guyot, N., Juliano, L., Moreau, T.,
968 Belaaouaj, A., Gauthier, F., 2009. Structural characterization of mouse neutrophil serine
969 proteases and identification of their substrate specificities: Relevance to mouse models of
970 human inflammatory diseases. *J. Biol. Chem.* 284, 34084–34091.
971 <https://doi.org/10.1074/jbc.M109.042903>

- 972 Käser, T., Renois, F., Wilson, H.L., Cnudde, T., Gerdts, V., Dillon, J.A.R., Jungersen, G., Agerholm, J.S.,
973 Meurens, F., 2018. Contribution of the swine model in the study of human sexually transmitted
974 infections. *Infect. Genet. Evol.* 66, 346–360. <https://doi.org/10.1016/j.meegid.2017.11.022>
- 975 Katalan, S., Falach, R., Rosner, A., Goldvaser, M., Brosh-Nissimov, T., Dvir, A., Mizrachi, A., Goren, O.,
976 Cohen, B., Gal, Y., Sapoznikov, A., Ehrlich, S., Sabo, T., Kronman, C., 2017. A novel swine model
977 of ricin-induced acute respiratory distress syndrome. *DMM Dis. Model. Mech.* 10, 173–183.
978 <https://doi.org/10.1242/dmm.027847>
- 979 Keller, I.E., Vosyka, O., Takenaka, S., Kloß, A., Dahlmann, B., Willems, L.I., Verdoes, M., Overkleeft,
980 H.S., Marcos, E., Adnot, S., Hauck, S.M., Ruppert, C., Günther, A., Herold, S., Ohno, S., Adler, H.,
981 Eickelberg, O., Meiners, S., 2015. Regulation of immunoproteasome function in the lung. *Sci.*
982 *Rep.* 5, 1–12. <https://doi.org/10.1038/srep10230>
- 983 Khatri, M., Dwivedi, V., Krakowka, S., Manickam, C., Ali, A., Wang, L., Qin, Z., Renukaradhya, G.J., Lee,
984 C.-W., 2010. Swine Influenza H1N1 Virus Induces Acute Inflammatory Immune Responses in Pig
985 Lungs: a Potential Animal Model for Human H1N1 Influenza Virus. *J. Virol.* 84, 11210–11218.
986 <https://doi.org/10.1128/JVI.01211-10>
- 987 Kim, H.M., Lee, Y.-W., Lee, K.-J., Kim, H.S., Cho, S.W., van Rooijen, N., Guan, Y., Seo, S.H., 2008.
988 Alveolar Macrophages Are Indispensable for Controlling Influenza Viruses in Lungs of Pigs. *J.*
989 *Virol.* 82, 4265–4274. <https://doi.org/10.1128/jvi.02602-07>
- 990 Kirschvink, N., Reinhold, P., 2008. Use of Alternative Animals as Asthma Models. *Curr. Drug Targets* 9,
991 470–484. <https://doi.org/10.2174/138945008784533525>
- 992 Klein, U., Dalla-Favera, R., 2008. Germinal centres: Role in B-cell physiology and malignancy. *Nat. Rev.*
993 *Immunol.* 8, 22–33. <https://doi.org/10.1038/nri2217>
- 994 Klingensmith, W.C., Tsan, M.-F., Wagner, H., 1976. Factors Affecting the Uptake of 99mTc-Sulfur
995 Colloid by the Lung and Kidney. *J. Nucl. Med.* 17, 681–685.
- 996 Klingensmith, W.C., Yang, S.L., Wagner, H.N., 1978. Lung uptake of Tc-99m sulfur colloid in liver and
997 spleen imaging. *J. Nucl. Med.* 19, 31–35.
- 998 Kobayashi, R., Nagaoka, K., Nishimura, N., Koike, S., Takahashi, E., Niimi, K., Murase, H., Kinjo, T.,
999 Tsukahara, T., Inoue, R., 2020. Comparison of the fecal microbiota of two monogastric
1000 herbivorous and five omnivorous mammals. *Anim. Sci. J.* 91, e13366.
1001 <https://doi.org/10.1111/asj.13366>
- 1002 Krejci, J., Nechvatalova, K., Blahutkova, M., Faldyna, M., 2013. The respiratory tract in pigs and its
1003 immune system: A review. *Vet. Med. (Praha).* 58, 206–220. [https://doi.org/10.17221/6759-](https://doi.org/10.17221/6759-VETMED)
1004 [VETMED](https://doi.org/10.17221/6759-VETMED)
- 1005 Kuebler, W.M., Goetz, A.E., 2002. The margined pool. *Eur. Surg. Res.* 34, 92–100.
1006 <https://doi.org/10.1159/000048894>
- 1007 Kuntz-Simon, G., Madec, F., 2009. Genetic and antigenic evolution of swine influenza viruses in
1008 Europe and evaluation of their zoonotic potential. *Zoonoses Public Health* 56, 310–325.
1009 <https://doi.org/10.1111/j.1863-2378.2009.01236.x>
- 1010 Lambrecht, B.N., Hammad, H., 2012. Lung Dendritic Cells in Respiratory Viral Infection and Asthma:
1011 From Protection to Immunopathology. *Annu. Rev. Immunol.* 30, 243–270.
1012 <https://doi.org/10.1146/annurev-immunol-020711-075021>
- 1013 Le Ludec, J.B., Debeer, S., Piras, F., Andréoni, C., Boudet, F., Laurent, P., Kaiserlian, D., Dubois, B.,
1014 2016. Intradermal vaccination with un-adjuvanted sub-unit vaccines triggers skin innate

- 1015 immunity and confers protective respiratory immunity in domestic swine. *Vaccine* 34, 914–922.
1016 <https://doi.org/10.1016/j.vaccine.2015.12.058>
- 1017 Lefèvre, F., Guillomot, M., D'Andréa, S., Battegay, S., La Bonnardière, C., 1998. Interferon-delta: The
1018 first member of a novel type I interferon family. *Biochimie* 80, 779–788.
1019 [https://doi.org/10.1016/S0300-9084\(99\)80030-3](https://doi.org/10.1016/S0300-9084(99)80030-3)
- 1020 Liebler-Tenorio, E., Pabst, R., 2006. MALT structure and function in farm animals. *Vet. Res.* 37, 257–
1021 280. <https://doi.org/10.1051/vetres:2006001>
- 1022 Liu, Q., Hu, W., Zhang, Y.L., Hu, S.P., Zhang, Z., He, X.J., Cai, X.H., 2018. Anti-viral immune response in
1023 the lung and thymus: Molecular characterization and expression analysis of
1024 immunoproteasome subunits LMP2, LMP7 and MECL-1 in pigs. *Biochem. Biophys. Res.*
1025 *Commun.* 502, 472–478. <https://doi.org/10.1016/j.bbrc.2018.05.190>
- 1026 Liu, Q., Zhang, Y.L., Hu, S.P., Ma, Z.L., Gao, S.L., Sun, B., Xiao, F., Zhang, Z., Cai, X.H., He, X.J., 2017.
1027 Expression of immunoproteasome subunits in the porcine lung: Alterations during normal and
1028 inflammatory conditions. *Vet. Microbiol.* 210, 134–141.
1029 <https://doi.org/10.1016/j.vetmic.2017.07.024>
- 1030 Lonardo, A., Nascimbeni, F., Ponz de Leon, M., 2017. Nonalcoholic fatty liver disease and COPD: is it
1031 time to cross the diaphragm? *Eur. Respir. J.* 49, 1700546.
1032 <https://doi.org/10.1183/13993003.00546-2017>
- 1033 Luna, C.M., Sibila, O., Agusti, C., Torres, A., 2009. Animal models of ventilator-associated pneumonia.
1034 *Eur. Respir. J.* 33, 182–188. <https://doi.org/10.1183/09031936.00046308>
- 1035 Lunney, J.K., 2007. Advances in swine biomedical model genomics. *Int. J. Biol. Sci.* 3, 179–184.
1036 <https://doi.org/10.7150/ijbs.3.179>
- 1037 Ma, W., Kahn, R., Richt, J., 2009. The pig as a mixing vessel for influenza viruses: Human and
1038 veterinary implications. *J Mol Genet Med* 3, 158–166. <https://doi.org/10.1038/nchina.2008.185>
- 1039 MacHuca, T.N., Cypel, M., Bonato, R., Yeung, J.C., Chun, Y.M., Juvet, S., Guan, Z., Hwang, D.M., Chen,
1040 M., Saito, T., Harmantas, C., Davidson, B.L., Waddell, T.K., Liu, M., Keshavjee, S., 2017. Safety
1041 and Efficacy of Ex Vivo Donor Lung Adenoviral IL-10 Gene Therapy in a Large Animal Lung
1042 Transplant Survival Model. *Hum. Gene Ther.* 28, 757–765.
1043 <https://doi.org/10.1089/hum.2016.070>
- 1044 Mair, K.H., Essler, S.E., Patzl, M., Storset, A.K., Saalmüller, A., Gerner, W., 2012. NKp46 expression
1045 discriminates porcine NK cells with different functional properties. *Eur. J. Immunol.* 42, 1261–
1046 1271. <https://doi.org/10.1002/eji.201141989>
- 1047 Mair, K.H., Sedlak, C., Käser, T., Pasternak, A., Levast, B., Gerner, W., Saalmüller, A., Summerfield, A.,
1048 Gerdts, V., Wilson, H.L., Meurens, F., 2014. The porcine innate immune system: An update. *Dev.*
1049 *Comp. Immunol.* 45, 321–343. <https://doi.org/10.1016/J.DCI.2014.03.022>
- 1050 Mair, K.H., Stadler, M., Talker, S.C., Forberg, H., Storset, A.K., Müllebner, A., Catharina Duvigneau, J.,
1051 Hammer, S.E., Saalmüller, A., Gerner, W., 2016. Porcine CD3+NKp46+lymphocytes have NK-cell
1052 characteristics and are present in increased frequencies in the lungs of influenza-infected
1053 animals. *Front. Immunol.* 7, 1–17. <https://doi.org/10.3389/fimmu.2016.00263>
- 1054 Maisonnasse, P., Bouguyon, E., Piton, G., Ezquerra, A., Urien, C., Deloizy, C., Bourge, M., Leplat, J.J.,
1055 Simon, G., Chevalier, C., Vincent-Naulleau, S., Crisci, E., Montoya, M., Schwartz-Cornil, I.,
1056 Bertho, N., 2016. The respiratory DC/macrophage network at steady-state and upon influenza
1057 infection in the swine biomedical model. *Mucosal Immunol.* 9, 835–849.
1058 <https://doi.org/10.1038/mi.2015.105>

- 1059 Man, W.H., De Steenhuijsen Piters, W.A.A., Bogaert, D., 2017. The microbiota of the respiratory tract:
1060 Gatekeeper to respiratory health. *Nat. Rev. Microbiol.* 15, 259–270.
1061 <https://doi.org/10.1038/nrmicro.2017.14>
- 1062 Mariscal, A., Caldarone, L., Tikkanen, J., Nakajima, D., Chen, M., Yeung, J., Cypel, M., Liu, M.,
1063 Keshavjee, S., 2018. Pig lung transplant survival model. *Nat. Protoc.* 13, 1814–1828.
1064 <https://doi.org/10.1038/s41596-018-0019-4>
- 1065 Marquet, F., Vu Manh, T.-P., Maisonnasse, P., Elhmouzi-Younes, J., Urien, C., Bouguyon, E., Jouneau,
1066 L., Bourge, M., Simon, G., Ezquerra, A., Lecardonnel, J., Bonneau, M., Dalod, M., Schwartz-
1067 Cornil, I., Bertho, N., Bertho, J., 2014. Pig Skin Includes Dendritic Cell Subsets Transcriptomically
1068 Related to Human CD1a and CD14 Dendritic Cells Presenting Different Migrating Behaviors and
1069 T Cell Activation Capacities. *J Immunol* 193, 5883–5893.
1070 <https://doi.org/10.4049/jimmunol.1303150>
- 1071 Martínez-Olondris, P., Sibila, O., Agustí, C., Rigol, M., Soy, D., Esquinas, C., Piñer, R., Luque, N.,
1072 Guerrero, L., Quera, M.Á., Marco, F., De La Bellacasa, J.P., Ramirez, J., Torres, A., 2010. An
1073 experimental model of pneumonia induced by methicillin-resistant *Staphylococcus aureus* in
1074 ventilated piglets. *Eur. Respir. J.* 36, 901–906. <https://doi.org/10.1183/09031936.00176709>
- 1075 Martini, V., Hinchcliffe, M., Blackshaw, E., Joyce, M., McNee, A., Beverley, P., Townsend, A.,
1076 MacLoughlin, R., Tchilian, E., 2020. Distribution of Droplets and Immune Responses After
1077 Aerosol and Intra-Nasal Delivery of Influenza Virus to the Respiratory Tract of Pigs. *Front.*
1078 *Immunol.* 11, 1–12. <https://doi.org/10.3389/fimmu.2020.594470>
- 1079 Martins, S., de Perrot, M., Imai, Y., Yamane, M., Quadri, S.M., Segall, L., Dutly, A., Sakiyama, S.,
1080 Chaparro, A., Davidson, B.L., Waddell, T.K., Liu, M., Keshavjee, S., 2004. Transbronchial
1081 administration of adenoviral-mediated interleukin-10 gene to the donor improves function in a
1082 pig lung transplant model. *Gene Ther.* 11, 1786–1796. <https://doi.org/10.1038/sj.gt.3302357>
- 1083 Matangkasombut, P., Pichavant, M., DeKruyff, R.H., Umetsu, D.T., 2009. Natural killer T cells and the
1084 regulation of asthma. *Mucosal Immunol.* 2, 383–392. <https://doi.org/10.1038/mi.2009.96>
- 1085 Mcfarlin, D.E., Binns, R.M., W, M.H.N., 1973. Lymph node function and lymphocyte circulation in the
1086 pig. *Adv Exp Med Biol* 29, 87–93.
- 1087 McLaughlin, R.F., Tyler, W.S., Canada, R.O., 1961. A study of the subgross pulmonary anatomy in
1088 various mammals. *Am. J. Anat.* 108, 149–165. <https://doi.org/10.1002/aja.1001080203>
- 1089 Meekins, D.A., Morozov, I., Trujillo, J.D., Gaudreault, N.N., Bold, D., Carossino, M., Artiaga, B.L.,
1090 Indran, S. V., Kwon, T., Balaraman, V., Madden, D.W., Feldmann, H., Henningson, J., Ma, W.,
1091 Balasuriya, U.B.R., Richt, J.A., 2020. Susceptibility of swine cells and domestic pigs to SARS-CoV-
1092 2. *Emerg. Microbes Infect.* 9, 2278–2288. <https://doi.org/10.1080/22221751.2020.1831405>
- 1093 Melzi, E., Caporale, M., Rocchi, M., Martín, V., Gamino, V., di Provvido, A., Marruchella, G., Entrican,
1094 G., Sevilla, N., Palmarini, M., 2016. Follicular dendritic cell disruption as a novel mechanism of
1095 virus-induced immunosuppression. *Proc. Natl. Acad. Sci.* 113, E6238–E6247.
1096 <https://doi.org/10.1073/pnas.1610012113>
- 1097 Mena, I., Nelson, M.I., Quezada-Monroy, F., Dutta, J., Cortes-Fernández, R., Lara-Puente, J.H., Castro-
1098 Peralta, F., Cunha, L.F., Trovão, N.S., Lozano-Dubernard, B., Rambaut, A., Van Bakel, H., García-
1099 Sastre, A., 2016. Origins of the 2009 H1N1 influenza pandemic in swine in Mexico. *Elife* 5, 1–21.
1100 <https://doi.org/10.7554/eLife.16777>
- 1101 Meurens, F., Summerfield, A., Nauwynck, H., Saif, L., Gerdts, V., 2012. The pig: A model for human
1102 infectious diseases. *Trends Microbiol.* 20, 50–57. <https://doi.org/10.1016/j.tim.2011.11.002>

- 1103 Mills, A.N., Lopez-Vidriero, M.T., Haworth, S.G., 1986. Development of the airway epithelium and
 1104 submucosal glands in the pig lung: Changes in epithelial glycoprotein profiles. *Br. J. Exp. Pathol.*
 1105 67, 821–829.
- 1106 Moreno, S., Alvarez, B., Poderoso, T., Revilla, C., Ezquerra, A., Alonso, F., Dominguez, J., 2010. Porcine
 1107 monocyte subsets differ in the expression of CCR2 and in their responsiveness to CCL2. *Vet. Res.*
 1108 41. <https://doi.org/10.1051/vetres/2010048>
- 1109 Morris, M.C., Kim, Y., Blakeman, T.C., Stevens-Topie, S., Jung, A.D., Cox, D.B., Robinson, B.B.R., Pritts,
 1110 T.A., Makley, A.T., Goodman, M.D., 2020. Early Identification of Acute Lung Injury in a Porcine
 1111 Model of Hemorrhagic Shock. *J. Surg. Res.* 247, 453–460.
 1112 <https://doi.org/10.1016/j.jss.2019.09.060>
- 1113 Moshous, D., Callebaut, I., De Chasseval, R., Corneo, B., Cavazzana-Calvo, M., Le Deist, F., Tezcan, I.,
 1114 Sanal, O., Bertrand, Y., Philippe, N., Fischer, A., De Villartay, J.P., 2001. Artemis, a novel DNA
 1115 double-strand break repair/V(D)J recombination protein, is mutated in human severe combined
 1116 immune deficiency. *Cell* 105, 177–186. [https://doi.org/10.1016/S0092-8674\(01\)00309-9](https://doi.org/10.1016/S0092-8674(01)00309-9)
- 1117 Murtaugh, M.P., Johnson, C.R., Xiao, Z., Scamurra, R.W., Zhou, Y., 2009. Species specialization in
 1118 cytokine biology: Is interleukin-4 central to the TH1-TH2 paradigm in swine? *Dev. Comp.*
 1119 *Immunol.* 33, 344–352. <https://doi.org/10.1016/j.dci.2008.06.014>
- 1120 Nayak, D.K., Zhou, F., Xu, M., Huang, J., Tsuji, M., Hachem, R., Mohanakumar, T., 2016. Long-Term
 1121 Persistence of Donor Alveolar Macrophages in Human Lung Transplant Recipients That
 1122 Influences Donor-Specific Immune Responses. *Am. J. Transplant.* 16, 2300–2311.
 1123 <https://doi.org/10.1111/ajt.13819>
- 1124 Ng, S.L., Teo, Y.J., Setiagani, Y.A., Karjalainen, K., Ruedl, C., 2018. Type 1 Conventional CD103+
 1125 Dendritic Cells Control Effector CD8+ T Cell Migration, Survival, and Memory Responses During
 1126 Influenza Infection. *Front. Immunol.* 9, 3043. <https://doi.org/10.3389/fimmu.2018.03043>
- 1127 Niederwerder, M.C., 2017. Role of the microbiome in swine respiratory disease. *Vet. Microbiol.* 209,
 1128 97–106. <https://doi.org/10.1016/j.vetmic.2017.02.017>
- 1129 Pabst, R., Binns, R.M., 1995. Lymphocytes migrate from the bronchoalveolar space to regional
 1130 bronchial lymph nodes. *Am. J. Respir. Crit. Care Med.* 151, 495–499.
 1131 <https://doi.org/10.1164/ajrccm.151.2.7842212>
- 1132 Parker, C.M., Cepek, K.L., Russell, G.J., Shaw, S.K., Posnett, D.N., Schwarting, R., Brenner, M.B., 1992.
 1133 A family of $\beta 7$ integrins on human mucosal lymphocytes. *Proc. Natl. Acad. Sci. U. S. A.* 89, 1924–
 1134 1928. <https://doi.org/10.1073/pnas.89.5.1924>
- 1135 Phan, T.G., Green, J.A., Xu, Y., Cyster, J.G., 2009. Immune complex relay by subcapsular sinus
 1136 macrophages and noncognate B cells drives antibody affinity maturation. *Nat. Immunol.* 10,
 1137 786–796. <https://doi.org/10.1038/ni.1745>
- 1138 Picarda, G., Chéneau, C., Humbert, J.-M., Bériou, G., Pilet, P., Martin, J., Duteille, F., Perrot, P., Bellier-
 1139 Waast, F., Heslan, M., Haspot, F., Guillon, F., Josien, R., Halary, F.A., 2016. Functional Langerin
 1140 high -Expressing Langerhans-like Cells Can Arise from CD14 high CD16 – Human Blood
 1141 Monocytes in Serum-Free Condition . *J. Immunol.* 196, 3716–3728.
 1142 <https://doi.org/10.4049/jimmunol.1501304>
- 1143 Pickering, B., Smith, S., Pinette, M., Embury-Hyatt, C., Moffat, E., Marszal, P., Lewis, C., 2020.
 1144 Susceptibility of domestic swine to experimental infection with SARS-CoV-2 2. *bioRxiv.*
- 1145 Platzer, B., Baker, K., Vera, M.P., Singer, K., Panduro, M., Lexmond, W.S., Turner, D., Vargas, S.O.,
 1146 Kinet, J.P., Maurer, D., Baron, R.M., Blumberg, R.S., Fiebiger, E., 2015. Dendritic cell-bound IgE

- 1147 functions to restrain allergic inflammation at mucosal sites. *Mucosal Immunol.* 8, 516–532.
1148 <https://doi.org/10.1038/mi.2014.85>
- 1149 Ploeg, I. Van Der, Alving, K., Sylvén, H., Dahlbäck, M., 2002. The tryptase inhibitor APC-366 reduces the
1150 acute airway response to allergen in pigs sensitized to *Ascaris suum* 967–971.
- 1151 Poulin, L.F., Salio, M., Griessinger, E., Anjos-Afonso, F., Craciun, L., Chen, J.-L., Keller, A.M., Joffre, O.,
1152 Zelenay, S., Nye, E., Le Moine, A., Faure, F., Donckier, V., Sancho, D., Cerundolo, V., Bonnet, D.,
1153 Reis e Sousa, C., 2010. Characterization of human DNCR-1⁺ BDCA3⁺ leukocytes as putative
1154 equivalents of mouse CD8α⁺ dendritic cells. *J. Exp. Med.* 207, 1261–1271.
1155 <https://doi.org/10.1084/jem.20092618>
- 1156 Powell, E.J., Cunnick, J.E., Knetter, S.M., Loving, C.L., Waide, E.H., Dekkers, J.C.M., Tuggle, C.K., 2016.
1157 NK cells are intrinsically functional in pigs with Severe Combined Immunodeficiency (SCID)
1158 caused by spontaneous mutations in the Artemis gene. *Vet. Immunol. Immunopathol.* 175, 1–6.
1159 <https://doi.org/10.1016/j.vetimm.2016.04.008>
- 1160 Rajao, D.S., Loving, C.L., Waide, E.H., Gauger, P.C., Dekkers, J.C.M., Tuggle, C.K., Vincent, A.L., 2017.
1161 Pigs with Severe Combined Immunodeficiency Are Impaired in Controlling Influenza A Virus
1162 Infection. *J. Innate Immun.* 9, 193–202. <https://doi.org/10.1159/000451007>
- 1163 Rajao, D.S., Vincent, A.L., 2015. Swine as a model for influenza avirus infection and immunity. *ILAR J.*
1164 56, 44–52. <https://doi.org/10.1093/ilar/ilv002>
- 1165 Ram-Mohan, S., Bai, Y., Schaible, N., Ehrlicher, A.J., Cook, D.P., Suki, X.B., Stoltz, D.A., Solway, J., Ai,
1166 X., Krishnan, R., Krishnan, R., 2020. Tissue traction microscopy to quantify muscle contraction
1167 within precision-cut lung slices. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* 318, L323–L330.
1168 <https://doi.org/10.1152/AJPLUNG.00297.2019>
- 1169 Ramos, L., Obregon-Henao, A., Henao-Tamayo, M., Bowen, R., Lunney, J.K., Gonzalez-Juarrero, M.,
1170 2017. The minipig as an animal model to study *Mycobacterium tuberculosis* infection and
1171 natural transmission. *Tuberculosis* 106, 91–98. <https://doi.org/10.1016/j.tube.2017.07.003>
- 1172 Rendas, A., Branthwaite, M., Reid, L., 1978. Growth of pulmonary circulation in normal pig: Structural
1173 analysis and cardiopulmonary function. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 45, 806–
1174 817. <https://doi.org/10.1152/jappl.1978.45.5.806>
- 1175 Renukaradhya, G.J., Manickam, C., Khatri, M., Rauf, A., Li, X., Tsuji, M., Rajashekara, G., Dwivedi, V.,
1176 2011. Functional invariant NKT cells in pig lungs regulate the airway hyperreactivity: A potential
1177 animal model. *J. Clin. Immunol.* 31, 228–239. <https://doi.org/10.1007/s10875-010-9476-4>
- 1178 Reutner, K., Leitner, J., Müllebner, A., Ladinig, A., Essler, S.E., Duvigneau, J.C., Ritzmann, M.,
1179 Steinberger, P., Saalmüller, A., Gerner, W., 2013. CD27 expression discriminates porcine T
1180 helper cells with functionally distinct properties. *Vet. Res.* 44, 1–14.
1181 <https://doi.org/10.1186/1297-9716-44-18>
- 1182 Reznikov, L.R., Liao, Y.S.J., Gu, T., Davis, K.M., Kuan, S.P., Atanasova, K.R., Dadural, J.S., Collins, E.N.,
1183 Guevara, M. V., Vogt, K., 2019. Sex-specific airway hyperreactivity and sex-specific
1184 transcriptome remodeling in neonatal piglets challenged with intra-airway acid. *Am. J. Physiol. -
1185 Lung Cell. Mol. Physiol.* 316, L131–L143. <https://doi.org/10.1152/ajplung.00417.2018>
- 1186 Ricklin, M.E., Python, S., Vielle, N.J., Brechbühl, D., Zumkehr, B., Posthaus, H., Zimmer, G., Ruggli, N.,
1187 Summerfield, A., 2017. Virus replicon particle vaccines expressing nucleoprotein of influenza A
1188 virus mediate enhanced inflammatory responses in pigs. *Sci. Rep.* 7, 1–12.
1189 <https://doi.org/10.1038/s41598-017-16419-w>
- 1190 Rodríguez-Gómez, I.M., Talker, S.C., Käser, T., Stadler, M., Hammer, S.E., Saalmüller, A., Gerner, W.,

- 1191 2016. Expression of T-bet, Eomesodermin and GATA-3 in porcine $\alpha\beta$ T cells. *Dev. Comp.*
1192 *Immunol.* 60, 115–126. <https://doi.org/10.1016/j.dci.2016.02.022>
- 1193 Rodríguez-Gómez, I.M., Talker, S.C., Käser, T., Stadler, M., Reiter, L., Ladinig, A., Milburn, J. V.,
1194 Hammer, S.E., Mair, K.H., Saalmüller, A., Gerner, W., 2019. Expression of T-bet, eomesodermin,
1195 and GATA-3 correlates with distinct phenotypes and functional properties in porcine $\gamma\delta$ T cells.
1196 *Front. Immunol.* 10. <https://doi.org/10.3389/fimmu.2019.00396>
- 1197 Rogers, C.S., Abraham, W.M., Brogden, K.A., Engelhardt, J.F., Fisher, J.T., McCray, P.B., McLennan, G.,
1198 Meyerholz, D.K., Namati, E., Ostedgaard, L.S., Prather, R.S., Sabater, J.R., Stoltz, D.A., Zabner, J.,
1199 Welsh, M.J., 2008. The porcine lung as a potential model for cystic fibrosis. *Am. J. Physiol. - Lung*
1200 *Cell. Mol. Physiol.* 295, 240–263. <https://doi.org/10.1152/ajplung.90203.2008>
- 1201 Rupa, P., Schmied, J., Wilkie, B.N., 2009. Porcine allergy and IgE. *Vet. Immunol. Immunopathol.* 132,
1202 41–45. <https://doi.org/10.1016/j.vetimm.2009.09.013>
- 1203 Saade, G., Ménard, D., Hervet, C., Renson, P., Hue, E., Zhu, J., Dubreil, L., Paillot, R., Pronost, S.,
1204 Bourry, O., Simon, G., Dupont, J., Bertho, N., Meurens, F., 2020. Porcine reproductive and
1205 respiratory syndrome virus interferes with swine influenza a virus infection of epithelial cells.
1206 *Vaccines* 8, 1–22. <https://doi.org/10.3390/vaccines8030508>
- 1207 Saponaro, F., Rutigliano, G., Sestito, S., Bandini, L., Storti, B., Bizzarri, R., Zucchi, R., 2020. ACE2 in the
1208 Era of SARS-CoV-2: Controversies and Novel Perspectives. *Front. Mol. Biosci.* 7, 1–25.
1209 <https://doi.org/10.3389/fmolb.2020.588618>
- 1210 Sasaki, K., Pabst, R., Rothkötter, H.J., 1994. The unique ultrastructure of high-endothelial venules in
1211 inguinal lymph nodes of the pig. *Cell Tissue Res.* 276, 85–90.
1212 <https://doi.org/10.1007/BF00354787>
- 1213 Sassu, E.L., Ladinig, A., Talker, S.C., Stadler, M., Knecht, C., Stein, H., Frömbling, J., Richter, B.,
1214 Spergser, J., Ehling-Schulz, M., Graage, R., Hennig-Pauka, I., Gerner, W., 2017. Frequency of
1215 Th17 cells correlates with the presence of lung lesions in pigs chronically infected with
1216 *Actinobacillus pleuropneumoniae*. *Vet. Res.* 48, 1–16. [https://doi.org/10.1186/s13567-017-](https://doi.org/10.1186/s13567-017-0411-z)
1217 [0411-z](https://doi.org/10.1186/s13567-017-0411-z)
- 1218 Schäfer, A., Hühr, J., Schwaiger, T., Dorhoi, A., Mettenleiter, T.C., Blome, S., Schröder, C., Blohm, U.,
1219 2019. Porcine Invariant Natural Killer T Cells : Functional Profiling and Dynamics in Steady State
1220 and Viral Infections 10, 1–19. <https://doi.org/10.3389/fimmu.2019.01380>
- 1221 Schlitzer, A., Ginhoux, F., 2014. Organization of the mouse and human DC network. *Curr. Opin.*
1222 *Immunol.* 26, 90–99. <https://doi.org/10.1016/j.coi.2013.11.002>
- 1223 Schneberger, D., Aharonson-Raz, K., Singh, B., 2012. Pulmonary intravascular macrophages and lung
1224 health: What are we missing? *AJP Lung Cell. Mol. Physiol.* 302, L498–L503.
1225 <https://doi.org/10.1152/ajplung.00322.2011>
- 1226 Schneider, C., Nobs, S.P., Heer, A.K., Kurrer, M., Klinke, G., van Rooijen, N., Vogel, J., Kopf, M., 2014.
1227 Alveolar Macrophages Are Essential for Protection from Respiratory Failure and Associated
1228 Morbidity following Influenza Virus Infection. *PLoS Pathog.* 10.
1229 <https://doi.org/10.1371/journal.ppat.1004053>
- 1230 Schulz, C., Perdiguero, E.G., Chorro, L., Szabo-Rogers, H., Cagnard, N., Kierdorf, K., Prinz, M., Wu, B.,
1231 Jacobsen, S.E.W., Pollard, J.W., Frampton, J., Liu, K.J., Geissmann, F., 2012. A lineage of myeloid
1232 cells independent of myb and hematopoietic stem cells. *Science (80-)*. 335, 86–90.
1233 <https://doi.org/10.1126/science.1219179>
- 1234 Schwaiger, T., Sehl, J., Karte, C., Schäfer, A., Hühr, J., Mettenleiter, T.C., Schröder, C., Köllner, B.,

- 1235 Ulrich, R., Blohm, U., 2019. Experimental H1N1pdm09 infection in pigs mimics human seasonal
1236 influenza infections. *PLoS One* 14, 1–21. <https://doi.org/10.1371/journal.pone.0222943>
- 1237 ScloTTau, K., Rissmann, M., Graaf, A., Schön, J., Sehl, J., Wylezich, C., Höper, D., Mettenleiter, T.,
1238 Balkema-Buschmann, A., Harder, T., Grund, C., Hoffmann, D., Breithaupt, A., Beer, M., 2020.
1239 SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *Lancet*
1240 1, e218–e225.
- 1241 Shi, J., Wen, Z., Zhong, G., Yang, H., Wang, C., Huang, B., Liu, R., He, X., Shuai, L., Sun, Z., Zhao, Y., Liu,
1242 P., Liang, L., Cui, P., Wang, J., Zhang, X., Guan, Y., Tan, W., Wu, G., Chen, H., Bu, Z., Bu, Z., 2020.
1243 Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2.
1244 *Science* (80-.). 368, 1016–1020. <https://doi.org/10.1126/science.abb7015>
- 1245 Shields, L.E., Jennings, J., Liu, Q., Lee, J., Ma, W., Blecha, F., Miller, L.C., Sang, Y., 2019. Cross-species
1246 genome-wide analysis reveals molecular and functional diversity of the unconventional
1247 interferon- ω subtype. *Front. Immunol.* 10. <https://doi.org/10.3389/fimmu.2019.01431>
- 1248 Sieck, G.C., Dogan, M., Young-soo, H., Valencia, S.O., Delmotte, P., 2019. Mechanisms underlying TNF
1249 α -induced enhancement of force generation in airway smooth muscle.
1250 <https://doi.org/10.14814/phy2.14220>
- 1251 Singh, B., Pearce, J.W., Gamage, L.N., Janardhan, K., Caldwell, S., 2004. Depletion of pulmonary
1252 intravascular macrophages inhibits acute lung inflammation. *Am J Physiol Lung Cell Mol Physiol*
1253 286, L363-72. <https://doi.org/10.1152/ajplung.00003.2003>
- 1254 Sinkora, M., Butler, J.E., 2016. Progress in the use of swine in developmental immunology of B and T
1255 lymphocytes. *Dev. Comp. Immunol.* 58, 1–17. <https://doi.org/10.1016/j.dci.2015.12.003>
- 1256 Sinkora, M., Sinkorova, J., 2014. B Cell Lymphogenesis in Swine Is Located in the Bone Marrow. *J.*
1257 *Immunol.* 193, 5023–5032. <https://doi.org/10.4049/jimmunol.1401152>
- 1258 Sinkora, M., Stepanova, K., Sinkorova, J., 2013. Different anti-CD21 antibodies can be used to
1259 discriminate developmentally and functionally different subsets of B lymphocytes in circulation
1260 of pigs. *Dev. Comp. Immunol.* 39, 409–418. <https://doi.org/10.1016/j.dci.2012.10.010>
- 1261 Sisson, S., Grossman, J., R, G., 1975. Porcine osteology: skull, in: Saunders (Ed.), *Anatomy of the*
1262 *Domestic Animal*. pp. 1231–1252.
- 1263 Soldevila, F., Edwards, J.C., Graham, S.P., Stevens, L.M., Crudgington, B., Crooke, H.R., Werling, D.,
1264 Steinbach, F., 2018. Characterization of the Myeloid Cell Populations' Resident in the Porcine
1265 Palatine Tonsil. *Front. Immunol.* 9, 1800. <https://doi.org/10.3389/fimmu.2018.01800>
- 1266 Speir, M., Hermans, I.F., Weinkove, R., 2017. Engaging Natural Killer T Cells as 'Universal Helpers' for
1267 Vaccination. *Drugs* 77, 1–15. <https://doi.org/10.1007/s40265-016-0675-z>
- 1268 Starbæk, S.M.R., Brogaard, L., Dawson, H.D., Smith, A.D., Heegaard, P.M.H., Larsen, L.E., Jungersen,
1269 G., Skovgaard, K., 2018. Animal Models for Influenza A Virus Infection Incorporating the
1270 Involvement of Innate Host Defenses: Enhanced Translational Value of the Porcine Model. *ILAR*
1271 *J.* 59, 323–337. <https://doi.org/10.1093/ilar/ily009>
- 1272 Steenhard, N.R., Jungersen, G., Kokotovic, B., Beshah, E., Dawson, H.D., Urban, J.F., Roepstorff, A.,
1273 Thamsborg, S.M., 2009. *Ascaris suum* infection negatively affects the response to a *Mycoplasma*
1274 *hyopneumoniae* vaccination and subsequent challenge infection in pigs. *Vaccine* 27, 5161–
1275 5169. <https://doi.org/10.1016/j.vaccine.2009.05.075>
- 1276 Stepanova, K., Sinkora, M., 2013. Porcine T Lymphocytes Can Be Categorized into Two Functionally
1277 and Developmentally Distinct Subsets according to Expression of CD2 and Level of TCR. *J.*

- 1278 Immunol. 190, 2111–2120. <https://doi.org/10.4049/jimmunol.1202890>
- 1279 Stylianou, E., Paul, M.J., Reljic, R., McShane, H., 2019. Mucosal delivery of tuberculosis vaccines: a
1280 review of current approaches and challenges. *Expert Rev. Vaccines* 18, 1271–1284.
1281 <https://doi.org/10.1080/14760584.2019.1692657>
- 1282 Su, S., Wong, G., Shi, W., Liu, J., Lai, A.C.K., Zhou, J., Liu, W., Bi, Y., Gao, G.F., 2016. Epidemiology,
1283 Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends Microbiol.* 24, 490–502.
1284 <https://doi.org/10.1016/j.tim.2016.03.003>
- 1285 Summerfield, A., Meurens, F., Ricklin, M.E., 2015. The immunology of the porcine skin and its value
1286 as a model for human skin. *Mol. Immunol.* 66, 14–21.
1287 <https://doi.org/10.1016/j.molimm.2014.10.023>
- 1288 Sun, S.H., Chen, Q., Gu, H.J., Yang, G., Wang, Y.X., Huang, X.Y., Liu, S.S., Zhang, N.N., Li, X.F., Xiong, R.,
1289 Guo, Y., Deng, Y.Q., Huang, W.J., Liu, Q., Liu, Q.M., Shen, Y.L., Zhou, Yong, Yang, X., Zhao, T.Y.,
1290 Fan, C.F., Zhou, Yu Sen, Qin, C.F., Wang, Y.C., 2020. A Mouse Model of SARS-CoV-2 Infection and
1291 Pathogenesis. *Cell Host Microbe* 28, 124-133.e4. <https://doi.org/10.1016/j.chom.2020.05.020>
- 1292 Sung, S.-S.J., Fu, S.M., Rose, C.E., Gaskin, F., Ju, S.-T., Beaty, S.R., 2006. A Major Lung CD103 (E)- 7
1293 Integrin-Positive Epithelial Dendritic Cell Population Expressing Langerin and Tight Junction
1294 Proteins. *J. Immunol.* 176, 2161–2172. <https://doi.org/10.4049/jimmunol.176.4.2161>
- 1295 Swindle, M.M., Makin, A., Herron, A.J., Clubb, F.J., Frazier, K.S., 2012. Swine as Models in Biomedical
1296 Research and Toxicology Testing. *Vet. Pathol.* 49, 344–356.
1297 <https://doi.org/10.1177/0300985811402846>
- 1298 Szebeni, J., Bawa, R., 2020. Human clinical relevance of the porcine model of pseudoallergic infusion
1299 reactions. *Biomedicines* 8, 1–24. <https://doi.org/10.3390/biomedicines8040082>
- 1300 Szelenyi, I., 2000. Animal models of bronchial asthma. *Inflamm. Res.* 49, 639–654.
1301 <https://doi.org/10.1007/s000110050642>
- 1302 Talker, S.C., Koinig, H.C., Stadler, M., Graage, R., Klingler, E., Ladinig, A., Mair, K.H., Hammer, S.E.,
1303 Weissenböck, H., Dürrwald, R., Ritzmann, M., Saalmüller, A., Gerner, W., 2015. Magnitude and
1304 kinetics of multifunctional CD4+ and CD8β+ T cells in pigs infected with swine influenza A virus.
1305 *Vet. Res.* 46, 1–16. <https://doi.org/10.1186/s13567-015-0182-3>
- 1306 Talker, S.C., Stadler, M., Koinig, H.C., Mair, K.H., Rodríguez-Gómez, I.M., Graage, R., Zell, R., Dürrwald,
1307 R., Starick, E., Harder, T., Weissenböck, H., Lamp, B., Hammer, S.E., Ladinig, A., Saalmüller, A.,
1308 Gerner, W., 2016. Influenza A Virus Infection in Pigs Attracts Multifunctional and Cross-Reactive
1309 T Cells to the Lung. *J. Virol.* 90. <https://doi.org/10.1128/jvi.01211-16>
- 1310 Tane, S., Noda, K., Shigemura, N., 2017. Ex Vivo Lung Perfusion: A Key Tool for Translational Science
1311 in the Lungs. *Chest* 151, 1220–1228. <https://doi.org/10.1016/j.chest.2017.02.018>
- 1312 Tungatt, K., Dolton, G., Morgan, S.B., Attaf, M., Fuller, A., Whalley, T., Hemmink, J.D., Porter, E.,
1313 Szomolay, B., Montoya, M., Hammond, J.A., Miles, J.J., Cole, D.K., Townsend, A., Bailey, M.,
1314 Rizkallah, P.J., Charleston, B., Tchilian, E., Sewell, A.K., 2018. Induction of influenza-specific local
1315 CD8 T-cells in the respiratory tract after aerosol delivery of vaccine antigen or virus in the
1316 Babraham inbred pig, *PLoS Pathogens*. <https://doi.org/10.1371/journal.ppat.1007017>
- 1317 Tyler, W.S., 1983. Small Airways and Terminal Units Comparative Subgross Anatomy of Lungs. *Am.*
1318 *Rev. Respir. Dis.* 128, S32–S36.
- 1319 Ugolini, M., Gerhard, J., Burkert, S., Jensen, K.J., Georg, P., Ebner, F., Volkers, S.M., Thada, S., Dietert,
1320 K., Bauer, L., Schäfer, A., Helbig, E.T., Opitz, B., Kurth, F., Sur, S., Dittrich, N., Gaddam, S.,

- 1321 Conrad, M.L., Benn, C.S., Blohm, U., Gruber, A.D., Hutloff, A., Hartmann, S., Boekschoten, M. V.,
 1322 Müller, M., Jungersen, G., Schumann, R.R., Suttorp, N., Sander, L.E., 2018. Recognition of
 1323 microbial viability via TLR8 drives TFHcell differentiation and vaccine responses. *Nat. Immunol.*
 1324 19, 386–396. <https://doi.org/10.1038/s41590-018-0068-4>
- 1325 Ulmer, J.B., Donnelly, J.J., Parker, S.E., Rhodes, G.H., Felgner, P.L., Dwarki, V.J., Gromkowski, S.H.,
 1326 Deck, R.R., DeWitt, C.M., Friedman, A., Hawe, L.A., Leander, K.R., Martinez, D., Perry, H.C.,
 1327 Shiver, J.W., Montgomery, D.L., Liu, M.A., 1993. Heterologous protection against influenza by
 1328 injection of DNA encoding a viral protein. *Science (80-)*. 259, 1745–1749.
 1329 <https://doi.org/10.1126/science.8456302>
- 1330 Van Merode, T., Maas, T., Twellaar, M., Kester, A., Van Schayck, C., 2007. PEDIATRIC ALLERGY AND
 1331 Gender-specific differences in the prevention of asthma-like symptoms in high-risk infants 196–
 1332 200. <https://doi.org/10.1111/j.1399-3038.2006.00513.x>
- 1333 Volk, T., Pannicke, U., Reisli, I., Bulashevskaya, A., Ritter, J., Björkman, A., Schäffer, A.A., Fliegau, M.,
 1334 Sayar, E.H., Salzer, U., Fisch, P., Pfeifer, D., Di Virgilio, M., Cao, H., Yang, F., Zimmermann, K.,
 1335 Keles, S., Caliskaner, Z., Güner, Ş., Schindler, D., Hammarström, L., Rizzi, M., Humme, M., Pan-
 1336 Hammarström, Q., Schwarz, K., Grimbacher, B., 2015. DCLRE1C (ARTEMIS) mutations causing
 1337 phenotypes ranging from atypical severe combined immunodeficiency to mere antibody
 1338 deficiency. *Hum. Mol. Genet.* 24, 7361–7372. <https://doi.org/10.1093/hmg/ddv437>
- 1339 Wang, Q., Vlasova, A.N., Kenney, S.P., Saif, L.J., 2019. Emerging and re-emerging coronaviruses in
 1340 pigs. *Curr. Opin. Virol.* 34, 39–49. <https://doi.org/10.1016/j.coviro.2018.12.001>
- 1341 Ware, L.B., 2006. Pathophysiology of acute lung injury and the acute respiratory distress syndrome.
 1342 *Semin. Respir. Crit. Care Med.* 27, 337–349. <https://doi.org/10.1055/s-2006-948288>
- 1343 Wernersson, R., Schierup, M.H., Jørgensen, F.G., Gorodkin, J., Panitz, F., Stærfeldt, H.H., Christensen,
 1344 O.F., Mailund, T., Hornshøj, H., Klein, A., Wang, Jun, Liu, B., Hu, S., Dong, W., Li, W., Wong,
 1345 G.K.S., Yu, J., Wang, Jian, Bendixen, C., Fredholm, M., Brunak, S., Yang, H., Bolund, L., 2005. Pigs
 1346 in sequence space: A 0.66X coverage pig genome survey based on shotgun sequencing. *BMC*
 1347 *Genomics* 6, 1–7. <https://doi.org/10.1186/1471-2164-6-70>
- 1348 Westerhof, L.M., McGuire, K., MacLellan, L., Flynn, A., Gray, J.I., Thomas, M., Goodyear, C.S.,
 1349 MacLeod, M.K.L., 2019. Multifunctional cytokine production reveals functional superiority of
 1350 memory CD4 T cells. *Eur. J. Immunol.* 49, 2019–2029. <https://doi.org/10.1002/eji.201848026>
- 1351 Wilson, H.L., Obradovic, M.R., 2020. Evidence for a common mucosal immune system in the pig. *Mol.*
 1352 *Immunol.* 66, 22–34.
- 1353 Winkler, G.C., Cheville, N.F., 1984. The neonatal porcine lung: Ultrastructural morphology and
 1354 postnatal development of the terminal airways and alveolar region. *Anat. Rec.* 210, 303–313.
 1355 <https://doi.org/10.1002/ar.1092100205>
- 1356 Woolcock, A., Macklem, P., 1971. Mechanical factors influencing collateral ventilation in human, dog,
 1357 and pig lungs. *J Appl Physiol* 30, 99–115.
- 1358 Xiao, L., Estellé, J., Kiilerich, P., Ramayo-Caldas, Y., Xia, Z., Feng, Q., Liang, S., Pedersen, A., Kjeldsen,
 1359 N.J., Liu, C., Maguin, E., Doré, J., Pons, N., Le Chatelier, E., Prifti, E., Li, J., Jia, H., Liu, X., Xu, X.,
 1360 Ehrlich, S.D., Madsen, L., Kristiansen, K., Rogel-Gaillard, C., Wang, J., 2016. A reference gene
 1361 catalogue of the pig gut microbiome. *Nat. Microbiol.* 1, 1–6.
 1362 <https://doi.org/10.1038/nmicrobiol.2016.161>
- 1363 Yang, G., Artiaga, B.L., Lewis, S.T., Driver, J.P., 2017a. Characterizing porcine invariant natural killer T
 1364 cells: A comparative study with NK cells and T cells. *Dev. Comp. Immunol.* 76, 343–351.

- 1365 <https://doi.org/10.1016/j.dci.2017.07.006>
- 1366 Yang, G., Artiaga, B.L., Lomelino, C.L., Jayaprakash, A.D., Sachidanandam, R., Mckenna, R., Driver, J.P.,
1367 2019. Next Generation Sequencing of the Pig $\alpha\beta$ TCR Repertoire Identifies the Porcine Invariant
1368 NKT Cell Receptor. *J. Immunol.* 202, 1981–1991. <https://doi.org/10.4049/jimmunol.1801171>
- 1369 Yang, G., Richt, J., Driver, J., Yang, G., Richt, J.A., Driver, J.P., 2017b. Harnessing Invariant NKT Cells to
1370 Improve Influenza Vaccines: A Pig Perspective. *Int. J. Mol. Sci.* 19, 68.
1371 <https://doi.org/10.3390/ijms19010068>
- 1372 Yao, Y., Jeyanathan, M., Haddadi, S., Barra, N.G., Vaseghi-Shanjani, M., Damjanovic, D., Lai, R.,
1373 Afkhami, S., Chen, Y., Dvorkin-Gheva, A., Robbins, C.S., Schertzer, J.D., Xing, Z., 2018. Induction
1374 of Autonomous Memory Alveolar Macrophages Requires T Cell Help and Is Critical to Trained
1375 Immunity. *Cell* 175, 1634-1650.e17. <https://doi.org/10.1016/j.cell.2018.09.042>
- 1376 Yu, C.I., Becker, C., Wang, Y., Marches, F., Helft, J., Leboeuf, M., Anguiano, E., Pourpe, S., Goller, K.,
1377 Pascual, V., Banchereau, J., Merad, M., Palucka, K., 2013. Human cd1c+dendritic cells drive the
1378 differentiation of cd103+cd8+mucosal effector t cells via the cytokine tgf- β . *Immunity* 38, 818–
1379 830. <https://doi.org/10.1016/j.immuni.2013.03.004>
- 1380 Zhang, C., Liu, Y., Chen, S., Qiao, Y., Zheng, Y., Xu, M., Wang, Z., Hou, J., Wang, J., Fan, H., 2019.
1381 Effects of intranasal pseudorabies virus AH02LA infection on microbial community and immune
1382 status in the ileum and colon of piglets. *Viruses* 11. <https://doi.org/10.3390/v11060518>
- 1383 Zhang, H., Li, H., Wang, W., Wang, Y., Han, G.Z., Chen, H., Wang, X., 2020. A unique feature of swine
1384 ANP32A provides susceptibility to avian influenza virus infection in pigs. *PLoS Pathog.* 16.
1385 <https://doi.org/10.1371/journal.ppat.1008330>
- 1386

Table I: Immune-related respiratory tract remarkable features of mouse, human and pig

	<i>Mouse</i>	<i>Human</i>	<i>Pig</i>	<i>Comments</i>
<i>Airways</i>	-Muzzle	-Nose	-Snout	-hm ≠ pg ≠ ms ≠ in higher respiratory tract size and structure
	-Small airways and lungs	-Large airways and lungs	-Large airways and lungs	-hm & pg ≠ ms ≠ in lower respiratory tract accessibility
<i>Lymph nodes</i>	-Centripetal circulation	-Centripetal circulation	-Centrifugal circulation	-hm & ms ≠ pg no known consequences
	-Lymphocytes exit through efferent lymphatics	-Lymphocytes exit through efferent lymphatics	-Lymphocytes exit through HEV	-hm & ms ≠ pg no known consequences
<i>Granulocytes</i>	-Neutrophil serine proteases not-sensitive to human inhibitors	-Neutrophil serine proteases sensitive to human inhibitors	-Neutrophil serine proteases sensitive to human inhibitors	-hm & pg ≠ ms ≠ in tissue inflammation handling
<i>AM</i>	-Embryonic origin	-Probably of embryonic origin	-Probably of embryonic origin	-hm=ms=pg
	-Self-renew -Inflammation, replacement by moAM	-No data -Large part of moAM	-No data -No data	
<i>PIM</i>	-No information	-Induced PIM upon liver conditions?	-Constitutive inflammatory PIM	-hm & ms ≠ pg high susceptibility of pigs to infused particles
<i>moM0/ interstitial M0</i>	-2 populations: nerve-associated blood vessel associated	-2 populations: nerve-associated blood vessel associated	-2 populations: location not explored	-hm=ms=pg
<i>DC</i>	-TLR3 expressed on cDC1	-TLR3 expressed on cDC1	-TLR3 expressed on pDC	-hm & ms ≠ pg ≠ dsRNA response?
	-cDC1 mainly involved in T CD8 activation	-cDC1 and cDC2 involved in T CD8 activation	-cDC1 and cDC2 involved in T CD8 activation	-hm & pg ≠ ms ≠ T CD8 induction?
	-cDC2 FcεRIα-, Lang-, CD103-, interstitial location	-cDC2 FcεRIα+, Lang+, CD103+, subepithelial location	-cDC2 FcεRIα+, Lang+, CD103+, subepithelial location	-hm & pg ≠ ms ≠ allergy induction?
<i>NK/NKT cells</i>	-NKp46 specifically expressed on NK cells	-NKp46 specifically expressed on NK cells	-NK subset NKP46-, T lymphocyte subset NKP46+	-hm & ms ≠ pg no known consequences
<i>T Lymphocytes</i>	-CD4 T cells devoid of CD8 expression	-CD4 T cells devoid of CD8 expression	-CD8α on memory CD4 T cells	-hm & ms ≠ pg no known consequences
	-Low proportion of γδ T cells	-Low proportion of γδ T cells	-High proportion of γδ T cells	-hm & ms ≠ pg ≠ innate immune response?
<i>B Lymphocytes</i>	-Naïve B cells transporting antigens to fDC	-Not described	-Centroblasts transporting antigens to fDC?	-hm & ms ≠ pg no known consequences

HEV: High-Endothelial Venules; AM: Alveolar Macrophage; PIM: Pulmonary Intravascular Macrophage; moAM: monocyte-derived AM; M0: Macrophages, moM0: monocyte derived Macrophage; DC: Dendritic Cell; cDC: conventional Dendritic Cell; pDC: plasmacytoid Dendritic Cell; fDC: follicular Dendritic Cell; NK: Natural Killer; hm: human; ms: mouse; pg: pig. Bibliographic references are in the main text.

Table 2: Non-transmissible respiratory diseases pro and con of the porcine model

	PROS	CONS	
RESPIRATORY INFECTIONS	<i>B. pertussis</i>	-naturally infected	
	<i>Mycobacterium sp</i>	-naturally infected	
	<i>S. aureus</i>	-naturally infected	
	<i>P. aeruginosa</i>	-naturally infected	
	<i>Influenzavirus</i>	-natural sensitivity to human and swine strains with mild clinical signs	-no example of highly pathogenic infections
	<i>Coronavirus</i>	-natural porcine <i>αCoronavirus</i> (PrCoV) similar to the human seasonal HCoV-229E	-disputed sensitivity to SARS-CoV-2
	<i>Orthopneumovirus</i>	-natural porcine <i>Orthopneumovirus?</i>	-not sensitive to hRSV
VACCINATION	Delivery	-skin structure -upper respiratory tract structure	
	Adjuvant	-functional TLR8 -cDC1 interstitial and cDC2 intra-subepithelial locations -αGalSer response	-TLR3 expression on pDC not on cDC1
MICROBIOTA		-omnivorous species -intestinal microbiota similar to herbivore - data paucity on porcine respiratory microbiota	
RESPIRATORY ALLERGY AND ASTHMA	Induction	-cDC2 location -cDC2 expressing IgE receptor (FcεRIα)	-no stable chronic allergic asthma porcine model
	Clinical signs	-IgE similar to human -lung smooth muscles porcine model -efficacy of tryptase inhibitors	
ACUTE AND CHRONIC INFLAMMATIONS		-model of early inflammatory ARDS responses -constitutive PIM	-no infectious model of ARDS induction -constitutive PIM
	LUNG GRAFT	-size allowing similar surgical intervention -primary graft dysfunction model - <i>ex vivo</i> lung perfusion model	

SARS: Severe Acute Respiratory Syndrome; hRSV: human Respiratory Syncytial Virus; cDC: conventional Dendritic Cell; pDC: plasmacytoid Dendritic Cell; ARDS: Acute respiratory distress syndrome; PIM: Pulmonary Intravascular Lymphocyte. Bibliographic references are in the main text.