

## First report of Tomato leaf curl New Delhi virus infecting courgette in France

Cecile Desbiez, P. Gentit, Pascaline Cousseau-Suhard, I. Renaudin, Eric  
Verdin

► **To cite this version:**

Cecile Desbiez, P. Gentit, Pascaline Cousseau-Suhard, I. Renaudin, Eric Verdin. First report of Tomato leaf curl New Delhi virus infecting courgette in France. *New Disease Reports, British Society for Plant Pathology*, 2021, 42 (2), 10.1002/ndr2.12006 . hal-03230529

**HAL Id: hal-03230529**

**<https://hal.inrae.fr/hal-03230529>**

Submitted on 20 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## NEW DISEASE REPORT

# First report of *Tomato leaf curl New Delhi virus* infecting courgette in France

C. Desbiez<sup>1</sup>  | P. Gentit<sup>2</sup> | P. Cousseau-Suhard<sup>2</sup> | I. Renaudin<sup>2</sup> | E. Verdin<sup>1</sup><sup>1</sup> Inrae, Pathologie Végétale, Montfavet F-84140, France<sup>2</sup> Anses, LSV-BVO, Angers F-49044, France

## Correspondence

Email: [Cecile.desbiez@inrae.fr](mailto:Cecile.desbiez@inrae.fr)

## KEYWORDS

begomovirus, cucurbit

Viruses constitute important problems for cucurbit production. In France, aphid-transmitted viruses remain the most important (Desbiez *et al.*, 2020), but whitefly-transmitted viruses, particularly begomoviruses, now emerging in the Mediterranean Basin (Leccoq & Desbiez, 2012) constitute major threats.

A few plants showing leaf crumpling, yellowing and downward curling were observed in a courgette (*Cucurbita pepo*) plot in Bouches-du-Rhône department (southeastern France) in September 2020. The symptoms were similar to those of *Tomato leaf curl New Delhi virus* (ToLCNDV), an emerging bipartite begomovirus affecting cucurbits in the Mediterranean Basin (Fortes *et al.*, 2016). After DNA extraction, PCR was performed on a symptomatic sample (CD20001) with specific primers TLCNDV-CP-5 and TLCNDV-CP-3, and ToLCNDV-B-900-5 and ToLCNDV-B-1940-3 (Romay *et al.*, 2019) targeting ToLCNDV DNA-A and DNA-B respectively. The amplified fragments (760 bp and 1072 bp) displayed 99.1% and 99.6% identity respectively to ToLCNDV isolate MU.12.ZU/1/2 (MH577761) from Spain. To complete the sequence of DNA-A, PCR amplification was performed with primers TLCNDV-fCP-F (5'-GTGACGGGAGGAACRTATGC-3') and TLCNDV-dCP-R (5'-CTAACACACATGACTTTGCC-3') before sequencing. For DNA-B, rolling circle amplification (RCA) was performed on the extracted DNA with the TempliPhi kit (GE Healthcare). A 2.7 Kb fragment was obtained after digestion with *Bam*HI and cloned in linearised pBlueScript(KS)+ before sequencing. Two clones were sequenced and were identical. The complete sequences of DNA-A and DNA-B (GenBank Accession Nos. MW310624 and MW310625, respectively) were 98–99% identical to isolates from the “Mediterranean” ToLCNDV clade (Fortes *et al.*, 2016).



**FIGURE 1** Foliar symptoms in a zucchini plot in which *Tomato leaf curl New Delhi virus* was detected in September 2021 in Bouches-du-Rhône, southeastern France

Five other symptomatic courgette plants were collected in the same plot and 12 plants in two other plots in Bouches-du-Rhône, at a distance of 10 km (8 plants) and 6 km (4 plants) respectively from the first plot (Figure 1). Four courgette samples with similar symptoms were also collected in Gard department, 30 km away from the first plot. The samples were tested by PCR with primers Beg-CP-F and Beg-580-R (Saison *et al.*, 2015). A 580 bp product was obtained for 17 of the 21 tested samples (3/5 in the first plot, 8/8 and 4/4 respectively in the two

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *New Disease Reports* published by British Society for Plant Pathology and John Wiley & Sons



other plots in Bouches-du-Rhône and 2/4 in Gard) and determined as ToLCNDV by sequencing, confirming the presence of the virus in all four areas. The sequences displayed 99.3-100% identity with isolate CD20001.



**FIGURE 2** Foliar symptoms in a zucchini plot in which *Tomato leaf curl New Delhi virus* was detected in September 2021 in Bouches-du-Rhône, southeastern France



**FIGURE 3** Fruit symptoms in a zucchini plot in which *Tomato leaf curl New Delhi virus* was detected in September 2021 in Bouches-du-Rhône, southeastern France

This is the first report of ToLCNDV in France. Since ToLCNDV is highly damaging in cucurbits and is a quarantine pest in Europe, it is important to prevent its emergence in France, particularly in the southeast which constitutes France's most important cucurbit production area. The presence of ToLCNDV in four locations within a 30 km range suggests that it has begun to spread efficiently or that multiple introductions have occurred. It is now important to prevent virus overwintering in weeds (Juárez *et al.*, 2019) and be careful to detect and eradicate potential infections in the next growing season.

#### ORCID

C. Desbiez  <https://orcid.org/0000-0001-8691-7423>

#### References

- Desbiez C, Wipf-Scheibel C, Millot P, Berthier K, Girardot G, Gognalons P, Hirsch J, Moury B, Nozeran K, Piry S, Schoeny A, Verdin E, 2020. Distribution and evolution of the major viruses infecting cucurbitaceous and solanaceous crops in the French Mediterranean area. *Virus Research* 286, 198042. <https://doi.org/10.1016/j.virusres.2020.198042>
- Fortes IM, Sánchez-Campos S, Fiallo-Olivé E, Díaz-Pendón JA, Navas-Castillo J, Moriones E, 2016. A novel strain of tomato leaf curl New Delhi virus has spread to the Mediterranean basin. *Viruses* 8, 307. <https://doi.org/10.3390/v8110307>
- Juárez M, Rabadán MP, Martínez LD, Tayahi M, Grande-Pérez A, Gómez P, 2019. Natural hosts and genetic diversity of the emerging tomato leaf curl New Delhi virus in Spain. *Frontiers in Microbiology* 10, 140. <https://doi.org/10.3389/fmicb.2019.00140>
- Lecoq H, Desbiez C, 2012. Viruses of cucurbit crops in the Mediterranean region: an ever-changing picture. *Advances in Virus Research* 84, 67–126. <https://doi.org/10.1016/B978-0-12-394314-9.00003-8>
- Romay G, Pitrat M, Lecoq H, Wipf-Scheibel C, Millot P, Girardot G, Desbiez C, 2019. Resistance against melon chlorotic mosaic virus and tomato leaf curl New Delhi virus in melon. *Plant Disease* 103, 2913–2919. <https://doi.org/10.1094/PDIS-02-19-0298-RE>
- Saison A, Gentit P, 2015. Development of a polyvalent detection method for Begomoviruses presenting a threat to the European tomato industry. Testa – EPPo conference on diagnostics for plant pests, Angers, France, 30/11/2015 to 02/12/2015. [https://www.eppo.int/media/uploaded\\_images/MEETINGS/Meetings\\_2015/testa\\_angers/21\\_Gentit.pdf](https://www.eppo.int/media/uploaded_images/MEETINGS/Meetings_2015/testa_angers/21_Gentit.pdf). Accessed 2 December 2020.

**How to cite this article:** Desbiez C, Gentit P, Cousseau-Suhard P, Renaudin I, Verdin E, First report of *Tomato leaf curl New Delhi virus* infecting courgette in France. *New Dis Rep.* 2021;43, e12006. <https://doi.org/10.1002/ndr2.12006>