



**HAL**  
open science

# Halving food-related greenhouse gas emissions can be achieved by redistributing meat consumption: progressive optimization Results of the NutriNet-Santé cohort

Emmanuelle Kesse-Guyot, H el ene Fouillet, Julia Baudry, Alison Dussot, Brigitte Langevin, Benjamin All es, Pauline Rebouillat, Jos ephine Brunin, Mathilde Touvier, Serge Hercberg, et al.

## ► To cite this version:

Emmanuelle Kesse-Guyot, H el ene Fouillet, Julia Baudry, Alison Dussot, Brigitte Langevin, et al.. Halving food-related greenhouse gas emissions can be achieved by redistributing meat consumption: progressive optimization Results of the NutriNet-Sant  cohort. *Science of the Total Environment*, 2021, 789, pp.147901. 10.1016/j.scitotenv.2021.147901 . hal-03238402

**HAL Id: hal-03238402**

**<https://hal.inrae.fr/hal-03238402v1>**

Submitted on 27 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

**Halving food-related greenhouse gas emissions can be achieved by redistributing meat consumption: progressive optimization Results of the NutriNet-Santé cohort**

Emmanuelle Kesse-Guyot<sup>1</sup>, H el ene Fouillet<sup>2</sup>, Julia Baudry<sup>1</sup>, Alison Dussot<sup>2</sup>, Brigitte Langevin<sup>5</sup>, Benjamin All es<sup>1</sup>, Pauline Rebouillat<sup>1</sup>, Jos ephine Brunin<sup>1,6</sup>, Mathilde Touvier<sup>1</sup>, Serge Hercberg<sup>1,3</sup>, Denis Lairon<sup>4</sup>, Fran ois Mariotti<sup>2</sup>, Philippe Pointereau<sup>5</sup>

<sup>1</sup> Sorbonne Paris Nord University, Inserm, INRAE, Cnam, Nutritional Epidemiology Research Team (EREN), Epidemiology and Statistics Research Center – University of Paris (CRESS), 93017 Bobigny, France

<sup>2</sup> UMR PNCA, AgroParisTech, INRAE, Universit  Paris-Saclay, 75005, Paris, France

<sup>3</sup> D partement de Sant  Publique, H pital Avicenne, 93017 Bobigny, France

<sup>4</sup> Aix Marseille Universit , INSERM, INRAE, C2VN, 13005 Marseille, France

<sup>5</sup> Solagro, 75, Voie TOEC, CS 27608, F-31076 Toulouse Cedex 3, France

<sup>6</sup> ADEME, Agence de l'Environnement et de la Ma trise de l'Energie), 49004 Angers, France

**Correspondence:** Email: [e.kesse@eren.smbh.univ-paris13.fr](mailto:e.kesse@eren.smbh.univ-paris13.fr)

Equipe de Recherche en Epid miologie Nutritionnelle (EREN)

SMBH Universit  Sorbonne Paris Nord, 74 rue Marcel Cachin, 93017 Bobigny, France

**Running title:** Low-Greenhouse Gas Emissions optimized diets

**Keywords:** diet sustainability, greenhouse gas emissions, optimization modelling, co-product trade-offs

**Number of tables:** 3/**Number of figures:** 3

**Appendix:** 4 tables and 3 figures

1 **Abstract 300 words**

2 **Background:** Diet-related greenhouse gas emissions (GHGe) mainly comes from animal-sourced  
3 foods. As progressive changes are more acceptable for a sustainable food transition, we aimed to  
4 identify nutritionally adequate and culturally acceptable optimized diets ensuring a gradual reduction  
5 in GHGe, using observed diet from a large sample of French adults, while considering the mode of  
6 food production (organic vs conventional farming) and the co-production link between milk and beef.

7 **Material and method:** Based on the consumption of 257 organic and conventional foods among  
8 29,413 participants (75% women, age: 53.5±14.0y) of the NutriNet-Santé study, we modelled optimal  
9 diets according to GHGe reduction scenarios in 5% steps, from 0 to 50% with nutritional,  
10 acceptability, and coproduct constraints, for men, premenopausal and menopausal women separately.

11 **Results:** Gradual GHGe decrease under these constraints led to optimal diets with an overall decrease  
12 in animal foods, with marked reductions in dairy products (up to -83%), together with a stable but  
13 largely redistributed meat consumption in favor of poultry (up to +182%) and pork (up to +46%) and  
14 at the expense of ruminant meat (down to -92%). Amounts of legumes increases dramatically (up to  
15 +238%). The greater the reduction in diet-related GHGe, the lower the cumulative energy demand  
16 (about -25%) and land use (about -43%). The proportion of organic food increased from ~30% in the  
17 observed diets to ~70% in the optimized diets.

18 **Conclusion:** Our results suggest that meeting both nutrient reference value and environmental  
19 objectives of up to 50% GHGe reduction requires the reduction of animal foods together with  
20 important substitutions between animal food groups, which result in drastic reductions in beef and  
21 dairy products. Further research is required to explore alignment with long-term health value and  
22 conflict with acceptability, in particular for even greater GHGe reductions.

## 23 1. Introduction

24 The current environmental crisis, beyond the irreversible damage to natural resources, is characterized  
25 by climate change and global warming (defined as the increase of both air and sea surface temperature  
26 over a long period of time (>30 years)). Thus, anthropogenic global warming in 2017 was +1°C  
27 compared to pre-industrial levels (1850-1900), *i.e.* about 0.2°C per decade (1).

28 Food systems are responsible for about 30% of global greenhouse gas emissions (GHGe) (2) and are  
29 major users of fresh water, therefore largely contributing to climate change (3). Unless action is taken  
30 in the next decades, various prospective scenarios have estimated that, by 2050, unsustainable diets  
31 will lead to an additional 80% rise in GHGe compared to the current situation (4). To mitigate climate  
32 change below 1.5°C, some scenarios have documented that halving agricultural carbon footprint by  
33 2050 would be necessary (5) and this would require strong dietary changes on a global scale (6).

34 The scientific literature about GHGe related to dietary patterns, based on optimization-based  
35 modelling and observational data, is growing and is consistently reporting that plant-based diets  
36 exhibit lower GHGe compared to those rich in animal products (7–10). Plant-based dietary patterns  
37 can also help to prevent chronic diseases (3,11–14), underlying co-benefits of plant-based diets for  
38 climate mitigation and human health promotion. Clark et al. (14) have documented, based on  
39 metadata, that beneficial foods for health, apart from fish, generally exhibit lower environmental  
40 pressures, encompassing GHGe, acidification, eutrophication, land and water use. Conversely, some  
41 foods which could be detrimental for health, such as red meat and processed meat (associated with  
42 increased risk for various health outcomes, including mortality and morbidity due to coronary heart  
43 diseases, stroke, diabetes and colorectal cancer), are also the highest contributors to diet-related GHGe  
44 and large variations in GHGe exist across food groups (14).

45 However, the question of how to achieve changes in dietary behavior including a reduction in meat  
46 consumption and more generally animal-based food has not been resolved (15). Indeed, food choices  
47 are diverse and based on multiple influencing factors which may constitute barriers (15,16). Thus, the  
48 strategies accompanying the transitions towards greater sustainability and in particular lower GHGe  
49 should be multiple and adapted to different types of consumers (17). We may hypothesize that among  
50 traditional high meat consumers, a first step of the transition can be based on intra-food group  
51 substitutions, especially due to cultural traits that hinder large reduction in meat consumption (18,19).  
52 For instance, in France, a previous study modelled a gradual reduction in GHGe and showed that a  
53 30% reduction was possible without drastically deviating from the current diets while respecting  
54 nutritional constraints and diet cost (20). Among the gaps in existing studies we can mention the  
55 following. First, most modelling studies used GHGe as constraints or objective function (10) but few  
56 have considered other environmental indicators in their analysis - as descriptors, constraints or  
57 objective - despite the fact that some conflicts are known to occur among the different environmental  
58 dimensions, which are related to the general organization of the food system, such as energy demand  
59 and land occupation (21). Besides, few studies have distinguished conventional and eco-friendly

60 production systems, as the data are generally based on life cycle assessment (LCA) for generic foods.  
61 Although organic farming has not been systematically related to lower GHGe, energy demand is lower  
62 while land use is higher compared to conventional agriculture (22,23). In a previous work based on  
63 observational data distinguishing between organic and conventional diets, we observed that diets with  
64 high GHGe were higher in animal-based food, more caloric and nutritionally less healthy (24). Thus,  
65 the role of various food systems on environmental pressure has not been yet considered enough (25).  
66 Finally, food systems also include some important structural determinants of food production, such as  
67 the fact that co-productions rules are often operating, as for milk and beef meat productions, but is  
68 rarely considered (26–28).

69 We can hypothesize that transitions to sustainable diets will require to activate all the levers and  
70 substitution is one of them. Thus, the objective of this study was to test whether the possibility to  
71 reduce GHGe of production by 50% as defined in the Paris agreement (29) in a set of culturally  
72 acceptable diets. We modelled dietary pattern characteristics with gradual decreasing GHGe under  
73 nutritional, coproduction and acceptability constraints and to relate these dietary patterns to other  
74 environmental indicators while considering two different food production systems, organic and  
75 conventional.

## 76 **2. Methods**

### 77 **2.1 Population**

78 The population included adults participating in the ongoing web-based prospective NutriNet-Santé  
79 cohort initiated in France in May 2009 (and on-going) whose aim is to investigate relationships  
80 between nutrition and health (30). Participants are recruited on a voluntary basis from the general  
81 French population. This study is conducted in accordance with the Declaration of Helsinki, and all  
82 procedures were approved by the Institutional Review Board of the French Institute for Health and  
83 Medical Research (IRB Inserm 0000388FWA00005831) and the National Commission on Informatics  
84 and Liberty (Commission Nationale de l'Informatique et des Libertés, CNIL 908450 and 909216).  
85 Electronic informed consent was obtained from all participants. The NutriNet-Santé study is registered  
86 in ClinicalTrials.gov (NCT03335644). At baseline and every year thereafter, participants provide data  
87 about their sociodemographic and economic status, anthropometrics, lifestyles and dietary intakes  
88 through self-administered questionnaires. They are also regularly invited to fill in complementary  
89 questionnaires.

### 90 **2.2 Sociodemographic and lifestyle characteristics**

91 Participants completed regularly validated questionnaires about sociodemographic and lifestyle  
92 features (31,32), thus data from the sociodemographic questionnaires that were the closest to the  
93 dietary questionnaire were used. Sociodemographic and lifestyle characteristics encompassed gender,  
94 age, education (<high school diploma, high school diploma, and post-secondary graduate), smoking

95 status (former, current, or never-smoker), and physical activity assessed using the International  
96 Physical Activity questionnaire (33).

### 97 2.3 Dietary data assessment

98 The present study is based on dietary data collected through a self-administered validated semi-  
99 quantitative food frequency questionnaire (FFQ), administered from June to December 2014. This  
100 questionnaire has been enriched by a five-point ordinal scale for each of the 264 food and beverage  
101 items to evaluate the share of organic food consumption (under official labels) (34). Organic  
102 production is one of the official signs of quality and origin in France. This method is governed by a  
103 European regulation since 1991, with the overall objectives of ensuring respect of the environment,  
104 biodiversity and animal welfare (35) recently updated (EU regulations 2018/848 and 2020/464 coming  
105 into force in January 2022. More specifically, for each item, participants were asked ‘How often was  
106 the product of organic origin?’ with the following answer modalities: “never”, “rarely”, “half-of-  
107 time”, “often” or “always”. This questionnaire developed within the frame of the BioNutriNet project  
108 has been extensively described elsewhere (22). Organic food consumption was estimated by allocating  
109 the respective weights: 0, 25, 50, 75 and 100 to the modalities. For clarity purpose, food and beverage  
110 items were grouped into 16 food groups as presented in **Appendix A**. Nutrient intake were calculated  
111 using a published food composition database (36).

## 112 2.4 Environmental pressure indicators

113 Environmental indicators were assessed using the DIALECTE tool, established by Solagro  
 114 (Toulouse, France) (37), whose aim is to evaluate environmental impacts of French farming  
 115 systems using attributional life cycle analysis (LCA) without considering land use change. The  
 116 perimeter of LCA was restricted to the agricultural production phase (conditioning, transport,  
 117 processing, storage or recycling stages were not considered). Upstream processes were  
 118 therefore included in the assessment, such as input production or energy supply. Three  
 119 environmental indicators were considered at the farm level: the GHGe measured as kg of CO<sub>2</sub>  
 120 equivalents (CO<sub>2</sub>eq), the cumulative energy demand (CED) in MJ, and the land occupation  
 121 expressed in m<sup>2</sup> for >60 raw products (24). The original database has been completed by other  
 122 data sources that have been previously listed (24), to obtain environmental pressures in organic  
 123 and conventional of 92 raw agricultural products covering the 264 food items. Data have been  
 124 published elsewhere (24). Environmental pressures of the FFQ food items as consumed were  
 125 retrieved from the 92 raw agricultural products by using a set of conversion coefficients  
 126 (economic allocation (accounting for co-products) and cooking and edibility coefficients).

### 127 Coproduct factors for ruminant products

128 We considered a meat to carcass weight ratio of 68% (38), and further yields of 90% during  
 129 distribution (due to 10% distribution losses) and 68% during consumption (due to 32% losses by  
 130 cooking, bones and wastes) (38).

131 In 2010 in France, 25 million tons of milk and 1.52 million tons of beef (expressed in carcass weight)  
 132 (5) were produced, of which 41% was from dairy herd, i.e., 0.62 million tons of beef (39). Thus, 1L of  
 133 milk corresponded to 10g of beef when applying the equation (1):

$$134 \quad (1) \quad 25 \text{ million tons of Milk (L)} = 1.52 \text{ million tons of beef} \times 41\% \times 68\%_{\text{carcass yield}} \times$$

$$135 \quad \quad \quad 90\%_{\text{distribution yield}} \times 68\%_{\text{preparation yield}}$$

136 Furthermore, we considered that 8L of milk are required to make 1kg of cheese and 1L of milk to  
 137 make 1kg of fresh dairy products, using the average figures from French processing chains.

## 138 2.6 Diets modelling and optimization

139 As nutrient requirements vary according to population subgroups, participants were grouped as men,  
 140 premenopausal women and menopausal women and diets were modeled for each subgroup to account  
 141 for differences in iron intake requirements. Postmenopausal women were considered to have a low  
 142 iron requirement and premenopausal women have a high iron requirement (the highest reference  
 143 value, i.e. the reference covering 97.5% of the women requirements; of note most women (80%) have  
 144 much lower requirements). Data related to observed food consumption as well as attributes of food  
 145 items, *i.e.* nutritional composition, environmental pressures and production mode (conventional or

146 organic), were used to define optimized diets being nutritionally adequate, acceptable, and more  
 147 sustainable.

148 Nutritional adequacy was ensured by a set of nutritional constraints by considering, in particular,  
 149 nutrient bioavailability for iron and zinc, as described in **Appendix B**.

150 The list of constraints was as follows:

- 151 - Nutritional constraints on total energy and 31 nutrients, with upper and/or lower bounds based  
 152 on nutrient reference intakes. Lower bounds were taken as recommended dietary allowance  
 153 (population reference intake) or adequate intake, or lower bound of reference intake range for  
 154 the French population (40) as mostly derived from the EFSA opinion (41). For some nutrients,  
 155 when the adequate intake was based on the observed average intake, the lower bound was set  
 156 as the value of the 5<sup>th</sup> percentile. Reference intakes also included upper levels, as tolerable  
 157 upper intakes for vitamins and minerals, when identified, and upper bound of reference intake  
 158 range.
- 159 - Acceptability constraints on some food groups, with upper bounds set as the population-  
 160 specific 95<sup>th</sup> percentiles for 37 *ad-hoc* food groups. Additional moderation constraints on  
 161 some food groups (dairy products  $\leq 2$  portions/d, fish  $\leq 2$  portions/week with 1 of fatty fish, and  
 162 red meat  $< 500$  g/week), to comply with national public health moderation recommendations  
 163 for animal products, as prescribed in French food-based dietary guidelines (42).
- 164 - Co-production constraint limiting the consumption of milk to a proportion of that of beef,  
 165 using the factor between milk and beef defined above in Eq. (1).
- 166 - Environmental constraint for a given (from 0 to -50 % by 5% decrement) reduction in GHGe  
 167 compared to the observed situation. For each food, during diet optimization, the model  
 168 selected the production option (conventional or organic) exhibiting lower GHGe.

169  
 170 The objective corresponded to the maximization of acceptability, i.e. minimizing the total departure  
 171 (D) from the observed diet (initial condition), as follows:

$$172 \quad \text{Min } D = \sum_i^{257} \left[ \frac{\text{Obs}_i - \text{Opt}_i}{SD_i} \right]^2$$

173 where  $\text{Obs}_i$  and  $\text{Opt}_i$  denote the daily consumption of food item (i) in the observed and optimized  
 174 diets, respectively and  $SD_i$  is the standard deviation of the observed daily consumption of food item  
 175 (i).

176 The climatic improvement approach was examined using scenarios of 5% gradual decreases in GHGe,  
 177 by using a GHGe constraint in each scenario from 0% (basal model:  $\text{GHGe} \leq \text{Observed situation}$ ) to -  
 178 50%.

179 The optimization process was performed using the procedure SAS/OR ® *optmodel* (version 9.4; SAS  
 180 Institute, Inc.) using the *nlp* non-linear optimization algorithm (since the objective and some



181 nutritional constraints were non-linear) and *multistart* option (to ensure that solutions were not only  
182 local optimums).

183 During diet optimization, we estimated the standardized dual values (i.e., the dual values associated  
184 with each constraint that has been standardized by its limiting bound), which represent the potential  
185 gain in objective for a 100% relaxation of each constraint's limiting bound. This allowed to identify  
186 the active (*vs.* inactive) constraints and compare their relative influences on the results. To conduct  
187 this sensitivity analysis even further, some alternative models were also tested, with either introducing  
188 some flexibility in some constraints (like the bioavailable zinc and iron nutritional constraints) or the  
189 suppression of some other (like the co-production constraint).

190 Consumptions of food groups, animal- and plant-based products, nutrient intakes, percentage of  
191 organic production mode per food group, monetary costs, environmental pressures (GHGe, cumulative  
192 energy demand and land occupation) and the pReCiPe, as previously described (22), were calculated  
193 for each optimized diet.

194 The pReCiPe (partial ReCiPe), a synthetic estimate of environmental impact based on GHGe,  
195 cumulative energy demand and land occupation, which enables to consider potential trade-offs  
196 between indicators (43), was calculated as follows:

$$197 \quad pReCiPe = 0.0459 \times GHGe + 0.0025 \times CED + 0.0439 \times LO$$

198  
199 with GHGe, in kg of CO<sub>2</sub>eq/d, CED, in MJ/d and LO, in m<sup>2</sup>/d. The higher the pReCiPe, the higher the  
200 environmental impact.

## 201 2.7 Statistical analysis

202 For the present study, we considered participants of the NutriNet-Santé study who had completed the  
203 Org-FFQ between June and December 2014 (N=37,685), with no missing covariates (N=37,305), not  
204 detected as under- or over-energy reporter (N=35,196), living in mainland France (N=34,453), and  
205 with available data regarding the place of purchase for the computation of the dietary monetary cost,  
206 leading to a final sample of 29,413 participants. The sociodemographic, lifestyle and dietary  
207 characteristics were presented by subgroup (men, premenopausal and menopausal women).

208 Dietary consumptions per subgroup were presented as observed mean (SD) or optimized values for  
209 scenarios for the main 16 food groups and further specifically detailed among both animal and plant-  
210 based foods.

211 All statistical analyses were performed using SAS (version 9.4; SAS Institute, Inc., Cary, NC, USA).

## 212 3. Results

### 213 3.1 Sample characteristics.

214 The sociodemographic characteristics of the sample are presented in **Table 1**. The sample included in  
215 the present analysis was constituted of 7,416 men, 9,710 premenopausal women and 12,287

216 menopausal women. The mean age was 53.5y (SD=14.0). Most of the sample was postgraduate (64%)  
217 and few individuals were current smokers (11%) or exhibited a low physical activity level (19%).  
218 For each population (men, premenopausal and menopausal women), the food group consumptions for  
219 the observed diet and the optimized diet under each model and by each scenario of GHGe reduction  
220 are presented in **Appendix C**.

### 221 3.2 Overall dietary changes

222 The overall food group composition of optimized diets meeting the set of nutritional, acceptability,  
223 moderation and coproduct constraints without (basal scenario, 0% reduction in GHGe) or with gradual  
224 GHGe reduction (following scenarios, up to 50% of the observed pressures), are presented in **Figure**  
225 **1**. In the basal scenario (0% reduction in GHGe, where the optimized diet was the closest diet to the  
226 observed diet that meet the nutritional, acceptability, moderation and coproduct constraints), nutrient  
227 constraints resulted in decreases in butter (up to -80% vs observed diet), dairy products (up to -64%),  
228 extra-foods (up to -75%), non-alcoholic beverages (up to -54%) and fish (up to -45%) and in contrast  
229 increases in soya-based food (up to +390%), eggs (up to +140%) and mixed dishes (up to +156%),  
230 with also some sex-specific effects (whole starchy foods 45% decrease in men but 70% increase in  
231 women, starchy foods and vegetable oils 54% and 145% respectively increase in men). In the  
232 following scenarios, as detailed in the **Appendix D**, fulfilment of the environmental constraints of  
233 gradual up to 50% decrease in GHGe was ensured by gradual further increases in soya-based food in  
234 women (up to +68% vs basal scenario) and eggs in all groups (up to +24%) and by gradual further  
235 decreases in extra-foods in all groups (up to -68%), whole grains & starchy foods, mixed dishes and  
236 dairy products in men (-95%, -32% and -32% respectively), in meat and vegetables oils in women (up  
237 to -27% and -29%, respectively).

### 238 3.3 Animal-based foods consumption

239 **Figure 2** presents the detailed intakes of animal-based foods in observed diets and in the optimized  
240 diets for the basal (0% reduction in GHGe) and following (up to 50% gradual decrease in GHGe)  
241 scenarios. Compared to the observed diets, in the basal scenario meeting the nutritional requirements,  
242 all optimized diets (whatever the population) were characterized by a reduction in total animal  
243 products (up to -44%), with suppression of milk and reductions in dairy products and cheese (up to -  
244 66% and -30%, respectively) and fish (up to -45%, to be reduced to its maximal recommendation), and  
245 in contrast increases in eggs and poultry (up to +140% and +182%, respectively). Moreover, in the  
246 basal scenario compared to the observed situation, so as to ensure the nutrient requirements and  
247 animal-based food dietary guidelines, ruminant meat increased (up to +30% in postmenopausal  
248 women) while pork meat decreased (up to -89% in men). These trends for ruminant and pork meats  
249 were then reversed during the following scenarios of up to 50% reduction in GHGe, which were  
250 systematically characterized by concomitant and gradual decrease in ruminant meat (up to -91%  
251 compared to the basal model with no GHGe decrease) and increase in pork (up to +964%). The GHGe  
252 50% reduction was also ensured, to a lesser extent, by some sex-specific effects in line with those

253 already observed in the basal scenario, namely a further dairy products reduction in men and a further  
254 egg increase in women.

255 Finally, compared to the observed diets, ensuring both nutritional needs, acceptability, moderation and  
256 coproduction constraints and 50% GHGe reduction was achieved by strong reductions in the  
257 consumptions of fish and ruminant meat (up to -23 g/d and -52 g/d) together with strong reductions in  
258 the consumptions of milk and dairy products (up to -65 g/d and -115 g/d), while the consumptions of  
259 poultry, eggs and pork increased (up to +46 g/d, +18 g/d and +18 g/d). Overall, if the total meat  
260 consumption remained relatively similar between the observed and optimized diets, it was strongly  
261 redistributed between meat types, as the contribution of poultry to total meat consumption greatly  
262 increased from 18%-24% in observed diets to 43%-50% in optimized diets, while the contribution of  
263 pork more moderately increased from 39%-42% to 46%-54% and the contribution of ruminant meat  
264 greatly decreased from 34%-41% to 3%-5%.

### 265 3.4 Plant-based foods consumption

266 **Figure 3** presents the detailed intakes of plant-based foods in the observed and optimized diets for the  
267 basal (0% reduction in GHGe) and following (up to 50% gradual decrease in GHGe) scenarios.

268 Compared to the observed diets, in the basal scenario meeting the nutritional requirements, all  
269 optimized diets were characterized by strong increases in legumes (up to +238%, i.e., +45g/d) and  
270 decreases in soups, soya-based food and fruits in all groups (up to -97%, -81% and -34%, i.e., -73 g/d  
271 and -3.6 g/d and -91 g/d respectively). Whole grains and starchy foods decreased in men (-63%, i.e. -  
272 59 g/d) but increase in women (up to +52%, i.e. 34 g/d). These effects were similar or even slightly  
273 further strengthened in the following scenarios of up to 50% GHGe reduction. Indeed, whole grains  
274 and starchy foods decreased in premenopausal women, increased in menopausal women and were  
275 almost totally suppressed in men. The 50% GHGe reduction was also achieved by a reduction in  
276 potatoes (up to -69%, i.e., -21 g/d).

### 277 3.5 Environmental and cost characteristics

278 The evolution of environmental and monetary cost indicators across the different scenarios is  
279 presented in **Table 2**. Compared to the observed situation, the basal scenario meeting the nutrient  
280 reference values (without GHGe reduction) yielded an increase in almost all these indicators (energy  
281 demand and land occupation, pReCiPe and monetary cost of the diet). In the following scenarios of  
282 gradual GHGe reduction, all these indicators then gradually decreased and reached lower values than  
283 those observed for the environmental indicators, but not for the diet monetary cost. Indeed, compared  
284 to the observed situation, in the last scenario of 50% GHGe reduction, the energy demand was lowered  
285 by up to -29%, land occupation by -48% and pReCiPe by -47%, while the monetary cost of the diet  
286 increased between +9% and +20%.

287 The share of organic food, starting from 26-32% in the observed diets, increased greatly and rapidly  
288 from the basal scenario and then stabilized around 65-70% in all optimized diets. As detailed in the

289 **Appendix E**, while animal foods were consumed mostly as non-organic, plant-based food were  
290 consumed as organic in optimized diet.

### 291 3.6 Nutrient characteristics

292 The nutrient intakes according to the observed and optimized diets are presented in **Table 3** Notably,  
293 in all optimized diets, the intakes of fibers and bioavailable zinc, which were insufficient in the  
294 observed diets, were leveled up to their reference value and were then kept unchanged. We found  
295 similar results for the intakes of bioavailable iron and vitamin B12, except that while they also  
296 increased in all the optimized compared to observed diets, they nevertheless decreased among  
297 optimized diets along with GHGe reductions. The intake of phytates was also gradually decreased as  
298 GHGe was reduced, allowing meeting reference values for bioavailable iron and zinc. Calcium intake  
299 increased in all optimized diets, except in menopausal women.

### 300 3.7 Active constraints and sensitivity analysis

301 Analysis of the standardized dual values showed that the most limiting constraints were, in descending  
302 order, bioavailable zinc, EPA+DHA (eicosapentaenoic acid + docosahexaenoic acid), energy intake,  
303 sodium and saturated fatty acids in men and bioavailable zinc, EPA+DHA and sodium in women. The  
304 redistribution between ruminant meat and pork across modeling scenarios was driven by the  
305 compromise between satisfying the nutritional constraints for bioavailable zinc and iron and sodium  
306 requirements and the environmental constraint of GHGe reduction, as tested by alternative models  
307 where we allowed some flexibility in the requirements for each of these nutrients one by one (data not  
308 shown).

309 The sensitivity analysis also showed that the nutritional constraints for bioavailable zinc and iron were  
310 determinant for the distribution between meat and whole grains products having a phytate content that  
311 limit the zinc and iron bioavailabilities. Indeed, as shown in **Appendix F**, we verified that allowing  
312 some flexibility for bioavailable zinc led to meat reduction together with whole grains and starch  
313 foods increase (in men: 110 g/d *vs.* 2 g/d of whole grains and starch foods with *vs.* without flexibility,  
314 respectively, under the 50% reduction in GHGe scenario).

315 Moreover, as shown in **Appendix G**, the constraint on livestock co-products had little influence on the  
316 modeling results that were fairly similar with or without considering this constraint.

## 317 4. Discussion

318 In the present diet optimization study, the minimal changes in current French diets necessary to first  
319 meet nutrient reference values and then reduce GHGe by up to 50% were characterized by an overall  
320 decrease in the consumption of foods of animal origin with notably suppression of milk and strong  
321 reductions in dairy products and cheese, together with a stable but largely redistributed meat  
322 consumption in favor of poultry and pork and against ruminant meat, as well as marked increases in  
323 the consumptions of legumes. It should be noted that strong dietary changes were induced as soon as  
324 the first, basal stage consisting in modelling diets meeting nutrient reference values (under

325 acceptability and moderation constraints), without any reduction in GHGe (which were however  
326 constrained to avoid any increase). From this first stage, the consumption of animal products decreases  
327 and the model opted for organic plant products, which are more efficient than non-organic ones to  
328 limit GHGe. During the second stage, GHGe reductions by up to 50% mainly resulted from a  
329 redistribution between meat types against ruminant meat, within total consumptions of meat and  
330 animal products remaining relatively stable. Noteworthy, the model selects the most efficient farming  
331 practice for each food (organic or not) thus the entire optimized consumption of each food item is  
332 either organic or conventional which does not reflect the reality of consumer behavior.  
333 Notably, in addition to food behaviors, a major challenge to improve the sustainability of food systems  
334 is the reduction of losses and waste (44). The lack of quantitative data about waste for each food did  
335 not allow us to consider this dimension in our models. This is all the more complex, as waste occurs at  
336 each link of the food system chain and depends on both the production and processing methods  
337

338 This study, by considering environmental pressures associated with food production while accounting  
339 for farming practices, as well as numerous detailed food items, allowed intra-group substitutions by  
340 favoring less emitter foods. This brings new insights since nowadays most French consumers are  
341 unlikely to be ready to follow drastic plant-based diets such as vegetarian or vegan diets, that would  
342 represent a radical change in eating habits for the highest consumers of animal products, and would  
343 require steps over time. In the meantime, small, low-impact dietary changes for a large proportion of  
344 the population are probably more acceptable than substantial changes as strong changes may need  
345 more time (45). Overall, our results are coherent with literature findings comparing emissions from  
346 observed diets more or less rich in animal products, which have documented lower emissions for diets  
347 richer in plant foods (7,46–48). However, such observed diets do not necessarily meet the nutritional  
348 requirements.

349  
350 Optimization modelling enables to identify environmental-friendly diets in line with nutrient  
351 requirements, (e.g. by avoiding counter-productive effects such as increase in consumption of sweet  
352 and fat products) (15). Scientific literature using diet optimization for exploring potential GHGe  
353 reduction under nutrient constraints is plentiful (7–10,48). Overall, from these studies, it appears that a  
354 drastic and specific reduction of ruminant meat as well as dairy products consumption is the main  
355 lever for GHGe reduction from diet, which is in line with our results. We indeed found that dairy  
356 products and ruminant meat have to be drastically decreased, without being totally suppressed, which  
357 is somewhat different from the results of a recent diet optimization study that identified the need to  
358 completely eliminate ruminant meat while maintaining dairy products (excluding butter and cheese) to  
359 comply with the 2030 and 2050 GHGe reduction targets (being much stronger than those modelled  
360 here) in the Netherlands (49). However this study, as well as most of the others, did not take into  
361 account nutrient bioavailability in nutrient constraints and did not include coproduction constraints,

362 whereas these important parameters may shape the modelling results and the order of magnitude of  
363 potential decreases and increases according to food groups (9). Herein, as previously done (26–28), we  
364 have considered and controlled the bioavailability of iron and zinc using validated equations for their  
365 absorption. We have shown that considering the bioavailability of iron and zinc was crucial for the  
366 concomitant variations in meat and whole grains products, whereas considering beef-milk  
367 coproduction had little influence in our context.

368  
369 A wide heterogeneity exists in terms of methodological aspects across modelling studies (50). First,  
370 the number of food items can vary greatly and we worked with a relatively large number of food items  
371 (~250) (9), with the notable feature of allowing the choice (or a mix) between two modes of  
372 production (organic or conventional) for each food item. Second, contrary to what has been most often  
373 done, we have considered constraints on food groups but not on food items so as to allow intra-group  
374 substitutions. These acceptability constraints based on the 95<sup>th</sup> percentile of each population, including  
375 participants with healthier diets than the general population, allowed stronger increases in some food  
376 groups. Finally, we have adopted a quadratic rather than a linear formulation for the objective function  
377 of diet departure to minimize, so as to favor more numerous but smaller changes rather than fewer but  
378 larger changes during optimization (51). All these methodological choices have provided levers for  
379 optimized diets, since we had a wide inventory of food items and since intra-group substitutions were  
380 favored by different means as modeled here.

381  
382 In our particular context, under all the considered nutritional and acceptability constraints and by  
383 accounting for the influence of anti-nutritional factors like phytate on zinc and iron bioavailabilities,  
384 total meat was maintained relatively stable, because of a decrease in whole grains and starchy foods  
385 (and thus a decrease in phytate), although it was qualitatively remodeled in disfavor of ruminant meat  
386 so as to reduce GHGe. In line with our results, a diet optimization study among old Dutch adults with  
387 50% reduction in GHGs found unchanged total meat consumption with an increased contribution of  
388 poultry and pork and a decreased contribution of beef (52).

389  
390 Several options regarding plant foods merit further discussion. In the optimized diets, non-alcoholic  
391 beverages (including coffee and tea) were strongly reduced (up to -54%), as they are poor in nutrients  
392 and represent important environmental pressures at the post-production stage. However, as culturally  
393 deeply entrenched in our usual diets, such drops could be an important limitation and all the more so  
394 since positive health effects have been reported (53). It should be noted that whole grains and starchy  
395 foods, whose beneficial role on health is well-documented (54), were lowered in optimized diets  
396 (almost in men), and this may be ascribed to their phytates content limiting the iron and zinc  
397 bioavailability. Such a prominent role in optimized solutions raises the issue related to nutritional  
398 constraints relying on nutritional references which are based on calculated physiological requirements

399 as for zinc, for instance, reliable biomarkers are lacking (55). Thus, while we have defined the nutrient  
400 constraints according to the French nutritional references and the literature equations for  
401 bioavailability, the methods of definition may be highly conservative and slightly lower intakes may  
402 not result in clear adverse effect on health, such as over-deficiency. Of note, the nutritional reference  
403 for fiber (i.e. >30g/d) favors the increase in foods with high content such as legumes, while reducing  
404 the ruminant meat for reducing GHGe required a reduction in phytate intake to allow sufficient  
405 absorption of iron and zinc, which in turn has favored the reduction of starchy foods. This was clearly  
406 illustrated by an alternative model allowing flexibility on zinc, with which whole grains foods did  
407 increase. Moreover, in the optimized diets, fish consumption was limited by an upper value. Fish and  
408 seafood are the major supplier of EPA and DHA that should be consumed at the highest level of their  
409 reference value. This reflects the fact that EPA+DHA is a limiting constraint for more sustainable diets  
410 and suggest that other and presumably better sustainable diets might be identified when introducing  
411 other new sources of these fatty acids.

412  
413 Finally, the proportion of organic foods drastically increased (in weight) from ~30% in the observed  
414 diets to ~70% in all the optimized diets, which explained the monetary cost increase of optimized diets  
415 (56). At the individual diet level, we previously reported that regular organic food consumers exhibit  
416 diets with a lower impact regarding GHGe, land use and energy demand but dietary patterns (i.e.  
417 plant-based patterns) prevailed on the mode of production in this association (22). However, at the  
418 food item level, organic farming may play a substantial role in reducing GHGe, depending on the food  
419 considered (23,57,58). Our results illustrate that when optimizing diets by selecting specific products  
420 like in the present study, rather than by only substituting some conventional by organic products at  
421 constant diet, as some consumers can do, organic foods can greatly help to the reduction of GHGe, as  
422 previously shown in our observational studies (59). The consumption of organic foods increases from  
423 the first step (i.e. 0% reduction in GHGe corresponding to modeling diet with  $\text{GHGe} \leq \text{observed}$   
424 value) which means that the foods preferentially selected to respect the nutritional constraints are more  
425 efficient as organic to maintain GHGe. However, due to modeling, one food is selected 100% as  
426 organic or 100% as conventional what does not reflect the reality of behaviors. This is interesting as an  
427 increased consumption of organic foods can have beneficial consequences on two levels. First, on the  
428 environmental level as organic production systems also exhibit improved energy efficiency (60), better  
429 biophysics and biological quality of soils (57,61) and are valuable for plant and animal biodiversity  
430 (57,60,61). Second, on the sanitary level, as high plant-based diets based on organic agriculture may  
431 lead to much lower exposure to pesticide residues (22,62,63), motivating the promotion of plant  
432 products produced without synthetic pesticides in the new French food-based dietary guidelines (64).

433  
434 As regards the obtained solutions, optimized diets exhibited high consumption of fruits and vegetables  
435 (>500g/d), low consumption of red meat (<500g/week), processed meat (<150g/week), sweet

436 products, low intake in salt and moderated consumption of dairy products, in line with the French  
437 food-based guidelines (42). Importantly, consumption of legumes among menopausal women was  
438 somewhat low and the consumption of whole grains and starchy foods was very low in men. As  
439 scientific literature has documented a notable beneficial role of plant-food diets, beyond fiber intake  
440 which are controlled in the present study, for health and environment (3), further steps of the  
441 transition, probably further away from the observed diets would require to introduce a higher plant  
442 versus animal food ratio. For instance, in the same cohort we previously described low emitting  
443 dietary patterns (GHGe for production lower than about 2.2 kg CO<sub>2</sub> eq) that were richer in plant-based  
444 food than the present solution but nutritional adequacy was not assessed (24). Higher shares of healthy  
445 plant protein such as whole grains and lower consumption of animal protein as red meat are  
446 considered as part of a healthy diet as documented by epidemiological data (65,66) and may be  
447 warranted for a full sustainable transition together with GHGe. Finally, the nutritional values of highly  
448 plant-based diets should be tested in the future by deleting or relaxing acceptability constraints,  
449 considering alignment with healthy eating patterns as defined from epidemiological data or by using  
450 hierarchical optimization as we did in a recent study (67). Finally, it has been previously documented  
451 that healthier diets are often more expensive (68). In line with this, the optimized diets were more  
452 expensive than the observed diet, constituting a potential barrier for some consumers. Without  
453 appropriate policies, this may jeopardize food security due to inaccessibility and potential low  
454 availability for vulnerable populations.

455 Some limitations of our work should be highlighted. First, the NutriNet-Santé cohort included  
456 volunteers who were probably more concerned by health and diet issues than the general population,  
457 limiting extrapolation to the general population as these participants exhibit diet rich in plant-based  
458 food. For instance, lowering energy intake is a well-known lever for reducing GHGe (69,70) but in  
459 this population including “small eaters”, energy intake increased in the basal model to reach the  
460 requirement. Second, post-farm environmental pressures for organic agriculture are lacking, thus life  
461 cycle assessments were limited to farm activities which have most impacts in the food system.  
462 Therefore, our scenarios may be insufficient to meet the global climatic objective, since some steps  
463 following food production were not considered. Concerning environmental indicators, LCA were used  
464 while it is recognized that some ecosystem services related to agroecological practices are misestimate  
465 by this method (25). Third, beyond GHGe, we considered two other environmental pressures for  
466 descriptive purpose, those three allowing an acceptable representativeness of the overall  
467 environmental impact (43), but other dimensions such as water use or biodiversity should be studied.  
468 However, in further works, it would be very important to consider water use in particular in the  
469 context of vegetable and fruit and the production of corn, mainly for feeding monogastric livestock  
470 breeding. We observed a decrease in land occupation with the gradual reduction in GHGe.  
471 Reallocation of released land may induce important fluctuations in GHGe which are the results of  
472 carbon balance of managed forests, agricultural soil organic carbon stocks soil and reallocation



473 (grassland, deforestation, afforestation, artificialization etc.)(71). But, an important factor that was not  
474 considered is land use reallocation since this analysis used attributional LCAs. Thus, the change in the  
475 type of meat consumed would have also an effect on the demand for arable land and therefore on  
476 carbon stocks and on GHGs (72). Notably, in addition to food behaviors, a major challenge to improve  
477 the sustainability of food systems is the food losses and waste reduction (44). The lack of quantitative  
478 data about waste for each food did not allow us to consider this important dimension in our models.  
479 This is all the more complex, as waste occurs at each link of the food system chain and depends on  
480 both the type production and level of processing. The reallocation of permanent grasslands is also an  
481 issue We have also assumed, as in most diet optimization studies (9), that the most acceptable diets are  
482 those the closest to the observed diets. While this classical assumption makes it possible to define a  
483 simple and very restrictive metric of cultural acceptability, it is known to account only very  
484 imperfectly for true acceptability as stronger dietary changes may occur, at least in certain segments of  
485 the population. Besides, this study integrates many strengths such as the level of detail for food  
486 consumption, the detailed and reliable consideration of the updated nutritional recommendations  
487 (including bioavailability of the micronutrients of concern in our context, iron and zinc, which is  
488 seldom done), the consideration of different food production methods and the corresponding  
489 environmental indicators.

490

491 In conclusion, this study in adults provides detailed results on the possible dietary changes that can be  
492 implemented to mitigate GHGe up to 50% with minimal departure from the observed diet. We were  
493 here able to identify more sustainable diets, being nutritionally adequate and culturally acceptable, and  
494 from which meat was not excluded. Because the present optimized nutrition model preferentially  
495 allowed intra-category substitutions, the plant/animal food ratio was not noticeably altered. Although  
496 adequate according to a large set of lower and upper nutrient reference values, the modelled diets may  
497 be sub-optimal for long-term health, which may benefit from further decrease in red meat and higher  
498 increases in whole grains. Furthermore, reducing the consumption of foods of animal origin,  
499 particularly beef and lamb, as well as milk and dairies, is necessary not only for environmental or  
500 health reasons but also for animal welfare considerations. Lastly, future research will be needed to  
501 document even greater reductions as this 50% is unlikely to be sufficient and further research focusing  
502 on specific subgroups, e.g. according to age or socioeconomic status would be of interest to fine-tune  
503 the optimized diet.

504

### 505 **Acknowledgements**

506 We thank Cédric Agaesse (dietitian); Thi Hong Van Duong, Younes Esseddik (IT manager), Régis  
507 Gatibelza, Djamal Lamri, Jagatjit Mohinder and Aladi Timera (computer scientists); Julien Allegre,  
508 Nathalie Arnault, Laurent Bourhis and Fabien Szabo de Edelenyi, PhD (supervisor) (data-  
509 manager/statisticians) for their technical contribution to the NutriNet-Santé study and Nathalie

510 Druesne-Pecollo, PhD (operational coordination). We thank all the volunteers of the NutriNet-Santé  
511 cohort.

## 512 **5. The authors' contributions are as follows:**

513 EKG, BA, MT, and SH conducted the study.

514 EKG, JB, BL, SH, DL and PP conducted the research and implemented databases.

515 EKG, HF, AD, and FM conducted the diet optimization.

516 EKG performed statistical analyses and drafted the manuscript.

517 All authors critically helped in the interpretation of results, revised the manuscript and provided relevant  
518 intellectual input. They all read and approved the final manuscript.

519 EKG had primary responsibility for the final content, she is the guarantor.

## 520 **6. Conflict of Interest**

521 No author declared conflict of interest.

## 522 **7. Funding**

523 The NutriNet-Santé study is funded by French Ministry of Health and Social Affairs, Santé Publique  
524 France, Institut National de la Santé et de la Recherche Médicale, Institut National de la Recherche  
525 Agronomique, Conservatoire National des Arts et Métiers, and Sorbonne Paris Nord University. The  
526 BioNutriNet project was supported by the French National Research Agency (Agence Nationale de la  
527 Recherche) in the context of the 2013 Programme de Recherche Systèmes Alimentaires Durables  
528 (ANR-13-ALID-0001). The funders had no role in the study design, data collection, analysis,  
529 interpretation of data, preparation of the manuscript, and decision to submit the paper.

## 530 8. References

- 531 1. The intergovernmental panel on climate change (IPCC). Global Warming of 1.5 °C [Internet].  
532 [cited 2019 Jul 18]. Available from: <https://www.ipcc.ch/sr15/>
- 533 2. Food production is responsible for one-quarter of the world's greenhouse gas emissions  
534 [Internet]. Our World in Data. [cited 2020 Jun 5]. Available from:  
535 <https://ourworldindata.org/food-ghg-emissions>
- 536 3. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the  
537 Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems.  
538 *Lancet*. 2019 Jan 16;393(10170):447–92.
- 539 4. Tilman D, Clark M. Global diets link environmental sustainability and human health. *Nature*.  
540 2014 Nov 27;515(1476-4687 (Electronic)):518–22.
- 541 5. Couturier C, Charru M, Doublet S, Pointereau P. The Afterres 2050 le scénario [Internet]. 2016  
542 [cited 2020 Jun 8]. Available from: [https://afterres2050.solagro.org/wp-](https://afterres2050.solagro.org/wp-content/uploads/2020/02/Afterres2050-eng.pdf)  
543 [content/uploads/2020/02/Afterres2050-eng.pdf](https://afterres2050.solagro.org/wp-content/uploads/2020/02/Afterres2050-eng.pdf)
- 544 6. Springmann M, Wiebe K, Mason-D'Croz D, Sulser TB, Rayner M, Scarborough P. Health and  
545 nutritional aspects of sustainable diet strategies and their association with environmental  
546 impacts: a global modelling analysis with country-level detail. *Lancet Planet Health*.  
547 2018;2(10):e451–61.
- 548 7. Aleksandrowicz L, Green R, Joy EJM, Smith P, Haines A. The Impacts of Dietary Change on  
549 Greenhouse Gas Emissions, Land Use, Water Use, and Health: A Systematic Review. Wiley AS,  
550 editor. *PLOS ONE*. 2016 Nov 3;11(11):e0165797.
- 551 8. Perignon M, Vieux F, Soler L-G, Masset G, Darmon N. Improving diet sustainability through  
552 evolution of food choices: review of epidemiological studies on the environmental impact of  
553 diets. *Nutrition Reviews*. 2017 Jan;75(1):2–17.
- 554 9. van Dooren C. A Review of the Use of Linear Programming to Optimize Diets, Nutritiously,  
555 Economically and Environmentally. *Front Nutr*. 2018;5:48.
- 556 10. Wilson N, Cleghorn CL, Cobiack LJ, Mizdrak A, Nghiem N. Achieving Healthy and Sustainable  
557 Diets: A Review of the Results of Recent Mathematical Optimization Studies. *Adv Nutr*. 2019  
558 01;10(Suppl\_4):S389–403.
- 559 11. Lindgren E, Harris F, Dangour AD, Gasparatos A, Hiramatsu M, Javadi F, et al. Sustainable  
560 food systems-a health perspective. *Sustain Sci*. 2018;13(6):1505–17.
- 561 12. Springmann M, Godfray HCJ, Rayner M, Scarborough P. Analysis and valuation of the health  
562 and climate change cobenefits of dietary change. *PNAS*. 2016 Apr 12;113(15):4146–51.
- 563 13. Hemler EC, Hu FB. Plant-Based Diets for Personal, Population, and Planetary Health. *Adv Nutr*.  
564 2019 Nov 1;10(Supplement\_4):S275–83.
- 565 14. Clark MA, Springmann M, Hill J, Tilman D. Multiple health and environmental impacts of  
566 foods. *PNAS*. 2019 Nov 12;116(46):23357–62.
- 567 15. Stoll-Kleemann S, Schmidt UJ. Reducing meat consumption in developed and transition  
568 countries to counter climate change and biodiversity loss: a review of influence factors. *Reg*  
569 *Environ Change*. 2017 Jun;17(5):1261–77.

- 570 16. Hartmann C, Siegrist M. Consumer perception and behaviour regarding sustainable protein  
571 consumption: A systematic review. *Trends in Food Science & Technology*. 2017 Mar 1;61:11–  
572 25.
- 573 17. Manners R, Blanco-Gutiérrez I, Varela-Ortega C, Tarquis AM. Transitioning European Protein-  
574 Rich Food Consumption and Production towards More Sustainable Patterns—Strategies and  
575 Policy Suggestions. *Sustainability*. 2020 Jan;12(5):1962.
- 576 18. Graça J, Godinho CA, Truninger M. Reducing meat consumption and following plant-based  
577 diets: Current evidence and future directions to inform integrated transitions. *Trends in Food  
578 Science & Technology*. 2019 Sep 1;91:380–90.
- 579 19. de Gavelle E, Davidenko O, Fouillet H, Delarue J, Darcel N, Huneau J-F, et al. Self-declared  
580 attitudes and beliefs regarding protein sources are a good prediction of the degree of transition to  
581 a low-meat diet in France. *Appetite*. 2019 Nov 1;142:104345.
- 582 20. Perignon M, Masset G, Ferrari G, Barré T, Vieux F, Maillot M, et al. How low can dietary  
583 greenhouse gas emissions be reduced without impairing nutritional adequacy, affordability and  
584 acceptability of the diet? A modelling study to guide sustainable food choices. *Public Health  
585 Nutrition*. 2016 Oct;19(14):2662–74.
- 586 21. Kurian M. The water-energy-food nexus: Trade-offs, thresholds and transdisciplinary  
587 approaches to sustainable development. *Environmental Science & Policy*. 2017 Feb 1;68:97–  
588 106.
- 589 22. Baudry J, Pointereau P, Seconda L, Vidal R, Taupier-Letage B, Langevin B, et al. Improvement  
590 of diet sustainability with increased level of organic food in the diet: findings from the  
591 BioNutriNet cohort. *Am J Clin Nutr*. 2019 Apr 1;109(4):1173–88.
- 592 23. Clark M, Tilman D. Comparative analysis of environmental impacts of agricultural production  
593 systems, agricultural input efficiency, and food choice. *Environmental Research Letters*. 2017  
594 Jun 1;12(6):064016.
- 595 24. Seconda L, Baudry J, Alles B, Boizot-Szantai C, Soler L-G, Galan P, et al. Comparing  
596 nutritional, economic, and environmental performances of diets according to their levels of  
597 greenhouse gas emissions. *Clim Change*. 2018 May;148(1–2):155–72.
- 598 25. van der Werf HMG, Knudsen MT, Cederberg C. Towards better representation of organic  
599 agriculture in life cycle assessment. *Nature Sustainability*. 2020 Jun;3(6):419–25.
- 600 26. Barré T, Perignon M, Gazan R, Vieux F, Micard V, Amiot M-J, et al. Integrating nutrient  
601 bioavailability and co-production links when identifying sustainable diets: How low should we  
602 reduce meat consumption? *PLoS ONE*. 2018;13(2):e0191767.
- 603 27. Ferguson EL, Darmon N, Briend A, Premachandra IM. Food-based dietary guidelines can be  
604 developed and tested using linear programming analysis. *J Nutr*. 2004 Apr;134(4):951–7.
- 605 28. Deptford A, Allieri T, Childs R, Damu C, Ferguson E, Hilton J, et al. Cost of the Diet: a method  
606 and software to calculate the lowest cost of meeting recommended intakes of energy and  
607 nutrients from local foods. *BMC Nutrition*. 2017 Mar 14;3(1):26.
- 608 29. European Commission. Paris Agreement [Internet]. Climate Action - European Commission.  
609 2016 [cited 2021 Feb 5]. Available from:  
610 [https://ec.europa.eu/clima/policies/international/negotiations/paris\\_en](https://ec.europa.eu/clima/policies/international/negotiations/paris_en)

- 611 30. Hercberg S, Castetbon K, Czernichow S, Malon A, Mejean C, Kesse E, et al. The Nutrinet-Sante  
612 Study: a web-based prospective study on the relationship between nutrition and health and  
613 determinants of dietary patterns and nutritional status. *BMC Public Health*. 2010;10(1471–  
614 2458):242.
- 615 31. Vergnaud A-C, Touvier M, Méjean C, Kesse-Guyot E, Pollet C, Malon A, et al. Agreement  
616 between web-based and paper versions of a socio-demographic questionnaire in the NutriNet-  
617 Santé study. *Int J Public Health*. 2011 Aug 1;56(4):407–17.
- 618 32. Touvier M, Mejean C, Kesse-Guyot E, Pollet C, Malon A, Castetbon K, et al. Comparison  
619 between web-based and paper versions of a self-administered anthropometric questionnaire.  
620 *EurJEpidemiol*. 2010 May;25:287–96.
- 621 33. Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a  
622 study of concurrent and construct validity. *Public Health Nutrition* [Internet]. 2006 Sep [cited  
623 2016 Apr 28];9(06). Available from:  
624 [http://www.journals.cambridge.org/abstract\\_S1368980006001261](http://www.journals.cambridge.org/abstract_S1368980006001261)
- 625 34. Baudry J, Méjean C, Allès B, Péneau S, Touvier M, Hercberg S, et al. Contribution of Organic  
626 Food to the Diet in a Large Sample of French Adults (the NutriNet-Santé Cohort Study).  
627 *Nutrients*. 2015;7(10):8615–32.
- 628 35. Council Regulation (EC) No 834/2007 of 28 June 2007 on organic production and labelling of  
629 organic products and repealing Regulation (EEC) No 2092/91 [Internet]. Available from:  
630 <https://eur-lex.europa.eu/eli/reg/2007/834/oj>
- 631 36. Etude Nutrinet-Santé. Table de composition des aliments de l'étude Nutrinet-Santé (Nutrinet-  
632 Santé Study Food Composition Database). Paris: Economica. 2013;
- 633 37. Pointereau P, Langevin B, Gimaret M. DIALECTE, a comprehensive and quick tool to assess  
634 the agro-environmental performance of farms. In 2012. Available from:  
635 <http://ifsa.boku.ac.at/cms/index.php?id=ifsa2012>
- 636 38. Idèle. Chiffres clés Bovins 2016. 2016.
- 637 39. France Agrimer. Filière bovine, quotas laitiers. 2012 [cited 2020 Sep 22];12. Available from:  
638 [https://www.franceagrimer.fr/fam/content/download/14506/document/filière bovine, quotas  
639 laitiers février 2012 A4.pdf](https://www.franceagrimer.fr/fam/content/download/14506/document/filière_bovine_quotas_laitiers_février_2012_A4.pdf)
- 640 40. French Agency for Food, Environmental and Occupational Health Safety (Anses). Actualisation  
641 des repères du PNNS : élaboration des références nutritionnelles [Internet]. Maison Alfort:  
642 ANSES; 2016 Dec. Available from: Available from:  
643 <https://www.anses.fr/fr/system/files/NUT2012SA0103Ra-2.pdf>
- 644 41. Dietary Reference Values | DRV Finder [Internet]. EFSA. [cited 2021 Jan 22]. Available from:  
645 <https://www.efsa.europa.eu/en/interactive-pages/drvs>
- 646 42. HCSP. Statement related to the revision of the 2017-2021 French Nutrition and Health  
647 Programme's dietary guidelines for adults [Internet]. Paris: Haut Conseil de la Santé Publique;  
648 2017 Feb [cited 2019 Feb 5]. Available from:  
649 <https://www.hcsp.fr/explore.cgi/avisrapportsdomaine?clefr=653>
- 650 43. Kramer GF, Tyszler M, Veer PV, Blonk H. Decreasing the overall environmental impact of the  
651 Dutch diet: how to find healthy and sustainable diets with limited changes. *Public Health Nutr*.  
652 2017 Jun;20(9):1699–709.

- 653 44. HLPE. Food losses and waste in the context of sustainable food systems. A report by the High  
654 Level Panel of Experts on Food Security and Nutrition of the Committee on World Food  
655 Security. Rome, Italy; 2014 p. 117.
- 656 45. Mariotti F, editor. Vegetarian and Plant-Based Diets in Health and Disease Prevention [Internet].  
657 Elsevier; 2017 [cited 2021 Feb 14]. Available from:  
658 <https://linkinghub.elsevier.com/retrieve/pii/C20150003999>
- 659 46. Chai BC, van der Voort JR, Grofelnik K, Eliasdottir HG, Klöss I, Perez-Cueto FJA. Which Diet  
660 Has the Least Environmental Impact on Our Planet? A Systematic Review of Vegan, Vegetarian  
661 and Omnivorous Diets. *Sustainability*. 2019 Jan;11(15):4110.
- 662 47. Macdiarmid JI, Whybrow S. Nutrition from a climate change perspective. *Proceedings of the*  
663 *Nutrition Society*. undefined/ed;1–8.
- 664 48. Auestad N, Fulgoni VL III. What current literature tells us about sustainable diets: emerging  
665 research linking dietary patterns, environmental sustainability, and economics. *Adv Nutr*. 2015  
666 Jan;6(2156-5376 (Electronic)):19–36.
- 667 49. Broekema R, Tyszler M, van 't Veer P, Kok FJ, Martin A, Lluch A, et al. Future-proof and  
668 sustainable healthy diets based on current eating patterns in the Netherlands. *Am J Clin Nutr*.  
669 2020 Nov 11;112(5):1338–47.
- 670 50. Gazan R, Brouzes CMC, Vieux F, Maillot M, Lluch A, Darmon N. Mathematical Optimization  
671 to Explore Tomorrow's Sustainable Diets: A Narrative Review. *Adv Nutr*. 2018 Sep 1;9(5):602–  
672 16.
- 673 51. Persson M, Fagt S, Pires SM, Poulsen M, Vieux F, Nauta MJ. Use of Mathematical Optimization  
674 Models to Derive Healthy and Safe Fish Intake. *The Journal of Nutrition*. 2018 Feb  
675 1;148(2):275–84.
- 676 52. Grasso AC, Olthof MR, van Dooren C, Broekema R, Visser M, Brouwer IA. Protein for a  
677 Healthy Future: How to Increase Protein Intake in an Environmentally Sustainable Way in Older  
678 Adults in the Netherlands. *The Journal of Nutrition*. 2021 Jan 4;151(1):109–19.
- 679 53. Poole R, Kenned O, Roderick P, Fallowfield J, Haye P, Parkes J. Coffee consumption and  
680 health: umbrella review of meta-analyses of multiple health outcomes. *BMJ* [Internet]. 2018 Jan  
681 12 [cited 2020 Dec 8];360. Available from:  
682 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5765813/>
- 683 54. McRae MP. Health Benefits of Dietary Whole Grains: An Umbrella Review of Meta-analyses. *J*  
684 *Chiropr Med*. 2017 Mar;16(1):10–8.
- 685 55. Bornhorst J, Kipp AP, Haase H, Meyer S, Schwerdtle T. The crux of inept biomarkers for risks  
686 and benefits of trace elements. *TrAC Trends in Analytical Chemistry*. 2018 Jul 1;104:183–90.
- 687 56. Boizot-Szantai C, Hamza O, Soler L-G. Organic consumption and diet choice: An analysis based  
688 on food purchase data in France. *Appetite*. 2017 01;117:17–28.
- 689 57. Tuomisto HL, Hodge ID, Riordan P, Macdonald DW. Does organic farming reduce  
690 environmental impacts? – A meta-analysis of European research. *Journal of Environmental*  
691 *Management*. 2012 Dec;112:309–20.
- 692 58. Meier MS, Stoessel F, Jungbluth N, Juraske R, Schader C, Stolze M. Environmental impacts of  
693 organic and conventional agricultural products--are the differences captured by life cycle  
694 assessment? *J Environ Manage*. 2015 Feb 1;149:193–208.

- 695 59. Lacour C, Seconda L, Allès B, Hercberg S, Langevin B, Pointereau P, et al. Environmental  
696 Impacts of Plant-Based Diets: How Does Organic Food Consumption Contribute to  
697 Environmental Sustainability? *Front Nutr.* 2018;5:8.
- 698 60. Reganold JP, Wachter JM. Organic agriculture in the twenty-first century. *Nature Plants.* 2016  
699 Feb 3;2(2):15221.
- 700 61. Gomiero T, Pimentel D, Paoletti MG. Environmental Impact of Different Agricultural  
701 Management Practices: Conventional vs. Organic Agriculture. *Critical Reviews in Plant  
702 Sciences.* 2011 Jan;30(1–2):95–124.
- 703 62. Fleury S, Rivière G, Allès B, Kesse-Guyot E, Méjean C, Hercberg S, et al. Exposure to  
704 contaminants and nutritional intakes in a French vegetarian population. *Food Chem Toxicol.*  
705 2017 Jul 25;109(Pt 1):218–29.
- 706 63. Rebouillat P, Vidal R, Cravedi J-P, Taupier-Letage B, Debrauwer L, Gamet-Payraastre L, et al.  
707 Estimated dietary pesticide exposure from plant-based foods using NMF-derived profiles in a  
708 large sample of French adults. *Eur J Nutr.* 2020 Jul 30;
- 709 64. Santé publique France - Santé publique France présente les nouvelles recommandations sur  
710 l'alimentation, l'activité physique et la sédentarité [Internet]. [cited 2019 Mar 1]. Available  
711 from: [https://www.santepubliquefrance.fr/Accueil-Presses/Tous-les-communiqués/Sante-  
712 publique-France-presente-les-nouvelles-recommandations-sur-l-alimentation-l-activite-physique-  
713 et-la-sedentarite](https://www.santepubliquefrance.fr/Accueil-Presses/Tous-les-communiqués/Sante-publique-France-presente-les-nouvelles-recommandations-sur-l-alimentation-l-activite-physique-et-la-sedentarite)
- 714 65. Mariotti F. Animal and Plant Protein Sources and Cardiometabolic Health. *Adv Nutr.* 2019  
715 01;10(Suppl\_4):S351–66.
- 716 66. Naghshi S, Sadeghi O, Willett WC, Esmailzadeh A. Dietary intake of total, animal, and plant  
717 proteins and risk of all cause, cardiovascular, and cancer mortality: systematic review and dose-  
718 response meta-analysis of prospective cohort studies. *BMJ.* 2020 Jul 22;370:m2412.
- 719 67. Seconda L, Fouillet H, Huneau J-F, Pointereau P, Baudry J, Langevin B, et al. Conservative to  
720 disruptive diets for optimizing nutrition, environmental impacts and cost in French adults from  
721 the NutriNet-Santé cohort. *Nature Food.* 2021 Mar 11;1:1–9.
- 722 68. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic  
723 disparities in diet quality and health: a systematic review and analysis. *Nutr Rev.* 2015  
724 Oct;73(10):643–60.
- 725 69. Nelson ME, Hamm MW, Hu FB, Abrams SA, Griffin TS. Alignment of Healthy Dietary  
726 Patterns and Environmental Sustainability: A Systematic Review<sup>12</sup>. *Adv Nutr.* 2016 Nov  
727 10;7(6):1005–25.
- 728 70. Vieux F, Darmon N, Touazi D, Soler L, Soler LG. Greenhouse gas emissions of self-selected  
729 individual diets in France: Changing the diet structure or consuming less? *EcolEcon.*  
730 2012;75:91–101.
- 731 71. Smith P, House JI, Bustamante M, Sobocká J, Harper R, Pan G, et al. Global change pressures  
732 on soils from land use and management. *Global Change Biology.* 2016;22(3):1008–28.
- 733 72. Garnett T. Livestock-related greenhouse gas emissions: impacts and options for policy makers.  
734 *Environmental Science & Policy.* 2009 Jun 1;12(4):491–503.

735

736 **Figure 1: Overall composition (g/d) of the observed and optimized diets according to the**  
737 **modelling scenario and population group (color figure)**

738 Main food groups intakes (g/d) in the observed diets and in the diets being nutritionally, culturally and  
739 environmentally optimized so as to ensure gradual reduction in greenhouse gas emissions (GHGe)  
740 from 0 to 50%. Abbreviations: Obs, observed diet. A: men, B: premenopausal women, C: menopausal  
741 women.

742 **Figure 2: Composition in animal-based foods in the observed and optimized diets**  
743 **according to the modelling scenario and population group (color figure)**

744 Detailed animal foods intakes (g/d) in the observed diets and in the diets being nutritionally, culturally  
745 and environmentally optimized so as to ensure gradual reduction in greenhouse gas emissions (GHGe)  
746 from 0 to 50%. Abbreviations: Obs, observed diet. A: men, B: premenopausal women, C: menopausal  
747 women.

748

749 **Figure 3: Composition in plant-based foods in the observed and optimized diets according**  
750 **to the modelling scenario and population group (color figure)**

751 Detailed plant-based foods intakes (g/d) in the observed diets and in the diets being nutritionally,  
752 culturally and environmentally optimized so as to ensure gradual reduction in greenhouse gas  
753 emissions (GHGe) from 0 to 50%. Abbreviations: Obs, observed diet. A: men, B: premenopausal  
754 women, C: menopausal women.