



**HAL**  
open science

# The thermal response of small and shallow lakes to climate change: new insights from 3D hindcast modelling

Francesco Piccioni, Céline Casenave, Bruno J. Lemaire, Patrick Le Moigne, Philippe Dubois, Brigitte Vinçon-Leite

## ► To cite this version:

Francesco Piccioni, Céline Casenave, Bruno J. Lemaire, Patrick Le Moigne, Philippe Dubois, et al.. The thermal response of small and shallow lakes to climate change: new insights from 3D hindcast modelling. 2021. hal-03247403

**HAL Id: hal-03247403**

**<https://hal.inrae.fr/hal-03247403v1>**

Preprint submitted on 3 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# The response of small and shallow lakes to climate change: new insights from hindcast modelling.

Francesco Piccioni<sup>1</sup>, Céline Casenave<sup>2</sup>, Bruno Jacques Lemaire<sup>1,3</sup>, Patrick Le Moigne<sup>4</sup>,  
Philippe Dubois<sup>1</sup>, and Brigitte Vinçon-Leite<sup>1</sup>

<sup>1</sup>LEESU, Ecole des Ponts ParisTech, Univ Paris Est Créteil, Marne-la-Vallée, France

<sup>2</sup>MISTEA, Université Montpellier, INRAE, Institut Agro, Montpellier, France

<sup>3</sup>AgroParisTech, Paris, France

<sup>4</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

**Correspondence:** Francesco Piccioni (francesco.piccioni@enpc.fr)

**Abstract.** Small and shallow water bodies are a dominant portion of inland freshwaters. However, the effects of climate change on such ecosystems have rarely been quantitatively addressed. We propose a methodology to evaluate the thermal response of a small and shallow lake to long-term changes in the meteorological conditions, through model simulations. To do so, a 3D hydrodynamic model is forced with meteorological data and used to hindcast the evolution of a urban lake in the Paris region between 1960 and 2017. Its thermal response is analyzed through the definition of a series of indices describing its thermal regime in terms of water temperature, thermal stratification and tendency to biomass production. Model results and meteorological forcing are analyzed over time to test the presence of monotonic trends and 3D simulations are exploited to highlight spatial patterns in the dynamics of stratification. The thermal regime of the study site underwent significant changes. Its response was highly correlated with three meteorological variables: air temperature, solar radiation and wind speed. Mean annual water temperature showed a considerable warming trend of 0.6°C/dec, accompanied by longer stratification and by an increase of thermal energy available for biomass production. Water warming was significant during all four seasons, with maxima in Spring and Summer, while stratification and energy for phytoplankton growth increased especially during Spring and Autumn. Stratification only established in the deeper areas of the water body, possibly inducing heterogeneity in the release of nutrient from the sediment and in the development of harmful algal blooms. Numerous similar ecosystems might be experiencing analogous changes, and appropriate management policies are needed to preserve their ecological value.

## 1 Introduction

Lakes and reservoirs represent 3.7% of the Earth's non-glaciated continental area (Verpoorter et al., 2014), and often act as "sentinels" of climate change (Adrian R. et al., 2009). They have experienced considerable warming along the past decades (O'Reilly et al., 2015; Schmid et al., 2014; Schneider and Hook, 2010; Piccolroaz et al., 2020), sometimes even accelerated in respect to the surrounding areas (Schneider et al., 2009). Climate change is expected to further deteriorate the ecological status of a number of lakes worldwide that already suffer from eutrophication. In particular, changes in water temperature and in the patterns of thermal stratification could have a strong influence on the development of harmful algal blooms. Warmer



water temperatures might favor the dominance of certain algal groups, such as cyanobacteria, whose increasing occurrence is a great concern in the management of water resources (Paerl and Huisman, 2008; Paerl and Paul, 2012; Wagner and Erickson, 2017). Furthermore, changes in the stratification and mixing regime could alter nutrients and light availability, sedimentation rates, and enhance the risk of hypolimnetic oxygen depletion (Song et al., 2013; Wilhelm and Adrian, 2008; Jankowski et al., 2006; Winder and Sommer, 2012).

The global areal extent of lakes and impoundments is dominated by millions of water bodies smaller than 1 km<sup>2</sup> (Downing et al., 2006). However, their role in climate change studies has often been overlooked. On the first hand, they might be relevant on a global scale in elemental budgets, such as the carbon budget (Mendonça et al., 2017), and should be taken into account in large-scale climate change analysis (Downing et al., 2006). On the other hand, the impact of climate change on small and shallow water bodies has rarely been quantitatively assessed, even though they play an important role for biodiversity and are prone to harmful algal blooms (Biggs et al., 2016; Wilkinson et al., 2020). Furthermore, with the advance of urbanization, the presence of aquatic environments has become a key feature for the improvement of life quality in the urban landscape (Frumkin et al., 2017; van den Bosch and Sang, 2017). Often small and shallow, urban lakes grant valuable ecosystem services and contribute to the preservation of biodiversity (Frumkin et al., 2017; Hill et al., 2017; Hassall, 2014; Higgins et al., 2019). However, to our best knowledge, only very few studies can be found on the effect of climate change on small and shallow water bodies (Biggs et al., 2016). This lack of scientific studies is mirrored in a general lack of long term *in situ* data, making it impossible to directly analyze how these environments respond to climate change solely through observations. Conversely, long term meteorological data are available for most regions of the globe (e.g. global or regional reanalysis), as a result of a network of systematic observations that developed consistently since the beginning of the 20th century.

Hydrodynamic models have been vastly used to simulate lake and reservoir dynamics over both short and long periods in order to test changes in systems subject to given meteorological and border conditions, often through one-dimensional simulations. Model results can be used to fill in gaps in sporadic series of observation or to gain knowledge outside of the observation period (Magee and Wu, 2017; Vinçon-Leite et al., 2014; Kerimoglu and Rinke, 2013; Hadley et al., 2014). However, most water bodies present a complex morphology, whose effects on the hydrodynamics can only be taken into account by three-dimensional models. This is crucial to study the presence of local patterns and spatial heterogeneity, and to reconstruct the lake dynamics not only in time but in space as well. The hydrodynamics of small and shallow lakes is complex and strongly influenced by meteorological conditions. They are usually polymictic and cannot be simply considered as completely mixed reactors (McEnroe et al., 2013). In fact, they alternate periods of complete mixing to periods of stable thermal stratification that, depending on the local meteorological conditions, can last up to a few weeks (Soulignac et al., 2017).

In this context, the objective of this paper is to evaluate the thermal response of a small and shallow lake to climate change, and to characterize the evolution of its thermal regime in relation to stratification phenology and tendency to biomass production. To do so, the hydrodynamics of a small urban lake was reconstructed and analyzed along the past six decades through a three-dimensional model, and the presence of long term trends and of spatial heterogeneity was tested. Furthermore, a series of indices that thoroughly describe the thermal regime of the water body, in terms of stratification dynamics and phytoplankton



growth was defined and analyzed. The proposed methodology was here applied to a study site located in the Paris region. However, it is generic and could be applied to other similar sites.

## 2 Materials and methods

### 60 2.1 Study site and in situ measurements

Lake Champs-sur-Marne is a sand-pit lake located in the East side of the Great Paris region, next to the Marne River. It is a small and shallow water body with a surface of 0.12 km<sup>2</sup>, mean depth of 2.5 m and maximum depth of 3.5 m. As shown in Fig. 1-b, the southern part of the lake is the deepest one, while depth decreases under 2 m around the island and in the northern part of the lake. Lake Champs-sur-Marne has no inflow or outflow and is fed primarily by groundwater and occasionally by rainfall runoff. Its water level varies weakly during the year, with monthly oscillations lower than 0.2 m on average.


The lake was originated in the 1940s by excavation and represents now a valuable and demanded recreational area. However, it suffers from strong eutrophic conditions and experiences severe harmful algal blooms, especially between late spring and early autumn. In particular, cyanobacteria such as *Microcystis* and *Aphanizomenon*, capable to produce toxins, often proliferate and become the dominant species in the lake. This leads regularly to bathing bans and to restrictions in the access to the lake. For these reasons, the lake is subject to a periodic monitoring. In order to provide a faster and more responsive survey of the main chemico-physical properties of the lake, a high-frequency (every 10 minutes) *in situ* measuring system has been installed at two different locations (A and B) during the spring 2015. Each measuring site is equipped with sensors at three depths: below the surface at 0.5 m depth, in the middle of the water column at 1.5 m and above the sediment at 2.5 m (Tran Khac et al., 2018). Water temperature measurements are recorded at the surface and bottom layers with a precision of 0.02 °C and a resolution of 0.05 °C through the thermal sensor SP2T10 (nke INSTRUMENT®), and through the MPx multi-parameter sensor (nke INSTRUMENT®) at the middle of the water column, with a precision and a resolution of 0.05°C (see Fig. 1-a). High-frequency water temperature observations are used here for the calibration and validation of the hydrodynamic model.

Lake Champs-sur-Marne is polymictic and its thermal behavior is strongly influenced by meteorological conditions. Between March and November periods of thermal stratification alternate with mixing and overturn of the water column. Depending on meteorological conditions, thermal stratification might form during the day and break up at night as well as last up to three or four consecutive weeks.

### 2.2 The model

#### 2.2.1 Presentation of Delft3D-FLOW

In order to simulate the thermal behavior of the lake and hindcast its evolution in the past decades, the FLOW module from the Delft3D modelling suite was used (Deltares, 2014). Delft3D-FLOW is a well known hydrodynamic model that has been applied in various contexts, from estuaries to rivers, lakes and reservoirs (Piccolroaz et al., 2019; Chanudet et al., 2012; Soullignac et al., 2017). It solves the Reynolds averaged Navier-Stokes equations for an incompressible fluid under the shallow water and



**Figure 1.** a) satellite picture of Lake Champs-sur-Marne (source: *géoportail.fr*) and sketch of the measuring system at the two locations (A and B); b) bathymetry and horizontal mesh of the study site as used in Delft3D.

the Boussinesq assumptions. The time integration of the partial differential equations is done through an Alternate Direction Implicit method (Deltare, 2014; Leendertse, 1967). For the spatial discretization of the horizontal advection terms the Cyclic  
90 scheme was selected (Stelling and Leendertse, 1992).

The bathymetry and the two-dimensional mesh of the domain representing the study site are shown in Fig. 1-b. The surface of the lake is divided in 255  $20\text{ m} \times 20\text{ m}$  cells. The Z-model was implemented for the discretization of the vertical axes, with 12 fixed parallel horizontal layers of 30 cm thickness. It is generally accepted that horizontal layers help avoiding artificial mixing, improving model results in terms of thermal stratification (Hodges, 2014). Turbulent eddy viscosity and diffusivity  
95 were computed through the  $k-\varepsilon$  turbulence closure model. Background values were set to zero  $[\text{m}^2 \cdot \text{s}^{-1}]$  for vertical viscosity and diffusivity, while they were set to  $0.01\text{ m}^2 \cdot \text{s}^{-1}$ , after Soullignac et al. (2017) and according to the grid size, for horizontal viscosity and diffusivity. Bottom roughness was computed through Chézy's formulation with the default value for the Chézy coefficient of  $65\text{ m}^{1/2} \cdot \text{s}^{-1}$ .

The computation of the heat exchange at the air-water interface is done through Murakami's model (Murakami et al., 1985).  
100 It requires as input time series of relative humidity [-], air temperature [ $^{\circ}\text{C}$ ], net solar radiation [ $\text{J} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$ ], wind speed [ $\text{m} \cdot \text{s}^{-1}$ ] and direction [ $^{\circ}\text{N}$ ], as well as constant values for sky cloudiness [-] and Secchi depth [m]. The heat flux model computes the heat budget at the air-water interface by taking into account the net incident solar radiation ( $Q_s$ ), the heat losses due to back radiation (long wave,  $Q_b$ ) and evaporation (latent heat flux,  $Q_e$ ), and the sensible convective heat flux ( $Q_c$ ). The total upward



heat flux through the air-water interface ( $Q$ ) is therefore:

$$105 \quad Q = -Q_s + Q_b + Q_e + Q_c \quad (1)$$

Finally, evaporative mass flux is here neglected and water volume and depth are therefore considered as constant. This assumption makes it possible to analyze exclusively the impact of changes in the climatic forcing.

## 2.2.2 Meteorological input data

The meteorological forcing for this study comes from the spatialized SAFRAN (Système d'Analyse Fournissant des Ren-  
110 seignements Atmosphériques à la Neige) meteorological analysis system (Durand et al., 1993; Quintana-Seguí et al., 2008; Raimonet et al., 2017). SAFRAN is part of the SAFRAN-ISBA-MODCOU chain of reanalysis that covers the hydrological cycle over France, from meteorology to snow and ice formation to hydrology, respectively (Habets et al., 2008). SAFRAN integrates spatialized data from meteorological models with various sources of observations through data assimilation techniques, in order to create a consistent and spatially detailed record of meteorological data over the french territory. The data  
115 are spatialized on a regular square grid (8 km between each cell center) that covers the entire French Territory. The location of Lake Champs-sur-Marne falls midway on the axes connecting the centers of SAFRAN cells number 1457 (North of the lake) and 1566 (South of the lake). The average of these two cells was therefore considered representative of the conditions over the study site and used as input for the hydrodynamic model.

Data are available from the SAFRAN suite in terms of: air temperature [ $^{\circ}\text{C}$ ], specific humidity [-], solar radiation (direct  
120 and diffused) [ $\text{W}\cdot\text{m}^{-1}$ ] and wind speed [ $\text{m}\cdot\text{s}^{-1}$ ]. They were downloaded at a hourly time step, in order to accurately simulate the daily variability of the thermal profile and improve the model performance. This is crucial in shallow water bodies, where thermal stratification and mixing can alternate between day and night. Specific humidity (SH) data had to be converted into relative humidity (RH) to match the input data set needed by Delft3D. This was done through the following formula:

$$RH = 100 \cdot \frac{w}{w_s} \approx 100 \cdot \frac{SH}{w_s} \quad (2)$$

125 where  $w$  is the mixing ratio of water with dry air [ $\text{kg}\cdot\text{kg}^{-1}$ ], the subscript  $s$  stands for saturation conditions and SH is the specific humidity, numerically very close to the mixing ratio value. The saturation mixing ratio can be calculated as follows:

$$w_s = \frac{R_a}{R_v} \cdot \frac{e_s}{p_{atm} - e_s} \quad (3)$$

where the atmospheric pressure ( $p_{atm}$ ) was considered to be constant and equal to the global average:  $p_{atm} = 1013$  hPa. The ratio between the air and vapor ideal gas constants ( $R_a$  and  $R_v$ , respectively) is equal to 0.622. The partial vapor pressure at  
130 saturation ( $e_s$ ) is temperature dependent and can be estimated (in hPa) through the Magnus equation:

$$e_s = 6.1094 \cdot \exp\left(\frac{17.625 \cdot T}{T + 243.04}\right) \quad (4)$$

where  $T$  is air temperature [ $^{\circ}\text{C}$ ]. The numerical coefficients in Eq. (4) were issued from (Alduchov and Eskridge, 1997). Finally, in order to complete the set of meteorological input for Dleft3D, daily wind direction data were downloaded from the



closest available MétéoFrance station (ID: 78621001 located in Trappes, roughly 40 km West of the study site), through the  
135 INRAE CLIMATIK platform (<https://intranet.inrae.fr/climatik/>, in French) managed by the AgroClim laboratory of Avignon,  
France.

### 2.2.3 Calibration and validation

Delft3D-FLOW stands on a robust mathematical and physical structure and only few parameters have to be calibrated. Here,  
only those directly involved in the heat-flux model and in the wind module were calibrated: the Secchi depth [m], the mean  
140 cloud cover [-] and the wind drag coefficient [-]. The Secchi depth ( $H_S$ ) is the parameter that defines water transparency. It  
is correlated with the penetration of solar radiation in water through the light extinction coefficient ( $\gamma = 1.7/H_S$  (Poole and  
Atkins, 1929)) and therefore has a strong influence on the stratification of the water column. In order to get a first estimate for  
the sky cloudiness parameter, cloud cover data from the MétéoFrance station in Trappes (ID: 78621001) were averaged over the  
calibration period. The wind drag coefficient was calibrated in order to take into account the presence of tall trees surrounding  
145 the contour of the lake, locally reducing wind speed. The calibration was done through a trial and error procedure based on  
high-frequency water temperature data at the surface, middle and bottom layers (0.5, 1.5 and 2.5 m depth, respectively) during  
the year 2016. The model was then run for validation over the whole period during which both meteorological data and *in situ*  
observations were available, i.e. from the 15th May 2015 to the 31st December 2017.

Model results were compared to water temperature data at three depths (surface, middle and bottom of the water column)  
150 and two different locations (A and B). Root mean square error (RMSE) and mean bias error (MBE) were calculated to evaluate  
model performances. For this purpose, high-frequency data were first averaged every hour to match the model output time step  
and cleaned from the outliers originated by periodic sensor maintenance. The latter were defined as sudden water temperature  
variations ( $> 1^\circ\text{C}$ ) over the 10 minutes separating two successive measurements, and consequently erased.

### 2.3 Indices for the characterization of the lake thermal regime

155 The thermal regime of the lake was assessed directly through the analysis of model results in terms of water temperature, as  
well as through a series of indices that explore the phenology of stratification and highlight the relation between temperature  
and phytoplankton production. The indices characterizing the thermal regime of the lake are described here-after.

#### 2.3.1 Stratification indices

In order to thoroughly characterize the phenology of stratification in Lake Champs-sur-Marne, two indices for the stability  
160 of the water column have been calculated: the Schmidt stability index and an index based on temperature difference between  
surface and bottom layers. The Schmidt stability index is a well known parameter often used in limnological studies to estimate  
the resistance of a water body to mixing, and therefore its stability. It has been extensively used in scientific literature to describe  
the strength of stratification in lakes and, more recently, to analyze its evolution over time in relation to climate change (Vinçon-  
Leite et al., 2014; Niedrist et al., 2018; Kraemer et al., 2015; Livingstone, 2003) and algal blooms (Wagner and Adrian, 2009).



165 The Schmidt stability index ( $S$ ) represents the amount of work per unit area that would be required to mix the lake water column at one time instant. It has been here calculated following Idso's formulation (Idso, 1973), in which the vertical axis  $z$  is considered positive downwards from the surface to the maximum lake depth  $z_M$  [m]:

$$S = \frac{g}{A_0} \int_0^{z_M} (z_v - z)(\rho_i - \rho_v)A(z)dz \quad [J.m^{-2}] \quad (5)$$

where:

170  $z_v = \frac{1}{V} \int_0^{z_M} zA(z)dz \quad (6)$

is the depth of the center of volume of the lake,  $\rho_v$  [ $kg.m^{-3}$ ] is water density at the depth of the center of volume  $z_v$ ,  $\rho_i$  is the mean uniform density,  $g$  [ $m.s^{-2}$ ] is the acceleration of gravity,  $V$  [ $m^3$ ] and  $A_0$  [ $m^2$ ] are respectively the volume and the surface area of the lake, and  $A(z)$  is the surface of the horizontal section of the lake at depth  $z$ . Computed for each time step, the Schmidt stability can also be averaged over each year or season.

175 Water resistance to mixing as estimated by the Schmidt stability index is closely correlated to temperature stratification. However, universal thresholds for the onset and breakdown of stratification are difficult to define based on this index and cannot be found in the literature, especially for shallow polymictic lakes. For these reasons, a second index based on temperature difference between surface and bottom layers ( $\Delta T$ ) is proposed. In order to assess the succession of stratification events in a polymictic water body, after Kerimoglu and Rinke (2013) and Magee and Wu (2017), the lake was considered to be stably  
180 stratified during one day if the minimum of  $\Delta T$  is greater than 1 °C during the whole day. This allows to identify all stably stratified days. The sum over a year of the stably stratified days (SSD) is called the annual number of SSD. The sum can also be evaluated on a seasonal basis (according to the definitions in section 2.4), leading to the seasonal number of SSD.

### 2.3.2 Growing degree days (GDD) and number of growing days (NGD)

In order to quantify how changes in the thermal regime might impact biomass production and in particular phytoplankton  
185 growth, we introduce two indices: the growing degree days (GDD) and the number of growing days (NGD). Growing degree days is a weather based indicator for biological growth, widely used in the field of agronomy. Based on air temperature, it gives an estimate of the rate of development and of the span of its phase for terrestrial plants and insects during the growing season. It is a useful indicator capable to link global warming and biology (Grigorieva et al., 2010; Schlenker et al., 2007). Approaches based on GDD have been increasingly applied to phytoplankton communities and fisheries (e.g. Gillooly, 2000; Neuheimer  
190 and Taggart, 2007; Ralston et al., 2014; Dupuis and Hann, 2009), in order to correlate water temperature and phytoplankton growth while taking into account interannual variability. After Dupuis and Hann (2009), GDD were calculated as follows:

$$GDD(t) = \sum_{i=t_0}^t \max\{0, (T_i - T_{base}) \cdot \Delta t\} \quad (7)$$





where  $t$  is the time (here in days) with  $t_0$  the reference day to start the calculation,  $\Delta t$  is the time step (equal to 1 day),  $T_i$  is the daily average of the modeled surface water temperature of day  $i$  and  $T_{base}$  is a physiological threshold below which growth does not occur. GDD can be calculated on an annual or a seasonal basis by adjusting the values of  $t_0$  and  $t$ . Annual GDD are calculated starting from the first of January until the 31st of December, while seasonal GDD can be obtained by adjusting  $t_0$  and  $t$  to the seasons defined in section 2.4. As in Dupuis and Hann (2009), the value of  $4^\circ\text{C}$  was selected for  $T_{base}$ . Such value was chosen to be a representative baseline for the growth of the whole phytoplankton community in Lake Champs-sur-Marne, generally composed of cyanobacteria, green algae and diatoms. This results in a succession of algal blooms that spans almost the whole year, usually starting in February when water temperature in the study site surpasses  $4^\circ\text{C}$ , until the end of Autumn.

The number of growing days (NGD) at day  $t$  is defined as the number of days during which  $(T_i - T_{base}) > 0$  from day  $i = t_0$  to  $i = t$ . It represents a proxy for the duration of the period favorable to growth for the phytoplankton community. Annual NGD and seasonal NGD can be calculated by adjusting the values of  $t_0$  and  $t$  as for the GDD. However, in a warm temperate climate water temperature never drops under the threshold of  $4^\circ\text{C}$  during Spring and Summer. This impedes any variability in the counts of NGD during these two seasons, that were therefore not addressed.

## 2.4 Long-term analysis

In the present paper we aim at hindcasting the long-term dynamics of a small and shallow urban lake in order to test the influence of climate change on such ecosystems. To do this we focus on 58 years, from 1960 to 2017, for which meteorological data are available from the SAFRAN reanalysis.

### 2.4.1 Long-term trends

The long-term hydrodynamic simulation starts on the first of January 1960. No data were available to set the initial conditions of the model, neither in terms of water temperature, nor in terms of current velocities. However, the model is strongly driven by the meteorological data and the influence of the initial condition vanishes after only a few days (Piccolroaz et al., 2019). The effect on model results of small perturbations in water temperature initial conditions ( $\pm 2^\circ\text{C}$ ) was tested and resulted to vanish in 5 to 7 days. The model was therefore initialized with water at rest and with a uniform water temperature of  $7^\circ\text{C}$ . This value was issued from the average of the water temperature recorded on the lake on the first of January in 2016, 2017, 2018 and 2019. Model results are stored at a hourly time step on every element of the mesh.

Model results at measuring site A are analyzed on an annual and seasonal basis for long-term trends. They are exploited directly in terms of water temperature (averaged over the water column) and to calculate the indices defined in section 2.3. Site A is located in one of the deepest part of the lake and therefore considered as representative of the general behavior of the water body. This assumption will be further analyzed through a spatial analysis of the three-dimensional model results (see section 2.4.2). The presence of long term trends is tested (with a threshold for significance  $\alpha = 0.05$ ) through the Mann-Kendall test (Mann, 1945; Kendall, 1975), a non-parametric test for the individuation of overall monotonic trends performed here through the MATLAB software (Burkey, 2020). The Mann-Kendall test is often preferred to simple linear regression in the analysis of meteorological and hydrological time series (Tímea et al., 2017; Wang et al., 2020), as it does not require any assumption on the



distribution of the analyzed dataset. Once a trend is detected, its strength is evaluated through the Sen's slope estimator (Sen, 1968), that uses a linear model to evaluate the intensity of the trend. In order to analyze both annual and seasonal trends, the year was divided into four groups of three months each: (i) January, February and March (Winter), (ii) April, May, June (Spring), (iii) July, August, September (Summer), (iv) October, November, December (Autumn). Meteorological forcing is crucial for this work, as it drives the hydrodynamic model and represents the only source of variability in our modelling configuration. The presence of long-term trends in the meteorological dataset was also evaluated by applying the Mann-Kendall test and the Sen's slope estimator to their annual averages.

#### 2.4.2 Spatial analysis of stratification

The phenology of stratification in Lake Champs-sur-Marne was further assessed exploiting the three-dimensional model simulations. The analysis was extended to the whole computational domain, with the objective of investigating the relation between climate change, water depth and thermal structure in a shallow and polymictic water body. The annual number of stably stratified days (SSD) was therefore calculated for every computational cell over the complete simulation period, using temperature values from the surface and local bottom layers. In order to detect the presence of heterogeneity in the overall spatial distribution of stratification, the annual SSD calculated for each computational cell were averaged over the 58 years of the simulation to obtain a space-dependent (but time-averaged) representation of thermal stratification.


The evolution of stratification in different areas of the water body can also be evaluated over time. To do so, the domain was divided into eight groups of cells depending on the local water depth, here considered constant in time. For instance, the first group includes the shallowest computational cells (depth between 1.2 and 1.5 m), the second one includes cells with depth between 1.2 and 1.5 m, and so forth until the last group of cells, with depth between 3.3 and 3.6 m. For each simulated year, the annual number of SSD was then averaged over each of the cell-groups, in order to represent the evolution over time of different areas of the lake with homogeneous depth.

### 3 Results

#### 3.1 Model calibration and validation

The model was calibrated on the year 2016 and validated from May 2015 to the end of 2017. Records from field campaigns show that the Secchi depth in Lake Champs-sur-Marne varies between 0.5 and 3 m; using this range of values, it was calibrated and finally set to 1.2 m. Sky cloudiness was calibrated and set to 80%, and a uniform wind drag coefficient was set to 0.005 [-].

During the calibration periods, the model tends to slightly overestimate water temperature during summer heat peaks (with residuals always lower than 2°C) and to slightly underestimate water temperature during winter (again, with residuals always lower than 2°C). However, overall model performances are satisfactory for all three layers, with RMSE values between simulated and observed water temperature of 0.85°C, 0.78°C and 0.81°C at site A respectively for the surface (0.5 m), middle (1.5 m) and bottom (2.5 m) layers. Model results are spatially robust and satisfactory also for the validation period, with similar




**Figure 2.** Water temperature during the validation period. a) Model output (black) and high-frequency observations (grey) for the middle layer (1.5 m depth) at measuring site A; b) residuals between model output and *in situ* data.

RMSE values for sites A and B (ranging between 0.96 and 1.00°C). Fig. 2-a shows simulation results and observations during the whole validation period at 1.5 m depth at site A, where the data series is the longest, together with the related residuals (Fig. 2-b); results were very similar for the two remaining sites. The model fits the data very well, especially between spring and autumn. During summer (respectively, winter), similarly to what was observed during calibration, it tends to slightly overestimate (respectively, underestimate) water temperature, with residuals always lower in module than 2°C. Over the validation period the model has a low but non negligible bias (MBE) ranging between -0.2 and -0.3 °C for sites A and B. Eventually, the number of SSD was calculated as defined in section 2.3.1 during the whole validation period at sites A and B for both the observed and simulated dataset. In case of gaps in the observation series, also simulation output were discarded from the calculation. The number of SSD between 2015 and 2017 calculated using water temperature observations is 122 for site A and 128 for site B. Using Delft3D simulations we found very similar results: 125 SSD for site A and 132 for site B. Overall, the model excellently reproduces high-frequency water temperature data, including the diurnal cycle, at both measuring sites.

## 3.2 Long-term trend analysis

### 3.2.1 Meteorological input data

Annual averages were calculated from 1960 to 2017 for the five meteorological variables used as inputs to the hydrodynamic model and tested with the Mann-Kendall test. Strongly significant monotonic trends ( $p \ll 0.05$ ) were found for the air temperature, solar radiation and wind speed, as shown in Fig. 3. The Sen's slope estimator was used to test the intensity of the significant monotonic trends. Air temperature displays a considerable warming trend of 0.3°C per decade; solar radiation also shows a significant increasing trend, with an intensity of 3.5 W.m<sup>-2</sup> per decade. Wind speed decreases quite sharply over time, at an estimated rate of 0.2 m.s<sup>-1</sup> per decade. While the increase in air temperature appears extremely linear (see Fig. 3-a), a



**Figure 3.** Annual averages of the three meteorological variables input to Delft3D (solid lines) that showed significant monotonic trends, and relative trend intensity evaluated through Sen's slope estimator (dashed lines). a) Air temperature; b) solar radiation; c) wind speed.

sharp shift in the behavior of both solar radiation and wind speed appears around the year 1989 (Fig. 3-b and -c, respectively). Solar radiation and wind speed appear to be piecewise functions of time that can be roughly divided into two sections with sensibly different mean values, one until 1989 and the second one starting from 1990. Despite this non linear behavior, the presence of monotonic increasing (for solar radiation) or decreasing (for wind speed) trends is confirmed by the very low p-values  
280 obtained for these variables through the Mann-Kendall test. Finally, no significant trend was found for relative humidity and wind direction. The two variables appear to be stationary, the former fluctuating around an annual average of roughly 80% and the latter around an annual prevailing wind direction of 200°N (South-West). Three of the five meteorological variables forcing the hydrodynamic model were therefore characterized by strongly significant monotonic trends along the past six decades, corroborating the idea of a changing climate in the region around the study site.

### 285 3.2.2 Model results

Long-term monotonic trends have been researched at site A on an annual and seasonal basis for: water temperature (vertically averaged), number of stably stratified days (SSD), mean Schmidt stability, growing degree days (GDD) and number of growing days (NGD). Figure 4 shows all the significant monotonic trends found from this analysis. On an annual basis, the Mann-Kendall test highlighted the presence of strongly significant increasing trends ( $p \ll 0.05$ ) for all variables.

290 Mean annual water temperature shows a very sharp warming tendency of 0.6°C per decade (see Fig. 4-a), even greater than what was found for air temperature (0.3°C). The Pearson correlation coefficient ( $r$ ) was calculated between water temperature and the five meteorological input variables in terms of annual averages in order to explain this behavior. Air temperature, solar radiation and wind speed were all strongly correlated with modeled water temperature, with correlation coefficients of 0.8 for solar radiation and air temperature and -0.9 for wind speed. Water temperature showed significant increase during all seasons, with higher slopes during Spring and Summer (0.8 and 0.7°C per decade, respectively), and a lower yet considerable intensity  
295 during Autumn and Winter (respectively 0.4 and 0.5°C per decade).

The warming trend is accompanied by reinforced stratification. An increase in water column stability was highlighted on an annual basis by both stratification indices: the annual number of SSD increased on average of around three days per decade,



while the Schmidt stability index increased of  $0.9 \text{ J}\cdot\text{m}^{-2}$  per decade (Fig. 4-b and -c, respectively). Despite a warming trend  
300 being present in all seasons, both stratification related indices showed significant increasing trends only during Winter (1 d and  
 $0.4 \text{ J}\cdot\text{m}^{-2}$  per decade) and Spring (2 d and  $2.6 \text{ J}\cdot\text{m}^{-2}$  per decade, for the seasonal SSD and the Schmidt index, respectively).  
The sharpest increase in stable stratification was therefore found during Spring.


The analysis of the growing degree days and of the number of growing days shows the progressive improvement of conditions  
for biomass production. The black line in Fig. 4-d shows the evolution over time of annual GDD. Its pattern is highly correlated  
305 to that of water temperature and shows an increasing rate of  $190^\circ\text{C}\cdot\text{d}$  per decade, with a shift around the year 1989. Even  
though this behavior cannot be regarded as linear, it corroborates the idea of a greater amount of thermal energy reaching the  
ecosystem. Furthermore, as shown by the increase in annual NGD (Fig. 4-e), this energy is spread over a wider period of time,  
now encompassing almost the whole year. The NGD increased by 10 days per decade on an annual basis, with significant  
seasonal trends for Winter (5 d per decade) and Autumn (3 d per decade). However, the amount of thermal energy available for  
310 biomass production did grow during all four seasons. Seasonal GDD show an estimated increase of: 24, 75, 67 and  $27^\circ\text{C}\cdot\text{d}$   
per decade, going from Winter to Autumn.

The changes found in the meteorological forcing clearly had an impact on the dynamics of the study site. The lake has  
sensibly warmed and its tendency to thermal stability has increased. Spring showed the sharpest trends in terms of water  
temperature, water column stability (Schmidt and SSD) and growing degree days, and might ultimately be the season suffering  
315 the strongest changes in terms of biomass production and algal blooms.


### 3.3 Spatial analysis of stratification

In order to examine the presence of spatial patterns regarding water column stratification, the three-dimensional model results  
were exploited for each cell of the domain and as well as for different depth classes. Figure 5-a displays a map of the study site  
representing for each cell of the domain the average over time (from 1960 to 2017) of the annual number of SSD, similarly to  
320 the spatial representation of sediment disturbance by waves found in Bachmann et al. (2000). The map shows a pronounced  
spatial pattern, with stratification developing only in some regions. The deeper areas were stratified on average for more than  
60 days a year, while the shallower northern part never significantly stratified. A strong linear correlation was found between  
water depth and number of stably stratified days, with a correlation coefficient between these two variables of 0.98.

The evolution over time of the spatial pattern of stratification was also investigated. The computational domain was divided  
325 into eight disjointed groups of cells depending on local water depth, which varies between 1.2 m and 3.6 m. For each year in  
the simulation, the annual number of SSD was then averaged over each of the eight cell-groups, in order to obtain the eight  
time series of annual SSD shown in Fig. 5-b. Figure 5-b shows that, for this study site, no thermal stratification ever develops in  
areas with a depth lower than 1.8 m. The depth of the thermocline therefore appears to be always greater than 1.8 m. Figure 5-b  
also shows an evolution over time of stratification. All cells groups with a depth greater than 1.8 m tend to experience a higher  
330 annual number of SSD towards the end of the simulated period, especially starting from 1995. In order to confirm this result,  
the time series of annual SSD calculated over each same-depth cells group (shown in Fig. 5-b through the color chart) was  
analyzed with the Mann-Kendall test. All cell-groups deeper than 1.8 m showed statistically significant ( $p < 0.05$ ) increasing



**Figure 4.** Statistically significant climate change trends at monitoring site A for the five indices, both on an annual (black) and seasonal (other colors) basis. a) Water temperature (averaged on the water column); b) Number of stably stratified days (SSD); c) Schmidt stability. d) Growing degree days (GDD); e) Number of growing days (NGD). Blue lines represent the Winter season, green lines represent Spring, red lines are for Summer trends and yellow lines for Autumn; black lines represent annual values.



**Figure 5.** a) Map of the annual number of SSD calculated over every cell in the computational grid, and averaged over the 58 years of the simulation; b) evolution over time of the annual number of SSD divided by cell groups with homogeneous depth.

trends in terms of annual number of SSD, with intensity varying between 2 d per decades (for cells with 1.8 m water depths) to 4 days per decade (for cells in the deepest part of the lake).



#### 335 4 Discussion

Long-term climate change trends were researched between 1960 to 2017 through the Mann-Kendall test for both meteorological input variables and model results.

In terms of meteorological variables, air temperature and solar radiation showed increasing monotonic trends ( $0.3^{\circ}\text{C}$  and  $3.5 \text{ W.m}^{-2}$  per decade, respectively), while wind speed showed a decreasing monotonic trend of  $-0.2 \text{ m.s}^{-1}$  per decade. A few  
340 studies already assessed climate change in the Paris region, mainly in terms of air temperature (Perrier et al., 2005; Lemonsu et al., 2013). Compared to our result, a milder increasing trend of  $0.1^{\circ}\text{C.dec}^{-1}$  was found between 1900 to 1987 analyzing measurements, with a steeper increment of  $0.7^{\circ}\text{C.dec}^{-1}$  later on until 2005 (Perrier et al., 2005). Similarly, we also find a steeper trend of  $0.55^{\circ}\text{C.dec}^{-1}$  if we limit our analysis on the years from 1987 to 2005. Less information can be found for solar radiation and wind speed. A decrease in wind speed on land was found over Europe since 1980 (around  $-0.1 \text{ m.s}^{-1}.\text{dec}^{-1}$ )  
345 as part of a large-scale analysis of observations in the northern hemisphere (Vautard et al., 2010), as well as on a global scale through meteorological reanalysis (Torralba et al., 2017). An overall increase in surface solar radiation was recently found for Europe between 1983 and 2015, specifically of  $3 \text{ W.m}^{-2}.\text{dec}^{-1}$  for western Europe (Pfeifroth et al., 2018). Solar radiation and wind speed showed here a piecewise non-linear behavior (Fig. 3), with a shift around the late 1980s. SAFRAN is a coherent and spatialized data set, that still partially depends on the surface observation network and might be influenced by changes in  
350 the instrumentation (Vidal et al., 2010). However, the existence of a shift in global climate in the 1980s has been highlighted by a number of studies using different data sources (Reid et al., 2016; Mariani et al., 2012; Gallagher et al., 2013).

Meteorological reanalyses usually cover multi-decadal periods and have the great benefit of being spatialized over vast portions of the globe. Even though their use in limnological studies is quite recent, they have already been used to simulate water temperature (Layden et al., 2016; Piccolroaz et al., 2020), stratification dynamics (Frassl et al., 2018) and phytoplankton  
355 distribution (Soulignac et al., 2018). As shown in this work, their use as external forcing to hydrodynamic models can yield, provided that data are available for calibration and validation, to accurate simulations of the behavior of water bodies even in the absence of local meteorological observations. This could open to a great range of applications in the field of limnology and paleolimnology (e.g. Jenny et al., 2016). The proposed methodology allows to thoroughly reconstruct the behavior of a water body both in time and space, independently of its proximity to meteorological stations. Furthermore, the use of biogeochemical  
360 modules could give additional insights on oxygen, nutrients and phytoplankton dynamics.

The use of an extensive data set of high-frequency observations (recorded every 10 min, at three depths and two locations) allowed to test the ability of the model to reproduce not only the general seasonal water temperature pattern, but also daily and sub-daily dynamics. In this respect the model performed very well, with RMSE values always lower than  $1^{\circ}\text{C}$  and comparable to those obtained in other studies with similar applications (Kerimoglu and Rinke, 2013; Magee and Wu, 2017; Lee et al.,  
365 2018) through less frequent observations.

Significant increasing trends were detected for water temperature both on an annual and seasonal basis. The highest warming was found during Spring and Summer ( $0.8$  and  $0.7^{\circ}\text{C.dec}^{-1}$ , respectively), while it was weaker during Autumn and Winter (around  $0.4^{\circ}\text{C.dec}^{-1}$ ). Overall, mean annual water temperature increased at a considerable rate of  $0.6^{\circ}\text{C}$  per decade, greater



than that of air temperature ( $0.3^{\circ}\text{C}\cdot\text{dec}^{-1}$ ), a behavior also highlighted by other studies (Austin and Colman, 2007; Schneider  
370 et al., 2009). A large-scale analysis showed how trends in summer surface lake water temperature are globally highly variable,  
comprised between  $-0.7$  and  $1.3^{\circ}\text{C}$  per decade (O'Reilly et al., 2015). After this study, lakes subject to similar changes in the  
meteorological forcing to our study site, namely in terms of air temperature and solar radiation, display an average summer  
warming trend of  $0.53^{\circ}\text{C}$  per decade, only slightly smaller than the result of our work. Mean annual water temperature showed  
a non-linear behavior (see Fig. 4-a), probably caused by that of solar radiation and wind speed. In fact, similarly to what was  
375 found by Magee and Wu (2017), mean annual water temperature was highly correlated (i.e.  $r \geq |0.8|$ ) with air temperature,  
solar radiation and wind speed. This suggests that meteorological variables might have additive effects that concur to enhance  
the response of the dependent variable. These effects might be particularly intense for wind over small and shallow lakes, due  
to their low volume to surface ratio.

Both stratification related indices (Schmidt stability and SSD) showed a mean annual increasing trend, mainly driven by  
380 an increase during Spring (respectively 2 d and  $2.6 \text{ J}\cdot\text{m}^{-2}$  per decade). Similar values were recently found for shallow water  
bodies in other long-term studies (Magee and Wu, 2017; Moras et al., 2019). The alternation between stratification and mixing  
in polymictic water bodies strongly influences the distribution and availability of nutrients over the water column (Song et al.,  
2013). The length of stable stratification and the frequency of mixing events have effects on the dynamics of sedimentation and  
resuspension of both nutrients and organic matter. This in turn might have strong effects on phytoplankton productivity and on  
385 the occurrence of harmful algal blooms, with impacts on the ecological state of the water body. To address this issue, the GDD  
and the NGD were introduced to link the thermal regime to the biological productivity of the study site. Both the annual GDD  
and the NGD showed a significant increasing trend. The seasonal GDD were found to significantly increase during all seasons,  
while the seasonal NGD increased significantly only during Autumn and Winter.

The thermal regime is a key factor in the regulation of the biogeochemical cycle and in the development of algal blooms.  
390 Small and shallow lakes react rapidly to climatic changes, partially due to their low volume and heat capacity. Changes in  
water temperature can alter the composition of phytoplankton groups and their succession throughout the year. Warmer water  
temperatures favor species with higher optimum temperatures such as cyanobacteria, capable of producing toxins (Paerl and  
Huisman, 2008). Stratification induces a separation between the sediment and the surface layers, influencing the distribution of  
nutrients and biomass over the water column. Due to their ability to migrate upwards towards the water surface, an increase in  
395 water column stability might also favor cyanobacteria over other species (Dupuis and Hann, 2009). Furthermore, in polymictic  
water bodies, a mechanism of accumulation (during stratification) and release (during subsequent mixing) of nutrients from  
the sediment has been suggested, causing multiple pulses that act as an internal nutrient source (Song et al., 2013; Wilhelm  
and Adrian, 2008). The combination of increasing trends for water temperature, stable stratification and the widening of the  
growing season can favor the occurrence of harmful algal blooms, further deteriorating ecosystems that are often already  
400 eutrophic (Winder and Sommer, 2012; Jones and Brett, 2014; Noble and Hassall, 2015). A number of small urban lakes  
similar to our study site might be undergoing similar changes as Lake Champs-sur-Marne. These changes might be especially  
sharp during Spring, that showed here the greatest increase for water temperature, GDD and SSD. The timing, composition





and intensity of spring blooms are extremely important in determining the succession of blooms in the subsequent months (Townsend et al., 1994; Sommer and Lengfellner, 2008; Lewandowska and Sommer, 2010).

405 Stratification affects many aspects of lake productivity, including nutrient recycling, habitat conditions for microorganisms and their distribution over depth (Hanna, 1998; Wilhelm and Adrian, 2008; Song et al., 2013). The spatial analysis of stratification showed a strong linear correlation between the number of annual SSD and local water depth. The deeper part of the study site does experience considerable stable stratification throughout the year (up to around 60 days on average), even though for non-consecutive periods of time. The increase in the annual number of SSD found for site A is shared with all cell-groups  
410 with water depth greater than 1.8 m. However, this is not the case in the shallower northern part of the basin, that never show stable stratification. Depending on the bathymetry, these spatial patterns might be strong in shallow water bodies, inducing heterogeneity in nutrients, phytoplankton and oxygen concentrations. Such heterogeneity might be even stronger in large water bodies, and can only be thoroughly inferred through three dimensional models (Gong et al., 2016; Frassl et al., 2018).

## 5 Conclusions

415 In this work, the long-term hydrodynamics of a shallow urban lake were profitably reconstructed through model simulations from 1960 to 2017. A series of indices were proposed with the objective of thoroughly describing the thermal regime of shallow water bodies, in relation with stratification dynamics and biological productivity. The meteorological data set was derived from the SAFRAN reanalysis and showed a significant increase in air temperature and solar radiation and a significant decrease in wind speed, with a regime shift in the late 1980s. Simulation results show that small urban lakes react rapidly to  
420 external meteorological conditions, with only limited resilience to climatic shifts. The increase in water temperature cannot be explained by air warming only. The additive effect of increasing solar radiation and decreasing wind speed acts on different terms of the heat budget at the lake surface, enhancing the changes found in the lake. Water warming ( $0.6^{\circ}\text{C}/\text{dec}$ ) is much quicker than air warming ( $0.3^{\circ}\text{C}/\text{dec}$ ), and especially intense in Spring, as is the lengthening of stratification. This could have favored early phytoplankton blooms, the development of cyanobacteria and ultimately the degradation of the whole aquatic  
425 ecosystem. Furthermore, the heterogeneity found in the spatial distribution of thermal stratification might concur to locally create favorable conditions for algal blooms, in terms of nutrient availability or warmer surface water temperature. The use of three-dimensional models is needed to thoroughly infer the dynamics of a water body, including horizontal patterns. Small and shallow lakes are extremely widespread ecosystems, and a number of them might be experiencing analogous changes. Our results suggest the urgent need for appropriate management in order to preserve their ecological value.

430 *Code and data availability.* The model set-up for long-term simulations, as well as the corresponding results at site A are available on Mendeley (<https://data.mendeley.com/datasets/92kzf5t5xn/draft?a=11918779-ce63-4e72-aa69-9207e8445fdc>). Model results were obtained using the Delft3D software package (Delft3D-flow, version 4.01.01.rc.03). Matlab codes used to obtain the datasets for this paper are available upon request from the corresponding author.



435 *Author contributions.* FP: conceptualization, investigation, writing - original draft. CC: conceptualization, supervision, writing - review & editing, funding acquisition, project administration. BJL: conceptualization, writing - review & editing. PLM: data curation, writing - review & editing. PD: data curation. BVL: conceptualization, supervision, writing - review & editing, funding acquisition, project administration.

*Competing interests.* The authors declare that they have no conflict of interest.

440 *Acknowledgements.* The data set used for model calibration and validation was collected under the OSSCyano (ANR-13-ECOT-0001) and ANSWER (ANR-16-CE32-0009-02) projects. The environmental observatories OSU EFLUVE and OLA contributed to the financial support for equipment maintenance. The authors also acknowledge the Base de loisirs du lac de Champs/Marne (CD93) for their logistical support in the field campaigns. The first author's PhD grant is funded by Ecole des Ponts ParisTech and ANSWER project. The authors also thank Nicolas Clercin for the revision of the English text.



## References

- Adrian R., O'Reilly C. M., Zagarese H., Baines S. B., Hessen D. O., Keller W., Livingstone D. M., Sommaruga R., Straile D., Van  
445 Donk E., Weyhenmeyer G. A., and Winder M.: Lakes as sentinels of climate change, *Limnology and Oceanography*, 54, 2283–2297,  
[https://doi.org/10.4319/lo.2009.54.6\\_part\\_2.2283](https://doi.org/10.4319/lo.2009.54.6_part_2.2283), 2009.
- Alduchov, O. A. and Eskridge, R. E.: Improved Magnus' form approximation of saturation vapor pressure, Tech. Rep. DOE/ER/61011-T6,  
Department of Commerce, Asheville, NC (United States), <https://doi.org/10.2172/548871>, 1997.
- Austin, J. A. and Colman, S. M.: Lake Superior summer water temperatures are increasing more rapidly than regional air temperatures: A  
450 positive ice-albedo feedback, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL029021>, 2007.
- Bachmann, R., Hoyer, M., and Canfield, D.: The Potential For Wave Disturbance in Shallow Florida Lakes, *Lake and Reservoir Management*,  
16, 281–291, <https://doi.org/10.1080/07438140009354236>, 2000.
- Biggs, J., von Fumetti, S., and Kelly-Quinn, M.: The importance of small waterbodies for biodiversity and ecosystem services: implications  
for policy makers, *Hydrobiologia*, 793, <https://doi.org/10.1007/s10750-016-3007-0>, 2016.
- 455 Burkey, J.: Mann-Kendall Tau-b with Sen's Method (enhanced), MATLAB Central File Exchange. Retrieved February 12, 2020, 2020.
- Chanudet, V., Fabre, V., and van der Kaaij, T.: Application of a three-dimensional hydrodynamic model to the Nam Theun 2 Reservoir (Lao  
PDR), *Journal of Great Lakes Research*, 38, 260–269, <https://doi.org/10.1016/j.jglr.2012.01.008>, 2012.
- Deltares: Delft3D-FLOW - Simulation of multi-dimensional hydrodynamic flows and transport phenomena, including sediments. User Man-  
ual Version: 3.15.30059, Delft Hydraulics, Delft, 2014.
- 460 Downing, J. A., Prairie, Y. T., Cole, J. J., Duarte, C. M., Tranvik, L. J., Striegl, R. G., McDowell, W. H., Kortelainen, P., Caraco, N. F.,  
Melack, J. M., and Middelburg, J. J.: The global abundance and size distribution of lakes, ponds, and impoundments, *Limnology and  
Oceanography*, 51, 2388–2397, <https://doi.org/10.4319/lo.2006.51.5.2388>, 2006.
- Dupuis, A. P. and Hann, B. J.: Warm spring and summer water temperatures in small eutrophic lakes of the Canadian prairies: potential  
implications for phytoplankton and zooplankton, *J Plankton Res*, 31, 489–502, <https://doi.org/10.1093/plankt/fbp001>, 2009.
- 465 Durand, Y., Brun, E., Merindol, L., Guyomarc'h, G., Lesaffre, B., and Martin, E.: A meteorological estimation of relevant parameters for  
snow models, *Annals of Glaciology*, 18, 65–71, <https://doi.org/10.3189/S0260305500011277>, 1993.
- Frassl, M. A., Boehrer, B., Holtermann, P. L., Hu, W., Klingbeil, K., Peng, Z., Zhu, J., and Rinke, K.: Opportunities and Limits of Using  
Meteorological Reanalysis Data for Simulating Seasonal to Sub-Daily Water Temperature Dynamics in a Large Shallow Lake, *Water*, 10,  
594, <https://doi.org/10.3390/w10050594>, 2018.
- 470 Frumkin, H., Bratman, G. N., Breslow, S. J., Cochran, B., Kahn, P. H., Lawler, J. J., Levin, P. S., Tandon, P. S., Varanasi, U.,  
Wolf, K. L., and Wood, S. A.: Nature Contact and Human Health: A Research Agenda, *Environ. Health Perspect.*, 125, 075001,  
<https://doi.org/10.1289/EHP1663>, 2017.
- Gallagher, C., Lund, R., and Robbins, M.: Change-point Detection in Climate Time Series with Long-Term Trends, *J. Climate*, 26, 4994–5006,  
<https://doi.org/10.1175/JCLI-D-12-00704.1>, 2013.
- 475 Gillooly, J.: Effect of body size and temperature on generation time in zooplankton, *Journal of Plankton Research*, 22,  
<https://doi.org/10.1093/plankt/22.2.241>, 2000.
- Gong, R., Xu, L., Wang, D., Li, H., and Xu, J.: Water Quality Modeling for a Typical Urban Lake Based on the EFDC Model, *Environ Model  
Assess*, 21, 643–655, <https://doi.org/10.1007/s10666-016-9519-1>, 2016.



- Grigorieva, E., Matzarakis, A., and De Freitas, C.: Analysis of growing degree-days as climate impact indicator in a region with extreme  
480 annual air temperature amplitude, *Climate Research*, 42, 143–154, <https://doi.org/10.3354/cr00888>, 2010.
- Habets, F., Boone, A., Champeaux, J.-L., Etchevers, P., Franchisteguy, L., Leblois, E., Ledoux, E., Le Moigne, P., Martin, E., Morel, S.,  
Noilhan, J., Quintana Seguí, P., Rousset, F., and Viennot, P.: The SAFRAN-ISBA-MODCOU hydrometeorological model applied over  
France, *Journal of Geophysical Research*, 113, <https://doi.org/10.1029/2007JD008548>, 2008.
- Hadley, K. R., Paterson, A. M., Stainsby, E. A., Michelutti, N., Yao, H., Rusak, J. A., Ingram, R., McConnell, C., and Smol, J. P.: Climate  
485 warming alters thermal stability but not stratification phenology in a small north-temperate lake, *Hydrological Processes*, 28, 6309–6319,  
<https://doi.org/10.1002/hyp.10120>, 2014.
- Hanna, M.: Evaluation of Models Predicting Mixing Depth, *Canadian Journal of Fisheries and Aquatic Sciences*, 47, 940–947,  
<https://doi.org/10.1139/f90-108>, 1998.
- Hassall, C.: The ecology and biodiversity of urban ponds, *Wiley Interdisciplinary Reviews: Water*, 1, 187–206,  
490 <https://doi.org/10.1002/wat2.1014>, 2014.
- Higgins, S. L., Thomas, F., Goldsmith, B., Brooks, S. J., Hassall, C., Harlow, J., Stone, D., Völker, S., and White, P.: Urban  
freshwaters, biodiversity, and human health and well-being: Setting an interdisciplinary research agenda, *WIREs Water*, 6, e1339,  
<https://doi.org/10.1002/wat2.1339>, 2019.
- Hill, M. J., Biggs, J., Thornhill, I., Briers, R. A., Gledhill, D. G., White, J. C., Wood, P. J., and Hassall, C.: Urban ponds as an aquatic  
495 biodiversity resource in modified landscapes, *Global Change Biology*, 23, 986–999, <https://doi.org/10.1111/gcb.13401>, 2017.
- Hodges, B.: Hydrodynamical Modeling, in: *Encyclopedia of Inland Waters*, <https://doi.org/10.1016/B978-0-12-409548-9.09123-5>, 2014.
- Idso, S. B.: On the concept of lake stability, *Limnology and Oceanography*, 18, 681–683, <https://doi.org/10.4319/lo.1973.18.4.0681>, 1973.
- Jankowski, T., Livingstone, D. M., Bührer, H., Forster, R., and Niederhauser, P.: Consequences of the 2003 European heat wave for lake tem-  
perature profiles, thermal stability, and hypolimnetic oxygen depletion: Implications for a warmer world, *Limnology and Oceanography*,  
500 51, 815–819, <https://doi.org/10.4319/lo.2006.51.2.0815>, 2006.
- Jenny, J.-P., Francus, P., Normandeau, A., Lapointe, F., Perga, M.-E., Ojala, A., Schimmelmann, A., and Zolitschka, B.: Global spread of  
hypoxia in freshwater ecosystems during the last three centuries is caused by rising local human pressure, *Global Change Biology*, 22,  
1481–1489, <https://doi.org/10.1111/gcb.13193>, 2016.
- Jones, J. and Brett, M. T.: Lake Nutrients, Eutrophication, and Climate Change, in: *Global Environmental Change*, edited by Freedman, B.,  
505 *Handbook of Global Environmental Pollution*, pp. 273–279, Springer Netherlands, Dordrecht, [https://doi.org/10.1007/978-94-007-5784-4\\_109](https://doi.org/10.1007/978-94-007-5784-4_109), 2014.
- Kendall, M.: Rank correlation methods, Rank correlation methods, Griffin, Oxford, England, 1975.
- Kerimoglu, O. and Rinke, K.: Stratification dynamics in a shallow reservoir under different hydro-meteorological scenarios and operational  
strategies, *Water Resour. Res.*, 49, 7518–7527, <https://doi.org/10.1002/2013WR013520>, 2013.
- 510 Kraemer, B., Anneville, O., Chandra, S., Dix, M., Kuusisto, E., Livingstone, D., Rimmer, A., Schladow, S., Silow, E., Sitoki, L., Tamatamah,  
R., Vadeboncoeur, Y., and McIntyre, P.: Morphometry and average temperature affect lake stratification responses to climate change: Lake  
Stratification Responses to Climate, *Geophysical Research Letters*, 42, <https://doi.org/10.1002/2015GL064097>, 2015.
- Layden, A., MacCallum, S., and Merchant, C.: Determining lake surface water temperatures worldwide using a tuned one-dimensional lake  
model (FLake, v1), *Geoscientific Model Development*, 9, 2167–2189, <https://doi.org/10.5194/gmd-9-2167-2016>, 2016.
- 515 Lee, R., Biggs, T., and Fang, X.: Thermal and Hydrodynamic Changes under a Warmer Climate in a Variably Stratified Hypereutrophic  
Reservoir, *Water*, 10, 1–24, <https://doi.org/10.3390/w10091284>, 2018.



- Leendertse, J. J.: Aspects of a Computational Model for Long-Period Water-Wave Propagation:, 1967.
- Lemonsu, A., Kounkou-Arnaud, R., Desplat, J., Salagnac, J.-L., and Masson, V.: Evolution of the Parisian urban climate under a global changing climate, *Climatic Change*, 116, <https://doi.org/10.1007/s10584-012-0521-6>, 2013.
- 520 Lewandowska, A. and Sommer, U.: Climate change and the spring bloom: a mesocosm study on the influence of light and temperature on phytoplankton and mesozooplankton, *MEPS*, 405, 101–111, <https://doi.org/10.3354/meps08520>, 2010.
- Livingstone, D. M.: Impact of Secular Climate Change on the Thermal Structure of a Large Temperate Central European Lake, *Climatic Change*, 57, 205–225, <https://doi.org/10.1023/A:1022119503144>, 2003.
- Magee, M. R. and Wu, C. H.: Response of water temperatures and stratification to changing climate in three lakes with different morphometry, 525 *Hydrology and Earth System Sciences*, 21, 6253–6274, <https://doi.org/10.5194/hess-21-6253-2017>, 2017.
- Mann, H. B.: Nonparametric Tests Against Trend, *Econometrica*, 13, 245–259, <https://doi.org/10.2307/1907187>, 1945.
- Mariani, L., Parisi, S. G., Cola, G., and Failla, O.: Climate change in Europe and effects on thermal resources for crops, *Int J Biometeorol*, 56, 1123–1134, <https://doi.org/10.1007/s00484-012-0528-8>, 2012.
- McEnroe, N. A., Buttle, J. M., Marsalek, J., Pick, F. R., Xenopoulos, M. A., and Frost, P. C.: Thermal and chemical stratification of urban 530 ponds: Are they ‘completely mixed reactors’?, *Urban Ecosyst*, 16, 327–339, <https://doi.org/10.1007/s11252-012-0258-z>, 2013.
- Mendonça, R., Müller, R. A., Clow, D., Verpoorter, C., Raymond, P., Tranvik, L. J., and Sobek, S.: Organic carbon burial in global lakes and reservoirs, *Nature Communications*, 8, 1694, <https://doi.org/10.1038/s41467-017-01789-6>, 2017.
- Moras, S., Ayala, A. I., and Pierson, D. C.: Historical modelling of changes in Lake Erken thermal conditions, *Hydrology and Earth System Sciences*, 23, 5001–5016, <https://doi.org/10.5194/hess-23-5001-2019>, 2019.
- 535 Murakami, M., Oonishi, Y., and Kunishi, H.: A numerical simulation of the distribution of water temperature and salinity in the Seto Inland Sea, *Journal of the Oceanographical Society of Japan*, 41, 213–224, <https://doi.org/10.1007/BF02109271>, 1985.
- Neuheimer, A. B. and Taggart, C. T.: The growing degree-day and fish size-at-age: The overlooked metric, *Canadian Journal of Fisheries and Aquatic Sciences*, 64, 375–385, 2007.
- Niedrist, G., Psenner, R., and Sommaruga, R.: Climate warming increases vertical and seasonal water temperature differences, and inter- 540 annual variability in a mountain lake, *Climatic Change*, <https://doi.org/10.1007/s10584-018-2328-6>, 2018.
- Noble, A. and Hassall, C.: Poor ecological quality of urban ponds in northern England: causes and consequences, *Urban Ecosystems*, 18, 649–662, <https://doi.org/10.1007/s11252-014-0422-8>, 2015.
- O’Reilly, C. M. et al.: Rapid and highly variable warming of lake surface waters around the globe, *Geophysical Research Letters*, 42, 10,773–10,781, <https://doi.org/10.1002/2015GL066235>, 2015.
- 545 Paerl, H. and Huisman, J.: Blooms Like It Hot, *Science (New York, N.Y.)*, 320, 57–8, <https://doi.org/10.1126/science.1155398>, 2008.
- Paerl, H. W. and Paul, V. J.: Climate change: Links to global expansion of harmful cyanobacteria, *Water Research*, 46, 1349–1363, <https://doi.org/10.1016/j.watres.2011.08.002>, 2012.
- Perrier, F., Le Mouél, J., Poirier, J., and Shnirman, M.: Long-term climate change and surface versus underground temperature measurements in Paris, *Long-term climate change and surface versus underground temperature measurements in Paris*, 25, 1619–1631, 2005.
- 550 Pfeifroth, U., Sanchez-Lorenzo, A., Manara, V., Trentmann, J., and Hollmann, R.: Trends and Variability of Surface Solar Radiation in Europe Based On Surface- and Satellite-Based Data Records, *Journal of Geophysical Research: Atmospheres*, 123, 1735–1754, <https://doi.org/10.1002/2017JD027418>, 2018.
- Piccolroaz, S., Amadori, M., Toffolon, M., and Dijkstra, H. A.: Importance of planetary rotation for ventilation processes in deep elongated lakes: Evidence from Lake Garda (Italy), *Scientific Reports*, 9, 1–11, <https://doi.org/10.1038/s41598-019-44730-1>, 2019.



- 555 Piccolroaz, S., Woolway, R. I., and Merchant, C. J.: Global reconstruction of twentieth century lake surface water temperature reveals different warming trends depending on the climatic zone, *Climatic Change*, <https://doi.org/10.1007/s10584-020-02663-z>, 2020.
- Poole, H. H. and Atkins, W. R. G.: Photo-electric Measurements of Submarine Illumination throughout the Year, *Journal of the Marine Biological Association of the United Kingdom*, 16, 297–324, <https://doi.org/10.1017/S0025315400029829>, 1929.
- 560 Quintana-Seguí, P., Le Moigne, P., Durand, Y., Martin, E., Habets, F., Baillon, M., Canellas, C., Franchisteguy, L., and Morel, S.: Analysis of Near-Surface Atmospheric Variables: Validation of the SAFRAN Analysis over France, *J. Appl. Meteor. Climatol.*, 47, 92–107, <https://doi.org/10.1175/2007JAMC1636.1>, 2008.
- Raimonet, M., Oudin, L., Thieu, V., Silvestre, M., Vautard, R., Rabouille, C., and Le Moigne, P.: Evaluation of Gridded Meteorological Datasets for Hydrological Modeling, *J. Hydrometeorol.*, 18, 3027–3041, <https://doi.org/10.1175/JHM-D-17-0018.1>, 2017.
- 565 Ralston, D. K., Keafer, B. A., Brosnahan, M. L., and Anderson, D. M.: Temperature dependence of an estuarine harmful algal bloom: Resolving interannual variability in bloom dynamics using a degree day approach, *Limnol. Oceanogr.*, 59, 1112–1126, <https://doi.org/10.4319/lo.2014.59.4.1112>, 2014.
- Reid, P. C., Hari, R. E., Beaugrand, G., Livingstone, D. M., Marty, C., Straile, D., Barichivich, J., Goberville, E., Adrian, R., Aono, Y., Brown, R., Foster, J., Groisman, P., Hélaouët, P., Hsu, H.-H., Kirby, R., Knight, J., Kraberg, A., Li, J., Lo, T.-T., Myneni, R. B., North, R. P., Pounds, J. A., Sparks, T., Stübi, R., Tian, Y., Wiltshire, K. H., Xiao, D., and Zhu, Z.: Global impacts of the 1980s regime shift, 570 *Global Change Biology*, 22, 682–703, <https://doi.org/10.1111/gcb.13106>, 2016.
- Schlenker, W., Hanemann, W. M., and Fisher, A. C.: Water Availability, Degree Days, and the Potential Impact of Climate Change on Irrigated Agriculture in California, *Climatic Change*, 81, 19–38, <https://doi.org/10.1007/s10584-005-9008-z>, 2007.
- Schmid, M., Hunziker, S., and Wüest, A.: Lake surface temperatures in a changing climate: a global sensitivity analysis, *Climatic Change*, 124, 301–315, <https://doi.org/10.1007/s10584-014-1087-2>, 2014.
- 575 Schneider, P. and Hook, S. J.: Space observations of inland water bodies show rapid surface warming since 1985, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL045059>, 2010.
- Schneider, P., Hook, S. J., Radocinski, R. G., Corlett, G. K., Hulley, G. C., Schladow, S. G., and Steissberg, T. E.: Satellite observations indicate rapid warming trend for lakes in California and Nevada, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL040846>, 2009.
- 580 Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall’s Tau, *Journal of the American Statistical Association*, 63, 1379–1389, <https://doi.org/10.1080/01621459.1968.10480934>, 1968.
- Sommer, U. and Lengfellner, K.: Climate change and the timing, magnitude, and composition of the phytoplankton spring bloom, *Global Change Biology*, 14, 1199–1208, <https://doi.org/10.1111/j.1365-2486.2008.01571.x>, 2008.
- Song, K., Xenopoulos, M. A., Buttle, J. M., Marsalek, J., Wagner, N. D., Pick, F. R., and Frost, P. C.: Thermal stratification patterns in urban ponds and their relationships with vertical nutrient gradients, *Journal of Environmental Management*, 127, 317–323, 585 <https://doi.org/10.1016/j.jenvman.2013.05.052>, 2013.
- Soullignac, F., Vinçon-Leite, B., Lemaire, B. J., Scarati Martins, J. R., Bonhomme, C., Dubois, P., Mezemate, Y., Tchiguirinskaia, I., Schertzer, D., and Tassin, B.: Performance Assessment of a 3D Hydrodynamic Model Using High Temporal Resolution Measurements in a Shallow Urban Lake, *Environ Model Assess*, 22, 309–322, <https://doi.org/10.1007/s10666-017-9548-4>, 2017.
- 590 Soullignac, F., Danis, P.-A., Bouffard, D., Chanudet, V., Dambrine, E., Guénand, Y., Harmel, T., Ibelings, B. W., Trevisan, D., Uittenbogaard, R., and Anneville, O.: Using 3D modeling and remote sensing capabilities for a better understanding of spatio-temporal heterogeneities of phytoplankton abundance in large lakes, *Journal of Great Lakes Research*, 44, 756–764, <https://doi.org/10.1016/j.jglr.2018.05.008>, 2018.



- Stelling, G. S. and Leendertse, J. J.: Approximation of Convective Processes by Cyclic AOI Methods, pp. 771–782, ASCE, 1992.
- Torralba, V., Doblas-Reyes, F. J., and Gonzalez-Reviriego, N.: Uncertainty in recent near-surface wind speed trends: a global reanalysis  
595 intercomparison, *Environ. Res. Lett.*, 12, 114 019, <https://doi.org/10.1088/1748-9326/aa8a58>, 2017.
- Townsend, D. W., Cammen, L. M., Holligan, P. M., Campbell, D. E., and Pettigrew, N. R.: Causes and consequences of variability in the timing of spring phytoplankton blooms, *Deep Sea Research Part I: Oceanographic Research Papers*, 41, 747–765, [https://doi.org/10.1016/0967-0637\(94\)90075-2](https://doi.org/10.1016/0967-0637(94)90075-2), 1994.
- Tran Khac, V., Hong, Y., Plec, D., Lemaire, B. J., Dubois, P., Saad, M., and Vinçon-Leite, B.: An Automatic Monitoring System  
600 for High-Frequency Measuring and Real-Time Management of Cyanobacterial Blooms in Urban Water Bodies, *Processes*, 6, 11, <https://doi.org/10.3390/pr6020011>, 2018.
- Tímea, K., Kovács-Székely, I., and Anda, A.: Comparison of parametric and non-parametric time-series analysis methods on a long-term meteorological data set, *Central European Geology*, 60, 1–17, <https://doi.org/10.1556/24.60.2017.011>, 2017.
- van den Bosch, M. and Sang, A. o.: 2096 - Urban Natural Environments As Nature Based Solutions for Improved Public Health – a Systematic  
605 Review of Reviews, *Journal of Transport & Health*, 5, S79, <https://doi.org/10.1016/j.jth.2017.05.230>, 2017.
- Vautard, R., Cattiaux, J., Yiou, P., Thépaut, J.-N., and Ciais, P.: Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness, *Nature Geoscience*, 3, 756–761, <https://doi.org/10.1038/ngeo979>, 2010.
- Verpoorter, C., Kutser, T., Seekell, D. A., and Tranvik, L. J.: A global inventory of lakes based on high-resolution satellite imagery, *Geophysical Research Letters*, 41, 6396–6402, <https://doi.org/10.1002/2014GL060641>, 2014.
- 610 Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., and Soubeyroux, J.-M.: A 50-year high-resolution atmospheric reanalysis over France with the Safran system, *International Journal of Climatology*, 30, 1627–1644, <https://doi.org/10.1002/joc.2003>, 2010.
- Vinçon-Leite, B., Lemaire, B. J., Khac, V. T., and Tassin, B.: Long-term temperature evolution in a deep sub-alpine lake, Lake Bourget, France: how a one-dimensional model improves its trend assessment, *Hydrobiologia*, 731, 49–64, <https://doi.org/10.1007/s10750-014-1818-4>, 2014.
- 615 Wagner, C. and Adrian, R.: Cyanobacteria dominance: Quantifying the effects of climate change, *Limnology and Oceanography*, 54, 2460–2468, [https://doi.org/10.4319/lo.2009.54.6\\_part\\_2.2460](https://doi.org/10.4319/lo.2009.54.6_part_2.2460), 2009.
- Wagner, T. and Erickson, L.: Sustainable Management of Eutrophic Lakes and Reservoirs, *Journal of Environmental Protection*, 08, 436–463, <https://doi.org/10.4236/jep.2017.84032>, 2017.
- Wang, F., Shao, W., Yu, H., Kan, G., He, X., Zhang, D., Ren, M., and Wang, G.: Re-evaluation of the Power of the Mann-Kendall Test for  
620 Detecting Monotonic Trends in Hydrometeorological Time Series, *Front. Earth Sci.*, 8, <https://doi.org/10.3389/feart.2020.00014>, 2020.
- Wilhelm, S. and Adrian, R.: Impact of summer warming on the thermal characteristics of a polymictic lake and consequences for oxygen, nutrients and phytoplankton, *Freshwater Biology*, 53, 226–237, <https://doi.org/10.1111/j.1365-2427.2007.01887.x>, 2008.
- Wilkinson, A. A., Hondzo, M., and Guala, M.: Vertical heterogeneities of cyanobacteria and microcystin concentrations in lakes using a seasonal In situ monitoring station, *Global Ecology and Conservation*, 21, e00 838, <https://doi.org/10.1016/j.gecco.2019.e00838>, 2020.
- 625 Winder, M. and Sommer, U.: Phytoplankton response to a changing climate, *Hydrobiologia*, 698, 5–16, <https://doi.org/10.1007/s10750-012-1149-2>, 2012.