



**HAL**  
open science

## Spatial heterogeneity of soil quality within a Mediterranean alley cropping agroforestry system: Comparison with a monocropping system

Esther Guillot, Isabelle Bertrand, Cornelia Rumpel, Claudia Gomez, Didier Arnal, Josiane Abadie, Philippe Hinsinger

### ► To cite this version:

Esther Guillot, Isabelle Bertrand, Cornelia Rumpel, Claudia Gomez, Didier Arnal, et al.. Spatial heterogeneity of soil quality within a Mediterranean alley cropping agroforestry system: Comparison with a monocropping system. *European Journal of Soil Biology*, 2021, 105, pp.103330. 10.1016/j.ejsobi.2021.103330 . hal-03251274

**HAL Id: hal-03251274**

**<https://hal.inrae.fr/hal-03251274>**

Submitted on 13 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Spatial heterogeneity of soil quality within a Mediterranean alley cropping**  
2 **agroforestry system: comparison with a monocropping system**

3

4 - Esther Guillot <sup>a</sup>

5 - Isabelle Bertrand <sup>a</sup>

6 - Cornelia Rumpel <sup>b</sup>

7 - Claudia Gomez <sup>b</sup>

8 - Didier Arnal <sup>a</sup>

9 - Josiane Abadie <sup>a</sup>

10 - Philippe Hinsinger <sup>a</sup>

11

12 *<sup>a</sup>Eco&Sols, Univ Montpellier, CIRAD, INRAE, IRD, Institut Agro, Montpellier, France*

13 *<sup>b</sup>CNRS, UMR 7618, IESS, Bâtiment EGER, F-78850 Thiverval Grignon, France*

14

15 **Corresponding author:** Isabelle Bertrand, Eco&Sols, Univ Montpellier, CIRAD, INRAE,  
16 IRD, Institut Agro, Montpellier, France

17 E-mail: [isabelle.bertrand@inrae.fr](mailto:isabelle.bertrand@inrae.fr)

18

19 **Highlights**

20 - Temperate alley cropping systems induce spatial heterogeneity in soil quality

21 - Soil quality is improved until 2 m beyond the tree row in the interrow

22 - 21 years of agroforestry increased the soil quality by 20% compared with monocrop

23 **Abstract**

24

25 Alley cropping agroforestry systems are complex agroecosystems highlighted for their  
26 positive effects on soil quality. However, the potential spatial heterogeneity of soil quality  
27 created by tree rows at the plot scale has seldom been studied. The aim of this study was to  
28 evaluate soil quality at the plot scale, under tree rows and along transects perpendicular to the  
29 tree row and to compare alley cropping systems with monocropping systems. This study was  
30 performed on an alley cropping system that combined hybrid walnut trees (21 years old) and  
31 peas. Topsoil was sampled at tree rows between 1 and 2, 2 and 4 and 4 and 6.5 m from the  
32 tree row in the alley cropping system, as well as in a neighbouring monocropping plot.  
33 Physical, chemical and microbiological indicators of soil quality were measured. Tree row  
34 implantation induced spatial heterogeneity in the chemical indicators, microbial biomass,  
35 activities and community structure at the alley cropping plot scale. Alley cropping not only  
36 improved microbiological soil quality indicators within the tree rows but also in the interrows  
37 when compared to a monocropping system. These indicators were then integrated into one  
38 soil quality index (SQI) built through a statistical approach. The soil quality index was  
39 calculated for the monocropping plot and for each position within the alley cropping plot.  
40 After 21 years of agroforestry practice, tree rows and permanent grass cover improved the  
41 SQI until 2 m in the interrow. Weighted SQIs were calculated relative to the surface area of  
42 each location for the entire alley cropping plot (i.e., tree row + interrow positions) and for the  
43 entire alley cropping interrow (i.e., removing the tree row surface area). The weighted SQI of  
44 the entire alley cropping plot significantly increased compared with that of the monocropping  
45 plot.

46

47 **Keywords:** Alley cropping system, spatial heterogeneity, soil quality index, microbial  
48 activities, soil organic carbon

## 49 **1. Introduction**

50

51 Soil quality can be defined as “the continued capacity of the soil to function within  
52 ecological and land-use boundaries, to sustain productivity, to promote the quality of air and  
53 water, and to maintain plant, animal and human health” [1]. Soil quality indicators are  
54 parameters that are sensitive enough to be modified by land use or management practices [2],  
55 [3] and should provide key information concerning the composition, structure and function of  
56 the soil [4]. In addition, a combination of physical, chemical and biological indicators must be  
57 used to correctly interpret soil quality [5], [6].

58 Physical indicators are related to the “inherent quality”, which is influenced by soil  
59 age and past climates [7], while most chemical and all biological indicators can be associated  
60 with the “dynamic quality”, which is highly sensitive to land use and soil management [8].  
61 Inherent and dynamic soil quality indicators refer to changes over long and medium to short  
62 terms, respectively [9]. Among physical soil quality indicators, water storage, bulk density,  
63 aggregate stability and texture are the most frequently measured [2]. Chemical indicators can  
64 be evaluated to characterize soil fertility and nutrient availability (e.g., soil organic carbon,  
65 total N, Olsen P, CEC, pH, exchangeable ions, and N mineralization rate) [10]. Biological  
66 indicators of soil quality are necessary to connect abiotic soil properties to soil functioning  
67 [2], [11]. Soil microbial biomass, abundance, diversity and activity play key roles as the main  
68 drivers of soil organic matter decomposition and nutrient cycling [12], [13]. Soil  
69 microorganisms can be studied as soil quality indicators through their biomass (e.g., microbial  
70 biomass C), activities (e.g., enzyme activities and substrate-induced respiration) and  
71 community structure (e.g., PLFA and sequencing) [14]. A minimum dataset of indicators has

72 to be selected either through expert opinion [15] or through statistical approaches [5], [16] to  
73 assess soil quality, avoiding collinearity between indicators. Both methods allow the  
74 construction of a soil quality index (SQI), which accounts for a more or less comprehensive  
75 set of indicators [10], [17]. The use of aggregated SQIs is currently strongly developed and  
76 can improve the comparisons of land use or management effects.

77 Land management is recognized as one of the main drivers affecting soil quality  
78 improvement or degradation [18]. Temperate agroforestry systems comprising a combination  
79 of trees and crops are examples of agroecosystems that provide a sustainable alternative to  
80 conventional cropping systems, in which plant diversity is low; hereafter, these conventional  
81 cropping systems are called monocropping systems. A recent review of European agroforestry  
82 practices confirmed their positive effect on ecosystem services; however, this effect was  
83 context-dependent [19]. One of the most important ecosystem services promoted by  
84 agroforestry systems is their potential capacity for carbon (C) storage, both temporarily in tree  
85 biomass through photosynthesis [20] and in the long term in soil organic matter [21]. High  
86 inputs of above- and belowground litter in the tree row in alley cropping can provide higher  
87 amounts of C and nutrient resources and thereby modify soil microbial communities [22].

88 Even though a global positive effect of agroforestry on soil quality has been  
89 recognized [23], it should be noted that only a small proportion of studies have considered the  
90 spatial heterogeneity induced by tree rows. As highlighted by Cardinael et al. [24], future  
91 studies should integrate more spatial dynamics and investigate lateral spatial heterogeneities  
92 instead of concentrating on only the tree row. The few publications that have considered the  
93 potential spatial heterogeneity of soil quality have not allowed us to claim that agroforestry  
94 systems significantly improve soil quality at the plot scale or within the interrow. For  
95 instance, [25] did not detect differences between tree rows and interrows in microbial biomass  
96 C (MBC) and basal respiration or [26] in bacterial diversity or [27] in enzyme activities.

97 Mungai et al. [28] found significantly more enzyme activities in tree rows than in interrows  
98 but no difference within cultivated interrows. The spatial extent to which tree rows and their  
99 understorey vegetation strips affect soil quality should be investigated.

100 In the present work, we evaluated the effect of an alley cropping agroforestry system  
101 on the physical, chemical and biological components of soil quality compared to a  
102 neighbouring, monocropping system managed in the same way as the interrows in the alley  
103 cropping system. We first hypothesized that spatial gradients of soil quality occur between the  
104 tree row and middle of the interrow, with the greatest values of soil quality indicators on and  
105 close to the tree row and decreasing values with increasing distance from the tree row.  
106 Second, we hypothesized that the weighted soil quality of alley cropping, integrating its  
107 spatial heterogeneity throughout the plot, is higher than that of the neighbouring  
108 monocropping plot.

109

## 110 **2. Materials and methods**

### 111 *2.1. Site description*

112 The experimental site of Restinclières is located at Prades-le-Lez, 15 km north of  
113 Montpellier, southern France (43°42'15 N, 3°51'41 E). This location has a subhumid  
114 Mediterranean climate, with an average temperature of 15.4 °C and average annual rainfall of  
115 658 mm between 2000 and 2016 (Fréjorgues weather station). The soil is a deep Fluvisol  
116 (WRB, 2007) with a silty-clay texture. In February 1995, hybrid walnut trees (*Juglans regia* ×  
117 *nigra* cv. NG23) were planted 4 m apart along east-west rows, with an interrow distance of 13  
118 m. The initial planting density was 208 trees ha<sup>-1</sup>, but the plot was thinned in 2004 down to a  
119 present density of 110 trees ha<sup>-1</sup>. Adult walnut trees have not been pruned since 2014. The  
120 tree rows (2 m wide) were covered with spontaneous herbaceous vegetation, as they have not  
121 been ploughed or treated with herbicides since 1995. Each interrow was ploughed to a 20 cm

122 depth every year before the winter crop was sown and was fertilized at an approximate rate of  
123 150 kg N ha<sup>-1</sup> yr<sup>-1</sup> (as NH<sub>4</sub>NO<sub>3</sub>). Most of the time since 2000, durum wheat has been the  
124 major crop in rotation, but pea has been introduced every 4 years since 2010, and barley has  
125 been introduced since 2015 [29]. Over the 2014-2016 period, durum wheat (*Triticum*  
126 *turgidum durum* cv. Claudio)-barley (*Hordeum vulgare* cv. Augusta)-pea (*Pisum sativum* cv.  
127 Igloo) rotation was practised on this 4.6-ha alley cropping plot and on the adjacent 1.4-ha  
128 monocropping plot, which has been managed in the same way as the cultivated interrows in  
129 the alley cropping since the experimental site was established. All crops were sown between  
130 the end of October and the beginning of December and harvested by the end of June/early  
131 July.

132

## 133 2.2. Soil sampling

134 Soil sampling was conducted in April 2016, when pea was growing in the  
135 monocropping plot and in the interrows of the alley cropping plot. At this stage, the walnut  
136 trees had no leaves yet. In the 1.4 ha monocropping plot, five sampling areas of 1 m<sup>2</sup> each, at  
137 least 15 m apart, were identified for soil sampling. In each 1 m<sup>2</sup> surface area, one composite  
138 soil sample, based on 4 soil cores (8 cm in diameter) from each angle of the area, was  
139 collected from the 0-15 cm layer of topsoil using a root auger. For the alley cropping plot,  
140 five transects at least 40 m apart were sampled from the tree row to the middle of the interrow  
141 on both the north and south sides of the tree row. Each north-south transect corresponded to 7  
142 positions (35 soil samples in total) (**Fig. A.1**). Soils were sampled in the tree row, with the  
143 centre of the 2-m<sup>2</sup> sampling area located on the axis of the tree row, 1 m from the nearest tree.  
144 Other sampling areas were located in the interrow between 1 and 2 m from the middle of the  
145 tree row, between 2 and 4 m and between 4 and 6.5 m. Soil samples were collected from the  
146 0-15 cm layer of topsoil using a root auger at each angle of a centred subsurface (representing

147 half of the total surface area of the location) (**Fig. A.1**). Areas sampled on the north side are  
148 indicated by “N”, and those on the south side are indicated by “s”. Once collected, the fresh  
149 soil samples were passed through a 2-mm sieve, and subsamples were stored at 4 °C for  
150 microbial biomass and MicroResp® analyses, which were performed within two weeks after  
151 sampling and at -20 °C prior to freeze-drying for PLFA analyses.

152

### 153 *2.3. Physical and chemical analyses*

154 Physical and chemical indicators were measured on the north side of the trees  
155 (corresponding to 4 positions per transect) and in the monocropping system, i.e., for a total of  
156 25 samples. The moisture content was determined for all samples after drying for 48 h at 105  
157 °C. Bulk density was determined by dividing the soil dry mass by the volume of the root  
158 auger. Additional analyses were performed by the national routine soil testing lab of INRAE  
159 at Arras (LAS, France). Soil texture was determined after CaCO<sub>3</sub> dissolution, and pH was  
160 determined in a water extract. The cation exchange capacity (CEC) was determined using the  
161 cobalt hexamine chloride method and analysed using inductively coupled plasma-atomic  
162 emission spectrometry (ICP-AES) [30]. Soil organic C (SOC) and total N were determined by  
163 dry combustion. The available P content was determined colorimetrically after extraction  
164 using the Olsen method [31]. Roots were collected from each soil core (15 cm height × 8 cm  
165 diameter) and pooled into one composite sample (4 cores) for each area. They were separated  
166 into three size classes as follows: 0-2 mm (fine roots), 2-5 mm (medium roots) and > 5 mm  
167 (large roots).

168

#### 169 *2.3.1. Density and particle size fractionation*

170 Density fractionation was performed on the north side of the trees and in the  
171 monocropping system, i.e., for a total of 25 soil samples. The fractionation procedure was



172 derived from Roscoe et al. [32]. Briefly, 25 g of soil was shaken with 100 ml of a 1.6 Mg m<sup>-3</sup>  
173 dense sodium polytungstate solution (NaPT) and then centrifuged for 30 min at 6800 g. The  
174 suspended material was termed the free light fraction (f-LF < 1.6 Mg m<sup>-3</sup>) and was separated  
175 from the supernatant by filtration using a 0.7 µm glass fibre filter. The material recovered on  
176 the filter paper was washed with distilled water to remove residual NaPT. New NaPT (density  
177 1.6 Mg m<sup>-3</sup>) was added to the remaining material. The new suspension was subjected to  
178 ultrasonic dispersion (25 J ml) for 3 min. Afterwards, the suspension was centrifuged at 6800  
179 g for 30 min, and the supernatant was filtered to recover the occluded light fraction (o-LF <  
180 1.6 Mg m<sup>-3</sup>) (procedure see above). The remaining material was then separated into two  
181 particle-size fractions by wet sieving: 50-2000 µm (sand) and < 50 µm (silt and clay). Before  
182 C and N measurements, each fraction was decarbonated by acid fumigation using 12 M HCl  
183 in a desiccator for 6 h [33]. After removing calcium carbonate, C and N contents in the total  
184 soil, the f-LF o-LF and 50-2000 µm and < 50 µm fractions were analysed by dry combustion  
185 with an elemental analyser (Elemental Analyser Vario Pyro Cube). This density fractionation  
186 procedure allowed the separation of free soil organic matter (f-LF), soil organic matter  
187 associated with minerals (o-LF and 50-2000 µm) and physically protected soil organic matter  
188 (< 50 µm).

189

### 190 2.3.2. *Non-cellulosic neutral sugar determination*

191 Rhamnose, fucose, mannose, galactose, glucose, ribose, arabinose and xylose were  
192 quantified in the total soil, f-LF, o-LF and < 50 µm fractions following the protocol of  
193 Rumpel and Dignac [34] modified through the addition of 0.9 ml of 2 M EDTA after  
194 hydrolysis with trifluoroacetic acid (TFA) [35]. Briefly, 500 mg for total soil, between 60 and  
195 70 mg for f-LF and o-LF and 600 mg for the < 50 µm size fraction were added to 10 ml of  
196 TFA and hydrolysed at 105 °C for 4 h. After hydrolysis, myoinositol was added as an internal

197 standard. Samples were filtered through a glass fibre filter (0.7  $\mu\text{m}$ ), and TFA was eliminated  
198 by evaporation before derivatization. Aldoses were reduced to their corresponding alditols  
199 after the addition of 1 ml of  $\text{NaBH}_4$  dissolved in dimethyl sulfoxide (DMSO). Acetylation  
200 was performed by adding 2 ml of acetic anhydride and 2 ml of glacial acetic acid using  
201 methylimidazole (2 ml) as the catalyst. The reaction was stopped after 10 min by adding 7 ml  
202 ice-cold deionized water. The derivatised sugar monomers were extracted with 1 ml of  
203 dichloromethane. The analyses were performed using a gas chromatograph (HP 6890 GC-  
204 FID) equipped with an SGE BPX-70 column (60 m  $\times$  0.32 mm internal diameter, 0.25  $\mu\text{m}$   
205 film thickness). The gas chromatography oven temperature program was 200  $^\circ\text{C}$  to 250  $^\circ\text{C}$  at  
206 8  $^\circ\text{C min}^{-1}$  and isothermal at 250  $^\circ\text{C}$  for 15 min with helium as the carrier gas.

207 The sum of extracted monosaccharides from plant-derived hemicellulose and  
208 microbial products [36] is hereafter termed “sugars”. The proportion of microorganism-  
209 derived sugars in relation to plant-derived sugars can be estimated from the ratio of  
210 hexose:pentose sugars: (galactose + mannose):(arabinose + xylose), hereafter called GM:AX  
211 [37]. GM:AX ratios  $< 0.5$  and  $> 2$  are representative of carbohydrates predominantly derived  
212 from plants and microorganisms, respectively [37].

213

#### 214 *2.4. Soil microbiological indicators*

215 The study plot had an east-west tree row orientation, and we wanted to test whether  
216 this orientation could modify soil microbiological parameters, which are sensitive to changes  
217 in climatic conditions at the local scale. Accordingly, we measured microbiological  
218 parameters on both the north and south sides of the tree row., i.e., for a total of 40 samples.

219 Microbial biomass C (MBC), N (MBN) and P (MBP) contents were quantified using  
220 the chloroform fumigation-extraction technique [38]. Briefly, 10 g (for MBC and MBN) or 2  
221 g (for MBP) of equivalent dry soil was exposed to chloroform vapor for 24 h and then

222 extracted with 40 ml of 0.025 M K<sub>2</sub>SO<sub>4</sub> (for MBC and MBN) or with 40 ml 0.5 M NaHCO<sub>3</sub>  
223 (for MBP), shaken for 45 min, centrifuged (10 min at 2683 g) and then filtered through 0.22-  
224 µm PTFE filters. Soil organic C and N concentrations in the extracts were measured by a  
225 TOC/TON analyser (OI-Analytical, Aurora 1030, College Station, USA). Inorganic P  
226 concentrations in the extracts were quantified colorimetrically using the malachite green  
227 method [39]. Microbial biomass C, N, and P (MBC, MBN and MBP) contents were  
228 calculated from the difference between the chloroform-fumigated and non-fumigated samples.  
229 We applied a conversion factor of 0.45 for MBC [40], 0.54 for MBN [41] and a conversion  
230 factor of 0.4 for MBP [40]. Data are expressed in mg C, N or P kg soil<sup>-1</sup>. The total C of  
231 nonfumigated samples was used to represent dissolved organic carbon (DOC).

232 Basal and substrate-induced respiration was measured using the MicroResp<sup>TM</sup>  
233 approach [42]. The soil water holding capacity (WHC) was determined using Richard's  
234 membrane press at pF 2.5 [43]. Sieved soils were adjusted to 40% of their WHC and were  
235 preincubated for 7 days at a temperature of 23 °C ± 2 °C in the dark following the protocol of  
236 Bérard et al. [44]. We tested three different soluble C substrates: glucose, trehalose and  
237 alanine [44]. In a 96-deep-well microplate, 350 mg of soil sample was distributed, and then 25  
238 µl of each substrate was dispensed in each of the wells at 10 µg substrate mg<sup>-1</sup> soil. The 96-  
239 deep-well microplate was then sealed with a CO<sub>2</sub>-trap gel and incubated for 6 h in dark  
240 conditions at 23 ± 2 °C. The optical density of each well of the CO<sub>2</sub>-trap gel was measured on  
241 a fluorometric microplate reader (Victor 3, Perkin Elmer) at 570 nm.

242 The microbial community structure was assessed using the phospholipid fatty acid  
243 (PLFA) method described by [45]. All analyses of PLFAs were performed by Microbial iD,  
244 Inc. (Newark, USA). Briefly, lipids were extracted from 5 g of freeze-dried soil by using a  
245 modified Bligh-Dyer extraction with 19 ml of extractant. Lipids were separated on a solid-  
246 phase extraction column, and phospholipids were eluted with 5 ml of methanol. After

247 evaporation, phospholipids were transesterified to fatty acid methyl esters, extracted in 4 ml of  
248 hexan, evaporated again and then analysed using a gas chromatograph (Agilent 6890  
249 Technologies, Wilmington, USA) [45]. PLFA peaks were identified using the MIDI PLFAD1  
250 calibration mix and naming table (MIDI, Inc., Newark, USA). Individual PLFA markers were  
251 used to determine the number of peaks identified per position (tree row, 1-2 m<sub>N/S</sub>, 2-4 m<sub>N/S</sub>,  
252 4-6.5 m<sub>N/S</sub> for alley cropping and monocropping). Gram-positive (GP) bacteria were  
253 identified by 13:0 iso, 14:0 iso, 15:1 iso w6c, 15:1 anteiso w9c, 15:0 iso, 15:0 anteiso, 16:0  
254 iso, 16:0 anteiso, 17:1 iso w9c, 17:1 anteiso w9c, 17:1 anteiso w7c, 17:0 iso, 17:0 anteiso,  
255 18:0 iso, and 20:0 iso; gram-negative (GN) bacteria were identified by 13:f1 w5c, 13:1 w4c,  
256 14:1 w5c, 15:1 w6c, 16:1 w9c, 16:1 w7c, 17:1 w8c, 17:0 cyclo w7c, 18:1 w7c, 18:1 w5c, 19:1  
257 w8c, 19:0 cyclo w7c, 20:1 w9c, 20:1 w6c, 21:1 w8c, 21:1 w3c and 22:1 w3c (reference table  
258 from MIDI, Inc., Newark, DE). The peaks associated with 12:0, 14:0, 15:0, 16:0, 17:0, 18:0,  
259 20:0, 22:0, 23:0 and 24:0 were considered general indicators [46], and they were not  
260 integrated in analyses and were classified as “non-identified peaks” in **Table A.1**. The GP:GN  
261 ratio was calculated based on the previously described peaks. A relative measure of the  
262 fungi/bacteria (F:B) ratio was calculated by dividing the fungal PLFA marker: 18:2 w6c, 18:1  
263 w9c [46], [47], [48] and 16:1 w5c, the last one representing the arbuscular mycorrhizal fungi  
264 (AMF) by the sum of GN and GP bacteria. Actinobacteria were identified by 16:0 10-methyl,  
265 17:1 w7c 10-methyl, 17:0 10-methyl, 18:1 w7c 10-methyl, 18:0 10-methyl and 19:1 w7c 10-  
266 methyl. Other unknown peaks were associated with the “others” category and were  
267 considered only in the calculation of the relative abundance of each PLFA group. All PLFA  
268 markers are listed in **Table A.1**.

269

270 *2.5. Calculation of the soil quality index (SQI)*

271 We followed the method proposed by Obriot et al. [10] to build an aggregated index of  
272 soil quality. This method is based on 4 steps and starts with a dataset from different soil  
273 variables and leads to a unitless single score. The first step (step 1, [10]) consisted of the  
274 inventory of soil parameters. The second step (step 2, [10]) aimed to produce a minimum  
275 dataset (MDS) for each category. Only those parameters that significantly discriminated the  
276 various locations within the agroforestry plot (mixed model analysis,  $P < 0.05$ ) were  
277 considered relevant [10]. Then, a Pearson correlation analysis was performed to identify  
278 correlated parameters ( $> 0.8$ ) and to avoid redundant information in the aggregated index. For  
279 the third step (step 3, [10]), each indicator was normalized between 0 and 1 using the “more is  
280 better”, “less is better” or “optimum” response curves. For each parameter, the type of  
281 response curve was determined according to the literature [10], [49], [50]. For all parameters,  
282 we used the “more is better” response curves except for bulk density and Olsen P, for which  
283 the “optimum” response curve was chosen [51], [52]. Selected indicators from the MDS were  
284 then computed in a principal component analysis (PCA), and the contribution of each  
285 indicator to the dimensions of the PCA was calculated. Finally, the SQI was calculated  
286 according to Equation 1 and Equation 2 (step 4, [10]), with a weighting of the transformed  
287 variables using the PCA eigenvectors and the percentages of total variability explained by  
288 each principal component:

289

290 Equation 1:  $W_i = \sum_{j=1}^p \lambda_j \times f_j$

291

292 Equation 2:  $SQI = \sum_{i=1}^n S_i \times W_i$

293

294 where  $f_j$  = relative percentage of total variability attributed to each principal component,  $\lambda_j$  =  
295 sum of squared coordinates on each eigenvector,  $S_i$  = normalized indicator scores and  $W_i$  =

296 weighted factors. One SQI was calculated for each position within the alley cropping plot, and  
297 one for the entire alley cropping (hereafter called weighted alley cropping), weighting each  
298 surface area on the north side for the SQI. Another SQI was calculated for the alley cropping  
299 interrow excluding the tree row area in the weighted calculation (hereafter called weighted  
300 interrow). The last SQI was calculated for the monocropping system.

301

302

### 303 2.6. *Statistical analyses*

304 Mixed models with distance to the tree row as a fixed effect and transect number as a  
305 random effect were created for all soil parameters. Normal distribution of residuals and  
306 homogeneity of variance were tested by using the Shapiro and Bartlett tests. When necessary,  
307 data were Box-Cox transformed (MASS R package). If significant, a Tukey post hoc test was  
308 used for pairwise multiple comparisons ( $P < 0.05$ ). Differences between monocropping and  
309 each position from the spatial gradient were tested by a one sample t-test ( $P < 0.05$ )  
310 considering the monocropping system as a reference.

311 A Pearson correlation matrix was calculated to assess relationships between  
312 environmental parameters and the abundance of soil microbial functional groups. Redundancy  
313 analysis (RDA) was performed to assess the relationship between the soil microbial  
314 community composition, i.e., relative abundance of different microbial groups and  
315 environmental parameters. Most discriminating variables were selected using a forward  
316 procedure, and significance was also tested (100,000 permutations) (vegan R package). All  
317 statistical analyses were performed with R software v.3.2.3 (R development Core Team,  
318 2015).

319

## 320 **3. Results**

321 *3.1. Physical and chemical soil quality*

322 The soil was calcareous with more than 50% calcium carbonate, which explained the  
323 alkaline pH (**Table 1**). Levels of Olsen P were systematically low regardless of the position in  
324 the alley cropping or monocropping plots. The tree row showed significantly higher values  
325 than the interrow positions for the following variables: soil moisture, SOC, DOC, N<sub>tot</sub>,  
326 mineral N, Olsen P, CEC and exchangeable cations.

327 The largest C content was systematically found in the smallest fraction, i.e., < 50 µm  
328 fraction, contributing to more than 50% of the total soil C (**Table 2**). The highest C content  
329 was found in the tree rows regardless of the fraction, e.g., within the 50-2000 µm fraction, 3.6  
330 g C kg<sup>-1</sup> soil was measured in the tree rows versus 2.0 - 2.3 g C kg<sup>-1</sup> soil within the interrows  
331 and 1.5 g C kg<sup>-1</sup> soil in the monocropping plots. The C:N ratios of f-LF and o-LF were  
332 significantly lower under the tree row (18.7 and 16.5, respectively) than under the middle of  
333 the interrow (22.8 and 18.7, respectively) (**Table A.2**).

334 The highest sugar-C concentration was also found in the < 50 µm fraction (1.87 - 2.36 mg  
335 sugar C g<sup>-1</sup> soil) and did not significantly differ among positions in the alley cropping plot or  
336 compared with the monocropping plot (**Table 3**). The sugar-C concentration in f-LF under the  
337 tree row was significantly (two-fold) higher than that in the middle of the interrow or in the  
338 monocropping plot. In the same fraction, the proportion of microorganism-derived sugars in  
339 relation to plant-derived sugars (GM:AX ratio) decreased from 0.85 in the tree row down to  
340 0.67 in 4-6.5 m<sub>N</sub> and was significantly higher than that in the monocropping plot (0.62).

341 In the tree row, the total root biomass was more than 2-fold higher than anywhere  
342 within the interrow or compared with the monocropping plot (**Fig. A.2**). Significant  
343 differences were noticeable for fine- and medium-class roots.

344

345 *3.2. Biological soil quality indicators*

346 We observed a soil MBC close to 2-fold higher in the tree row (427 mg C kg<sup>-1</sup>) than in  
347 the middle of the interrow (219 mg C kg<sup>-1</sup> soil) and in the monocropping plot (199 mg C kg<sup>-1</sup>  
348 soil) (**Fig. 1**). The same difference was observed for MBN and MBP. Compared with  
349 monocropping, we systematically found a significant increase in MBC, MBN and MBP in the  
350 tree row and in 1-2 m<sub>N/S</sub>.

351 The MB-C:N ratio significantly increased with increasing distance from the tree row,  
352 from 6.0 in the tree row to 7.4 in 4-6.5 m<sub>N</sub> (**Table A.3**). The MB-C:P ratio was significantly  
353 lower under the tree row (24.5) than in 4-6.5 m<sub>N</sub> (41.4). Values of MB-C:N and values of  
354 MB-C:P were similar in the monocropping plot and close to those in the tree rows.

355 Basal respiration was significantly increased only in the 1-2 m<sub>S</sub> position compared  
356 with all others (**Table 4**). Compared with monocropping, glucose-induced respiration was  
357 significantly enhanced at all positions in the alley cropping plot except in 4-6.5 m<sub>N</sub>. Alanine-  
358 and trehalose-induced respiration was significantly higher in the tree row and close to it in the  
359 alley cropping plot than in the monocropping plot. Alanine-induced respiration was more than  
360 2-fold higher under the tree row than in the middle of the interrow. The metabolic quotient  
361 (qCO<sub>2</sub>) was similar across all positions ( $P > 0.05$ , **Table 4**).

362 The PLFA diversity, i.e., the number of detected peaks, decreased with increasing  
363 distance from the tree row (**Fig. A.3**). Actinobacteria abundance was significantly highest in  
364 the tree row (2.58 μg PLFAs g<sup>-1</sup> soil) and similar for the interrow positions and for the  
365 monocropping plot (**Fig. 2a**). For GP and GN bacteria, AMF and other fungi, soils under the  
366 tree row and in 1-2 m<sub>S</sub> showed significantly higher abundance than those in the middle of the  
367 interrow or in the monocropping plot (**Fig. 2a, b**). The F:B ratio did not show significant  
368 differences between positions within the alley cropping plot or compared with the  
369 monocropping plot: all values were approximately 0.30 ± 0.1 (data not shown). The GP:GN



370 ratio was significantly lower in and close to the tree row compared with the interrow or  
371 monocropping plot (**Fig. 2a**).

372

### 373 *3.3 Soil quality index*

374 Eight indicators, i.e., bulk density, SOC, N<sub>tot</sub>, CEC, Olsen P, basal respiration, and  
375 glucose- and alanine-induced respiration ultimately contributed to the construction of the SQI  
376 (**Fig. 3**).

377 We observed a gradual decrease in the SQI with increasing distance from the tree row (0.76 to  
378 0.53). The monocropping plot had an SQI similar (0.52) to that found in the middle of the  
379 interrow. The contribution of each individual indicator to SQI construction was approximately  
380 12%, except Olsen P, which contributed less than 3%. Individual contributions varied  
381 according to the position in the alley cropping plot following a same spatial pattern that was  
382 similar to the SQI. The weighted SQI of the entire alley cropping plot (0.64) was significantly  
383 20% higher than that of the monocropping plot (0.52), while an increase of 10% of the  
384 weighted SQI for the alley cropping interrow (0.57) was not significantly different from that  
385 of the monocropping system ( $P = 0.08$ ) (**Fig. 3**).

386

### 387 *3.4. Relationships between environmental parameters and microbial community structure*

388 To understand the link between soil environmental parameters and soil microbial  
389 community structure, we i) calculated a Pearson correlation matrix between different soil  
390 characteristics and diversity, and total PLFA biomass and abundance of each microbial group  
391 (**Table A.4**) and ii) performed RDA to identify the determinants of the relative abundance of  
392 PLFA markers (**Fig. 4**).

393 The two first axes explained 45 and 10% of the total variation, respectively (**Fig. 4**).

394 The first axis clearly distinguished the relative abundance of actinobacteria and other GP

395 bacteria, which were positively correlated with silt content, and the relative abundance of  
396 AMF, GN bacteria and fungi. The AMF and GN bacteria were positively correlated with  
397 sugar content and GM:AX ratio in the f-FL fraction, and fungi were positively correlated with  
398 sugar content in the o-LF fraction and negatively correlated with clay content.

399

400

401

## 402 **4. Discussion**

### 403 *4.1. Soil quality index in agroforestry and monocropping plots*

404 In the present work, the selected indicators for the minimum dataset were in line with  
405 expert opinion [2], [53], [54]. Bulk density and SOC content represent important weighted  
406 indicators in soil quality assessment for their role in water infiltration, storage and supply, as  
407 well as CEC, N and P contents for nutrient storage and supply [11]. Among the selected  
408 biological indicators, basal respiration is considered a physiological trait of microbial  
409 communities, and glucose-induced respiration is assumed to be proportional to active  
410 microbial biomass [44]. In this study, we focused on the microbial component of soil  
411 functioning, although we are aware that other biological indicators, such as those related to  
412 meso- or macro-fauna, could have been helpful for soil quality assessment [9], [55].

413 Our aim was to aggregate information obtained using different parameters into one  
414 integrative soil quality index based on several soil functions: water, C and nutrient storage  
415 and cycling and sustainability of biological activities. In the present work, we initially had a  
416 large number of indicators (30 in total), and several methods for SQI calculation offered to us.  
417 Comparing three different SQI constructions, i.e., additive SQI, weighted additive SQI and  
418 statistically modelled SQI, [56] showed that the last one was more efficient in time and cost.

419 The method for index calculation proposed by [10] allowed us to objectively select the most  
420 relevant indicators to assess changes in physical, chemical and biological soil quality in the  
421 studied field plots. Without this statistical approach, we could have overestimated the positive  
422 effect of tree rows on soil quality because of the co-variability of the measured indicators.  
423 However, this method considers as relevant for the first step of minimum dataset selection  
424 only parameters that significantly changed statistically according to the treatments, which  
425 could be considered a subjective choice.

426 This is the first time, to the our knowledge, that soil quality was assessed in temperate  
427 alley cropping through the use of an integrative and weighted SQI. We showed a strong  
428 positive effect from the 21-year-old trees and accompanying permanent grass cover in the tree  
429 rows of the studied alley cropping system on soil physicochemical indicators compared with  
430 the interrow and monocropping systems; this effect was restricted to the tree rows (**Table 1**).  
431 The increase in SOC content in the tree rows led to an increase in those soil properties that are  
432 usually related to it, such as CEC and exchangeable cation contents [10] (**Table 1**).

433 We showed that the SQI was significantly improved in an alley cropping system in  
434 comparison with a monocropping system, considering the spatial heterogeneity induced by  
435 these systems. This indicates here that the positive effect from the trees and herbaceous cover  
436 on the weighted entire alley cropping SQI was significant beyond the tree rows. The tree  
437 rows, which represent approximately 13% of the field plot surface area, yielded a significant  
438 beneficial effect on SQI over approximately 20% compared with monocropping (**Fig. 3**).  
439 These findings should be put in perspective, as the results depend on one site and one  
440 sampling date. Additional sub-annual sampling would be necessary to reject a possible  
441 seasonal effect. Even though not significant ( $P = 0.08$ ), there was a 10% increase in the SQI  
442 weighted interrow (i.e., 1-2 m + 2-4 m + 4-6.5 m) compared with the neighbouring

443 monocropping plot. Repeating these indicator measurements in time and increasing the  
444 number of replicates would allow us to see if this trend could be significant or not.

445

#### 446 *4.2. What explains the higher SOC content in the alley cropping system?*

447 In our study, the topsoil of tree rows showed SOC contents 20-30% higher than that of  
448 the interrow and 50% greater than that of the monocropping plot (**Table 2**), which is in line  
449 with former results at the same study site [57]. Within and close to tree rows, belowground  
450 litter, as aboveground litter can also be enhanced, as tree roots have been shown to colonize  
451 the topsoil at the same agroforestry site until 1.5 m in the interrow [58]. In addition, a  
452 substantial portion of roots can originate from (i) herbaceous cover root systems, which can  
453 colonize the topsoil close to the tree row, as shown by [59] in younger walnut tree alley  
454 cropping, or ii) weed colonization with low dispersal abilities, originating from herbaceous  
455 cover [60]. Perennial plants can indeed allocate up to 10-15% more C belowground than  
456 crops [61], [62]. The absence of tillage in the tree rows likely favoured some accumulation of  
457 both C and N [63].

458 The novelty of our study is that it provides greater insight into the pools of C found in  
459 the topsoil at various locations in the alley cropping system than in the monocropping system.  
460 We demonstrate that the additional C found in the topsoil comes from fractions with a higher  
461 C:N ratio (> 16) (**Table A.2**), which is more similar to that of plant debris than that of smaller  
462 size fractions with a C:N ratio < 10, i.e., close to the C:N ratio of soil microbial communities  
463 and soil organic matter [64], [65]. These coarser fractions can be more easily mineralized by  
464 microorganisms, suggesting a less stable C pool.

465 The increased GM:AX ratio in the f-LF fraction from the middle of the interrow to the  
466 tree row suggests that a higher proportion of microbial-derived carbohydrates was present in  
467 the tree row (**Table 3**). This may have induced higher and faster microbial turnover based on

468 more organic matter recycling through microorganisms and could indicate that the production  
469 of organic compounds is easier to stabilize [66], [67]. The RDA confirmed that the increased  
470 GM:AX ratio favours more fast-growing microorganisms, i.e., microbial communities with  
471 higher GN bacteria proportions (**Fig. 4**).

472

#### 473 *4.3. Spatial heterogeneity of microbial soil properties in the alley cropping system*

474 Our work demonstrated that an agroforestry system such as the one we studied here  
475 may considerably impact soil microbial communities through its biomass, activity and  
476 structure, resulting in the creation of within-plot spatial heterogeneity.

477 Higher microbial biomass and activities in and close to the tree row may represent a  
478 higher potential capacity to decompose organic matter and improve nutrient cycling.  
479 MicroResp® appeared not to be the most appropriate method to evaluate the activity of  
480 microorganisms in the carbonate-rich soil; however, as carbonate content was similar  
481 everywhere, it allowed us to compare positions. Our qCO<sub>2</sub> results indicate a similar  
482 physiological activity of the microbial biomass regardless of the position within the  
483 agroforestry plot [68]. This result was surprising because of the different tillage management  
484 practices between the tree row and interrow [69]. However, [70] showed in a meta-analysis  
485 that similar qCO<sub>2</sub> values were observed for tilled and no-tilled plots in long-term (10 years)  
486 plots. This might indicate that qCO<sub>2</sub> is not sensitive to soil tillage or that microorganisms from  
487 the tree row are, at the time of our sampling, as efficient in using C as those in the interrow or  
488 in the monocropping system.

489 In tree rows, the MBC/SOC ratio was significantly higher than that in the interrow or  
490 in the monocropping plot (i.e., 23 versus 17, data not shown). This indicates that the MBC  
491 increase in tree rows was not merely due to the increase in SOC but also to some properties of

492 the soil organic matter (e.g., greater C availability) within tree rows that favour the  
493 development of an abundant soil microbial biomass [71].

494 Compared with the monocropping system, the microbial biomass and substrate-  
495 induced respiration were more significantly higher on the south side of the tree row than on  
496 the north side in the alley cropping plot (**Fig. 1, Table 4**). This result underlines the potential  
497 legacy effect of microclimate created in this alley cropping agroforestry system. Dufour et al.  
498 [72] showed in the same plot that photosynthetically active radiation can be reduced by 38%  
499 at 3 m on the north side when the tree canopy is well developed (end of June). Our soil  
500 sampling was performed in April before walnut tree budburst began. At this stage, they  
501 showed that the reduction in incident light was negligible (10%) compared with that in June  
502 (50%). This suggests that the slight difference observed between the north and south sides of  
503 the tree rows was due to a legacy effect of the 21-year-old practice.

504 We found that trees and the herbaceous grass community (dominated by weeds,  
505 *Bromus* and *Torilis* species) in the tree rows increased PLFA diversity and biomass. These  
506 two parameters were highly correlated with pH (negatively), soil moisture, DOC and total N,  
507 as observed by [48] and [22]. In our case, an additional correlation with root biomass  
508 highlighted its potential key role in shaping microbial community dynamics (**Table A.4**).

509 In the present study, we observed that tree rows favour a higher proportion of GN  
510 bacteria than GP bacteria, leading to a lower GP:GN ratio. Environments rich in C and  
511 available N are known to promote soil GN bacteria [73], which have high affinity for recent  
512 plant-derived C compounds [74]. The RDA also suggested a correlation between the sugar  
513 content in f-LF and the GN bacterial proportion (**Fig. 4**). In contrast, GP bacteria have been  
514 shown to be less affected by low C availability [73]. Actinobacteria and other GP bacteria are  
515 known to be able to feed on more complex polymers, such as older organic matter [48]. This  
516 could explain their higher proportion in the middle of the interrow and in the monocropping

517 system. Variation in the GP:GN ratio suggests that microorganisms could be more  
518 copiotrophic in the tree row than in the interrow of the alley cropping system as the GP:GN  
519 ratio increases with decreasing C availability in soil [75] (**Fig. 2a**).

520 Moreover, high-recalcitrant litter inputs, such as walnut leaves, i.e., litter known to be  
521 rich in tannins and lignin [76], could have favoured the fungal pathway within or close to the  
522 tree row, while we did not observe this. We hypothesize that the herbaceous cover under the  
523 tree row adding labile litter and rhizodeposits buffers the recalcitrant effect of walnut litter.  
524 These results raise the question of the importance of considering herbaceous cover in future  
525 alley cropping studies.

526

## 527 **5. Conclusion**

528 Through the design of our sampling strategy, we evaluated changes in soil quality  
529 according to the distance from the tree rows in the interrow of an alley cropping system and  
530 compared them with a neighbouring monocropping system. The use of an integrative soil  
531 quality index related to water and nutrient storage and to the support of biological activity  
532 simplified the comparison of the various positions within the alley cropping plot and with the  
533 monocropping plot. After 21 years, soil quality was significantly improved in the tree rows  
534 compared with the interrow positions and monocropping. The tree rows and its permanent  
535 herbaceous cover enriched the soil in organic matter through an increased input of above- and  
536 belowground litters and possibly through enhanced rhizodeposition. The SQI was  
537 significantly greater under tree rows and beyond until a 2-m distance from the rows, which  
538 was clear evidence of spatial heterogeneity within the cultivated interrows. For the first time,  
539 we showed that by considering the SQI of the weighted entire plot of the alley cropping  
540 system that soil quality was significantly improved compared with that of the monocropping  
541 system.

542

543 **Acknowledgments**

544

545 This work was supported by “La Fondation de France” and the French National  
546 Program EC2CO Biohefect, France. The authors want to thank Annette Bérard (INRAE) at  
547 UMR EMMAH in Avignon and Josiane Abadie (INRAE) at UMR Eco&Sols in Montpellier  
548 for their help during MicroResp™ analyses, Delphine Mézière and Lydie Dufour (INRAE) at  
549 UMR SYSTEM in Montpellier for their knowledge of the study site and their advice on the  
550 sampling location and Paula Fernandes (CIRAD) at UR HORTSYS for her help with the  
551 PLFA.

552

553

554 **References**

555

556 [1] J.W. Doran, M. Safley, 1997. Defining and assessing soil health and sustainable  
557 productivity. Biological indicators of soil health. New York: CAB International.

558

559 [2] E.K. Bünemann, G. Bongiorno, Z. Bai, R.E. Creamer, G. De Deyn, R. Goede, L. Fleskens,  
560 V. Geissen, T.W. Kuyper, P. Mäder, M. Pulleman, W. Sukkel, J.W. van Groenigen,  
561 L. Brussaard, 2018. Soil quality – A critical review. Soil Biology and Biochemistry  
562 120:105–125. <https://doi.org/10.1016/j.soilbio.2018.01.030>.

563

564 [3] M. Rutgers, H.J. van Wijnen, A.J. Schouten, C. Mulder, A.M.P. Kuiten, L. Brussaard,  
565 A.M. Breure, 2012. A method to assess ecosystem services developed from soil  
566 attributes with stakeholders and data of four arable farms. Science of The Total  
567 Environment 415:39–48. <https://doi.org/10.1016/j.scitotenv.2011.04.041>.



568

569 [4] J. Paz-Ferreiro, S. Fu, 2016. Biological indices for soil quality evaluation: perspectives  
570 and limitations. *Land Degradation & Development*, 27(1), 14-25.  
571 <https://doi.org/10.1002/ldr.2262>

572

573 [5] S.S. Andrews, D.L. Karlen, C.A. Cambardella, 2004. The Soil Management Assessment  
574 Framework. *Soil Science Society of America Journal* 68:1945–1962.  
575 <https://doi.org/10.2136/sssaj2004.1945>.

576

577 [6] J.E. Herrick, J.R. Brown, A.J. Tugel, P.L. Shave, K.M. Havstad, 2002. Application of soil  
578 quality to monitoring and management: paradigms from rangeland ecology. *Agron*  
579 *J.*, 94, pp. 3-11. <https://doi.org/10.2134/agronj2002.3000>

580

581 [7] E. Dominati, M. Patterson, A. Mackay, 2010. A framework for classifying and quantifying  
582 the natural capital and ecosystem services of soils. *Ecological Economics* 69, 1858–  
583 1868. <https://doi.org/10.1016/j.ecolecon.2010.05.002>.

584

585 [8] L. Gianfreda, P. Ruggiero, 2006. Enzyme activities in soil. In *Soil biology*, volume 8.  
586 nucleic acids and proteins in soil, Nannipieri P, Smalla K (eds). Springer-Verlag:  
587 Berlin Heidelberg.

588 [9] P. Coll, E. Le Cadre, E. Blanchart, P. Hinsinger, C. Villenave, 2011. Organic viticulture  
589 and soil quality: A long-term study in Southern France. *Applied Soil Ecology*.  
590 <https://doi.org/10.1016/j.apsoil.2011.07.013>.

591

- 592 [10] F. Obriot, M. Stauffer, Y. Goubard, N. Cheviron, G. Peres, M. Eden, A. Revallier, L.  
593 Vieublé-Gonod, S. Houot, 2016. Multi-criteria indices to evaluate the effects of  
594 repeated organic amendment applications on soil and crop quality. *Agriculture,*  
595 *Ecosystems & Environment* 232:165–178.  
596 <https://doi.org/10.1016/j.agee.2016.08.004>.  
597
- 598 [11] A.C.R. Lima, L. Brussaard, M.R Totola, W.B. Hoogmoed, R.G.M. De Goede, 2013. A  
599 functional evaluation of three indicator sets for assessing soil quality. *Applied Soil*  
600 *Ecology*, 64, 194-200. <https://doi.org/10.1016/j.apsoil.2012.12.009>  
601
- 602 [12] J. Six, S.D. Frey, R.K. Thiet, K.M. Batten, 2006. Bacterial and fungal contributions to  
603 carbon sequestration in agroecosystems. *Soil Science Society of America*  
604 *Journal*, 70(2), 555-569. <https://doi.org/10.2136/sssaj2004.0347>  
605
- 606 [13] R.G. Joergensen, C. Emmerling, 2006. Methods for evaluating human impact on soil  
607 microorganisms based on their activity, biomass, and diversity in agricultural soils. *J*  
608 *Plant Nutr Soil Sci* 169:295–309. <https://doi.org/10.1002/jpln.200521941>  
609
- 610 [14] J. Bloem, D.W. Hopkins, A. Benedetti, Eds., 2005. *Microbiological methods for*  
611 *assessing soil quality*. Cabi).  
612
- 613 [15] J. Laishram, K.G. Saxena, R.K. Maikhuri, K.S. Rao, 2012. Soil quality and soil health: A  
614 review. *Int. J. Ecol. Environ. Sci.* 38, 19–37.  
615

- 616 [16] M. Shukla, R. Lal, M. Ebinger, 2006. Determining soil quality indicators by factor  
617 analysis. *Soil Tillage Res.* 87, 194–204. <https://doi.org/10.1016/j.still.2005.03.011>.  
618
- 619 [17] A. Thoumazeau, C. Bessou, M-S. Renevier, J. Trap, R. Marichal, L. Mareschal, T.  
620 Decaëns, N. Bottinelli, B. Jaillard, T. Chevallier, N. Suvannang, K. Sajjaphan, P.  
621 Thaler, F. Gay, A. Brauman, 2019. Biofunctool®: a new framework to assess the  
622 impact of land management on soil quality. Part A: concept and validation of the set  
623 of indicators. *Ecol. Indic.* 97, 100–110.  
624 <https://doi.org/10.1016/j.ecolind.2018.09.023>.  
625
- 626 [18] R. Lal, 2015. Restoring Soil Quality to Mitigate Soil Degradation. *Sustainability* 7:5875–  
627 5895. <https://doi.org/10.3390/su7055875>.  
628
- 629 [19] M. Torralba, N. Fagerholm, P.J. Burgess, G. Moreno, T. Plieninger, 2016. Do European  
630 agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis.  
631 *Agriculture, Ecosystems & Environment* 230:150–161.  
632 <https://doi.org/10.1016/j.agee.2016.06.002>.  
633
- 634 [20] K.R. Kirby, C. Potvin, 2007. Variation in carbon storage among tree species:  
635 Implications for the management of a small-scale carbon sink project. *Forest*  
636 *Ecology and Management* 246:208–221.  
637 <https://doi.org/10.1016/j.foreco.2007.03.072>.  
638
- 639 [21] F. Abbas, H.M. Hammad, S. Fahad, A. Cerdà, M. Rizwan, W. Farhad, S. Ehsan, H.F.  
640 Bakhat, 2017. Agroforestry: a sustainable environmental practice for carbon

641 sequestration under the climate change scenarios—a review. *Environ Sci Pollut Res*  
642 24:11177–11191. <https://doi.org/10.1007/s11356-017-8687-0>.

643

644 [22] S. Banerjee, M. Baah-Acheamfour, C.N. Carlyle, A. Bissett, A.E. Richardson, T.  
645 Siddique, E.W. Bork, S.X. Chang, 2016. Determinants of bacterial communities in  
646 Canadian agroforestry systems. *Environ Microbiol* 18:1805–1816.  
647 <https://doi.org/10.1111/1462-2920.12986>.

648

649 [23] J. Dollinger, S. Jose, 2018. Agroforestry for soil health. *Agroforest Syst* 1–7.  
650 <https://doi.org/10.1007/s10457-018-0223-9>.

651

652 [24] R. Cardinael, Z. Mao, C. Chenu, P. Hinsinger, 2020. Belowground functioning of  
653 agroforestry systems: recent advances and perspectives. *Plant Soil*.  
654 <https://doi.org/10.1007/s11104-020-04633-x>

655

656 [25] S. Nii-Annang, H. Grünewald, D. Freese, R.F. Hüttl, O. Dilly, 2009. Microbial activity,  
657 organic C accumulation and <sup>13</sup>C abundance in soils under alley cropping systems  
658 after 9 years of recultivation of quaternary deposits. *Biol Fertil Soils* 45:531–538.  
659 <https://doi.org/10.1007/s00374-009-0360-4>.

660

661 [26] S. Bardhan, S. Jose, R.P. Udawatta, F. Fritschi, 2013. Microbial community diversity in a  
662 21-year-old temperate alley cropping system. *Agrorest Syst* 87:1031–1041.  
663 <https://doi.org/10.1007/s10457-013-9617-x>.

664

- 665 [27] R. Beuschel, H-P. Piepho, R.G. Joergensen, C. Wachendorf, 2019. Similar spatial  
666 patterns of soil quality indicators in three poplar-based silvo-arable alley cropping  
667 systems in Germany. *Biol Fertil Soils* 55:1–14. [https://doi.org/10.1007/s00374-018-](https://doi.org/10.1007/s00374-018-1324-3)  
668 1324-3.
- 669
- 670 [28] N.W. Mungai, P.P. Motavalli, R.J. Kremer, K.A. Nelson, 2005. Spatial variation of soil  
671 enzyme activities and microbial functional diversity in temperate alley cropping  
672 systems. *Biology and Fertility of Soils* 42:129–136. [https://doi.org/10.1007/s00374-](https://doi.org/10.1007/s00374-005-0005-1)  
673 005-0005-1.
- 674
- 675 [29] L. Dufour, M. Gosme, J. Le Bec, C. Dupraz, 2020. Does pollarding trees improve the  
676 crop yield in a mature alley-cropping agroforestry system? *Journal of Agronomy and*  
677 *Crop Science*, 206(5), 640-649. <https://doi.org/10.1111/jac.12403>
- 678
- 679 [30] H. Ciesielski, T. Sterckeman, 1997. A comparison between three methods for the  
680 determination of cation exchange capacity and exchangeable cations in soils.  
681 *agronomie*, 17, 9-15.
- 682
- 683 [31] S.R. Olsen, C.V. Cole, F.S. Watanabe, L.A. Dean, 1954. Estimation of available  
684 phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939. U.S.  
685 Government Printing Office, Washington D.C.
- 686
- 687 [32] R. Roscoe, P. Buurman, E.J. Velthorst, C.A. Vasconcellos, 2001. Soil organic matter  
688 dynamics in density and particle size fractions as revealed by the <sup>13</sup>C/<sup>12</sup>C isotopic

689 ratio in a Cerrado's oxisol. *Geoderma* 104:185–202. [https://doi.org/10.1016/S0016-](https://doi.org/10.1016/S0016-7061(01)00080-5)  
690 7061(01)00080-5.

691

692 [33] D. Harris, W.R. Horwath, C. van Kessel, 2001. Acid fumigation of soils to remove  
693 carbonates prior to total organic carbon or CARBON-13 isotopic analysis. *Soil*  
694 *Science Society of America Journal* 65:1853–1856.  
695 <https://doi.org/10.2136/sssaj2001.1853>.

696

697 [34] C. Rumpel, M-F. Dignac, 2006. Gas chromatographic analysis of monosaccharides in a  
698 forest soil profile: Analysis by gas chromatography after trifluoroacetic acid  
699 hydrolysis and reduction–acetylation. *Soil Biology and Biochemistry* 38:1478–1481.  
700 <https://doi.org/10.1016/j.soilbio.2005.09.017>.

701

702 [35] E. Eder, S. Spielvogel, A. Kölbl, G. Albert, I. Kögel-Knabner, 2010. Analysis of  
703 hydrolysable neutral sugars in mineral soils: Improvement of alditol acetylation for  
704 gas chromatographic separation and measurement. *Organic Geochemistry* 41:580–  
705 585. <https://doi.org/10.1016/j.orggeochem.2010.02.009>.

706

707 [36] G. Guggenberger, B.T. Christensen, W. Zech, 1994. Land-use effects on the composition  
708 of organic matter in particle-size separates of soil: I. Lignin and carbohydrate  
709 signature. *European Journal of Soil Science* 45:449–458.  
710 <https://doi.org/10.1111/j.1365-2389.1994.tb00530.x>.

711

- 712 [37] J.M. Oades, 1984. Soil organic matter and structural stability: mechanisms and  
713 implications for management. *Plant Soil* 76:319–337.  
714 <https://doi.org/10.1007/BF02205590>.
- 715
- 716 [38] E.D. Vance, P.C. Brookes, D.S. Jenkinson, 1987. An extraction method for measuring  
717 soil microbial biomass C. *Soil Biology and Biochemistry* 19:703–707.  
718 [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6).
- 719
- 720 [39] [37] T. Ohno, L.M. Zibilske, 1991. Determination of Low Concentrations of Phosphorus  
721 in Soil Extracts Using Malachite Green. *Soil Science Society of America Journal*  
722 55:892–895. <https://doi.org/10.2136/sssaj1991.03615995005500030046x>.
- 723
- 724 [40] D. Jenkinson, 2004. Measuring soil microbial biomass. *Soil Biology and Biochemistry*  
725 36:5–7. <https://doi.org/10.1016/j.soilbio.2003.10.002>.
- 726
- 727 [41] P.C. Brookes, A. Landman, G. Pruden, D.S. Jenkinson, 1985. Chloroform fumigation  
728 and the release of soil nitrogen: a rapid direct extraction method to measure  
729 microbial biomass nitrogen in soil. *Soil biology and biochemistry*, 17(6), 837-842.  
730 [https://doi.org/10.1016/0038-0717\(85\)90144-0](https://doi.org/10.1016/0038-0717(85)90144-0)
- 731
- 732 [42] C.D. Campbell, S.J. Chapman, C.M. Cameron, M.S. Davidson, J.M. Potts, 2003. A  
733 Rapid Microtiter Plate Method To Measure Carbon Dioxide Evolved from Carbon  
734 Substrate Amendments so as To Determine the Physiological Profiles of Soil  
735 Microbial Communities by Using Whole Soil. *Appl Environ Microbiol* 69:3593–  
736 3599. <https://doi.org/10.1128/AEM.69.6.3593-3599.2003>.

737

738 [43] L.A. Richards, Pressure-membrane apparatus — Construction and use. *Agric. Eng.*,28,  
739 1947, pp. 451-454.

740

741 [44] A. Bérard, M.B. Sassi, P. Renault, R. Gros, 2012. Severe drought-induced community  
742 tolerance to heat wave. An experimental study on soil microbial processes. *J Soils*  
743 *Sediments* 12:513–518. <https://doi.org/10.1007/s11368-012-0469-1>.

744

745 [45] J.S. Buyer, M. Sasser, 2012. High throughput phospholipid fatty acid analysis of soils.  
746 *Applied Soil Ecology* 61:127–130. <https://doi.org/10.1016/j.apsoil.2012.06.005>.

747

748 [46] N. Fanin, I. Bertrand, 2016. Aboveground litter quality is a better predictor than  
749 belowground microbial communities when estimating carbon mineralization along a  
750 land-use gradient. *Soil Biology and Biochemistry* 94:48–60.  
751 <https://doi.org/10.1016/j.soilbio.2015.11.007>.

752

753 [47] Y. Feng, A.C. Motta, D.W. Reeves, C.H. Burmester, E. Van Santen, J.A. Osborne, 2003.  
754 Soil microbial communities under conventional-till and no-till continuous cotton  
755 systems. *Soil Biology and Biochemistry* 35:1693–1703.  
756 <https://doi.org/10.1016/j.soilbio.2003.08.016>.

757

758 [48] Q. Zhang, J. Wu, F. Yang, Y. Lei, Q. Zhang, X. Cheng, 2016. Alterations in soil  
759 microbial community composition and biomass following agricultural land use  
760 change. *Scientific Reports* 6:36587. <https://doi.org/10.1038/srep36587>.

761



- 762 [49] M.S. Askari, N.M. Holden, 2015. Quantitative soil quality indexing of temperate arable  
763 management systems. *Soil Tillage Res.*, 150 (2015), pp. 57-67.  
764 <https://doi.org/10.1016/j.still.2015.01.010>  
765
- 766 [50] J.D. Glover, J.P. Reganold, P.K. Andrews, 2000. Systematic method for rating soil  
767 quality of conventional, organic, and integrated apple orchards in Washington State.  
768 *Agric. Ecosyst. Environ.*, 80 (2000), pp. 29-45. [https://doi.org/10.1016/S0167-](https://doi.org/10.1016/S0167-8809(00)00131-6)  
769 [8809\(00\)00131-6](https://doi.org/10.1016/S0167-8809(00)00131-6)  
770
- 771 [51] M.P. Lefebvre, 2010. Spatialisation de modèles de fonctionnement hydromécanique des  
772 sols appliquée à la prévision des risques de tassement à l'échelle de la France (PhD  
773 thesis). Université d'Orléans.  
774
- 775 [52] UNIFA, [http://www.unifa.fr/raisonner-la-fertilisation/calculer-les-apports/fertilisation-p-](http://www.unifa.fr/raisonner-la-fertilisation/calculer-les-apports/fertilisation-p-a-k.html)  
776 [a-k.html](http://www.unifa.fr/raisonner-la-fertilisation/calculer-les-apports/fertilisation-p-a-k.html), 2016  
777
- 778 [53] D.L. Karlen, N.C. Wollenhaupt, D.C. Erbach, E.C. Berry, J.B. Swan, N.S. Eash, J.L.  
779 Jordhal, 1994. Crop residue effects on soil quality following 10-years of no-till corn.  
780 *Soil Tillage Res.*, 31, pp. 149-167. [https://doi.org/10.1016/0167-1987\(94\)90077-9](https://doi.org/10.1016/0167-1987(94)90077-9)  
781
- 782 [54] C.H. Lee, M.Y. Wu, V.B. Asio, C.U. Zueng-Sang, 2006. Using a soil quality index to  
783 assess the effects of applying swine manure compost on soil quality under a crop  
784 rotation system in Taiwan. *Soil Sci.*, 171, pp. 210-222.  
785 <https://doi.org/10.1097/01.ss.0000199700.78956.8c>  
786

- 787 [55] S. Yan, A.N. Singh, S. Fu, C. Liao, S. Wang, Y. Li, L. Hu, 2012. A soil fauna index for  
788 assessing soil quality. *Soil Biology and Biochemistry*, 47, 158-165.  
789 <https://doi.org/10.1016/j.soilbio.2011.11.014>  
790
- 791 [56] A. Mukherjee, R. Lal, 2014. Comparison of soil quality index using three methods. *PloS*  
792 *one*, 9(8), e105981. <https://doi.org/10.1371/journal.pone.0105981>
- 793 [57] R. Cardinael, T. Chevallier, B.G. Barthès, N.P.A. Saby, T. Parent, C. Dupraz, M.  
794 Bernoux, C. Chenu, 2015a. Impact of alley cropping agroforestry on stocks, forms  
795 and spatial distribution of soil organic carbon — A case study in a Mediterranean  
796 context. *Geoderma* 259–260:288–299.  
797 <https://doi.org/10.1016/j.geoderma.2015.06.015>.  
798
- 799 [58] R. Cardinael, Z. Mao, I. Prieto, A. Stokes, C. Dupraz, J.H. Kim, C. Jourdan, 2015b.  
800 Competition with winter crops induces deeper rooting of walnut trees in a  
801 Mediterranean alley cropping agroforestry system. *Plant Soil* 391:219–235.  
802 <https://doi.org/10.1007/s11104-015-2422-8>.  
803
- 804 [59] P. Battie-Laclau, E. Taschen, C. Plassard, D. Dezette, J. Abadie, D. Arnal, P. Bezenech,  
805 M. Duthoit, A.L. Pablo, C. Jourdan, J.P. Laclau, I. Bertrand, A. Taudière, P.  
806 Hinsinger, 2019. Role of trees and herbaceous vegetation beneath trees in  
807 maintaining arbuscular mycorrhizal communities in temperate alley cropping  
808 systems. *Plant Soil*. <https://doi.org/10.1007/s11104-019-04181-z>.  
809

- 810 [60] S. Boinot, G. Fried, J. Storkey, H. Metcalfe, K. Barkaoui, P.E. Lauri, D. Mézière, 2019.  
811 Alley cropping agroforestry systems: Reservoirs for weeds or refugia for plant  
812 diversity? *Agriculture, Ecosystems & Environment*, 284, 106584.  
813 <https://doi.org/10.1016/j.agee.2019.106584>.
- 814
- 815 [61] J. Pausch, Y. Kuzyakov, 2018. Carbon input by roots into the soil: Quantification of  
816 rhizodeposition from root to ecosystem scale. *Glob Change Biol* 24:1–12.  
817 <https://doi.org/10.1111/gcb.13850>.
- 818
- 819 [62] S.J. Grayston, D. Vaughan, D. Jones, 1997. Rhizosphere carbon flow in trees, in  
820 comparison with annual plants: the importance of root exudation and its impact on  
821 microbial activity and nutrient availability. *Applied Soil Ecology* 5:29–56.  
822 [https://doi.org/10.1016/S0929-1393\(96\)00126-6](https://doi.org/10.1016/S0929-1393(96)00126-6).
- 823
- 824 [63] W.R. Cookson, D.V. Murphy, M.M. Roper, 2008. Characterizing the relationships  
825 between soil organic matter components and microbial function and composition  
826 along a tillage disturbance gradient. *Soil Biology and Biochemistry* 40:763–777.  
827 <https://doi.org/10.1016/j.soilbio.2007.10.011>.
- 828
- 829 [64] E. Guillot, P. Hinsinger, L. Dufour, J. Roy, I. Bertrand, 2019. With or without trees:  
830 Resistance and resilience of soil microbial communities to drought and heat stress in  
831 a Mediterranean agroforestry system. *Soil Biology and Biochemistry* 129:122–135.  
832 <https://doi.org/10.1016/j.soilbio.2018.11.011>.
- 833

- 834 [65] I. Bertrand, M. Sauvadet, E. Guillot, C. d’Hervilly, C. Plassard, E. Taschen, C. Marsden,  
835 M. Hedde, 2019. Relations entre modes de gestion des agroécosystèmes et  
836 biodiversité fonctionnelle des sols. *Innov. Agron.* 75, 107–124.  
837 <https://doi.org/10.15454/hl4c8y>.
- 838
- 839 [66] M.F. Cotrufo, M.D. Wallenstein, C.M. Boot, K. Deneff, E. Paul, 2013. The Microbial  
840 Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter  
841 decomposition with soil organic matter stabilization: do labile plant inputs form  
842 stable soil organic matter? *Global Change Biology* 19:988–995.  
843 <https://doi.org/10.1111/gcb.12113>.
- 844
- 845 [67] M.F. Cotrufo, J.L. Soong, A.J. Horton, E.E. Campbell, M.L. Haddix, D.H. Wall, W.J.  
846 Parton, 2015. Formation of soil organic matter via biochemical and physical  
847 pathways of litter mass loss. *Nature Geoscience* 8:776–779.  
848 <https://doi.org/10.1038/ngeo2520>.
- 849
- 850 [68] T.H. Anderson, K.H. Domsch, 1985. Determination of ecophysiological maintenance  
851 carbon requirements of soil microorganisms in a dormant state. *Biology and fertility*  
852 *of soils*, 1(2), 81-89.
- 853
- 854 [69] E.L. Balota, A. Colozzi Filho, D.S. Andrade, R.P. Dick, 2004. Long-term tillage and  
855 crop rotation effects on microbial biomass and C and N mineralization in a Brazilian  
856 Oxisol. *Soil Tillage Res.*, 77 (2004), pp. 137-145.  
857 <https://doi.org/10.1016/j.still.2003.12.003>
- 858

- 859 [70] S.M. Zuber, M.B. Villamil, 2016. Meta-analysis approach to assess effect of tillage on  
860 microbial biomass and enzyme activities. *Soil Biology and Biochemistry*, 97, 176-  
861 187. <https://doi.org/10.1016/j.soilbio.2016.03.011>  
862
- 863 [71] T.H. Anderson, K.H. Domsch, 1989. Ratios of microbial biomass carbon to total organic  
864 carbon in arable soils. *Soil biology and biochemistry*, 21(4), 471-479.  
865 [https://doi.org/10.1016/0038-0717\(89\)90117-X](https://doi.org/10.1016/0038-0717(89)90117-X)  
866
- 867 [72] L. Dufour, A. Metay, G. Talbot, C. Dupraz, 2013. Assessing Light Competition for  
868 Cereal Production in Temperate Agroforestry Systems using Experimentation and  
869 Crop Modelling. *Journal of Agronomy and Crop Science* 199:217–227.  
870 <https://doi.org/10.1111/jac.12008>.  
871
- 872 [73] N. Fierer, J.P. Schimel, P.A. Holden, 2003. Variations in microbial community  
873 composition through two soil depth profiles. *Soil Biology and Biochemistry* 35:167–  
874 176. [https://doi.org/10.1016/S0038-0717\(02\)00251-1](https://doi.org/10.1016/S0038-0717(02)00251-1).  
875
- 876 [74] C. Kramer, G. Gleixner, 2008. Soil organic matter in soil depth profiles: Distinct carbon  
877 preferences of microbial groups during carbon transformation. *Soil Biol. Biochem.*  
878 40, 425–433. <https://doi.org/10.1016/j.soilbio.2007.09.016>.  
879
- 880 [75] N. Fanin, P. Kardol, M. Farrell, M.C. Nilsson, M.J. Gundale, D.A. Wardle, 2019. The  
881 ratio of Gram-positive to Gram-negative bacterial PLFA markers as an indicator of  
882 carbon availability in organic soils. *Soil Biology and Biochemistry*, 128, 111-114.  
883 <https://doi.org/10.1016/j.soilbio.2018.10.010>

884

885 [76] V. Nour, I. Trandafir, S. Cosmulescu, 2013. HPLC Determination of Phenolic Acids,

886 Flavonoids and Juglone in Walnut Leaves. *J Chromatogr Sci* 51:883–890.

887 <https://doi.org/10.1093/chromsci/bms180>.

888

889


890

891


892

893

**Figure 1**




**Figure 2a**






**Figure 2b**





**Figure 4**



**Table 1**

Parameter	Tree row	1-2 m <sub>N</sub>	2-4 m <sub>N</sub>	4-6.5 m <sub>N</sub>	MC
Clay (< 2 μm) (g kg <sup>-1</sup> soil)	178 ± 9 <sup>a</sup>	173 ± 24 <sup>a</sup>	186 ± 7 <sup>a</sup>	184 ± 13 <sup>a</sup>	177 ± 10
Silt (2-50 μm) (g kg <sup>-1</sup> soil)	157 ± 24 <sup>a*</sup>	165 ± 43 <sup>a*</sup>	157 ± 23 <sup>a*</sup>	158 ± 17 <sup>a*</sup>	208 ± 16
Sand (50-2000 μm) (g kg <sup>-1</sup> soil)	109 ± 18 <sup>a*</sup>	101 ± 22 <sup>a</sup>	105 ± 25 <sup>a</sup>	103 ± 22 <sup>a</sup>	76 ± 20
Bulk density (g cm <sup>-3</sup> )	1.24 ± 0.1 <sup>a*</sup>	1.41 ± 0.1 <sup>a</sup>	1.23 ± 0.1 <sup>a*</sup>	1.30 ± 0.1 <sup>a</sup>	1.43 ± 0.1
CaCO <sub>3</sub> (g kg <sup>-1</sup> soil)	539 ± 11 <sup>a</sup>	545 ± 22 <sup>a</sup>	540 ± 26 <sup>a</sup>	539 ± 18 <sup>a</sup>	527 ± 20
Moisture	14.3 ± 1.8 <sup>c*</sup>	13.3 ± 1.1 <sup>bc*</sup>	11.8 ± 0.7 <sup>a</sup>	11.9 ± 0.9 <sup>b</sup>	11.4 ± 0.6
pH <sub>water</sub>	8.40 ± 0.05 <sup>a*</sup>	8.52 ± 0.04 <sup>b*</sup>	8.55 ± 0.03 <sup>b</sup>	8.57 ± 0.05 <sup>b</sup>	8.58 ± 0.03
SOC (g kg <sup>-1</sup> soil)	19.0 ± 3.5 <sup>b*</sup>	16.4 ± 4.5 <sup>ab</sup>	14.6 ± 3.0 <sup>a</sup>	13.8 ± 2.6 <sup>a</sup>	11.8 ± 1.0
DOC (mg kg <sup>-1</sup> soil)	34.5 ± 5.6 <sup>b*</sup>	23.4 ± 2.8 <sup>a</sup>	17.4 ± 4.9 <sup>a</sup>	18.1 ± 3.2 <sup>a</sup>	20.3 ± 2.6
N <sub>tot</sub> (g kg <sup>-1</sup> soil)	1.8 ± 0.2 <sup>c*</sup>	1.4 ± 0.1 <sup>b</sup>	1.3 ± 0.1 <sup>ab</sup>	1.2 ± 0.1 <sup>a</sup>	1.1 ± 0.1
C:N	10.9 ± 1.6 <sup>a</sup>	11.4 ± 2.0 <sup>a</sup>	11.2 ± 2.3 <sup>a</sup>	11.5 ± 3.9 <sup>a</sup>	10.2 ± 3.1
Mineral N (mg kg <sup>-1</sup> soil)	31 ± 5 <sup>b*</sup>	16 ± 4 <sup>a*</sup>	11 ± 2 <sup>a</sup>	11 ± 2 <sup>a</sup>	11 ± 2
Olsen P (mg kg <sup>-1</sup> soil)	8.3 ± 2.1 <sup>b</sup>	5.8 ± 1.2 <sup>a</sup>	5.2 ± 1.6 <sup>a</sup>	5.6 ± 2.3 <sup>a</sup>	5.8 ± 1.3
CEC (cmol <sup>+</sup> kg <sup>-1</sup> soil)	14.7 ± 1.9 <sup>b*</sup>	13.0 ± 1.6 <sup>a</sup>	12.5 ± 1.4 <sup>a</sup>	12.4 ± 1.3 <sup>a</sup>	11.8 ± 0.4
Exch. Ca (cmol <sup>+</sup> kg <sup>-1</sup> soil)	14.7 ± 1.6 <sup>b*</sup>	13.3 ± 1.5 <sup>a</sup>	13.1 ± 1.1 <sup>a</sup>	12.8 ± 1.1 <sup>a</sup>	12.5 ± 0.5
Exch. Mg (cmol <sup>+</sup> kg <sup>-1</sup> soil)	0.73 ± 0.12 <sup>b*</sup>	0.47 ± 0.10 <sup>a*</sup>	0.39 ± 0.09 <sup>a*</sup>	0.38 ± 0.07 <sup>a*</sup>	0.29 ± 0.02
Exch. K (cmol <sup>+</sup> kg <sup>-1</sup> soil)	0.51 ± 0.08 <sup>b*</sup>	0.42 ± 0.08 <sup>ab*</sup>	0.36 ± 0.08 <sup>a</sup>	0.35 ± 0.07 <sup>a</sup>	0.31 ± 0.02

**Table 2**

Fraction	Tree row	1-2 m <sub>N</sub>	2-4 m <sub>N</sub>	4-6.5 m <sub>N</sub>	MC
f-LF (g C kg <sup>-1</sup> soil)	3.6 ± 1.3 <sup>b*</sup>	2.1 ± 0.9 <sup>a</sup>	1.5 ± 0.5 <sup>a</sup>	1.3 ± 0.5 <sup>a</sup>	1.2 ± 0.3
o-LF (g C kg <sup>-1</sup> soil)	2.2 ± 2.2 <sup>b*</sup>	1.7 ± 1.7 <sup>ab*</sup>	1.4 ± 1.4 <sup>a</sup>	1.2 ± 1.2 <sup>a</sup>	1.0 ± 1.0
50-2000 μm (g C kg <sup>-1</sup> soil)	3.6 ± 0.9 <sup>b</sup>	2.3 ± 0.3 <sup>ab</sup>	2.0 ± 0.2 <sup>a</sup>	2.0 ± 0.1 <sup>a</sup>	1.5 ± 0.1
<50μm (g C kg <sup>-1</sup> soil)	12.0 ± 2.1 <sup>b*</sup>	9.0 ± 1.8 <sup>a</sup>	8.6 ± 1.9 <sup>a</sup>	7.9 ± 1.8 <sup>a</sup>	7.8 ± 1.0

**Table 3**

Fraction	Tree row	1-2 m <sub>N</sub>	2-4 m <sub>N</sub>	4-6.5 m <sub>N</sub>	MC
Total (mg sugar C g <sup>-1</sup> soil)	4.4 ± 0.2 <sup>b</sup>	4.1 ± 0.1 <sup>a</sup>	4.0 ± 0.2 <sup>a</sup>	4.0 ± 0.2 <sup>a</sup>	4.0 ± 0.3
f-LF (mg sugar C g <sup>-1</sup> soil)	0.57 ± 0.19 <sup>b*</sup>	0.35 ± 0.11 <sup>a</sup>	0.32 ± 0.07 <sup>a</sup>	0.28 ± 0.04 <sup>a</sup>	0.29 ± 0.08
o-LF (mg sugar C g <sup>-1</sup> soil)	0.38 ± 0.05 <sup>a*</sup>	0.31 ± 0.07 <sup>a*</sup>	0.29 ± 0.20 <sup>a</sup>	0.22 ± 0.09 <sup>a</sup>	0.16 ± 0.06
< 50µm (mg sugar C g <sup>-1</sup> soil)	2.26 ± 0.4 <sup>a</sup>	2.15 ± 0.2 <sup>a</sup>	1.87 ± 0.7 <sup>a</sup>	2.10 ± 0.2 <sup>a</sup>	2.36 ± 0.1
GM:AX Total	0.95 ± 0.06 <sup>a</sup>	0.96 ± 0.04 <sup>a</sup>	1.00 ± 0.04 <sup>a</sup>	0.99 ± 0.04 <sup>a</sup>	0.97 ± 0.05
GM:AX f-LF	0.85 ± 0.15 <sup>b*</sup>	0.73 ± 0.09 <sup>ab</sup>	0.71 ± 0.09 <sup>ab</sup>	0.67 ± 0.09 <sup>a</sup>	0.62 ± 0.08
GM:AX o-LF	0.95 ± 0.03 <sup>a</sup>	0.91 ± 0.07 <sup>a</sup>	0.97 ± 0.04 <sup>a*</sup>	0.95 ± 0.04 <sup>a</sup>	0.90 ± 0.03
GM:AX < 50µm	1.05 ± 0.01 <sup>a</sup>	1.03 ± 0.02 <sup>a</sup>	1.04 ± 0.02 <sup>a</sup>	1.02 ± 0.02 <sup>a</sup>	1.03 ± 0.01

**Table 4**

Substrate	Tree row	1-2m <sub>N</sub>	1-2m <sub>S</sub>	2-4m <sub>N</sub>	2-4m <sub>S</sub>	4-6.5m <sub>N</sub>	4-6.5m <sub>S</sub>	MC
Water	0.57 ± 0.09 <sup>ab</sup>	0.59 ± 0.20 <sup>ab</sup>	0.72 ± 0.25 <sup>b</sup>	0.45 ± 0.12 <sup>ab</sup>	0.38 ± 0.11 <sup>a</sup>	0.36 ± 0.10 <sup>a</sup>	0.35 ± 0.06 <sup>a</sup>	0.43 ± 0.16
Glucose	2.65 ± 0.47 <sup>b*</sup>	2.12 ± 0.50 <sup>ab*</sup>	2.49 ± 0.50 <sup>b*</sup>	1.97 ± 0.41 <sup>ab*</sup>	1.98 ± 0.31 <sup>ab*</sup>	1.72 ± 0.41 <sup>a</sup>	1.81 ± 0.30 <sup>ab*</sup>	1.30 ± 0.21
Trehalose	1.99 ± 0.57 <sup>a*</sup>	1.80 ± 0.61 <sup>a*</sup>	1.98 ± 0.51 <sup>a*</sup>	1.39 ± 0.35 <sup>a</sup>	1.44 ± 0.22 <sup>a*</sup>	1.17 ± 0.52 <sup>a</sup>	1.11 ± 0.14 <sup>a</sup>	1.05 ± 0.19
Alanine	1.50 ± 0.56 <sup>c*</sup>	1.03 ± 0.61 <sup>abc*</sup>	1.44 ± 0.50 <sup>bc*</sup>	0.95 ± 0.20 <sup>abc</sup>	0.99 ± 0.34 <sup>abc</sup>	0.62 ± 0.16 <sup>a</sup>	0.72 ± 0.17 <sup>ab</sup>	0.73 ± 0.12
qCO <sub>2</sub>	0.22 ± 0.02 <sup>a</sup>	0.28 ± 0.05 <sup>a</sup>	0.29 ± 0.08 <sup>a</sup>	0.24 ± 0.09 <sup>a</sup>	0.19 ± 0.05 <sup>a</sup>	0.21 ± 0.03 <sup>a</sup>	0.20 ± 0.03 <sup>a</sup>	0.33 ± 0.11