



**HAL**  
open science

## Characterization of key aroma compounds in Burgundy truffle

Isabelle Andriot, Karine Gourrat, Rémy Reynaud, Sylvie Cordelle, Caroline Peltier, Olivier Berdeaux, Géraldine Lucchi

► **To cite this version:**

Isabelle Andriot, Karine Gourrat, Rémy Reynaud, Sylvie Cordelle, Caroline Peltier, et al.. Characterization of key aroma compounds in Burgundy truffle. The 16. Weurman Flavour Research Symposium, May 2021, Virtual meeting (Dijon), France. , 2021. hal-03253287

**HAL Id: hal-03253287**

**<https://hal.inrae.fr/hal-03253287v1>**

Submitted on 25 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Characterization of key aroma compounds in Burgundy truffle

Isabelle Andriot, Karine Gourrat, Rémy Reynaud, Sylvie Cordelle, Caroline Peltier, Olivier Berdeaux, Géraldine Lucchi



CSGA, ChemoSens Platform, INRAE Dijon, France



## Introduction



Truffles are a fungus of the genus *Tuber*. Some of them have an important economic value due to their gastronomic qualities appreciated in "grande cuisine". Périgord truffle (*Tuber melanosporum* or Black truffle) and White Alba truffle (*Tuber magnatum pico*) are actually the noblest ones. While these two species are well-valued, Burgundy truffle (*Tuber uncinatum*) is less well-characterized in its production area and truffle producers encounter some difficulties to sell their harvest at its fair value. The aroma compounds in truffles are investigated with sensory evaluation and two physico-chemical methods.

## Methodology

**French Fresh truffles**  
 ✓ Burgundy truffles (summer, autumn)  
 ✓ Truffles of south region (summer)

**Diced truffles**

### Sensory analysis


- ✓ 16 panellists were trained with 8 sessions
- ✓ Truffles were evaluated only by olfaction

### Classical descriptive profile (QDA)

Descriptor		Weak	Strong
Button mushroom	Spicy		
Cep	Smoked		
Girolle mushroom	Animal		
Undergrowth	Pungent		
Earthy	Pepper		
Mouldy	Fruity		
Nut	Vanilla		
Alcohol, pharmaceutical	Bread		

### Grounded truffles

### Dynamic Headspace (DHS)




### Gas Chromatography – Mass Spectrometry (GC-MS)


GC-MS parameters	
GC-MS	7890A - 8975 C TAD (Agilent Technologies)
Column	DB-HeavyWax (30 m x 0.25 mm x 0.5 µm)
T°C	40 to 240°C at 4°C/min
He	40 cm/s
Injection	Split/splitless 1 min 25 mL/min
Ionisation	Electronic impact
Detector	Quadrupole

Retention index

m/z  
**Aroma compounds identification**  
 (NIST08, INRAMass, WILEY11N17)



**In vitro dilution system :**  
 - 25 mg of truffles  
 - Incubation 36°C 2h  
 - Pre-analysis (blank): 1 min  
 - Analysis (sample): 2 min 15 s  
 - Post-analysis (calibration): 15 s



### Proton Transfer Reaction – Mass Spectrometry (PTR-MS)


PTR-MS parameters	
MS	PTR-TOF 8000 (Ionicon)
Mode	Funnel
Ionisation	H <sub>3</sub> O <sup>+</sup>
H <sub>2</sub> O	5.6 sccm
Transfer line	110°C
E/N	92 Td
Inlet flow	65 mL/min
Scans	500 ms

m/z  
**Aroma compounds mass fingerprint**  
 (statistical analysis)


## Results

### Sensory analysis

CVA of scores (55.98%)



**Figure 1: Canonic variables analysis of truffles (QDA)**  
 Truffles of Lot (south of France, summer)  
 Burgundy Truffles (summer) / Burgundy Truffles (autumn)




**Figure 2: Hierarchical classification analysis of truffles**  
 Truffles of Lot (south of France, summer)  
 Burgundy Truffles (summer) / Burgundy Truffles (autumn)


The figures 1 and 2 showed that:  
 - The Burgundy truffles (summer) are weakly odorant,  
 - The Lot truffles are characterised by forest odours and sweet odours,  
 - The Burgundy truffles (autumn) are characterised by negative odours and spicy odours.

### PTR-MS analysis

Observations (axes F1 et F2 : 55.83%)



**Figure 3: PCA of truffles from different harvests and places**



**Figure 4: Dendrogram of truffles (the same truffles as PCA)**

The results presented in figure 3 and 4 show that it is possible to discriminate the truffles according to places and seasons (1,2,3). However, the analyses of frozen mix truffles as a control have shown a sample dispersion according to time from June to November (data not shown). We can suspect truffle conservation problems and / or an instrumental bias.

### GC-MS analysis

The results obtained by GC-MS confirm the results obtained by PTR-MS and sensory analyses.

The truffles are separated according to harvest places and seasons (Figure 5).


A volcano plot with a foldchange higher than 3 and p-value<10<sup>-5</sup> allows compounds to be discriminated depending on their peak area in the TIC of the chromatogram. The results are presented in Table 1.

**Table 1: Compounds over-expressed, and the corresponding truffles**

Compounds which the quantities are more important	Truffles
butan-2-one	Bonnieres – Daix / November
butan-2-ol	Bonnieres – Daix / November
pentane-2,3-dione	Bonnieres – Daix / November
3-hydroxypentane-2-one	Bonnieres – Daix / November
2-hydroxypentane-3-one	Bonnieres – Daix / November
4-hydroxyhexan-3-one	Vauderben / June

Pentane-2,3-dione and butan-2-one are characteristic of the truffle freshness (4).

The same results are found with the PTR-MS for the truffles of october and november.



**Figure 5: PCA of truffles from different places in Burgundy, the same truffles analysed by sensory analyses and PTR-MS analyses. Unfortunately, the Lot truffles are not enough for the GC-MS**

## Conclusion & Perspectives


Sensory analyses, PTR-MS, GC-MS are, three of them, good methods to differentiate truffles according to harvest places and the seasons. The Burgundy truffles harvested in summer are less odorant than Burgundy truffles harvested in autumn. These results will have to be combined to microbiota and genetic analyses. This work will bring a scientific contribution to the creation of an IGP (Indication Géographique Protégée) request.

## Contact

**Centre des Sciences du Goût et de l'Alimentation**

UMR1324 INRA – 6265 CNRS – Université de Bourgogne - Franche Comté  
 – AgroSupDijon, Dijon, France, **contact:** [isabelle.andriot@inrae.fr](mailto:isabelle.andriot@inrae.fr)

This study is part of the research project BIJOU, supported by FEADER and Bourgogne Franche-Comté Region.



(1). Vita, F., et al. (2015). "Sci Rep 5(12629).  
 (2). Giocchini Anna, M., et al. (2005). " Rapid Communications in Mass Spectrometry 19(17): 2365-2370.  
 (3). March, R. E., et al. (2006). International Journal of Mass Spectrometry 249-250: 60-67.  
 (4). Vahdatzadeh M, et al. (2019). Food Microbiology 84:103251

