



HAL
open science

Stilbenes in grape berries and wine and their potential role as anti-obesity agents: A review

Nawel Benbougerra, Ruth Hornedo-Ortega, François Garcia, Toni El Khawand, Cédric Saucier, Tristan Richard

► To cite this version:

Nawel Benbougerra, Ruth Hornedo-Ortega, François Garcia, Toni El Khawand, Cédric Saucier, et al.. Stilbenes in grape berries and wine and their potential role as anti-obesity agents: A review. Trends in Food Science and Technology, 2021, 112, pp.362-381. 10.1016/j.tifs.2021.03.060 . hal-03256032

HAL Id: hal-03256032

<https://hal.inrae.fr/hal-03256032>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

1 **Stilbenes in grape berries and wine and their potential role as anti-obesity agents: a**
2 **review**

3 Nawel Benbouguerra^{a,1}, Ruth Hornedo-Ortega^{b,1}, François Garcia^a, Toni El Khawand^b, Cédric
4 Saucier^{a*}, Tristan Richard^{b*}

5 ^a *SPO, Univ. Montpellier, INRA SupAgro. 34093, Montpellier Cedex 5, France*

6 ^b *Unité de Recherche Oenologie, EA4577, USC 1366 INRA, ISVV, Université de Bordeaux,*
7 *Villenave d'Ornon Cedex, France*

8 *Corresponding authors: cedric.saucier@umontpellier.fr; tristan.richard@u-bordeaux.fr

9 ¹ These authors contributed equally to the manuscript.

10

11 **Abstract**

12 **Background**

13 Stilbenes are a group of naturally occurring phenolic compounds. These compounds are
14 synthesized by plants in response of biotic or abiotic stress situations. The major dietary
15 sources of stilbenes for humans are grape berries and wine. To accurately identify and
16 quantify these compounds, the research community has undertaken considerable efforts to
17 optimise samples preparations and analytical methods. In addition, stilbenes are well-known
18 to possess a wide range of biological activities for human health. One of the most recent and
19 promising properties demonstrated by stilbenes are their anti-obesity effects.

20 **Scope and Approach**

21 The principal objectives of this review to address and discuss about: i) stilbenes in grape
22 berries and wine, the factors that can modulate their concentrations and the most efficient
23 sample preparation techniques and analytical methods used for their detection; ii) the most
24 important investigations related to the *in vitro* and *in vivo* anti-obesity effects of grape and
25 wine stilbenes and the associated molecular mechanisms.

26 **Key Findings and Conclusions**

27 Stilbene concentration in grape and wines can vary substantially. The composition of stilbenes
28 in red wine is much more complex than in white wine. Until today, more than 30 stilbenes
29 have been identified in grapes and wines. The liquid chromatography coupled to mass
30 spectrometry is the most efficient method to investigate stilbene content. Regarding anti-
31 obesity properties of stilbenes, a great number of *in vitro* and *in vivo* studies have allowed to
32 demonstrate not only the positive implications of these bioactives but also the underlying
33 mechanisms of the observed effects.

34 **Keywords:** stilbenes; resveratrol; grape, wine; *Vitis vinifera*; obesity

- 35 **Abbreviation list**
- 36 **ACC:** acetylCoA carboxylase
- 37 **ACO:** acyl-coenzyme A oxidase
- 38 **aP2:** adipocyte protein 2
- 39 **ATAD3:** ATPase family AAA Domain-containing protein 3
- 40 **ATGL:** adipose triglyceride lipase
- 41 **BAT:** brown adipose tissue
- 42 **Bmp2:** bone morphogenetic protein 2
- 43 **CE:** capillary electrophoresis
- 44 **C/EBPs:** CCAAT/enhancer binding proteins
- 45 **CHS:** chalcone synthase
- 46 **CLA:** conjugated linoleic acid
- 47 **COX:** cyclooxygenase
- 48 **CPT-1:** carnitine palmitoyltransferase 1
- 49 **CRP:** C-reactive protein
- 50 **Cyc-D:** cyclin D
- 51 **DAD:** diode array detector
- 52 **Ddit3:** DNA-damage inducible transcript 3
- 53 **DLLME:** dispersive liquid-liquid microextraction
- 54 **ERK:** extracellular receptor kinase
- 55 **FABP4:** fatty acid binding protein 4
- 56 **FAS:** fatty acid synthase
- 57 **Fgf10:** fibroblast growth factor 10
- 58 **Fiaf:** fasting-induced adipose factor
- 59 **FLRD:** fluorescence detector
- 60 **FW:** fresh weight
- 61 **GC:** gas chromatography
- 62 **G6PDH:** glucose-6-phosphate dehydrogenase
- 63 **GLUT4:** glucose transporter 4
- 64 **HFCSD:** high fructose corn syrup diet
- 65 **HFD:** high fat diet
- 66 **HO-1:** heme-oxygenase 1
- 67 **HPD:** high protein diet

- 68 **HPLC:** high performance liquid chromatography
- 69 **HSL:** hormone-sensitive lipase
- 70 **IFN:** interferon
- 71 **IL:** interleukin
- 72 **IR:** insulin receptor
- 73 **KLF9:** Kruppel-like factor 9
- 74 **KR:** β -ketoacyl reductase
- 75 **LDL:** low density lipoprotein
- 76 **LPL:** lipoprotein lipase
- 77 **LXR α :** liver X Receptor alpha
- 78 **MS:** mass spectrometry
- 79 **MCP-1:** monocyte chemoattractant protein 1
- 80 **ME:** malic enzyme
- 81 **MEPS:** microextraction by packed sorbents
- 82 **Mfn2:** mitofusin 2
- 83 **miRNA:** microRNA
- 84 **MRM:** multiple reaction monitoring
- 85 **MS:** mass spectrometry
- 86 **mTOR:** mammalian target of rapamycin
- 87 **MTTP:** microsomal triglyceride transfer protein
- 88 **NF-KB:** nuclear Factor Kappa B
- 89 **Nrf1:** nuclear respiratory factor 1
- 90 **PDA:** photodiode array detector
- 91 **PDE3B:** phosphodiesterase 3B
- 92 **PKA:** protein kinase A
- 93 **PKC δ :** Protein Kinase C delta
- 94 **PPAR γ :** peroxisome proliferator-activated receptor gamma
- 95 **PRDM16:** PR domain-containing 16
- 96 **PVPP:** polyvinylpolypyrrolidone
- 97 **QqQ:** triple quadrupole
- 98 **QTOF:** quadrupole time-of-flight
- 99 **Rip 140:** receptor interacting protein 140
- 100 **ROS:** reactive oxygen species

101 **SBSE:** stir-bar sorptive extraction
102 **SCD-1:** stearyl-CoA desaturase-1
103 **SD:** standard diet
104 **SIRT1-AMPK-FOXO1:** sirtuin 1-AMP-activated protein kinase-Forkhead box protein O1
105 **SPE:** solid phase microextraction
106 **SPME:** solid phase microextraction
107 **SREBP1:** sterol regulatory element binding transcription factor 1
108 **STS:** stilbene synthase
109 **TG:** triglycerides
110 **TLR:** Toll-like receptor
111 **TNF- α :** tumor necrosis factor alpha
112 **UCP-1:** uncoupling protein 1
113 **UV:** ultraviolet
114 **WAT:** white adipose tissue
115 **WHO:** world health organization
116

117 **Highlights**

118 • Grape berries and wines are among the major sources of stilbenes in human nutrition

119 • Red wines contain more complex stilbenes than white wines

120 • Stilbenes could reduce obesity by regulating different pathways

121 • Some stilbenes show better anti-obesity activities than resveratrol

122 • Combination of stilbenes with others polyphenols give promising results

123

124 **Introduction**

125 Stilbenes (1,2-diphenylethylene) are phenolic compounds derived from the general
126 phenylalanine pathway (**Figure 1**). Among the commonly identified stilbenes, resveratrol is
127 the most popular compound (Gambini et al., 2015). The structural unit is constituted by two
128 phenyl rings linked together by an ethylene bridge forming a C6-C2-C6 chain. This double
129 bond allows stilbenes to exist in the *trans* (E) and in the *cis* form (Z) (Rivière et al., 2012).
130 The aromatic rings could be substituted by different functional groups such as hydroxyl,
131 methoxyl, prenyl or geranyl groups. Moreover, monomeric units can also be coupled, leading
132 to the formation of oligomers. Over a thousand of natural stilbenes have been described in the
133 literature. Despite this chemical diversity, only a limited number of plant families produce
134 these secondary metabolites, such as Polygonaceae, Cyperaceae, Pinaceae or Vitaceae
135 (Rivière et al., 2012). Furthermore, stilbenes are considered as phytoalexins since they are
136 associated with the resistance of plants to diseases and their synthesis is often a response to an
137 attack by phytopathogenic agents and other stress factors like UV irradiation, ozone, heavy
138 metal ions, injury or frost (Błaszczuk et al., 2019).

139 Grapes and red wine are among the major dietary sources of stilbenes for human nutrition,
140 especially in European countries (Guerrero et al., 2009; Weiskirchen and Weiskirchen, 2016).
141 However, it should be noted that stilbenes are also present in minor quantities in other
142 foodstuffs such as peanut, pistachio, almonds, berries, banana, pineapple, apple, peach,
143 passion fruit or dark chocolate (Neveu et al., 2010; Weiskirchen and Weiskirchen, 2016).
144 Focusing only on grapes berries and wines, it is noteworthy that stilbene amounts can be
145 extremely variable (Guerrero et al., 2016; Hasan and Bae, 2017). Several factors could
146 influence their quantities in grapes such as the stage of ripening, the grape varieties or various
147 external stimuli. The duration of maceration, yeast activity, and other winemaking processes

148 are also important factors that contribute to the final stilbene amounts described in wines.

149 For a long time, stilbene analyses in grape berries and wines were limited to resveratrol and
150 piceid (Lamuela-Raventós et al., 1995; Mattivi et al., 1995). Today, thanks to the analytical
151 methods such as ultra-high performance liquid chromatography tandem mass spectrometry
152 (UHPLC–MS/MS), more than thirty-three stilbenes have been identified and quantified in
153 grape berries and wines (**Figure 2**). In addition, it must be emphasized that not only the
154 selection of the most adapted analytical method but also the sample preparation techniques
155 are crucial to have reliable and accurate results.

156 Stilbenes have demonstrated to possess a great range of biological activities potentially
157 beneficial for human health such as neuroprotective, antioxidant and antitumor effects
158 (Weiskirchen and Weiskirchen, 2016). Among the more recent research lines, stilbenes are
159 gaining considerable interest as potential anti-obesity agents. Obesity is the most common
160 nutritional disorder in the world. According to the World Health Organization (WHO),
161 obesity is defined as an abnormal or excessive fat accumulation that may impact health.
162 Prevalence data estimate that 650 millions of adults were obese in 2016, representing about
163 13% of the world adult population. Moreover, alarming figures are presented regarding young
164 population. In fact, more than 124 millions of children and adolescents were obese in 2016
165 (WHO, 2018, <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>). It is
166 well known that the fundamental cause of obesity is an energy imbalance between calories
167 intake and loss, as a consequence globally, of a high intake of energy-fat foods and a
168 sedentary style of life. When a positive energy imbalance occurs, triglycerides (TG) are
169 accumulated in adipocytes producing an increase in the number of adipocyte (hyperplasia) or
170 an increase of its size (hypertrophy) (Hausman et al., 2001). Additionally, obesity can also
171 bring a potential risk to develop cardiovascular diseases, diabetes and some types of cancer,

172 endometrial, breast, ovarian, prostate, liver, gallbladder, kidney, and colon, among others
173 (Gallagher and LeRoith, 2015). Due to the increase of the prevalence and the associated
174 complications, several global government policies, laws and regulations have been developed
175 to halt and reverse the obesity epidemic. The first solution to stem obesity is focused in a
176 change of lifestyle by decreasing the energy/fat/sugar intake and increasing physical activity.
177 However, the implementation and effectiveness of these recommendations are usually
178 unsatisfactory. The research community tries then to find anti-obesity bioactive molecules
179 such as stilbenes that can be combined with other recommendations and treatments in order to
180 improve the results.

181 It has been shown that stilbenes may reduce obesity by regulating different pathways related
182 to fat metabolism as adipogenesis, lipogenesis, lipolysis and thermogenesis (Chou et al.,
183 2018; Fernández-Quintela et al., 2017). Resveratrol has been the most studied bioactive
184 compound, but a considerable number of works are already indicating that other stilbenes are
185 promising anti-obesity agents and may even be more potent than resveratrol.

186 For all above explained, the aim of this review is to address and discuss about: i) stilbene
187 compounds in grape berries and wines, the principal factors that can modulate their
188 concentrations and the most efficient sample preparation techniques and analytical methods
189 used for their detection; ii) the most important investigations related to the *in vitro* and *in vivo*
190 anti-obesity effects of grape and wine stilbenes and the associated molecular mechanisms.

191 **Stilbenes content in grape berries and wine**

192 ***Grape berries stilbenes***

193 In grape berries, stilbenes as other polyphenols are mainly concentrated in skins (Babazadeh
194 et al., 2017). They present a great variability in composition and content depending on

195 different biotic and abiotic factors including grape variety, stage of ripening, viticultural
196 factors and practices (Bavaresco et al., 2012; Błaszczuk et al., 2019).

197 The stilbene content offers sharp contrasts due to many potential external stimuli (Vincenzi et
198 al., 2013). Red varieties seem to present higher stilbene content than white ones (Guerrero et
199 al., 2010; Viñas et al., 2011). Concerning monomers, glucosides (piceid and astringin) are the
200 main compounds identified in grape berries (**Table 1**). The piceid (mean 1.36 mg/kg, *cis*- and
201 *trans*-isomers) is the main compound followed by astringin (mean 0.83 mg/kg), and
202 resveratrol (mean 0.68 mg/kg). Both isomeric forms, *trans*- and *cis*-isomers, were identified
203 in similar levels for piceid and mainly *trans* forms for astringin and resveratrol. These
204 compounds were observed in all berry growth stages in different concentrations (Jeandet et
205 al., 1991). They are subject to enzymatic transformations leading to the formation of a pool of
206 different compounds (Chong et al., 2009). The hydroxylation of resveratrol and piceid leads
207 to the formation of piceatannol and its glucoside, astringin (Bavaresco et al., 2002). In
208 addition to these three monomers some other minor monomers were identified such as
209 pterostilbene (Pezet et al., 1994), and isorhapontigenin (Fernández-Marín et al., 2012). The
210 oxidative coupling of resveratrol induces the formation of more complex oligomers including
211 dimers, trimers and tetramers (Takaya et al., 2005). Significant levels of oligomers were
212 reported (Rosso et al., 2016). In *Vitis vinifera* cultivars, dimers compounds were identified
213 such as pallidol (Vrhovsek et al., 2012), ϵ -viniferin and δ -viniferin (Flamini et al., 2013). In
214 addition to these dimers, trimers were identified, like miyabenol C and α -viniferin, and a pool
215 of tetramers including hopeaphenol and isohopeaphenol (Flamini et al., 2016; Rosso et al.,
216 2016). Unfortunately, few studies were focused on the quantification of these complex
217 compounds in grape berries. In contrast, wild *Vitis* species appear to contain a greater number
218 and diversity of oligomeric stilbenes (He et al., 2009; Jiang et al., 2012).

219 Different biotic and abiotic factors affect the stilbene content in grape berries, including the
220 grape varieties or species, stage of ripening, environmental conditions or postharvest
221 treatments (Hasan and Bae, 2017; Błaszczyk et al., 2019). The main factors that can have an
222 impact on stilbene quantities in grapes were described below.

223 ***Stage of ripening***

224 The development of grape berries is a dynamic process that involves a complex series of
225 changes divided into three major phases: before veraison, veraison and after veraison. Before
226 veraison, the cell division is rapid and all cells are established within two weeks of flowering,
227 followed by a subsequent sigmoidal increase in berry size (Deluc et al., 2007). At the end of
228 this period, the STS is accumulated and reaches its maximum two weeks before veraison. The
229 stage after veraison is characterized by the initiation of colour development (accumulation of
230 anthocyanins in red grapes) until maturity and the accumulation of stilbenes (Wang et al.,
231 2016). Even if the STS its maximum level two weeks before veraison, the accumulation of
232 stilbenes only starts in the second week after veraison and then increases until maturity from 5
233 to 35 mg/kg (Wang et al., 2016).

234 ***Grape varieties and species***

235 The accumulation of stilbenes seems to depend on the *Vitis vinifera* grape cultivar. In a
236 comparative study between three different cultivars, Syrah, Tempranillo and Merlot, Guerrero
237 et al. have shown that Syrah had the highest stilbene content 4.02 mg/kg followed by Merlot
238 and Tempranillo with 3.59 and 0.45 mg/kg; respectively (Guerrero et al., 2010). This result
239 was confirmed by Gatto et al. that compared 78 different cultivars during three years (Gatto et
240 al., 2008). Firstly, they observed higher stilbene content in red varieties than in white or pink
241 ones. Secondly, Pinots and related cultivars (Pinot Gris, Tête de Negre, and Noir) exhibited

242 the highest stilbene levels. Biochemical and transcriptomic analyses have shown the genotype
243 influence on stilbene accumulation in healthy grapevine berries. In the high yielding varieties,
244 glucosylated forms such as piceid were accumulated preferentially, resveratrol being mainly
245 produced after external stimuli such as fungal infection.

246 Nevertheless, these data have to be substantiated by consistent results because external
247 stimuli, such as light or pathogen infection could have higher effect than the intrinsic
248 differences between cultivars (Hasan and Bae, 2017). As for other secondary metabolites
249 (Hilbert et al., 2015), it is likely that wild species present higher content and diversity of
250 stilbene than the cultivated one (Jiang et al., 2012). In a recent article, Gabaston et al.
251 compared the stilbene content of nine wild *Vitis* species to *Vitis vinifera* showing that stilbene
252 content in wild *Vitis* was generally richest and more complex than the cultivated *Vitis*
253 (Gabaston et al., 2020). For example, *Vitis champinii*, an American species, may contain up to
254 ten times more stilbenes than *Vitis vinifera* (208.8 versus 20.5 mg/kg, respectively). These
255 results are in agreement with the data concerning other parts of the grapevine, such stem or
256 roots. Wild *Vitis* species, that have higher resistance to external stress, seem to contain more
257 stilbenes than cultivated vine (Pawlus et al., 2013b).

258 ***External stimuli***

259 Several external stimuli could modulate the stilbene content in grape berries including
260 pathogen infection, elicitor applications, or UVC treatments. The effects of stimuli could
261 increase the stilbene content more than ten times the normal. As stilbenes are well-known
262 phytoalexins (Langcake and Pryce, 1977), the effect of different pathogens on stilbenes
263 accumulation in grape was widely investigated. Infection by *Botrytis cinerea* induce the
264 accumulation of resveratrol mainly in its glucosidic form from 2.06 mg/kg before infection to
265 63.60 mg/kg after induction (Roldán et al., 2003). Similar results were observed for

266 *Plasmopara viticola*, *Uncinula necator* and *Rhizopus stolonifera* (Hasan and Bae, 2017), with
267 accumulation rate ranging between three and twelve times than control. Remarkably, the
268 increase of the resveratrol level in berries due to pathogen infection does not lead to a
269 significant modification of the stilbene content in wines (Jeandet et al., 1995; Roldán et al.,
270 2003). Authors suppose that grape berries with a high *Botrytis* infestation presented lower
271 resveratrol contents due to the oxidation of resveratrol by the laccase enzymes secreted by
272 fungi. In the same manner, chemical compounds such as elicitors are able to induce the
273 production of stilbenes (Krisa et al., 1999). This effect was mainly demonstrated in grape cell
274 cultures using different chemicals such as methyl jasmonate, salicylic acid, glucan, or
275 chitosan (Vuong et al., 2014). The induction could increase stilbene production ten times
276 compared to the control. This observation was confirmed in grape berries, where methyl
277 jasmonate induced a significant increase of resveratrol (from 0.78 to 1.40 mg/kg), piceatannol
278 (from 0 to 0.20 mg/kg), isorhapontigenin (from 0.15 to 0.21 mg/kg) and ϵ -viniferin (from 0.14
279 to 0.29 mg/kg) content up to two times in comparison to the control (Fernández-Marín et al.,
280 2014). Physical processes could also stimulate the production of stilbenes. Pre- and post-
281 harvest treatments by UVC induce an increase of the stilbene content in grape berries (Adrian
282 et al., 2000; Guerrero et al., 2010). The values were ranged between one and twenty times the
283 normal resveratrol concentration (Błaszczuk et al., 2019). In addition to the increase in the
284 amount of resveratrol, levels of other stilbenes rise in the same proportion. This effect was
285 observed on different stilbenes including piceid, piceatannol, ϵ -viniferin, δ -viniferin and
286 hopeaphenol (Guerrero et al., 2016).

287 To conclude, several biotic and abiotic factors can strongly influence the stilbenes content in
288 grape berries including species and various external stimuli. It is difficult to draw conclusions
289 on the relative impact of each of these factors. But some of these external stimuli such

290 elicitors and UVC treatments could be combined to increase artificially the stilbene content in
291 grape berries before and/or after harvest.

292 *Wine stilbenes*

293 Wine is an important dietary source of resveratrol (Kasiotis et al., 2013), in which it can be
294 found at concentrations up to 20 mg/L (Ribeiro De Lima et al., 1999). As for grape berries,
295 the monomeric forms are the most abundant stilbenes in wine, while oligomeric forms were
296 mainly identified in red wines (Guerrero et al., 2020). The main compounds identified in
297 wines were piceid (*cis*- and *trans*-isomers), resveratrol (*cis*- and *trans*-isomers), astringin (*cis*-
298 and *trans*-isomers) and piceatannol (**Table 1**). In addition to these compounds several other
299 stilbenes were identified including dimers such as pallidol (Landrault et al., 2002), ϵ -viniferin
300 (Amira-Guebailia et al., 2009; Landrault et al., 2002), δ -viniferin (Moss et al., 2013), ω -
301 viniferin (Guerrero et al., 2020), parthenocissin A (Vitrac et al., 2001), quadrangularin A
302 (Pawlus et al., 2013a); trimers such as miyabenol C (Guerrero et al., 2020), and α -viniferin
303 (Arraki et al., 2017); and tetramers such as hopeaphenol (Guebailia et al., 2006),
304 isohopeaphenol, and r2-viniferin (Guerrero et al., 2020). Glucosidic and diglucosidic forms of
305 these compounds were also identified including ϵ -viniferin diglucosides, pallidol 3-O-
306 glucoside, and pallidol diglucosides (Baderschneider and Winterhalter, 2000). In addition,
307 analyses by mass spectrometry indicated the presence of several other oligomers in wines
308 (Moss et al., 2013).

309 Red wines have a higher content in stilbenes than white wines (**Table 1**). Piceid is the main
310 stilbene in wines (means 8.26 and 0.88 mg/L in red and white wines, respectively).
311 Surprisingly, the *cis*-isomer levels are higher in both cases (means 6.20 and 0.68 mg/L in red
312 and white wines, respectively). In red wines, the piceid is followed by resveratrol (mean
313 2.29 mg/L). The derivatives of piceatannol are present at a concentration of about 1.73 mg/L.

314 Oligomers are minor compounds except isohopeaphenol with a mean value of 1.39 mg/L in
315 red wines (**Table 1**). No oligomers were reported in white wines. Several parameters are able
316 to modulate the stilbene content in wine including mainly the extraction from grape, but also
317 yeast strains, and enological practices.

318 *Duration of maceration*

319 Red wines contain higher stilbene levels than white wines that are obtained from a limited
320 maceration with the pomaces (**Tables 1**). In fact, the stilbene content in wines is strongly
321 modulated by the duration of the maceration and the solubility of these compounds in alcohol
322 (Mattivi et al., 1995; Kostadinović et al., 2012). Due to their polarity, monomeric glucosides
323 are extracted before their aglycone forms (Mattivi et al., 1995). While oligomers such as
324 δ -viniferin are observed in wines after few days of maceration (Poussier et al., 2003). After a
325 long time of maceration the concentration of stilbenes decrease mainly that of glucosides such
326 as *cis*- and *trans*-piceid (Poussier et al., 2003). The decrease of piceid could be attributed to
327 the β -glucosidase activity of the different yeasts (Jeandet et al., 1994). In addition to this
328 phenomenon, the *cis/trans* isomerization of stilbenes could explain the formation of some of
329 these compounds such as *cis*-resveratrol. The levels of this compound are similar to those of
330 *trans*-resveratrol in red wines and significantly lower in grapes (**Table 1**). Finally, even if few
331 studies about resveratrol degradation during maceration were conducted, yeasts seem to be
332 able to induce a degradation of resveratrol during wine fermentation (Vacca et al., 1997).

333 *Yeast activities*

334 As previously mentioned, yeast activities influence the stilbene content in wines. A
335 comparison between French and Macedonian yeasts demonstrated a variation of resveratrol
336 content up to four times depending on the yeast and maceration times (Kostadinović et al.,

337 2012). The β -glucosidase activities of yeasts induce the hydrolyze of glucosides which leads
338 to the formation of non glucosylated stilbenes (Mattivi et al., 1995). Thus, the selection of
339 exogenous yeasts could significantly affect the stilbene content in wines. The use of specific
340 yeasts enriched in β -glucosidase could positively increase the level of resveratrol in wines
341 (González-Candelas et al., 2000). Nevertheless more studies are needed to better understand
342 the impact of yeast activities on stilbene content in wines.

343 *Winemaking processes and wine ageing*

344 Few studies were focused on the impact of winemaking processes on the stilbene content in
345 wines. Grapes exposure to UVC radiations before winemaking induces an increase of stilbene
346 production in wines (Threlfall et al., 1999; Cantos et al., 2003). This positive effect is
347 observed on monomeric stilbenes such as resveratrol and piceatannol whereas other stilbenes
348 such as ϵ -viniferin are not affected (Fernández-Marín et al., 2014). On the contrary, the use of
349 some fining agents, such as polyvinylpyrrolidone (PVPP), could reduce the stilbene
350 content in wines (Threlfall et al., 1999; Vrhovsek et al., 1997).

351 As for winemaking processes few studies were focusing on the stilbene stability during wine
352 ageing. Stilbenes are known to be relatively stable compounds (Bavaresco et al., 2012;
353 Gaudette and Pickering, 2011). Nevertheless, even if the total stilbene content is not affected,
354 the ratio between resveratrol and piceid isomers could be impacted during wine ageing (Sun
355 et al., 2006; Favre et al., 2020). In addition some specific ageing processes such as those used
356 in Sherry wines could reduce the resveratrol content (Roldán et al., 2010). Finally, heat could
357 induce oxidative coupling between monomeric stilbenes inducing the formation of oligomers
358 in wines (El Khawand et al., 2020).

359 **Analytical methods for stilbenes analysis in grapes and wines**

360 The relatively low concentration of stilbenes, their structural diversity, and the complexity of
361 grape and wine matrices restrain their identification and quantification. Today, the studies are
362 mainly focused on the monomeric stilbenes. Several analytical procedures were developed in
363 order to determine the stilbene content in grapevines and wines from monomers to complex
364 oligomers. These strategies often require multi-steps sample preparation (Baderschneider and
365 Winterhalter, 2000; Romero-Pérez et al., 2001; Liu et al., 2013).

366 Concerning grapevines, the first step consists of the extraction of stilbenes from the raw
367 material. Several extraction procedures were applied including classical solvent extraction
368 from fresh, frozen, dried or lyophilized berries (Romero-Pérez et al., 2001), or more recently
369 ultrasonication-assisted extraction (Cho et al., 2006). The classical solvent extraction remains
370 the most applied technique the main used methodology using acetone, chloroform, ethanol,
371 ethyl acetate, methanol, and some mixed solvents. Methanol, pure or mixed, is the most
372 widely used solvent for stilbene extraction from grapevine berries. Several studies were
373 focused on optimizing the extraction conditions form grapevine berries (Liu et al., 2013;
374 Romero-Pérez et al., 2001; Sun et al., 2006). The main parameters analyzed were the solvent,
375 the extraction time, the temperature, and the ratio solid to solvent. Based on Liu et al. (2013)
376 studies (Liu et al., 2013), the best extraction solutions were: methanol or methanol/ethyl
377 acetate (1:1 (v/v)) with a ratio solid to solvent of 1 g/10 mL at 25°C for 24 h. In addition, in
378 order to avoid *cis/trans* isomerization, extraction should be carried out in the dark (Careri et
379 al., 2003). Interestingly, innovative extraction techniques such as ultrasonication-assisted
380 extraction seem to be able to increase the stilbene extraction rate while reducing time and
381 solvent consumption (Cho et al., 2006).

382 As for grape berries, several methods were proposed to extract stilbenes in wines, from direct

383 analyses (Lamuela-Raventós et al., 1995) to more complex procedures using solid phase
384 extraction (Mattivi et al., 1995), stir-bar sorptive extraction (Viñas et al., 2008),
385 microextraction by packed sorbents (Gonçalves and Câmara, 2011), liquid–liquid extraction
386 (Rabesiaka et al., 2011), dispersive liquid–liquid microextraction (Rodríguez-Cabo et al.,
387 2012), semi preparative liquid chromatography (Amira-Guebailia et al., 2009), or counter
388 current chromatography (Fernandez-Marin et al., 2012). The main drawback of these methods
389 is that they are mainly focused on resveratrol and piceid. In the research conducted by
390 Baderschneider and Winterhalter a combination of solid phase extraction and counter current
391 chromatography were used to identify new stilbene skeletons in wines (Baderschneider and
392 Winterhalter, 2000).

393 Several methods were used for stilbene analysis in grape berries and wines including capillary
394 electrophoresis (CE), gas chromatography (GC), high performance liquid chromatography
395 (HPLC) and ultra-high performance liquid chromatography (UHPLC).

396 The capillary electrophoresis is appropriate for the separation of small molecules and could
397 separate some optical isomeric forms. This method allows separating positional isomers as
398 well as optical isomers. This method has different advantages such as fast and accurate
399 analysis, low amount and consumption (Coelho et al., 2016).

400 Gas chromatography (GC) was successfully used for resveratrol quantification in wines
401 (Barbanti et al., 1996), and grapes (Viñas et al., 2009). Today, GC analyses were coupled with
402 mass spectrometry detector (GC–MS) in order to increase the accuracy and sensibility of the
403 detection (Rodríguez-Cabo et al., 2016). These methods provide limits of detection close to
404 ng/L in wines. However, GC analyses have different drawbacks such as isomerization,
405 degradation of analytes.

406 High-performance liquid chromatography (HPLC) is considered among the most commonly
407 applied method for stilbenes analysis in grapes (Błaszczuk et al., 2019) and wines
408 (Fabjanowicz et al., 2018), using different detection systems such as UV-visible or diode
409 array detectors (Mattivi et al., 1995; Sun et al., 2006), or fluorescence detectors in order to
410 increase sensitivity and specificity (Vitrac et al., 2002). Nowadays, HPLC combined with
411 mass spectrometry (HPLC-MS) has become the most used technique for the determination of
412 phenolic compounds in general and stilbenes in particular (Pugajeva et al., 2018). This
413 method allows the identification of all stilbenes (free and conjugated) without any
414 derivatization or hydrolysis of samples. Different mass detectors were used including triple
415 quadrupole (QqQ) and quadrupole time-of-flight (QTOF). Information on liquid
416 chromatography methods developed for stilbenes identification and quantification in grape
417 berries and wines are summarized in **Table 2**.

418 In recent years, UHPLC coupled with mass spectrometry (UHPLC-MS) has been widely used
419 as an alternative for other methods due to its higher sensitivity, accuracy, rapidity, and low
420 solvent consumption for grape berry (Flamini et al., 2013), and wine analyses (Moss et al.,
421 2013; Guerrero et al., 2020). For example, based on a targeted metabolomics approach using
422 UHPLC-QTOF mass spectrometer, Flamini et al. have identified and quantified eighteen
423 stilbenes in grape berries including oligomers. In wines, an UHPLC-QqQ-MS method was
424 designed to identify and quantify fifteen stilbenes in white and red wines with limit of
425 detection ranging between 4 and 28 µg/L (Guerrero et al., 2020). Similarly, using a UHPLC-
426 QTOF mass spectrometer, forty-one stilbenes were identified in red wines, including six
427 monomers, twenty-three dimers, eight trimers and four tetramers (**Table 3**).

428 Due to the complexity and the diversity of stilbene structures, pure compounds are needed as
429 standard to obtain accurate results. Unfortunately, since pure samples of complex stilbene

430 oligomers are often unavailable, results are expressed in resveratrol or piceid equivalent
431 (Flamini et al., 2013). This approach could lead to severe underestimation of the oligomeric
432 stilbene content (Biais et al., 2017). For example, the quantification of ϵ -viniferin using
433 resveratrol by a HPLC-DAD method underestimated the concentration ϵ -viniferin, R-viniferin
434 and isohopeaphenol by a factors upper than two, five and ten, respectively.

435 **Stilbenes, *in vitro* and *in vivo* anti-obesity effects and molecular mechanisms**

436 *Anti-obesity in vitro effects of stilbenes*

437 As was noted in the Introduction section, stilbenes might act as anti-obesity agents by
438 regulating different fat metabolism pathways such as adipogenesis, lipogenesis, lipolysis and
439 thermogenesis (Chou et al., 2018; Fernández-Quintela et al., 2017).

440 Adipogenesis is defined as the differentiation process of preadipocytes to fully mature to
441 adipocytes. This process can be divided in different stages: growth phase and growth arrest,
442 clonal expansion, gene expression of lipogenic proteins inducing triglyceride (TG)
443 accumulation and differentiation and cell death (apoptosis) (Esteve Ràfols, 2014). Some
444 transcription factors like the peroxisome proliferator-activated receptor gamma (PPAR γ), the
445 sterol regulatory element binding protein 1c (SREBP-1c) and CCAAT-enhancer binding
446 proteins (C/EBPs) are key elements in production of the fully mature adipocytes (Rosen et al.,
447 2002). Moreover, others proteins such as glucose transporter 4 (GLUT4), adipocyte protein 2
448 (aP2) or lipoprotein lipase (LPL) are also involved in this process (MacDougald and
449 Mandrup, 2002).

450 Concerning the anti-adipogenic effects of stilbenes, the most investigated molecule was
451 resveratrol. Several papers demonstrated that resveratrol is able to inhibit preadipocyte and
452 adipocyte differentiation through the decrease of gene and protein expressions of PPAR γ ,

453 C/EBP α and C/EBP β at concentrations ranging between 0.03-400 μ M in mouse, bovine and
454 human cells (**Table 4**). In addition to suppressing the expression of several key lipogenic
455 genes, resveratrol can interfere by diminishing preadipocyte proliferation and inhibiting the
456 clonal expansion stage or the cell cycle entry to G2/M phase. Actually, a decrease of the
457 expression of cell cycle genes, such as cyclin D1 and A2, cyclin-dependent kinase 2 and 4,
458 and DNA-damage inducible transcript 3 (Ddit3, also known as Chop-10), was observed
459 (Kwon et al., 2012b; Mitterberger and Zwerschke, 2013; Santos et al., 2014). Furthermore,
460 resveratrol interacted with the insulin receptor (IR) in 3T3-L1 preadipocyte cells and inhibited
461 the insulin signaling pathway in the early phase of adipogenesis (Kwon et al., 2012b).

462 More recently, a very interesting work of Eseberri et al. showed that some resveratrol
463 metabolites, and specifically resveratrol 3'- and 4'-glucuronide (at 25 μ M concentration), were
464 able to regulate and inhibit the expression of C/EBP β and Krüppel-like factor (KLF9) that
465 mediates both the early and late stages of the differentiation program (Eseberri et al., 2017).

466 Some of these studies, also showed that resveratrol decrease or attenuates the production of
467 other key adipogenic proteins such as fatty acid binding protein 4 (FABP4) that regulates
468 adipogenesis by downregulating PPAR γ (Garin-Shkolnik et al., 2014). Santos et al. also
469 reported an important modulation role of resveratrol on the gene expression of Adipogenic,
470 Bone morphogenetic protein 2 (Bmp2), fatty acid synthase (FAS), fibroblast growth factor 10
471 (Fgf10) and leptin (Santos et al., 2014). Likewise, this lipid-lowering effect has been
472 associated and depends on the sirtuin 1-AMP-activated protein kinase-Forkhead box protein
473 O1 (SIRT1-AMPK-FOXO1) pathway (Liu et al., 2018). It is well know that SIRT 1 is
474 responsible of fat mobilization in mature adipocytes and its activation give rise to the
475 inhibition of PPAR γ expression (Picard et al., 2004). Additionally, AMPK pathway plays an
476 important role on the control of body fat stress. In fact, some studies have demonstrated that

477 hormones, such as leptin and adiponectin, adrenergic agonists, and metformin, activate
478 AMPK in adipocytes (Rossmeisl et al., 2004). AMPK negatively regulates white
479 adipogenesis, specifically blocking the clonal expansion of preadipocytes by attenuating
480 adipocyte differentiation (Kang et al., 2005). In addition, AMPK activation in the early phase
481 of differentiation inhibits PPAR γ and C/EBP α expression as well as late adipogenic markers
482 such as FAS and acetylCoA carboxylase (ACC). For these reasons, it is well accepted that
483 AMPK activity is inversely related to white adipogenesis.

484 Moreover, resveratrol can induce the cell death in both mice 3T3-L1 cells (Rayalam et al.,
485 2008), and human pre-adipocytes (Liu et al., 2018). In fact, a modulation of the expression of
486 caspase-3, and the pro-apoptotic protein Bax was demonstrated after resveratrol treatment that
487 also implicated the SIRT1-AMPK-FOXO1 pathway (Liu et al., 2018).

488 Concerning other stilbenes, r- and r2-viniferin (tetrameric stilbenes) that were recently
489 identified in wine (Guerrero et al., 2020), are also thought to be able to inhibit adipocytes
490 differentiation and reduce lipid accumulation in 3T3-L1 cells by decreasing the expression of
491 PPAR γ , C/EBP α and FABP4 genes (Tie et al., 2018). In accordance with these results, r2-
492 viniferin suppressed the adipogenic process by blocking the cell cycle at the G1-S phase
493 through p21- (CDK inhibitor) and Rb-dependent suppression of transcription in 3T3-L1 cells
494 (Kim et al., 2008). Long-term treatment at low concentrations (5-10 μ M) by pterostilbene, a
495 methylated derivative of resveratrol, inhibited adipocyte differentiation in 3T3-L1
496 preadipocytes and 3T3-F442A cells (Gomez-Zorita et al., 2017; Hsu et al., 2012). This
497 compound induced heme-oxygenase 1 (HO-1) expression which acts as a regulator of
498 Chop10, suppressing in consequence the initiation of mitotic clonal expansion (Seo et al.,
499 2017). Piceatannol acts in the early phase of adipogenesis delaying the cell cycle entry into
500 G2/M phase at 24 h after initiation of adipogenesis and suppressing the mitotic clonal and the

501 activation of the insulin-signaling pathways (Kwon et al., 2012a). Furthermore, ϵ -viniferin, a
502 resveratrol dimer, showed anti-adipogenic effects by downregulating PPAR γ mRNA levels at
503 50 μ M concentration (Ohara et al., 2015).

504 Additionally, the combination of stilbenes with other polyphenols may present synergistic or
505 additional effects. Some studies proved that the combination of resveratrol with genistein,
506 quercetin or epigallocatechin gallate enhanced the resveratrol efficiency by inhibiting
507 adipogenesis and decreasing the lipid accumulation and TG content in 3T3-L1 and human
508 primary adipocytes cells (**Table 4**). More specifically, the treatment with these compounds
509 (30 μ M) decreased the protein expression of PPAR γ , C/EBP α , FABP4 and perilipin (Ahmed
510 et al., 2017). Additionally, an increase of the apoptotic process has also been observed in
511 early- and mid-phase maturing and lipid-filled mature human primary adipocytes after
512 resveratrol, genistein and quercetin treatment (Park et al., 2008).

513 Lipogenesis process involves *de novo* fatty acid and TG synthesis from glucose metabolism
514 products (Wang et al., 2004). In human body this pathway is active in liver and adipose tissue.
515 Once glucose is incorporated into the cells, it undergoes a series of biochemical
516 transformations (glycolysis) to produce acetyl-CoA. This last compound is transformed in
517 malonyl CoA by acetyl-CoA carboxylase (ACC) which is a substrate for fatty acid synthase
518 (FAS) to synthesize fatty acids. Two other enzymes, malic enzyme (ME) and glucose-6-
519 phosphate dehydrogenase (G6PDH) supply NADPH to fatty acid synthesis reactions. SREBP-
520 1c has also an important role as a transcription factor by regulating the expression of ACC
521 and FAS (Vázquez-Vela et al., 2008).

522 In addition, at adipose tissue level, the fatty acids involved in triacylglycerol synthesis can be
523 transported via triacylglycerol-rich lipoproteins (chylomicrons and low density lipoproteins
524 (LDL)). The enzyme lipoprotein lipase (LPL), which is located in the luminal surface of

525 endothelial cells, hydrolyses lipoprotein triacylglycerols into two free fatty acids and one
526 monoacylglycerol. In this case, PPAR γ is the transcriptional factor that controls the
527 expression of this enzyme. AMPK plays again an important role because its phosphorylation
528 is related with the decrease of fatty acid synthesis and the activation of ACC and also with the
529 downregulation of SREBP-1c through the mammalian target of rapamycin (mTOR) and Liver
530 X Receptor alfa (LXR α) (Zhang et al., 2009).

531 Resveratrol showed anti-lipogenic effects at both hepatic and adipose levels (**Table 4**).
532 Treatment with this bioactive compound at low doses down regulates, PPAR γ , SREBP-1c,
533 ACC, FAS gene expression in adipocyte cells (Chen et al., 2011; Liang et al., 2013). A
534 similar effect has also been described for pterostilbene (5-10 μ M) that is able to reduce the
535 G6PDH activity in 3T3-L1 cells (Hsu et al., 2012). The ability of stilbenes to increase the
536 phosphorylation of APMK and their capacity to bond to the ketoacyl reductase (KR) domain
537 of FAS are two mechanisms implicated in the anti-lipogenic effects of these bioactives (Chen
538 et al., 2011; Li et al., 2016; Liang et al., 2013).

539 Alternatively, by using different hepatocytes cell lines, resveratrol and oxyresveratrol are able
540 to decrease the hepatic lipogenesis by suppression of SREBP-1, FAS, ACC and stearyl- CoA
541 desaturase-1 (SCD-1) (Gnoni and Paglialonga, 2009; Jin et al., 2013; Choi et al., 2014; Lee et
542 al., 2018). This last protein is a rate-limiting enzyme that catalyzes the synthesis of
543 monounsaturated fatty acids and it is essential for the assembly of VLDL particles, which
544 transport triacylglycerol (TG) from liver to adipose tissue and other sites (Li et al., 2009).
545 AMPK activation and the inhibition of LXR α which activate the SREBP-1 have been the
546 molecular pathways related (Jin et al., 2013; Choi et al., 2014; Lee et al., 2018).

547 Lipolysis and β -oxidation occur when the body requires energy. In adipocytes the TG are
548 metabolized giving rise to glycerol and fatty acids by the catabolic action of adipose

549 triglyceride lipase (ATGL) and hormone-sensitive lipase (HSL). HSL activation depends on
550 protein kinase A (PKA) phosphorylation, which is mediated via the accumulation of cAMP
551 (Duncan et al., 2007). With regard to the lipolytic effects of stilbenes, resveratrol was the
552 main compound studied. Some published articles proved that resveratrol, at different doses
553 (0.03-400 μ M), is able to enhance the free fatty acid and the glycerol release by increasing
554 HSL and ATGL expression in some cellular models (bovine intramuscular adipocytes, 3T3-
555 L1 and SGBS adipocytes) (Rayalam et al., 2007; Chang et al., 2016; Lee et al., 2018).
556 Interestingly, resveratrol had also synergistic action with genistein, CLA and epinephrine
557 enhancing its lipolytic capacity (Rayalam et al., 2007; Szkudelska et al., 2009; Lasa et al.,
558 2011).

559 An increment of β -oxidation capacity in adipocyte cells was demonstrated by resveratrol and
560 oxyresveratrol. In fact, the upregulation of carnitine palmitoyltransferase 1 (CPT-1) which is
561 necessary for mitochondrial import of fatty acids and the repression of receptor interacting
562 protein 140 (rip 140), a suppressor of oxidative metabolism, was involved (Mercader et al.,
563 2011; Lee et al., 2018).

564 Finally, another important pathway related to fat metabolism is thermogenesis that is literally
565 defined as heat production. The white adipose tissue (WAT) and the brown adipose tissue
566 (BAT) are the main types of adipose tissues with antagonistic functions. WAT stores the
567 excess of energy in the form of TG and BAT is specialized in heat production. BAT is
568 specialized in dissipate energy thanks to the high number of mitochondria. Mitochondria
569 membranes contain high amounts of Uncoupling protein 1 (UCP-1), an inner membrane
570 protein that uncouples the electron transport chain from ATP synthesis resulting in energy
571 dissipation rather than ATP synthesis (Madden, 2017). Under certain conditions (e.g. extreme
572 low temperature or β -3 adrenergic agonist) the number of mitochondria increased drastically;

573 this process is named “browning” and the type of WAT is called “beige” (Bartelt and Heeren,
574 2014). Thus, an increase of BAT thermogenesis is considered nowadays as a potential
575 strategy to reduce obesity.

576 Several recent *in vitro* studies have revealed that resveratrol (10 μ M) can activate the
577 “browning” process. In particular, an increase of UCP-1 expression in stromal vascular cells
578 from interscapular WAT and BAT has been observed (Wang et al., 2015; Wang et al., 2017).
579 Beyond this, these studies have also demonstrated an increase of expression of others
580 transcriptional factors of brown adipogenic program as PR domain-containing 16 (PRDM16)
581 that leads to the activation of mitochondrial biogenesis. This effect was related with AMPK
582 signaling pathway activation. Other papers showed that resveratrol at 100 μ M concentration
583 up-regulated the gene expression of some mitochondrial activity regulators such as SIRT3
584 that influences the mitochondrial function by reducing membrane potential, mitofusin 2
585 (Mfn2) that participates in mitochondrial fusion in mammalian cells, and ATPase family
586 AAA Domain-containing protein 3 (ATAD3), a protein that regulates mitochondrial
587 biogenesis (Rayalam et al., 2008; Li et al., 2016).

588 To conclude, resveratrol but also others stilbenes and the combinations or these with other
589 bioactive compounds have proved be able to counteract the adipogenic and the lipogenic
590 (adipose and hepatic level) processes by reducing the gen and protein expression of several
591 transcription factors, avoiding the adypocytes differentiation/proliferation or promoting the
592 apoptosis program with a direct implication of SIRT, AMPK and FOXO pathway.
593 Additionally, these compounds can enhance not only the lipolysis and β -oxidation but also the
594 thermogenesis and mitochondrial biogenesis although more studies are necessary to be able to
595 claim the mechanism of action.

596 *Anti-obesity in vivo effects of stilbenes*

597 Regarding the *in vivo* effects and similarly with the *in vitro* studies the most examined
598 stilbene molecule was resveratrol.

599 Several animal models were used to investigate the anti-obesity effects of stilbenes, mice
600 (Swiss, C57BL/6J, FVB/N, CD1, kunming) and rats (Sprague-Dawley, Wistar, Zucker (*fa/fa*))
601 being the most commonly used. In addition, other animal models such as *Megalobrama*
602 *amblycephala* (fish) or *Caenorhabditis elegans* were used. Experimentally, these species were
603 treated with standard (SD), high fat (HFD), high protein (HPD), high fructose corn syrup
604 (HFCSD) diets supplemented with stilbenes (between 1 to 300 mg/kg/day) during different
605 periods of time varying from some hours to 20 weeks (Table 5). This large numbers of
606 conditions (animal model, dose, diet and period of treatment) makes sometimes difficult the
607 understanding of results found by these studies. Nevertheless, several studies showed that
608 resveratrol is able to mitigate the body weight gain in different animal models. In some
609 instances, even a 50% reduction was described (Choi et al., 2014; Jeon et al., 2014; Majumdar
610 et al., 2014; Qiao et al., 2014; Mendes et al., 2016).

611 In addition to weight reduction, an attenuation of lipid deposition on internal adipose tissues
612 such as epididymal and intraperitoneal tissues was observed after resveratrol treatment
613 (Gómez-Zorita et al., 2013; Jeon et al., 2014; De Almeida Pinheiro et al., 2017; Zhang et al.,
614 2018). This fact was related to the capacity of resveratrol to control and reduce the
615 adipogenesis process. Particularly, resveratrol (0.4%) reversed the HFD-induced up-
616 regulation of key adipogenic genes such as PPAR γ , C/EBP α , SREBP-1c, FAS, LPL, aP2, and
617 leptin in mice adipose tissues (Kim et al., 2011). The results reported by Kim et al.
618 highlighted that galanin-mediated signaling molecules are also implicated on the anti-
619 adipogenesis effects of resveratrol. Galanin is a neuropeptide that plays a role in food intake

620 and its circulating serum levels are elevated in obese individuals (Kim and Park, 2010).
621 Resveratrol has demonstrated its capacity to significantly reverse the HFD-induced up-
622 regulation of galanin and its receptors along with increased expression and/or activation of
623 downstream molecules related to adipogenesis, such as Protein Kinase C Delta (PKC δ),
624 cyclin D (Cyc-D), transcriptional factor E2F1, and Extracellular Receptor Kinase (ERK)
625 (Kim et al., 2011).

626 There is evidence that microbiota has an important role in obesity. In fact, the presence of
627 certain bacteria and overall, the relative proportions and composition of microbial
628 communities is key for energy homeostasis (Tennyson and Friedman, 2008). Although this
629 research line is recent and remains not fully explored, some studies were published indicating
630 that in obese people exposed to low calorie diet, *Bacteroidetes* level increased while
631 *Firmicutes* levels decreased (Ley et al., 2006). In addition, it seems that gut microbiota can
632 abolish the expression of fasting-induced adipose factor (Fiaf) that belongs to the family of
633 fibrinogen/angiopoietin-like proteins. When Fiaf was suppressed an increase of LPL and a
634 higher deposition of TG in adipose tissues have been observed (Bäckhed et al., 2004). For this
635 reason, the study of gut microbiota in obesity research is gaining more and more relevance.

636 HFD was related with microbiota dysbiosis (promoting the growth of endotoxin producers)
637 producing a decrease of *Lactobacillus* and an increase of *Enterococcus faecalis*. In this
638 context it is worth mentioning that resveratrol long term (12 weeks) treatment in Kunming
639 mice's demonstrated that this stilbene increased the *Bacteroidetes* to *Firmicutes* ratio and
640 diminished the growth of *Enterococcus faecalis* (Qiao et al., 2014). Similarly, the numbers of
641 *Lactobacillus* and *Bifidobacterium* were significantly increased. At intestinal level, this work
642 showed that resveratrol increases the Fiaf expression which can be linked to the suppression
643 of LPL and SCD-1 expressions in the liver, and with the expression of adipogenesis/

644 lipogenesis genes (PPAR γ , ACC1, and FAS) in adipose tissues (Qiao et al., 2014).

645 Likewise, some of the above-mentioned papers proved that resveratrol can also be able to
646 reduce lipogenesis process. A reduction of expression levels of FAS, ACC, SCD-1, and
647 G6PDH, and therefore, a suppression of fatty acid uptake and TG synthesis at adipose and
648 hepatic level has been observed (**Table 5**). Furthermore, resveratrol causes the increase of
649 GLUT2 mRNA expression allowing to restore the normal glucose fluxes induced by HFD
650 (Zhang et al., 2018). Another recent work showed for the first time that resveratrol modifies
651 the microRNA (miRNA) profile in WAT. Actually, the reduction of protein levels of FAS,
652 SREBP-1 and SP-1 (acts together with SREBP1 to synergistically activate the promoter of
653 FAS) has been linked to the up-regulation of miR-539-5p (Gracia et al., 2016).

654 An inhibition of adipogenesis by resveratrol at hepatic level has also been evidenced. Some
655 works have mentioned a decrease of mRNA expression of PPAR γ , SREBP-1c and FAS after
656 resveratrol supplementation with HFD, HFSCD or atherogenic diets (Ahn et al., 2008; Shang
657 et al., 2008; Alberdi et al., 2013; Andrade et al., 2014; Sadi et al., 2015). Furthermore, this
658 action is mediated by the activation of AMPK/SIRT1 axis (Shang et al., 2008; Alberdi et al.,
659 2013). Indeed, Shang et al. showed an increase of 164% of AMPK phosphorylation level in
660 liver after the oral administration of resveratrol (100 mg/Kg/day) during 10 weeks (Shang et
661 al., 2008). Furthermore, higher gene and protein expressions levels of a great number of
662 insulin-signaling molecules including IR, IRS-1/2, eNOS as well as SIRT1 have been outlined
663 after resveratrol supplementation (Sadi et al., 2015).

664 One of the key developments in obesity research is the recognition that this disorder is also
665 characterized by chronic mild inflammation. Indeed, an increase of circulating levels of
666 inflammatory markers in obese people such as CRP (C-reactive protein), tumor necrosis
667 factor alpha (TNF- α), interleukin (IL)-6 and IL-18, was described (Festa et al., 2001;

668 Monteiro and Azevedo, 2010). In this context, some papers indicated that a reduction of
669 inflammatory markers (TNF- α , interferon IFN α , IFN β , IL-6, monocyte chemoattractant
670 protein 1 (MCP-1), and CRP) was observed in the adipose tissues of mice and rat after
671 resveratrol oral treatment, relating that with the repression of Toll-like receptor (TLR)2- and
672 TLR4-mediated pro-inflammatory signaling cascades (as Nuclear Factor Kappa B (NF- κ B)
673 pathway) in the adipose tissues of mice and rat after resveratrol oral treatment (Kim et al.,
674 2011; Gómez-Zorita et al., 2013). This action was also observed at hepatic level and
675 evidenced by the reduction of TNF- α , IL-1 β , IL-6 and NF- κ B expression in liver in
676 association with the up-regulation of SIRT1 in mice treated with an HFD and resveratrol
677 (Andrade et al., 2014).

678 Additionally, resveratrol is able to improve fatty acid oxidation in liver and WAT.
679 Particularly, Enzyme activities involved in fatty acid oxidation as CPT-1a, a marker of
680 mitochondrial oxidation, and acyl-coenzyme A oxidase (ACO), a marker of peroxisome
681 oxidation, were significantly increased by resveratrol in liver (Gómez-Zorita et al., 2012;
682 Alberdi et al., 2013). It was furthermore demonstrated by other authors that the increase of
683 expression levels of both CPT-1 and UCP-2 may exert a protective effect of resveratrol on
684 mitochondrial dysfunction not only by inhibition of fatty acid oxidation but also in association
685 with reactive oxygen species (ROS) generation (Khaleel et al., 2018). By using a fish animal
686 model (*M. amblycephala*), the supplementation with different doses of resveratrol (0.04, 0.36
687 and 1.08%) resulted in a significant reduction of ATGL, CPT-1 and the microsomal
688 triglyceride transfer protein (MTTP) implying up-regulation of lipolysis and β -oxidation
689 (Zhang et al., 2018). Finally, an augmentation of HSL, without affecting the ATGL levels,
690 was observed in Zucker (*falfa*) rats after oral supplementation of 15 mg/Kg/day of resveratrol
691 during 6 weeks (Gómez-Zorita et al., 2013).

692 Finally, a strengthening of “browning” and mitochondrial biogenesis has been observed by
693 increasing protein contents of UCP1, PRDM16, and Cytochrome C along with an increase of
694 AMPK in CD1 mice treated with 0.1% of resveratrol (Wang et al., 2015).

695 As for *in vitro* tests, the combination of resveratrol with other polyphenols, bioactives and
696 drugs has been object of certain studies (Table 5). Two works published by the same research
697 group have proved that the mixture of resveratrol (15 mg/Kg/day) and quercetin (30
698 mg/Kg/day) for a 6 weeks period of time decreases the weight of liver and all the fats depots.
699 This effect has been related with the reduction of LPL activity and ACC and the increase of
700 ATGL and CPT-1 (Arias et al., 2015; Arias et al., 2016). However, non-synergistic effects
701 have been found when resveratrol is combined with conjugated linoleic acid (CLA) (Arias et
702 al., 2014). Melatonin is a neurohormone related with the circadian rhythms but also a
703 bioactive found ubiquously in several foods and also in wine (Hornedo-Ortega et al., 2016).
704 The supplementation of melatonin (3 mg/kg/day) with resveratrol in ovariectomized rats
705 reduced the body weight by 16% and body mass index (BMI) by 19%. Moreover, this
706 combination is able to reduce the insulin resistance and macrophage infiltration in liver
707 (Majumdar et al., 2014).

708 With regard to other stilbenes, some *in vivo* studies pointed out that piceatannol, pterostilbene
709 and oxyresveratrol are interesting molecules to combat obesity (Table 5). Starting with the
710 first one and by utilizing *C. elegans* as *in vivo* model, authors demonstrated that this
711 compound reduced the fat accumulation induced by high glucose conditions; an attenuation of
712 SBP-1 (encode SREBP-1c) and FAS and a reduction on HOSL-1 expression (encodes HSL)
713 prove that piceatannol can prevent the lipid synthesis and stimulate the lipolysis (Shen et al.,
714 2017). Furthermore, Tung et al. showed that piceatannol (0.1 and 0.25%; 18 weeks) can
715 decrease the C/EBP α , PPAR γ , FAS and CPT-1 and consequently promote the mitochondrial

716 FA, oxidation and lipid accumulation in adipocytes and liver. As was displayed for
717 resveratrol, this compound can also alter the composition of gut microbiota specifically by
718 increasing of *Firmicutes/Lactobacillus* and decreasing *Bacteroidetes* (Tung et al., 2016).

719 Pterostilbene at low doses (15 and 30 mg/Kg/day) can interfere in *de novo* lipogenesis at
720 adipose and hepatic level by reducing the activity of: ME, FAS, ACC, G6PDH, CPT-1, which
721 is in part explained by the increase of *p*-AMPK levels (Gómez-Zorita et al., 2014). This paper
722 highlighted that pterostilbene was more efficient than resveratrol at a dose of 15 mg/kg/day
723 while at 30 mg/kg/day both stilbenes had similar responses. This fact can be explained, by the
724 higher bioavailability of pterostilbene (Gómez-Zorita et al., 2014). Using the same doses of
725 pterostilbene, a thermogenic and oxidative capacity by increasing of brown adipose tissue
726 markers (UCP-1, CPT-1b, nuclear respiratory factor 1 (Nrf1), cyclooxygenase 2 (COX2)) was
727 observed (Aguirre et al., 2016).

728 Finally, an improvement of insulin resistance and hyperglycemia accompanied by the
729 reduction in body weight (up to 26%), liver weight (up to 28%), and visceral fat (up to 51%)
730 was observed for oxyresveratrol in C57BL/6 mouse experiments. The down regulation of
731 G6PDH expression was interpreted as the repression of free glucose production in liver. They
732 also observed an increase of GLUT4 and IRS1 that also plays a vital role in intracellular
733 glucose uptake (Tan et al., 2017).

734 In summary, there is seemingly compelling *in vivo* evidence that demonstrate that overall
735 resveratrol but also other stilbenes possess anti-adipogenic and anti-lipogenic effects with
736 SIRT and AMPK as implicated molecular pathways. In addition, a significant reduction on
737 body and adipose tissue weight, accompanied with a reduction of cholesterol, TG and glucose
738 blood levels has been confirmed. The enhancement of the browning process or mitochondrial
739 biogenesis and the possible control of the microbiota dysbiosis displayed by resveratrol is

740 gaining attention as promising molecular approaches to combat obesity even though more
741 studies are still necessary.

742 An important concern that should be taken in account after examined the *in vivo* animal anti-
743 obesity effects of resveratrol and other stilbenes is the quantity that should be consumed to
744 reach the bioactivity. In fact and although wine and grapes are the richest sources of stilbenes
745 in diet, the required quantities cannot be reached with a normal diet. If we take into account
746 only the consumption of red grapes and wine (resveratrol mean of 0.67 mg/kg and 1.41 mg/L,
747 respectively) more than one hundred kg of red grapes or more than fifty liters of wine per day
748 should be consumed. It should not be forgotten that the high consumption of grapes and wine
749 can be exacerbates the obesity and the related disorders due to the high content of sugar or
750 ethanol. Consequently, it is therefore essential to turn to food supplements in order to achieve
751 these concentrations. One of the objectives of this review has been to examine the literature
752 about the anti-obesity effects of resveratrol in combination with other polyphenols, nutrients
753 or other bioactives. Several of these such as quercetin, CLA or melatonin are ubiquitously
754 present in food and consequently, the potential effects of resveratrol and other stilbenes can
755 be improved. Moreover, this experimental approach represents undoubtedly a more realistic
756 situation.

757 Even if it is not one of the goals of this work it can be mentioned that until recently there was
758 a considerable controversy about the effects of resveratrol as anti-obesity agent proved by
759 human intervention studies because the majority of these studies had not noted any effect of
760 resveratrol concerning body weight or body composition. Multiple reasons might explain the
761 lack of link between animal and human studies. For example, in human studies participants
762 consume generally their usual diet and not a high-fat or high-calorie diet in contrast to the
763 animal studies. Concerning doses, in human trials a relatively low dose of resveratrol (75-

764 3000 mg/day) have been used, however at these doses in animals no effect is observed.
765 Another fact that should not be overlooked is that the digestion, metabolism or absorption
766 differs between humans and animals. Furthermore, the duration of the resveratrol
767 supplementation could also have been too short in the human intervention studies in
768 comparison with animal studies. However, two recent systematic reviews and meta-analysis
769 has concluded by using random-effects model that resveratrol supplementation significantly
770 decrease body weight, body mass index, fat mass and waist circumference in trials using
771 resveratrol at the dosage under 500 mg/day, those with long-term interventions (≥ 3 month),
772 and performed on people with obesity (Akbari et al., 2020; Mousavi et al., 2019).

773 **Challenges and future trends**

774 Both, the analytical determination of grapes and wine monomeric/oligomeric stilbenes and,
775 the study on their biological activities represent nowadays a major challenge for the research
776 community. Consideration must be given concerning the deep matrix complexity of grapes
777 and wine, which makes the stilbene analysis more exigent. Thanks to the widespread
778 development on sample preparation, extraction techniques and detection instrumentation such
779 as mass spectrometry, more accurate stilbene identification is possible nowadays. However, it
780 should be kept in mind that a generalized problem is the lacks of commercial standards
781 (overall of oligomers) that force to express the results in terms of monomers (resveratrol or
782 piceid equivalent). This lack of precision could severally underestimate the oligomeric
783 stilbene content.

784 Regarding anti-obesity properties of grapes and wine stilbenes, a great number of *in vitro* and
785 *in vivo* studies have allowed to demonstrate the positive implications of these bioactives and
786 the underlying mechanisms of the observed effects. The understanding of these mechanisms is
787 an essential task since the scientists seeks the development of more specific dietary

788 recommendations to prevent or combat obesity. Apart from resveratrol, other stilbenes
789 (pterostilbene, oxyresveratrol, viniferins and vitisins) and the combination of these bioactives
790 with others polyphenols or drugs, are showing promising results. However, more studies are
791 necessary to allow reliable conclusions. To finish, it is important to emphasize that a great
792 number of published papers use in general high doses, and without reflecting important
793 processes such as absorption and metabolism or even the role of gut microbiota. Furthermore,
794 intervention studies are still necessary in order to prove these beneficial effects in humans.

795 **References**

- 796 Adrian, M., Jeandet, P., Douillet-Breuil, A.C., Tesson, L., Bessis, R. (2000). Stilbene content
797 of mature *Vitis vinifera* berries in response to UV-C elicitation. *Journal of Agricultural and*
798 *Food Chemistry* 48(12), 6103-6105.
- 799 Aguirre, L., Milton-Laskibar, I., Hijona, E., Bujanda, L., Rimando, A.M., Portillo, M.P.
800 (2016). Effects of pterostilbene in brown adipose tissue from obese rats. *Journal of*
801 *Physiology and Biochemistry* 73(3), 457-464.
- 802 Ahmed, B., Liu, S., Si, H. (2017). Antiadipogenic Effects and Mechanisms of Combinations
803 of Genistein, Epigallocatechin-3-Gallate, and/or Resveratrol in Preadipocytes. *Journal of*
804 *Medicinal Food* 20(2), 162-170.
- 805 Ahn, J., Cho, I., Kim, S., Kwon, D., Ha, T. (2008). Dietary resveratrol alters lipid
806 metabolism-related gene expression of mice on an atherogenic diet. *Journal of Hepatology*
807 49(6), 1019-1028.
- 808 Akbari, M., Tamtaji, O.R., Lankarani, K.B., Tabrizi, R., Dadgostar, E., Haghghat, N.,
809 Kolahdooz, F., Ghaderi, A., Mansournia, M.A., Asemi, Z. (2020). The effects of resveratrol
810 on lipid profiles and liver enzymes in patients with metabolic syndrome and related disorders:
811 a systematic review and meta-analysis of randomized controlled trials. *Lipids in Health and*
812 *Disease* 19(1), 25.
- 813 Alberdi, G., Rodríguez, V.M., Macarulla, M.T., Miranda, J., Churrua, I., Portillo, M.P.
814 (2013). Hepatic lipid metabolic pathways modified by resveratrol in rats fed an obesogenic
815 diet. *Nutrition* 29(3), 562-567.
- 816 Amira-Guebailia, H., Valls, J., Richard, T., Vitrac, X., Monti, J.P., Delaunay, J.C., Mérillon,
817 J.M. (2009). Centrifugal partition chromatography followed by HPLC for the isolation of cis-
818 ϵ -viniferin, a resveratrol dimer newly extracted from a red Algerian wine. *Food Chemistry*
819 113(1), 320-324.
- 820 Andrade, J.M.O., Paraíso, A.F., de Oliveira, M.V.M., Martins, A.M.E., Neto, J.F., Guimarães,
821 A.L.S., de Paula, A.M., Qureshi, M., Santos, S.H.S. (2014). Resveratrol attenuates hepatic
822 steatosis in high-fat fed mice by decreasing lipogenesis and inflammation. *Nutrition* 30(7-8),
823 915-919.
- 824 Arias, N., Macarulla, M.T., Aguirre, L., Milton, I., Portillo, M.P. (2016). The combination of
825 resveratrol and quercetin enhances the individual effects of these molecules on triacylglycerol
826 metabolism in white adipose tissue. *European Journal of Nutrition* 55(1), 341-348.

827 Arias, N., Macarulla, M.T., Aguirre, L., Miranda, J., Portillo, M.P. (2015). Liver delipidating
828 effect of a combination of resveratrol and quercetin in rats fed an obesogenic diet. *Journal of*
829 *Physiology and Biochemistry* 71(3), 569-576.

830 Arias, N., Miranda, J., Macarulla, M.T., Aguirre, L., Fernández-Quintela, A., Andres-
831 Lacueva, C., Urpi-Sarda, M., Portillo, M.P. (2014). The combination of resveratrol and
832 conjugated linoleic acid attenuates the individual effects of these molecules on triacylglycerol
833 metabolism in adipose tissue. *European Journal of Nutrition* 53(2), 575-582.

834 Arraki, K., Renouf, E., Waffo Teguo, P., Merillon, J.M., Richard, T., Decendit, A. (2017).
835 Identification and quantification of stilbenes in some Tunisian red wines using UPLC-MS and
836 HPLC-DAD. *OENO One* 51(2-3), 231-236.

837 Babazadeh, A., Taghvimi, A., Hamishehkar, H., Tabibiazar, M. (2017). Development of new
838 ultrasonic-solvent assisted method for determination of trans-resveratrol from red grapes:
839 Optimization, characterization, and antioxidant activity (ORAC assay). *Food Bioscience* 20,
840 36-42.

841 Bäckhed, F., Ding, H., Wang, T., Hooper, L.V., Gou, Y.K., Nagy, A., Semenkovich, C.F.,
842 Gordon, J.I. (2004). The gut microbiota as an environmental factor that regulates fat storage.
843 *Proceedings of the National Academy of Sciences of the United States of America* 101(44),
844 15718-15723.

845 Baderschneider, B., Winterhalter, P. (2000). Isolation and characterization of novel stilbene
846 derivatives from Riesling wine. *Journal of Agricultural and Food Chemistry* 48(7), 2681-
847 2686.

848 Barbanti, D., Galassi, S., Versari, A., Burattini, R. (1996). Gas chromatography and high-
849 pressure liquid chromatography determination of resveratrol in Italian red wines. *International*
850 *Journal of Phytoremediation* 21(1), 5-11.

851 Bartelt, A., Heeren, J. (2014). Adipose tissue browning and metabolic health. *Nature Reviews*
852 *Endocrinology* 10(1), 24-36.

853 Bavaresco, L., Fregoni, M., Trevisan, M., Mattivi, F., Vrhovsek, U., Falchetti, R. (2002). The
854 occurrence of the stilbene piceatannol in grapes. *Vitis* 41(3), 133-136.

855 Bavaresco, L., Mattivi, F., de Rosso, M., Flamini, R. (2012). Effects of elicitors, viticultural
856 factors, and enological practices on resveratrol and stilbenes in Grapevine and Wine. *Mini-*
857 *Reviews in Medicinal Chemistry* 12(13), 1366-1381.

858 Biais, B., Krisa, S., Cluzet, S., Da Costa, G., Waffo-Teguo, P., Méillon, J.M., Richard, T.
859 (2017). Antioxidant and cytoprotective activities of grapevine stilbenes. *Journal of*
860 *Agricultural and Food Chemistry* 65(24), 4952-4960.

861 Błaszczuk, A., Sady, S., Sielicka, M. (2019). The stilbene profile in edible berries.
862 *Phytochemistry Reviews* 18(1), 37-67.

863 Brillante, L., De Rosso, M., Dalla Vedova, A., Maoz, I., Flamini, R., Tomasi, D. (2018).
864 Insights on the stilbenes in Raboso Piave grape (*Vitis vinifera* L.) as a consequence of
865 postharvest vs on-vine dehydration. *Journal of the Science of Food and Agriculture* 98(5),
866 1961-1967.

867 Buiarelli, F., Coccioli, F., Jasionowska, R., Merolle, M., Terracciano, A. (2007). Analysis of
868 some stilbenes in Italian wines by liquid chromatography/tandem mass spectrometry. *Rapid*
869 *Communications in Mass Spectrometry* 21(18), 2955-2964.

870 Cantos, E., Espín, J.C., Fernández, M.J., Oliva, J., Tomás-Barberán, F.A. (2003). Postharvest
871 UV-C-Irradiated Grapes as a Potential Source for Producing Stilbene-Enriched Red Wines.
872 *Journal of Agricultural and Food Chemistry* 51(5), 1208-1214.

873 Careri, M., Corradini, C., Elviri, L., Nicoletti, I., Zagnoni, I. (2003). Direct HPLC analysis of
874 quercetin and trans-resveratrol in red wine, grape, and winemaking byproducts. *Journal of*
875 *Agricultural and Food Chemistry* 51(18), 5226-5231.

876 Careri, M., Corradini, C., Elviri, L., Nicoletti, I., Zagnoni, I. (2004). Liquid chromatography-
877 electrospray tandem mass spectrometry of cis-resveratrol and trans-resveratrol: Development,
878 validation, and application of the method to red wine, grape, and winemaking byproducts.
879 *Journal of Agricultural and Food Chemistry* 52(23), 6868-6874.

880 Chang, C.C., Lin, K.Y., Peng, K.Y., Day, Y.J., Hung, L.M. (2016). Resveratrol exerts anti-
881 obesity effects in high-fat diet obese mice and displays differential dosage effects on
882 cytotoxicity, differentiation, and lipolysis in 3T3-L1 cells. *Endocrine Journal* 63(2), 169-178.

883 Chen, S., Li, Z., Li, W., Shan, Z., Zhu, W. (2011). Resveratrol inhibits cell differentiation in
884 3T3-L1 adipocytes via activation of AMPK. *Canadian Journal of Physiology and*
885 *Pharmacology* 89(11), 793-799.

886 Cho, Y.J., Hong, J.Y., Chun, H.S., Lee, S.K., Min, H.Y. (2006). Ultrasonication-assisted
887 extraction of resveratrol from grapes. *Journal of Food Engineering* 77(3), 725-730.

888 Choi, Y.J., Suh, H.R., Yoon, Y., Lee, K.J., Kim, D.G., Kim, S., Lee, B.H. (2014). Protective
889 effect of resveratrol derivatives on high-fat diet induced fatty liver by activating AMP-
890 activated protein kinase. *Archives of Pharmacal Research* 37(9), 1169-1176.

891 Chong, J., Poutaraud, A., Hugueney, P. (2009). Metabolism and roles of stilbenes in plants.
892 *Plant Science* 177(3), 143-155.

893 Chou, Y.C., Ho, C.T., Pan, M.H. (2018). Stilbenes: Chemistry and Molecular Mechanisms of
894 Anti-obesity. *Current Pharmacology Reports* 4(3), 202-209.

895 Coelho, C., Bagala, F., Gougeon, R.D., Schmitt-Kopplin, P. (2016). Capillary electrophoresis
896 in wine science, *Methods in Molecular Biology*, pp. 509-523.

897 Costa, C.d.S., Rohden, F., Hammes, T.O., Margis, R., Bortolotto, J.W., Padoin, A.V., Mottin,
898 C.C., Guaragna, R.M. (2011). Resveratrol Upregulated SIRT1, FOXO1, and Adiponectin and
899 Downregulated PPAR γ 1-3 mRNA Expression in Human Visceral Adipocytes. *Obesity*
900 *Surgery* 21(3), 356-361.

901 De Almeida Pinheiro, T., De Almeida Pinheiro, T., Feltenberger, J.D., Andrade, J.M.O.,
902 Ferreira, E.C.N., De Farias Lelis, D., Guimarães, A.L.S., De Paula, A.M.B., Caldeira, A.P.,
903 Santos, S.H.S. (2017). Effects of resveratrol and ACE inhibitor enalapril on glucose and lipid
904 profiles in mice. *Protein and Peptide Letters* 24(9), 854-860.

905 Deluc, L.G., Grimplet, J., Wheatley, M.D., Tillett, R.L., Quilici, D.R., Osborne, C., Schooley,
906 D.A., Schlauch, K.A., Cushman, J.C., Cramer, G.R. (2007). Transcriptomic and metabolite
907 analyses of Cabernet Sauvignon grape berry development. *BMC Genomics* 8.

908 Duncan, R.E., Ahmadian, M., Jaworski, K., Sarkadi-Nagy, E., Sul, H.S. (2007). Regulation of
909 lipolysis in adipocytes, *Annual Review of Nutrition*, pp. 79-101.

910 El Khawand, T., Valls Fonayet, J., Da Costa, G., Hornedo-Ortega, R., Jourdes, M., Franc, C.,
911 de Revel, G., Decendit, A., Krisa, S., Richard, T. (2020). Resveratrol transformation in red
912 wine after heat treatment. *Food Research International* 132.

913 Eseberri, I., Lasa, A., Miranda, J., Gracia, A., Portillo, M.P. (2017). Potential miRNA
914 involvement in the anti-adipogenic effect of resveratrol and its metabolites. *PloS One* 12(9).

915 Esteve Ràfols, M. (2014). Adipose tissue: Cell heterogeneity and functional diversity.
916 *Endocrinologia y Nutricion* 61(2), 100-112.

917 Fabjanowicz, M., Płotka-Wasyłka, J., Namieśnik, J. (2018). Detection, identification and
918 determination of resveratrol in wine. Problems and challenges. *TRAC Trends in Analytical*
919 *Chemistry* 103, 21-33.

920 Favre, G., Piccardo, D., Sergio, G.-A., Pérez-Navarro, J., García-Romero, E., Mena-Morales,
921 A., González-Neves, G. (2020). Stilbenes in grapes and wines of Tannat, Marselan and Syrah
922 from Uruguay. *OENO One* 54(1), 27-36.

923 Fernández-Marín, M.I., Guerrero, R.F., García-Parrilla, M.C., Puertas, B., Ramírez, P.,
924 Cantos-Villar, E. (2013). Terroir and variety: Two key factors for obtaining stilbene-enriched
925 grapes. *Journal of Food Composition and Analysis* 31(2), 191-198.

926 Fernandez-Marin, M.I., Guerrero, R.F., Garcia-Parrilla, M.C., Puertas, B., Richard, T.,
927 Rodriguez-Werner, M.A., Winterhalter, P., Monti, J.P., Cantos-Villar, E. (2012).
928 Isorhapontigenin: A novel bioactive stilbene from wine grapes. *Food Chemistry* 135(3), 1353-
929 1359.

930 Fernández-Marín, M.I., Guerrero, R.F., García-Parrilla, M.C., Puertas, B., Richard, T.,
931 Rodríguez-Werner, M.A., Winterhalter, P., Monti, J.P., Cantos-Villar, E. (2012).
932 Isorhapontigenin: A novel bioactive stilbene from wine grapes. *Food Chemistry* 135(3), 1353-
933 1359.

934 Fernández-Marín, M.I., Puertas, B., Guerrero, R.F., García-Parrilla, M.C., Cantos-Villar, E.
935 (2014). Preharvest Methyl Jasmonate and Postharvest UVC Treatments: Increasing Stilbenes
936 in Wine. *Journal of Food Science* 79(3), C310-C317.

937 Fernández-Quintela, A., Milton-Laskibar, I., González, M., Portillo, M.P. (2017). Antiobesity
938 effects of resveratrol: Which tissues are involved?, *Annals of the New York Academy of*
939 *Sciences*, pp. 118-131.

940 Festa, A., D'Agostino Jr, R., Williams, K., Karter, A.J., Mayer-Davis, E.J., Tracy, R.P.,
941 Haffner, S.M. (2001). The relation of body fat mass and distribution to markers of chronic
942 inflammation. *International Journal of Obesity* 25(10), 1407-1415.

943 Fischer-Posovszky, P., Kukulius, V., Tews, D., Unterkircher, T., Debatin, K.-M., Fulda, S.,
944 Wabitsch, M. (2010). Resveratrol regulates human adipocyte number and function in a Sirt1-
945 dependent manner. *The American Journal of Clinical Nutrition* 92(1), 5-15.

946 Flamini, R., De Rosso, M., De Marchi, F., Dalla Vedova, A., Panighel, A., Gardiman, M.,
947 Maoz, I., Bavaresco, L. (2013). An innovative approach to grape metabolomics: stilbene
948 profiling by suspect screening analysis. *Metabolomics* 9(6), 1243-1253.

949 Flamini, R., Zanzotto, A., de Rosso, M., Lucchetta, G., Vedova, A.D., Bavaresco, L. (2016).
950 Stilbene oligomer phytoalexins in grape as a response to *Aspergillus carbonarius* infection.
951 *Physiological and Molecular Plant Pathology* 93, 112-118.

952 Gabaston, J., Valls Fonayet, J., Franc, C., Waffo-Teguo, P., de Revel, G., Hilbert, G., Gomes,
953 E., Richard, T., Mérillon, J.-M. (2020). Characterization of Stilbene Composition in Grape
954 Berries from Wild *Vitis* Species in Year-To-Year Harvest. *Journal of Agricultural and Food*
955 *Chemistry* in press.

956 Gallagher, E.J., LeRoith, D. (2015). Obesity and Diabetes: The Increased Risk of Cancer and
957 Cancer-Related Mortality. *Physiological Reviews* 95(3), 727-748.

958 Gambini, J., Inglés, M., Olaso, G., Lopez-Grueso, R., Bonet-Costa, V., Gimeno-Mallench, L.,
959 Mas-Bargues, C., Abdelaziz, K. M., Gomez-Cabrera, M. C., Vina, J., & Borrás, C. (2015).
960 Properties of Resveratrol: In Vitro and In Vivo Studies about Metabolism, Bioavailability,
961 and Biological Effects in Animal Models and Humans. *Oxidative Medicine and Cellular*
962 *Longevity*, 2015, 837042.

963 Garin-Shkolnik, T., Rudich, A., Hotamisligil, G.S., Rubinstein, M. (2014). FABP4 attenuates
964 PPAR γ and adipogenesis and is inversely correlated with PPAR γ in adipose tissues. *Diabetes*
965 63(3), 900-911.

966 Gatto, P., Vrhovsek, U., Muth, J., Segala, C., Romualdi, C., Fontana, P., Pruefer, D.,
967 Stefanini, M., Moser, C., Mattivi, F., Velasco, R. (2008). Ripening and genotype control
968 stilbene accumulation in healthy grapes. *Journal of Agricultural and Food Chemistry* 56(24),
969 11773-11785.

970 Gaudette, N.J., Pickering, G.J. (2011). Sensory and chemical characteristics of trans-
971 resveratrol-fortified wine. *Australian Journal of Grape and Wine Research* 17(2), 249-257.

972 Gnoni, G.V., Paglialonga, G. (2009). Resveratrol inhibits fatty acid and triacylglycerol
973 synthesis in rat hepatocytes. *European Journal of Clinical Investigation* 39(3), 211-218.

974 Gomez-Zorita, S., Belles, C., Briot, A., Fernández-Quintela, A., Portillo, M.P., Carpené, C.
975 (2017). Pterostilbene Inhibits Lipogenic Activity similar to Resveratrol or Caffeine but
976 Differently Modulates Lipolysis in Adipocytes. *Phytotherapy Research* 31(8), 1273-1282.

977 Gómez-Zorita, S., Fernández-Quintela, A., Lasa, A., Aguirre, L., Rimando, A.M., Portillo,
978 M.P. (2014). Pterostilbene, a dimethyl ether derivative of resveratrol, reduces fat
979 accumulation in rats fed an obesogenic diet. *Journal of Agricultural and Food Chemistry*
980 62(33), 8371-8378.

981 Gómez-Zorita, S., Fernández-Quintela, A., Lasa, A., Hijona, E., Bujanda, L., Portillo, M.P.
982 (2013). Effects of resveratrol on obesity-related inflammation markers in adipose tissue of
983 genetically obese rats. *Nutrition* 29(11-12), 1374-1380.

984 Gómez-Zorita, S., Fernández-Quintela, A., MacArulla, M.T., Aguirre, L., Hijona, E.,
985 Bujanda, L., Milagro, F., Martínez, J.A., Portillo, M.P. (2012). Resveratrol attenuates
986 steatosis in obese Zucker rats by decreasing fatty acid availability and reducing oxidative
987 stress. *British Journal of Nutrition* 107(2), 202-210.

988 Gonçalves, J., Câmara, J.S. (2011). New method for determination of (E)-resveratrol in wine
989 based on microextraction using packed sorbent and ultra-performance liquid chromatography.
990 *Journal of Separation Science* 34(18), 2376-2384.

991 González-Candelas, L., Gil, J.V., Lamuela-Raventós, R.M., Ramón, D. (2000). The use of
992 transgenic yeasts expressing a gene encoding a glycosyl-hydrolase as a tool to increase
993 resveratrol content in wine. *International Journal of Food Microbiology* 59(3), 179-183.

994 Gracia, A., Miranda, J., Fernández-Quintela, A., Eseberri, I., Garcia-Lacarte, M., Milagro,
995 F.I., Martínez, J.A., Aguirre, L., Portillo, M.P. (2016). Involvement of miR-539-5p in the
996 inhibition of de novo lipogenesis induced by resveratrol in white adipose tissue. *Food and*
997 *Function* 7(3), 1680-1688.

998 Guebailia, H.A., Chira, K., Richard, T., Mabrouk, T., Furiga, A., Vitrac, X., Monti, J.P.,
999 Delaunay, J.C., Merillon, J.M. (2006). Hopeaphenol: The first resveratrol tetramer in wines
1000 from North Africa. *Journal of Agricultural and Food Chemistry* 54(25), 9559-9564.

1001 Guerrero, R.F., Cantos-Villar, E., Puertas, B., Richard, T. (2016). Daily Preharvest UV-C
1002 Light Maintains the High Stilbenoid Concentration in Grapes. *Journal of Agricultural and*
1003 *Food Chemistry* 64(25), 5139-5147.

1004 Guerrero, R.F., García-Parrilla, M.C., Puertas, B., Cantos-Villar, E. (2009). Wine, resveratrol
1005 and health: A review. *Natural Product Communications* 4(5), 635-658.

1006 Guerrero, R.F., Puertas, B., Fernández, M.I., Palma, M., Cantos-Villar, E. (2010). Induction
1007 of stilbenes in grapes by UV-C: Comparison of different subspecies of *Vitis*. *Innovative Food*
1008 *Science and Emerging Technologies* 11(1), 231-238.

1009 Guerrero, R.F., Valls-Fonayet, J., Richard, T., Cantos-Villar, E. (2020). A rapid quantification
1010 of stilbene content in wine by ultra-high pressure liquid chromatography – Mass
1011 spectrometry. *Food Control* 108.

1012 Hasan, M.M., Bae, H. (2017). An overview of stress-induced resveratrol synthesis in grapes:
1013 Perspectives for resveratrol-enriched grape products. *Molecules* 22(2).

1014 Hausman, D.B., DiGirolamo, M., Bartness, T.J., Hausman, G.J., Martin, R.J. (2001). The
1015 biology of white adipocyte proliferation. *Obesity Reviews* 2(4), 239-254.

1016 He, S., Jiang, L., Wu, B., Li, C., Pan, Y. (2009). Chunganenol: An Unusual Antioxidative
1017 Resveratrol Hexamer from *Vitis chunganensis*. *The Journal of Organic Chemistry* 74(20),
1018 7966-7969.

1019 Hilbert, G., Tamsamani, H., Bordenave, L., Pedrot, E., Chaher, N., Cluzet, S., Delaunay, J.C.,
1020 Ollat, N., Delrot, S., Merillon, J.M., Gomès, E., Richard, T. (2015). Flavonol profiles in
1021 berries of wild *Vitis* accessions using liquid chromatography coupled to mass spectrometry
1022 and nuclear magnetic resonance spectrometry. *Food Chemistry* 169, 49-58.

1023 Hornedo-Ortega, R., Cerezo, A.B., Troncoso, A.M., Garcia-Parrilla, M.C., Mas, A. (2016).
1024 Melatonin and other tryptophan metabolites produced by yeasts: Implications in
1025 cardiovascular and neurodegenerative diseases. *Frontiers in Microbiology* 6(JAN).
1026 Hsu, C.L., Lin, Y.J., Ho, C.T., Yen, G.C. (2012). Inhibitory effects of garcinol and
1027 pterostilbene on cell proliferation and adipogenesis in 3T3-L1 cells. *Food and Function* 3(1),
1028 49-57.
1029 Hu, P., Zhao, L., Chen, J. (2015). Physiologically achievable doses of resveratrol enhance
1030 3T3-L1 adipocyte differentiation. *European Journal of Nutrition* 54(4), 569-579.
1031 Jeandet, P., Bessis, R., Gautheron, B. (1991). The Production of Resveratrol (3,5,4'-
1032 trihydroxystilbene) by Grape Berries in Different Developmental Stages. *American Journal of*
1033 *Enology and Viticulture* 42(1), 41-46.
1034 Jeandet, P., Bessis, R., Maume, B.F., Meunier, P., Peyron, D., Trollat, P. (1995). Effect of
1035 Enological Practices on the Resveratrol Isomer Content of Wine. *Journal of Agricultural and*
1036 *Food Chemistry* 43(2), 316-319.
1037 Jeandet, P., Bessis, R., Sbaghi, M., Meunier, P. (1994). Occurrence of a resveratrol β -D-
1038 glucoside in wine: preliminary studies. *Vitis - Journal of Grapevine Research* 33(3), 183-184.
1039 Jeon, S.M., Lee, S.A., Choi, M.S. (2014). Antiobesity and vasoprotective effects of
1040 resveratrol in ApoE-deficient mice. *Journal of Medicinal Food* 17(3), 310-316.
1041 Jiang, L., He, S., Sun, C., Pan, Y. (2012). Selective 1O₂ quenchers, oligostilbenes, from *Vitis*
1042 *wilsonae*: Structural identification and biogenetic relationship. *Phytochemistry* 77, 294-303.
1043 Jin, S.H., Yang, J.H., Shin, B.Y., Seo, K., Shin, S.M., Cho, I.J., Ki, S.H. (2013). Resveratrol
1044 inhibits LXR α -dependent hepatic lipogenesis through novel antioxidant Sestrin2 gene
1045 induction. *Toxicology and Applied Pharmacology* 271(1), 95-105.
1046 Kang, H.K., Min, J.S., Chung, J., Park, H., Jae, B.K. (2005). Hypoxia inhibits adipocyte
1047 differentiation in a HDAC-independent manner. *Biochemical and Biophysical Research*
1048 *Communications* 333(4), 1178-1184.
1049 Kasiotis, K.M., Pratsinis, H., Kletsas, D., Haroutounian, S.A. (2013). Resveratrol and related
1050 stilbenes: Their anti-aging and anti-angiogenic properties. *Food and Chemical Toxicology* 61,
1051 112-120.
1052 Khaleel, E.F., Abdel-Aleem, G.A., Mostafa, D.G. (2018). Resveratrol improves high-fat diet
1053 induced fatty liver and insulin resistance by concomitantly inhibiting proteolytic cleavage of
1054 sterol regulatory element-binding proteins, free fatty acid oxidation, and intestinal triglyceride
1055 absorption. *Canadian Journal of Physiology and Pharmacology* 96(2), 145-157.
1056 Kim, A., Park, T. (2010). Diet-induced obesity regulates the galanin-mediated signaling
1057 cascade in the adipose tissue of mice. *Molecular Nutrition and Food Research* 54(9), 1361-
1058 1370.
1059 Kim, S., Jin, Y., Choi, Y., Park, T. (2011). Resveratrol exerts anti-obesity effects via
1060 mechanisms involving down-regulation of adipogenic and inflammatory processes in mice.
1061 *Biochemical Pharmacology* 81(11), 1343-1351.
1062 Kim, S.h., Park, H.S., Lee, M.s., Cho, Y.J., Kim, Y.S., Hwang, J.T., Sung, M.J., Kim, M.S.,
1063 Kwon, D.Y. (2008). Vitisin A inhibits adipocyte differentiation through cell cycle arrest in
1064 3T3-L1 cells. *Biochemical and Biophysical Research Communications* 372(1), 108-113.
1065 Kolouchová-Hanzlíková, I., Melzoch, K., Filip, V., Šmidrkal, J. (2004). Rapid method for
1066 resveratrol determination by HPLC with electrochemical and UV detections in wines. *Food*
1067 *Chemistry* 87(1), 151-158.
1068 Kostadinović, S., Wilkens, A., Stefova, M., Ivanova, V., Vojnoski, B., Mirhosseini, H.,
1069 Winterhalter, P. (2012). Stilbene levels and antioxidant activity of Vranec and Merlot wines
1070 from Macedonia: Effect of variety and enological practices. *Food Chemistry* 135(4), 3003-
1071 3009.

1072 Krisa, S., Larronde, F., Budzinski, H., Decendit, A., Deffieux, G., Mérillon, J.-M. (1999).
1073 Stilbene Production by *Vitis vinifera* Cell Suspension Cultures: Methyl Jasmonate Induction
1074 and ¹³C Biolabeling. *Journal of Natural Products* 62(12), 1688-1690.

1075 Kwon, J.Y., Seo, S.G., Heo, Y.S., Yue, S., Cheng, J.X., Lee, K.W., Kim, K.H. (2012a).
1076 Piceatannol, natural polyphenolic stilbene, inhibits adipogenesis via modulation of mitotic
1077 clonal expansion and insulin receptor-dependent insulin signaling in early phase of
1078 differentiation. *Journal of Biological Chemistry* 287(14), 11566-11578.

1079 Kwon, J.Y., Seo, S.G., Yue, S., Cheng, J.X., Lee, K.W., Kim, K.H. (2012b). An inhibitory
1080 effect of resveratrol in the mitotic clonal expansion and insulin signaling pathway in the early
1081 phase of adipogenesis. *Nutrition Research* 32(8), 607-616.

1082 Lamuela-Raventós, R.M., Romero-Pérez, A.I., Waterhouse, A.L., de la Torre-Boronat, M.C.
1083 (1995). Direct HPLC Analysis of cis- and trans-Resveratrol and Piceid Isomers in Spanish
1084 Red *Vitis vinifera* Wines. *Journal of Agricultural and Food Chemistry* 43(2), 281-283.

1085 Landrault, N., Larronde, F., Delaunay, J.C., Castagnino, C., Vercauteren, J., Merillon, J.M.,
1086 Gasc, F., Cros, G., Teissedre, P.L. (2002). Levels of stilbene oligomers and astilbin in French
1087 varietal wines and in grapes during noble rot development. *Journal of Agricultural and Food*
1088 *Chemistry* 50(7), 2046-2052.

1089 Langcake, P., Pryce, R.J. (1977). A new class of phytoalexins from grapevines. *Experientia*
1090 33(2), 151-152.

1091 Lasa, A., Miranda, J., Churruga, I., Simón, E., Arias, N., Milagro, F., Martínez, J.A., del Puy
1092 Portillo, M. (2011). The combination of resveratrol and CLA does not increase the
1093 delipidating effect of each molecule in 3T3-L1 adipocytes. *Nutricion Hospitalaria* 26(5), 997-
1094 1003.

1095 Lasa, A., Schweiger, M., Kotzbeck, P., Churruga, I., Simón, E., Zechner, R., Portillo, M.d.P.
1096 (2012). Resveratrol regulates lipolysis via adipose triglyceride lipase. *The Journal of*
1097 *Nutritional Biochemistry* 23(4), 379-384.

1098 Lee, J., Rennaker, C. (2007). Antioxidant capacity and stilbene contents of wines produced in
1099 the Snake River Valley of Idaho. *Food Chemistry* 105(1), 195-203.

1100 Lee, J.H., Baek, S.Y., Jang, E.J., Ku, S.K., Kim, K.M., Ki, S.H., Kim, C.E., Park, K.I., Kim,
1101 S.C., Kim, Y.W. (2018). Oxyresveratrol ameliorates nonalcoholic fatty liver disease by
1102 regulating hepatic lipogenesis and fatty acid oxidation through liver kinase B1 and AMP-
1103 activated protein kinase. *Chemico-Biological Interactions* 289, 68-74.

1104 Ley, R.E., Turnbaugh, P.J., Klein, S., Gordon, J.I. (2006). Microbial ecology: Human gut
1105 microbes associated with obesity. *Nature* 444(7122), 1022-1023.

1106 Li, S., Bouzar, C., Cottet-Rousselle, C., Zagotta, I., Lamarche, F., Wabitsch, M., Tokarska-
1107 Schlattner, M., Fischer-Posovszky, P., Schlattner, U., Rousseau, D. (2016). Resveratrol
1108 inhibits lipogenesis of 3T3-L1 and SGBS cells by inhibition of insulin signaling and
1109 mitochondrial mass increase. *Biochimica et Biophysica Acta - Bioenergetics* 1857(6), 643-
1110 652.

1111 Li, Z.Z., Berk, M., McIntyre, T.M., Feldstein, A.E. (2009). Hepatic lipid partitioning and liver
1112 damage in nonalcoholic fatty liver disease: role of stearoyl-CoA desaturase. *The Journal of*
1113 *biological chemistry* 284(9), 5637-5644.

1114 Liang, Y., Tian, W., Ma, X. (2013). Inhibitory effects of grape skin extract and resveratrol on
1115 fatty acid synthase. *BMC Complementary and Alternative Medicine* 13.

1116 Liu, C., Wang, L., Wang, J., Wu, B., Liu, W., Fan, P., Liang, Z., Li, S. (2013). Resveratrols in
1117 *Vitis* berry skins and leaves: Their extraction and analysis by HPLC. *Food Chemistry* 136(2),
1118 643-649.

1119 Liu, X., Zhao, H., Jin, Q., You, W., Cheng, H., Liu, Y., Song, E., Liu, G., Tan, X., Zhang, X.,
1120 Wan, F. (2018). Resveratrol induces apoptosis and inhibits adipogenesis by stimulating the

1121 SIRT1-AMPK α -FOXO1 signalling pathway in bovine intramuscular adipocytes. *Molecular*
1122 *and Cellular Biochemistry* 439(1-2), 213-223.

1123 Lukić, I., Radeka, S., Budić-Leto, I., Bubola, M., Vrhovsek, U. (2019). Targeted UPLC-QqQ-
1124 MS/MS profiling of phenolic compounds for differentiation of monovarietal wines and
1125 corroboration of particular varietal typicity concepts. *Food Chemistry* 300, 125251.

1126 MacDougald, O.A., Mandrup, S. (2002). Adipogenesis: Forces that tip the scales. *Trends in*
1127 *Endocrinology and Metabolism* 13(1), 5-11.

1128 Madden, C.J. (2017). Brown fat in obesity: Uncoupling protein-1 versus thermogenic activity.
1129 *Temperature* 4(2), 126-127.

1130 Majumdar, A.S., Giri, P.R., Pai, S.A. (2014). Resveratrol- and melatonin-abated ovariectomy
1131 and fructose diet-induced obesity and metabolic alterations in female rats. *Menopause* 21(8),
1132 876-885.

1133 Mattivi, F., Reniero, F., Korhammer, S. (1995). Isolation, Characterization, and Evolution in
1134 Red Wine Vinification of Resveratrol Monomers. *Journal of Agricultural and Food Chemistry*
1135 43(7), 1820-1823.

1136 Mendes, K.L., De Pinho, L., Andrade, J.M.O., Paraíso, A.F., Lula, J.F., MacEdo, S.M.,
1137 Feltenberger, J.D., Guimarães, A.L.S., De Paula, A.M.B., Santos, S.H.S. (2016). Distinct
1138 metabolic effects of resveratrol on lipogenesis markers in mice adipose tissue treated with
1139 high-polyunsaturated fat and high-protein diets. *Life Sciences* 153, 66-73.

1140 Mercader, J., Palou, A., Bonet, M.L. (2011). Resveratrol enhances fatty acid oxidation
1141 capacity and reduces resistin and Retinol-Binding Protein 4 expression in white adipocytes.
1142 *Journal of Nutritional Biochemistry* 22(9), 828-834.

1143 Mitterberger, M.C., Zwerschke, W. (2013). Mechanisms of resveratrol-induced inhibition of
1144 clonal expansion and terminal adipogenic differentiation in 3T3-L1 preadipocytes. *Journals of*
1145 *Gerontology - Series A Biological Sciences and Medical Sciences* 68(11), 1356-1376.

1146 Monteiro, R., Azevedo, I. (2010). Chronic inflammation in obesity and the metabolic
1147 syndrome. *Mediators of Inflammation* 2010.

1148 Moss, R., Mao, Q., Taylor, D., Saucier, C. (2013). Investigation of monomeric and oligomeric
1149 wine stilbenoids in red wines by ultra-high-performance liquid chromatography/electrospray
1150 ionization quadrupole time-of-flight mass spectrometry. *Rapid Communications in Mass*
1151 *Spectrometry* 27(16), 1815-1827.

1152 Mousavi, S.M., Milajerdi, A., Sheikhi, A., Kord-Varkaneh, H., Feinle-Bisset, C., Larijani, B.,
1153 Esmailzadeh, A. (2019). Resveratrol supplementation significantly influences obesity
1154 measures: a systematic review and dose-response meta-analysis of randomized controlled
1155 trials. *Obesity Reviews* 20(3), 487-498.

1156 Nagao, K., Jinnouchi, T., Kai, S., Yanagita, T. (2017). Pterostilbene, a dimethylated analog of
1157 resveratrol, promotes energy metabolism in obese rats. *The Journal of Nutritional*
1158 *Biochemistry* 43, 151-155.

1159 Neveu, V., Perez-Jiménez, J., Vos, F., Crespy, V., du Chaffaut, L., Mennen, L., Knox, C.,
1160 Eisner, R., Cruz, J., Wishart, D., Scalbert, A. (2010). Phenol-Explorer: an online
1161 comprehensive database on polyphenol contents in foods. *Database* 2010, bap024.

1162 Ohara, K., Kusano, K., Kitao, S., Yanai, T., Takata, R., Kanauchi, O. (2015). ϵ -Viniferin, a
1163 resveratrol dimer, prevents diet-induced obesity in mice. *Biochemical and Biophysical*
1164 *Research Communications* 468(4), 877-882.

1165 Park, H.J., Yang, J.Y., Ambati, S., Della-Fera, M.A., Hausman, D.B., Rayalam, S., Baile,
1166 C.A. (2008). Combined effects of genistein, quercetin, and resveratrol in human and 3T3-L1
1167 adipocytes. *Journal of Medicinal Food* 11(4), 773-783.

1168 Pawlus, A., Cantos-Villar, E., Richard, T., Bisson, J., Poupard, P., Papastamoulis, Y., Monti,
1169 J.-P., Teissedre, P.L., Waffo-Teguo, P., Merillon, J.-M. (2013a). Chemical dereplication of

1170 wine stilbenoids using high performance liquid chromatography-nuclear magnetic resonance
1171 spectroscopy. *Journal of Chromatography A* 1289, 19 - 26.

1172 Pawlus, A.D., Sahli, R., Bisson, J., Rivière, C., Delaunay, J.C., Richard, T., Gomès, E.,
1173 Bordenave, L., Waffo-Téguo, P., Mérillon, J.M. (2013b). Stilbenoid profiles of canes from
1174 *Vitis* and *Muscadinia* species. *Journal of Agricultural and Food Chemistry* 61(3), 501-511.

1175 Pezet, R., Pont, V., Cuenat, P. (1994). Method to determine resveratrol and pterostilbene in
1176 grape berries and wines using high-performance liquid chromatography and highly sensitive
1177 fluorimetric detection. *Journal of Chromatography A* 663(2), 191-197.

1178 Picard, F., Kurtev, M., Chung, N., Topark-Ngarm, A., Senawong, T., De Oliveira, R.M., Leid,
1179 M., McBurney, M.W., Guarente, L. (2004). Sirt1 promotes fat mobilization in white
1180 adipocytes by repressing PPAR- γ . *Nature* 429(6993), 771-776.

1181 Poussier, M., Guilloux-Benatier, M., Torres, M., Heras, E., Adrian, M. (2003). Influence of
1182 Different Maceration Techniques and Microbial Enzymatic Activities on Wine Stilbene
1183 Content. *American Journal of Enology and Viticulture* 54(4), 261-266.

1184 Pugajeva, I., Perkons, I., Górnas, P. (2018). Identification and determination of stilbenes by
1185 Q-TOF in grape skins, seeds, juice and stems. *Journal of Food Composition and Analysis* 74,
1186 44-52.

1187 Qiao, Y., Sun, J., Xia, S., Tang, X., Shi, Y., Le, G. (2014). Effects of resveratrol on gut
1188 microbiota and fat storage in a mouse model with high-fat-induced obesity. *Food and*
1189 *Function* 5(6), 1241-1249.

1190 Rabesiaka, M., Rakotondramasy-Rabesiaka, L., Mabilie, I., Porte, C., Havet, J.-L. (2011).
1191 Extraction of trans-resveratrol from red wine and optimization by response surface
1192 methodology. *Separation and Purification Technology* 81(1), 56-61.

1193 Rayalam, S., Della-Fera, M.A., Yang, J.Y., Hea, J.P., Ambati, S., Baile, C.A. (2007).
1194 Resveratrol potentiates genistein's antiadipogenic and proapoptotic effects in 3T3-L1
1195 adipocytes. *Journal of Nutrition* 137(12), 2668-2673.

1196 Rayalam, S., Yang, J.Y., Ambati, S., Della-Fera, M.A., Baile, C.A. (2008). Resveratrol
1197 induces apoptosis and inhibits adipogenesis in 3T3-L1 adipocytes. *Phytotherapy Research*
1198 22(10), 1367-1371.

1199 Ribeiro De Lima, M.T., Waffo-Téguo, P., Teissedre, P.L., Pujolas, A., Vercauteren, J.,
1200 Cabanis, J.C., Mérillon, J.M. (1999). Determination of stilbenes (trans-astringin, cis- and
1201 trans-piceid, and cis- and trans-resveratrol) in Portuguese wines. *Journal of Agricultural and*
1202 *Food Chemistry* 47(7), 2666-2670.

1203 Rivière, C., Pawlus, A.D., Mérillon, J.M. (2012). Natural stilbenoids: distribution in the plant
1204 kingdom and chemotaxonomic interest in Vitaceae. *Natural Product Reports* 29(11), 1317-
1205 1333.

1206 Rodríguez-Cabo, T., Rodríguez, I., Cela, R. (2012). Determination of hydroxylated stilbenes
1207 in wine by dispersive liquid-liquid microextraction followed by gas chromatography mass
1208 spectrometry. *Journal of Chromatography A* 1258, 21-29.

1209 Rodríguez-Cabo, T., Rodríguez, I., Ramil, M., Silva, A., Cela, R. (2016). Multiclass semi-
1210 volatile compounds determination in wine by gas chromatography accurate time-of-flight
1211 mass spectrometry. *Journal of Chromatography A* 1442, 107-117.

1212 Roldán, A., Palacios, V., Caro, I., Pérez, L. (2003). Resveratrol content of Palomino fino
1213 grapes: Influence of vintage and fungal infection. *Journal of Agricultural and Food Chemistry*
1214 51(5), 1464-1468.

1215 Roldán, A., Palacios, V., Caro, I., Pérez, L. (2010). Evolution of Resveratrol and Piceid
1216 Contents during the Industrial Winemaking Process of Sherry Wine. *Journal of Agricultural*
1217 *and Food Chemistry* 58(7), 4268-4273.

1218 Romero-Pérez, A.I., Lamuela-Raventós, R.M., Andrés-Lacueva, C., de la Torre-Boronat,
1219 M.C. (2001). Method for the Quantitative Extraction of Resveratrol and Piceid Isomers in

1220 Grape Berry Skins. Effect of Powdery Mildew on the Stilbene Content. *Journal of*
1221 *Agricultural and Food Chemistry* 49(1), 210-215.

1222 Rosen, E.D., Hsu, C.-H., Wang, X., Sakai, S., Freeman, M.W., Gonzalez, F.J., Spiegelman,
1223 B.M. (2002). C/EBPalpha induces adipogenesis through PPARgamma: a unified pathway.
1224 *Genes & Development* 16(1), 22-26.

1225 Rosenow, A., Noben, J.-P., Jocken, J., Kallendrusch, S., Fischer-Posovszky, P., Mariman,
1226 E.C.M., Renes, J. (2012). Resveratrol-Induced Changes of the Human Adipocyte Secretion
1227 Profile. *Journal of Proteome Research* 11(9), 4733-4743.

1228 Rossmeisl, M., Flachs, P., Brauner, P., Sponarova, J., Matejkova, O., Prazak, T., Ruzickova,
1229 J., Bardova, K., Kuda, O., Kopecky, J. (2004). Role of energy charge and amp-activated
1230 protein kinase in adipocytes in the control of body fat stores. *International Journal of Obesity*
1231 28, S38-S44.

1232 Rosso, M.D., Soligo, S., Panighel, A., Carraro, R., Vedova, A.D., Maoz, I., Tomasi, D.,
1233 Flamini, R. (2016). Changes in grape polyphenols (*V. vinifera* L.) as a consequence of post-
1234 harvest withering by high-resolution mass spectrometry: Raboso Piave versus Corvina.
1235 *Journal of Mass Spectrometry* 51(9), 750-760.

1236 Sadi, G., Ergin, V., Yilmaz, G., Pektas, M.B., Yildirim, O.G., Menevse, A., Akar, F. (2015).
1237 High-fructose corn syrup-induced hepatic dysfunction in rats: improving effect of resveratrol.
1238 *European Journal of Nutrition* 54(6), 895-904.

1239 Santos, J.C., Gotardo, E.M.F., Brianti, M.T., Pirae, M., Gambero, A., Ribeiro, M.L. (2014).
1240 Effects of yerba maté, a plant extract formulation ("YGD") and resveratrol in 3T3-L1
1241 adipogenesis. *Molecules* 19(10), 16909-16924.

1242 Sato, M., Suzuki, Y., Okuda, T., Yokotsuka, K. (1997). Contents of resveratrol, piceid, and
1243 their isomers in commercially available wines made from grapes cultivated in japan.
1244 *Bioscience, Biotechnology and Biochemistry* 61(11), 1800-1805.

1245 Seo, Y.J., Kim, K.J., Koh, E.J., Choi, J., Lee, B.Y. (2017). Anti-adipogenesis mechanism of
1246 pterostilbene through the activation of heme oxygenase-1 in 3T3-L1 cells. *Phytomedicine* 33,
1247 7-13.

1248 Shang, J., Chen, L.L., Xiao, F.X., Sun, H., Ding, H.C., Xiao, H. (2008). Resveratrol improves
1249 non-alcoholic fatty liver disease by activating AMP-activated protein kinase. *Acta*
1250 *Pharmacologica Sinica* 29(6), 698-706.

1251 Shen, Y., Xu, Z., Sheng, Z. (2017). Ability of resveratrol to inhibit advanced glycation end
1252 product formation and carbohydrate-hydrolyzing enzyme activity, and to conjugate
1253 methylglyoxal. *Food Chemistry* 216, 153-160.

1254 Sun, B., Ribes, A.M., Leandro, M.C., Belchior, A.P., Spranger, M.I. (2006). Stilbenes:
1255 Quantitative extraction from grape skins, contribution of grape solids to wine and variation
1256 during wine maturation. *Analytica Chimica Acta* 563(1), 382-390.

1257 Sun, L., Wang, Y., Song, Y., Cheng, X.-R., Xia, S., Rahman, M.R.T., Shi, Y., Le, G. (2015).
1258 Resveratrol restores the circadian rhythmic disorder of lipid metabolism induced by high-fat
1259 diet in mice. *Biochemical and Biophysical Research Communications* 458(1), 86-91.

1260 Szkudelska, K., Nogowski, L., Szkudelski, T. (2009). Resveratrol, a naturally occurring
1261 diphenolic compound, affects lipogenesis, lipolysis and the antilipolytic action of insulin in
1262 isolated rat adipocytes. *Journal of Steroid Biochemistry and Molecular Biology* 113(1-2), 17-
1263 24.

1264 Takaya, Y., Terashima, K., Ito, J., He, Y.-H., Tateoka, M., Yamaguchi, N., Niwa, M. (2005).
1265 Biomimic transformation of resveratrol. *Tetrahedron* 61(43), 10285-10290.

1266 Tan, H.Y., Tse, I.M.Y., Li, E.T.S., Wang, M. (2017). Oxyresveratrol supplementation to
1267 C57bl/6 mice fed with a high-fat diet ameliorates obesity-associated symptoms. *Nutrients*
1268 9(2).

1269 Tennyson, C.A., Friedman, G. (2008). Microecology, obesity, and probiotics. *Current*
1270 *Opinion in Endocrinology, Diabetes and Obesity* 15(5), 422-427.

1271 Threlfall, R.T., Morris, J.R., Mauromoustakos, A. (1999). Effect of variety, ultraviolet light
1272 exposure, and enological methods on the trans-resveratrol level of wine. *American Journal of*
1273 *Enology and Viticulture* 50(1), 57-64.

1274 Tie, F.F., Luan, G.X., Zhou, W.N., Wang, Z.H., Shi, X.B., Li, G., Wang, H.L. (2018). Effects
1275 of the oligostilbenes from *Iris lactea* Pall. var. *chinensis* (Fisch.) Koidz on the adipocytes
1276 differentiation of 3T3-L1 cells. *Pharmazie* 73(2), 98-103.

1277 Tung, Y.C., Lin, Y.H., Chen, H.J., Chou, S.C., Cheng, A.C., Kalyanam, N., Ho, C.T., Pan,
1278 M.H. (2016). Piceatannol exerts anti-obesity effects in C57BL/6 mice through modulating
1279 adipogenic proteins and gut microbiota. *Molecules* 21(11).

1280 Vacca, V., Leccis, L., Fenu, P., Pretti, L., Farris, G.A. (1997). Wine yeasts and resveratrol
1281 content. *Biotechnology Letters* 19(6), 497-498.

1282 Vázquez-Vela, M.E.F., Torres, N., Tovar, A.R. (2008). White Adipose Tissue as Endocrine
1283 Organ and Its Role in Obesity. *Archives of Medical Research* 39(8), 715-728.

1284 Viñas, P., Campillo, N., Hernández-Pérez, M., Hernández-Córdoba, M. (2008). A comparison
1285 of solid-phase microextraction and stir bar sorptive extraction coupled to liquid
1286 chromatography for the rapid analysis of resveratrol isomers in wines, musts and fruit juices.
1287 *Analytica Chimica Acta* 611(1), 119-125.

1288 Viñas, P., Campillo, N., Martínez-Castillo, N., Hernández-Córdoba, M. (2009). Solid-phase
1289 microextraction on-fiber derivatization for the analysis of some polyphenols in wine and
1290 grapes using gas chromatography-mass spectrometry. *Journal of Chromatography A* 1216(9),
1291 1279-1284.

1292 Viñas, P., Martínez-Castillo, N., Campillo, N., Hernández-Córdoba, M. (2011). Directly
1293 suspended droplet microextraction with in injection-port derivatization coupled to gas
1294 chromatography–mass spectrometry for the analysis of polyphenols in herbal infusions, fruits
1295 and functional foods. *Journal of Chromatography A* 1218(5), 639-646.

1296 Vincenzi, S., Tomasi, D., Gaiotti, F., Lovat, L., Giacosa, S., Torchio, F., Río Segade, S.,
1297 Rolle, L. (2013). Comparative study of the resveratrol content of twenty-one italian red grape
1298 varieties. *South African Journal of Enology and Viticulture* 34(1), 30-35.

1299 Vitrac, X., Castagnino, C., Waffo-Téguo, P., Delaunay, J.-C., Vercauteren, J., Monti, J.-P.,
1300 Deffieux, G., Mérillon, J.-M. (2001). Polyphenols Newly Extracted in Red Wine from
1301 Southwestern France by Centrifugal Partition Chromatography. *Journal of Agricultural and*
1302 *Food Chemistry* 49(12), 5934-5938.

1303 Vitrac, X., Monti, J.P., Vercauteren, J., Deffieux, G., Mérillon, J.M. (2002). Direct liquid
1304 chromatographic analysis of resveratrol derivatives and flavanonols in wines with absorbance
1305 and fluorescence detection. *Analytica Chimica Acta* 458(1), 103-110.

1306 Vrhovsek, U., Masuero, D., Gasperotti, M., Franceschi, P., Caputi, L., Viola, R., Mattivi, F.
1307 (2012). A versatile targeted metabolomics method for the rapid quantification of multiple
1308 classes of phenolics in fruits and beverages. *Journal of Agricultural and Food Chemistry*
1309 60(36), 8831-8840.

1310 Vrhovsek, U., Wendelin, S., Eder, R. (1997). Effects of various vinification techniques on the
1311 concentration of cis and trans-resveratrol and resveratrol glucoside isomers in wine. *American*
1312 *Journal of Enology and Viticulture* 48(2), 214-219.

1313 Vuong, T.V., Franco, C., Zhang, W. (2014). Treatment strategies for high resveratrol
1314 induction in *Vitis vinifera* L. cell suspension culture. *Biotechnology Reports* 1-2, 15-21.

1315 Wang, J., Wang, S., Liu, G., Edwards, E.J., Duan, W., Li, S., Wang, L. (2016). The Synthesis
1316 and Accumulation of Resveratrol Are Associated with Veraison and Abscisic Acid
1317 Concentration in Beihong (*Vitis vinifera* × *Vitis amurensis*) Berry Skin. *Frontiers in Plant*
1318 *Science* 7, 1605-1605.

1319 Wang, S., Liang, X., Yang, Q., Fu, X., Rogers, C.J., Zhu, M., Rodgers, B.D., Jiang, Q.,
1320 Dodson, M.V., Du, M. (2015). Resveratrol induces brown-like adipocyte formation in white
1321 fat through activation of AMP-activated protein kinase (AMPK) α 1. *International Journal of*
1322 *Obesity* 39(6), 967-976.

1323 Wang, S., Liang, X., Yang, Q., Fu, X., Zhu, M., Rodgers, B.D., Jiang, Q., Dodson, M.V., Du,
1324 M. (2017). Resveratrol enhances brown adipocyte formation and function by activating AMP-
1325 activated protein kinase (AMPK) α 1 in mice fed high-fat diet. *Molecular Nutrition and Food*
1326 *Research* 61(4).

1327 Wang, Y., Voy, B.J., Urs, S., Kim, S., Soltani-Bejnood, M., Quigley, N., Heo, Y.R.,
1328 Standridge, M., Andersen, B., Dhar, M., Joshi, R., Wortman, P., Taylor, J.W., Chun, J.,
1329 Leuze, M., Claycombe, K., Saxton, A.M., Moustaid-Moussa, N. (2004). The Human Fatty
1330 Acid Synthase Gene and De Novo Lipogenesis Are Coordinately Regulated in Human
1331 Adipose Tissue. *Journal of Nutrition* 134(5), 1032-1038.

1332 Weiskirchen, S., Weiskirchen, R. (2016). Resveratrol: How Much Wine Do You Have to
1333 Drink to Stay Healthy? *Advances in Nutrition* 7(4), 706-718.

1334 Zhang, B.B., Zhou, G., Li, C. (2009). AMPK: An Emerging Drug Target for Diabetes and the
1335 Metabolic Syndrome. *Cell Metabolism* 9(5), 407-416.

1336 Zhang, D., Yan, Y., Tian, H., Jiang, G., Li, X., Liu, W. (2018). Resveratrol supplementation
1337 improves lipid and glucose metabolism in high-fat diet-fed blunt snout bream. *Fish*
1338 *Physiology and Biochemistry* 44(1), 163-173.

1339 Zhang, X.-H., Huang, B., Choi, S.-K., Seo, J.-S. (2012). Anti-obesity effect of resveratrol-
1340 amplified grape skin extracts on 3T3-L1 adipocytes differentiation. *Nutrition research and*
1341 *practice* 6(4), 286-293.

1342 Zhao, W., Li, A., Feng, X., Hou, T., Liu, K., Liu, B., Zhang, N. (2016). Metformin and
1343 resveratrol ameliorate muscle insulin resistance through preventing lipolysis and
1344 inflammation in hypoxic adipose tissue. *Cellular Signalling* 28(9), 1401-1411.

1345

1346 **Figure legends**

1347 **Figure 1.** Biosynthesis of trans-resveratrol and its chemical diversification pathways.(Chong
1348 et al., 2009)

1349 **Figure 2.** Main stilbenes identified in grape berries and wines.

1350

Table 1. Stilbenes content in berries (in mg/kg fw) and wines (in mg/L) (mean, standard deviation, minimum and maximum, number of studies, and number of measures).

Compounds	Content	SD	Min.	Max.	n	N	Ref.
Berries							
Monomers							
<i>trans</i> -resveratrol	0.67	0.97	0.00	3.56	12	84	(Brillante et al., 2018; Rosso et al., 2016; Flamini et al., 2016; Fernández-Marín et al., 2013; Flamini et al., 2013; Vincenzi et al., 2013; Vrhovsek et al., 2012; Viñas et al., 2011; Guerrero et al., 2010; Viñas et al., 2009; Kolouchová-Hanzlíková et al., 2004; Bavaresco et al., 2002)
<i>cis</i> -resveratrol	0.01	0.07	0.00	0.40	5	38	
Total resveratrol	0.68						
piceatannol	0.12	0.22	0.00	1.35	11	76	
<i>trans</i> -piceid	0.65	1.30	0.00	6.87	7	32	
<i>cis</i> -piceid	0.71	1.61	0.00	6.77	7	32	
Total piceid	1.36						
<i>trans</i> -astringin	0.71	0.58	0.12	1.73	5	7	
<i>cis</i> -astringin	0.12	0.09	0.04	0.29	4	6	
Total astringin	0.83						
Red wines							
Monomers							
<i>trans</i> -resveratrol	1.41	1.17	0.00	3.75	8	103	(Guerrero et al., 2020; Arraki et al., 2017; Lukić et al., 2019; Viñas et al., 2009; Buiarelli et al., 2007; Rennaker, 2007; Careri et al., 2004; Kolouchová-Hanzlíková et al., 2004; Lee and Sato et al., 1997)
<i>cis</i> -resveratrol	0.88	1.07	0.00	6.08	7	96	
Total resveratrol	2.29						
piceatannol	0.68	0.94	0.00	5.22	3	35	
<i>trans</i> -piceid	2.06	2.19	0.00	9.31	6	82	
<i>cis</i> -piceid	6.20	8.09	0.00	38.47	6	75	
Total piceid	8.26						
<i>trans</i> -astringin	0.52	0.61	0.00	3.00	3	32	
<i>cis</i> -astringin	0.13	0.32	0.00	1.59	2	28	
Total astringin	0.65						
Dimers							
<i>trans</i> - ϵ -viniferin	0.06	0.21	0.00	0.81	2	15	
<i>trans</i> - ω -viniferin	0.03	0.10	0.00	0.30	1	10	
Others							
miyabenol C	0.14	0.45	0.00	1.41	1	10	
hopeaphenol	0.08	0.15	0.00	0.49	2	15	
Isohopeaphenol	1.39	1.80	0.00	7.47	3	19	
R2-viniferin	0.42	1.05	0.00	3.28	1	10	
White wines							
Monomers							
<i>trans</i> -resveratrol	0.08	0.20	0.00	1.16	6	55	(Guerrero et al., 2020; Lukić et al., 2019; Arraki et al., 2017; Viñas et al., 2009; Buiarelli et al., 2007; Lee and Rennaker, 2007; Sato et al., 1997)
<i>cis</i> -resveratrol	0.04	0.13	0.00	0.76	5	55	
Total resveratrol	0.12						
piceatannol	0.04	0.16	0.00	0.59	3	14	
<i>trans</i> -piceid	0.20	0.39	0.00	1.91	6	61	
<i>cis</i> -piceid	0.68	2.23	0.00	16.26	6	61	
Total piceid	0.88						
<i>trans</i> -astringin	0.05	0.16	0.00	0.72	3	20	
<i>cis</i> -astringin	0.11	0.35	0.00	1.32	2	14	
Total astringin	0.16						

Table 2. Fragmentation patterns and tentative assignments of stilbenes in red wines (Moss et al., 2013).

Compounds	RT (Min)	Formula	Calculated	Experimental	Mass error (ppm)	CE (eV)	MS/MS product ions
<i>trans</i> -resveratrol	20.005	C ₁₄ H ₁₁ O ₃	227.0714	227.0725	4.84	22.5	185; 143
<i>cis</i> -resveratrol	21.96	C ₁₄ H ₁₁ O ₃	227.0714	227.0717	1.32	22.5	185; 143
piceatannol	14.242	C ₁₄ H ₁₁ O ₄	243.0663	243.0659	-1.65	25	201; 159
<i>trans</i> -piceid	11.185	C ₂₀ H ₂₁ O ₈	389.1242	389.1235	-1.8	20	227
<i>cis</i> -piceid	13.561	C ₂₀ H ₂₁ O ₈	389.1242	389.1263	5.4	20	227
astringin	8.209	C ₂₀ H ₂₁ O ₉	405.1191	405.1206	3.7	15	243; 201; 159
pallidol	21.459	C ₂₈ H ₂₁ O ₆	453.1344	453.1347	0.66	20	359; 265
parthenocissin A	23.959	C ₈ H ₂₁ O ₆	453.1344	453.1344	-4.19	20	359; 289
ampelopsin D	26.765	C ₂₈ H ₂₁ O ₆	453.1344	453.1363	4.19	20	359; 289
<i>cis</i> - ϵ -viniferin	31.448	C ₂₈ H ₂₁ O ₆	453.1344	453.1345	0.22	20	435; 411; 369; 359; 347; 333; 225
<i>trans</i> - ϵ -viniferin	33.833	C ₂₈ H ₂₁ O ₆	453.1344	453.1376	7.06	20	435; 411; 369; 359; 347; 333; 225
<i>cis</i> - ω -viniferin	32.202	C ₂₈ H ₂₁ O ₆	453.1332	453.1344	-2.65	20	435; 411; 369; 359; 347; 333; 225
<i>trans</i> - ω -viniferin	34.34	C ₂₈ H ₂₁ O ₆	453.1344	453.1357	2.87	20	435; 411; 369; 359; 347; 333; 225
<i>cis</i> - δ -viniferin	39.705	C ₂₈ H ₂₁ O ₆	453.1344	453.1329	-3.31	25	453; 411; 369; 359; 333
<i>trans</i> - δ -viniferin	38.789	C ₂₈ H ₂₁ O ₆	453.1344	453.1356	1.77	25	435; 411; 369; 359; 333
<i>trans</i> -scirpusin A	27.93	C ₂₈ H ₂₁ O ₇	469.1293	469.1307	2.98	25	451; 427; 385; 375; 359; 347; 333; 241
restrisol A	11.303	C ₂₈ H ₂₃ O ₇	471.1449	471.1447	-0.42		377; 349; 255; 121
parthenostilbenin A	19.64	C ₂₉ H ₂₅ O ₇	485.1606	485.1635	5.98	15	453; 391; 359; 289; 255; 187
parthonostilbenin B	21.088	C ₂₉ H ₂₅ O ₇	485.1606	485.1594	-2.47	15	453; 391; 359; 289; 255; 187
ϵ -viniferin glucoside	25.202	C ₃₄ H ₃₁ O ₁₁	615.1872	615.1878	0.98	20	453; 411; 359; 347
ampelopsin C	30.044	C ₄₂ H ₃₂ O ₉	679.1974	679.1981	1.03	30	585; 573; 491; 479; 385
<i>trans</i> -miyabenol C	35.372	C ₄₂ H ₃₂ O ₉	679.1974	679.1978	0.59	30	661; 637; 585; 573; 555; 451; 479; 357; 345
<i>cis</i> -miyabenol C	36.322	C ₄₂ H ₃₂ O ₉	679.1974	679.1984	1.47	30	661; 637; 585; 573; 555; 479; 451; 357; 345
hopeaphenol	31.695	C ₅₆ H ₄₁ O ₁₂	905.2604	905.2573	-3.42	35	811; 717; 611; 451; 359; 265

Table 3. Information on liquid chromatography methods for the identification and quantification of stilbenes in grape berries and wines.

Compounds	Sample preparation	Analytical methods	LOD	LOQ	References
1,2	Wine: direct injection Grape skin: extraction ethanol (80%) Standard used: <i>trans</i> - and <i>cis</i> - resveratrol	HPLC- UV	3-15 µg/L	-	(Kolouchová-Hanzlíková et al., 2004)
1, 7, 9, 15, 24	Wine: extraction ethyl acetate Grape berries: water/acetonitrile (1:1, v/v) Standard used: pure compounds	HPLC-UV	wine: 5-8 ng	-	(Landrault et al., 2002)
1, 7, 8	Grape skin: extraction with various solvent Standard used: pure compounds	UPLC-MS/MS (identification) HPLC-DAD (quantification)	-	-	(Sun et al., 2006)
1	Wine: direct injection Grape skin and pomace: extraction methanol/ethanol (4:1, v/v) Standard used: <i>trans</i> -resveratrol	HPLC-UV HPLC-ESI-MS/MS	10 µg/L	16 µg/L	(Careri et al., 2004)
1, 3, 15, 20	Grape skin: extraction methanol Standard used: <i>trans</i> -resveratrol	UPLC-MS/MS (identification) HPLC-DAD (quantification)	0.01 mg/kg	0.04 mg/kg	(Guerrero et al., 2010)
1, 2, 9, 14, 15, 16, 17, 24, 27, 28, 29, 31	Wine: direct injection Grape berries: extraction water/methanol/chloroform (1:2:2, v/v/v) Standard used: pure compounds	UPLC-QqQ-MS/MS	-	8-400 pg	(Vrhovsek et al., 2012)
1, 2, 3, 7, 8, 9, 10, 12, 15, 17, 28, 30, 31, 32, 33	Wine: direct injection Standard used: pure compounds	UPLC-QqQ-MS/MS	5-28 µg/L	12-84 µg/L	(Guerrero et al., 2020)
1, 3, 7, 9	Wine: solid-phase extraction (SPE) protocol Standard used: <i>trans</i> -resveratrol, <i>trans</i> -piceid, <i>trans</i> -piceatannol	UPLC-QTOF-MS/MS (identification) UPLC-QqQ-MS/MS (quantification)	48-50 µg/L	160-167 µg/L	(Buiarelli et al., 2007)
1, 3, 7, 8, 9, 10, 11, 14, 15, 17, 19, 24, 27, 28, 29, 30, 31	Grape berries: extraction methanol Standard used: <i>trans</i> -resveratrol, <i>trans</i> -piceid, ϵ -viniferin	UPLC-QTOF-MS/MS	-	-	(Flamini et al., 2016; Rosso et al., 2016; Flamini et al., 2013)
1, 2, 3, 7, 8, 9, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 29, 30, 31	Wine: extraction diethyl ether	UPLC-QTOF-MS/MS	-	-	(Moss et al., 2013)

Table 4. *In vitro* effects of resveratrol, other stilbenes and combination of stilbenes and other bioactives on adipogenesis, apoptosis, lipogenesis, lipolysis and thermogenesis.;

ADIPOGENESIS & APOPTOSIS			
Effect/mechanism of action	Cell model	Doses	Reference
Resveratrol and metabolites			
↓ PPAR γ , C/EBP α , CEBP β , FAS, FGF10, leptin, LPL, BMP2, HSL			(Liu et al., 2018; Eseberri et al., 2017; Chang et al., 2016; Hu et al., 2015; Zhang et al., 2012; Rayalam et al., 2008; Costa et al., 2011)
↑ adiponectin	Bovine intramuscular adipocytes		(Costa et al., 2011)
↓ Preadipocyte proliferation and adipogenic differentiation			
↓ Cell cycle entry (↓ AKT, MAPK, cyclin D1)	Human SGBS cells		(Santos et al., 2014; Mitterberger and Zwerschke, 2013; Kwon et al., 2012b; Fischer-Posovszky et al., 2010)
↓ Clonal expansion (cyclin A2)			
↑ SIRT1, SIRT 2, AMPK α , FOXO1	Human visceral adipocytes	1-400 μ M	(Liu et al., 2018; Santos et al., 2014; Costa et al., 2011; Fischer-Posovszky et al., 2010)
↑ apoptosis (↑ Caspase-3, Bax)	3T3-L1 cells		(Liu et al., 2018; Rayalam et al., 2008)
↓ IR activity			(Kwon et al., 2012b)
↓ IL-6, IL-8			(Fischer-Posovszky et al., 2010)
Other stilbenes: Pterostilbene, R-viniferin, R2-viniferin, ϵ-viniferin			
↓ PPAR γ , C/EBP α , FABP4, aP2, CHOP10, leptin, resistin, FAS			(Tie et al., 2018; Seo et al., 2017; Ohara et al., 2015; Hsu et al., 2012; Kim et al., 2008)
↑ adiponectin	3T3-L1 cells	6-50 μ M	(Hsu et al., 2012)
↓ Preadipocyte proliferation and adipogenic differentiation			
↓ Cell cycle entry at the G1-S and G2/M phase			(Tie et al., 2018; Hsu et al., 2012; Kim et al., 2008)
↑ HO-1			(Seo et al., 2017)
Combination resveratrol with other polyphenols and bioactives (quercetin, genistein, CLA, EGCG)			
↓ PPAR γ , C/EBP α , FABP4, perilipin			(Yang et al., 2008; Rayalam et al., 2007)
↑ apoptosis	Primary human adipocytes (HAs)		(Park et al., 2008)
↓ Preadipocyte proliferation and adipogenic differentiation	3T3-L1 cells	10-50 μ M	(Ahmed et al., 2017; Lasa et al., 2011)
↓ ERK $\frac{1}{2}$, ↑ JNK phosphorylation			(Yang et al., 2008; Rayalam et al., 2007)

LIPOGENESIS (adipose and hepatic)			
Effect/mechanism of action	Cell model	Doses	Reference
Resveratrol			
↓Lipogenesis (adipose and hepatic)	Human SGBS cells		(Li et al., 2016; Choi et al., 2014; Liang et al., 2013; Chen et al., 2011; Mercader et al., 2011; Szkudelska et al., 2009; Jin et al., 2013; Gnoni and Paglialonga, 2009)
↓ACC ↓ FAS	3T3-L1 cells	1-100 μM	(Li et al., 2016; Choi et al., 2014; Liang et al., 2013; Gnoni and Paglialonga, 2009)
↓SREBP-1c, PPARγ, SCD-1	H4IIEC3 cells		(Choi et al., 2014; Chen et al., 2011; Mercader et al., 2011; Jin et al., 2013)
↓LXR-α	HepG2 cells		(Jin et al., 2013)
↓AMPK			(Chen et al., 2011)
Pterostilbene			
↓Lipogenesis (adipose)	3T3-L1 cells	5-10 μM	(Hsu et al., 2012)
↓PPARγ, FAS			
↓GPDH			
Oxyresveratrol			
↓Lipogenesis (hepatic)	HepG2 cells	30 μM	(Lee et al., 2018)
↓SREBP-1c, PPARγ, SCD-1			
↓LXR-α			

LIPOLYSIS			
Effect/mechanism of action	Cell model	Doses	Reference
Resveratrol			
↓ PPAR γ	Bovine intramuscular adipocytes		(Liu et al., 2018; Chang et al., 2016)
↑HSL, ATGL	3T3-L1 cells		(Liu et al., 2018; Lasa et al., 2012)
↓TG content	3T3-L1 cells		(Rosenow et al., 2012)
↑Glycerol release	Rat adipocytes (epididymal tissue)	0.03-400 μ M	(Gomez-Zorita et al., 2017; Chang et al., 2016; Rosenow et al., 2012; Szkudelska et al., 2009)
↑ β -oxidation	3T3-F442A cells		(Mercader et al., 2011)
↑ CPT-1	Human SGBS cells		(Chang et al., 2016)
↓TNF α -induced lipolysis	3T3-F442A cells		
Resveratrol + CLA or + genistein			
↓ PPAR γ			
↑HSL, ATGL	3T3-L1 cells	10-25 μ M	(Lasa et al., 2011; Rayalam et al., 2007)
↓TG content			
↑Glycerol release			
Pterostilbene			
↑Glycerol release	3T3-F442A cells	100 nM-10 μ M	(Gomez-Zorita et al., 2017)
Oxyresveratrol			
↑ β -oxidation	HepG2 cells	30 μ M	(Lee et al., 2018)
↑ CPT-1			
THERMOGENESIS/MITOCHONDRIAL BIOGENESIS			
Effect/mechanism of action	Cell model	Doses	Reference
Resveratrol			
↑ PRDM16	3T3-L1 cells		(Wang et al., 2017; Wang et al., 2015)
↑ UCP-1	Adipocytes derived from primary mouse embryonic fibroblasts (MEF)		(Wang et al., 2017; Wang et al., 2015; Santos et al., 2014; Mercader et al., 2011; Rayalam et al., 2008)
↑ ATAD3	Human SGBS cells	10-200 μ M	(Li et al., 2016)
↑ SIRT3	Human SGBS cells		(Rayalam et al., 2008)
↑ Mfn2	Vascular cells isolated from iBAT		(Rayalam et al., 2008)
↑AMPK α			(Wang et al., 2017; Wang et al., 2015)

Table 5. *In vivo* effects of resveratrol, other stilbenes and combination of stilbenes and other bioactives on adipogenesis, apoptosis, lipogenesis, lipolysis and thermogenesis.

Stilbene and dose	Diet/Model	Doses/Duration	Reference
Resveratrol			
↓Adipogenesis ↓PPAR γ , C/EBP α , SREBP-1c, ACC, FAS, leptin ↑SIRT1 ↓Galanin-mediated signaling molecules: GalR1/2, PKCd, Cyc-D, E2F1, p-ERK	HFD, HPD FVB/N mice Kunming mice C57BL/6J mice Wistar rats	30-300 mg/Kg/day 6-12 weeks	(Mendes et al., 2016; Qiao et al., 2014; Andrade et al., 2014; Alberdi et al., 2013; Kim et al., 2011)
↓Lipogenesis (adipose or hepatic) ↓ACC, FAS, ME, G6PDH ↓PPAR γ , SREBP-1c ↑SIRT1, eNOS ↑AMPK	SD, HFD, HFCSD, atherogenic diet Blunt snout bream (<i>Megalobrama amblycephala</i>) Swiss mice FVB/N mice Kunming mice C57BL/6J mice ICR mice Apo-E deficient mice SAMP10 mice Wistar rats Sprague-Dawley rats	15-300 mg/Kg/day 2 days-20 weeks	(Zhang et al., 2018; Khaleel et al., 2018; De Almeida Pinheiro et al., 2017; Mendes et al., 2016; Gracia et al., 2016; Sun et al., 2015; Sadi et al., 2015; Choi et al., 2014; Andrade et al., 2014; Qiao et al., 2014; Jeon et al., 2014; Alberdi et al., 2013; Gómez-Zorita et al., 2013; Kim et al., 2011; Ahn et al., 2008; Shang et al., 2008; Shiozaki et al., 2011)
↓ Body weight, BMI ↓ Body adiposity and weight adipose tissue ↓ Iwat index (iwat mass/body weight) ↓ Adipocyte diameter	SD, HFD, HFCSD Blunt snout bream (<i>Megalobrama amblycephala</i>) Swiss mice FVB/N mice C57BL/6J mice Kunming mice CD1 mice Apo-E deficient mice Wistar rats Zucker (<i>falfa</i>) rat	0.0125-0.4% 1-300 mg/Kg/day 4-12 weeks	(Zhang et al., 2018; De Almeida Pinheiro et al., 2017; Mendes et al., 2016; Chang et al., 2016; Sun et al., 2015; Sadi et al., 2015; Wang et al., 2015; Andrade et al., 2014; Majumdar et al., 2014; Jeon et al., 2014; Qiao et al., 2014; Kim et al., 2011; Ahn et al., 2008) (Gómez-Zorita et al., 2013)
↓Total cholesterol ↓TG, LDL ↑HDL-cholesterol ↓TG	HFD, HPD, HFCSD, atherogenic diet Blunt snout bream (<i>Megalobrama amblycephala</i>) FVB/N mice Apo-E deficient mice C57BL/6J mice ICR mice CD1 mice	0.0125-1.08 % 15-300 mg/Kg/day 2 days-12 weeks	(Zhang et al., 2018; Mendes et al., 2016; Wang et al., 2015; Andrade et al., 2014; Majumdar et al., 2014; Qiao et al., 2014; Jeon et al., 2014; Choi et al., 2014; Kim et al., 2011; Ahn et al., 2008)

	Sprague-Dawley rats		
↑Insulin sensitivity ↓Insulin plasma level ↓Blood glucose ↑Glucose transporter 2	SD, HFD Blunt snout bream (<i>Megalobrama amblycephala</i>) Swiss mice Kunming mice Wistar rats	0.04-1.08 % 20-200 mg/Kg/day 1-12 weeks	(Khaleel et al., 2018; Zhang et al., 2018; De Almeida Pinheiro et al., 2017; Andrade et al., 2014; Qiao et al., 2014; Shang et al., 2008)
↓ Lipolysis ↑ HSL, ATGL ↑ Beta-oxidation ↑ CPT-1, MTPP, UCP-1, UCP-2, ACO	SD, HFD Blunt snout bream (<i>Megalobrama amblycephala</i>) CD1 mice SAMP10 mice Wistar rats Sprague-Dawley rats	0.04-1.08 % 20-30 mg/Kg/day 4-10 weeks	(Zhang et al., 2018; Khaleel et al., 2018; Wang et al., 2015; Alberdi et al., 2013; Shiozaki et al., 2011)
↑Browning ↑Mitochondrial biogenesis ↑UCP-1, PRDM16, Cytochrome C ↑AMPK α	HFD CD1 mice	0.01 % 4 weeks	(Wang et al., 2015)
↓ Inflammation ↓TNF- α , IL-6, IFN α , IFN β , NF-kB, MCP-1, CRP ↓ Macrophage infiltration	SD, HFD FVB/N mice C57BL/6J mice Zucker (<i>fa/fa</i>) rat Sprague-Dawley rats	0.04-1.08 % 20-30 mg/Kg/day 6-20 weeks	(Andrade et al., 2014; Majumdar et al., 2014; Gómez-Zorita et al., 2013; Kim et al., 2011)
Gut microbiota ↑Fiaf ↑ <i>Bacteroidetes</i> to <i>Firmicutes</i> ratios ↓ <i>Enterococcus faecalis</i> ↑ <i>Bifidobacterium</i> , <i>Lactobacillus</i>	HFD Kunming mice	200 mg/Kg/day 12 weeks	(Qiao et al., 2014)

Continuation

Stilbene and dose	Diet/Model	Doses/Duration	Reference
Combination resveratrol with other polyphenols and bioactives (quercetin, CLA, melatonin)			
↓ Body weight, BMI ↓ Weight adipose tissue ↓ Blood glucose ↓ Adipogenesis (↓FAS, ACC) ↓ Lipogenesis (↓FAS, ACC) ↓ Lipolysis (↑ HSL, ATGL) ↓ Total cholesterol, TG	HFD, HFCSD ICR mice Wistar rats Sprague-Dawley rats	Resveratrol: 15-50 mg/Kg/day Others: 3-30 mg/Kg/day 10 days-12 weeks	(Arias et al., 2016; Zhao et al., 2016; Arias et al., 2014; Majumdar et al., 2014)
Piceatannol			
↓ Body weight ↓ Total cholesterol, LDL and HDL cholesterol ↓ Blood glucose ↑ AMPK ↑ <i>Firmicutes</i> , <i>Lactobacillus</i> ↓ <i>Bacteroidetes</i>	S, HG, HFD <i>Caenorhabditis elegans</i> C57BL/6 mice	0.1 and 0.25% 50 and 100 μM 48 hours-18 weeks	(Shen et al., 2017; Tung et al., 2016)
Pterostilbene			
↓ Lipolysis ↑ Beta-oxidation ↑ UCP-1 ↓ Lipogenesis (hepatic) ↓ ACC, FAS, ME, G6PDH ↑ AMPK	SD, HFD Zucker (<i>fafa</i>) rats OLETF rats Wistar rats	15 and 300 mg/kg/day 4-6 weeks	(Nagao et al., 2017; Aguirre et al., 2016; Gómez-Zorita et al., 2014)
Oxyresveratrol			
↓ Body weight, BMI ↓ Insulin resistance ↓ C/EBPα, SREBP-1c, ↓ Fasting glucose ↓ Total cholesterol, LDL cholesterol ↑ SIRT1 ↑ AMPK	SD HFD C57BL/6 mice	0.25 and 0.5 % 10-30 mg/kg/day 4-8 weeks	(Lee et al., 2018; Tan et al., 2017)














Figure 1.




Monomers














Glycosylated monomers




Dimers









Glycosylated dimer



Trimers






Figure 2.