



HAL
open science

Relative Weight of Organic Waste Origin on Compost and Digestate 16S rRNA Gene Bacterial Profilings and Related Functional Inferences

Axel Aigle, Emilie Bourgeois, Laurence Marjolet, Sabine Houot, Dominique Patureau, Emmanuel Doelsch, Benoit Cournoyer, Wessam Galia

► To cite this version:

Axel Aigle, Emilie Bourgeois, Laurence Marjolet, Sabine Houot, Dominique Patureau, et al.. Relative Weight of Organic Waste Origin on Compost and Digestate 16S rRNA Gene Bacterial Profilings and Related Functional Inferences. *Frontiers in Microbiology*, 2021, 12, 10.3389/fmicb.2021.667043 . hal-03259341

HAL Id: hal-03259341

<https://hal.inrae.fr/hal-03259341>

Submitted on 14 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Relative Weight of Organic Waste Origin on Compost and Digestate 16S rRNA Gene Bacterial Profilings and Related Functional Inferences

Axel Aigle^{1†}, Emilie Bourgeois¹, Laurence Marjolet¹, Sabine Houot², Dominique Patureau³, Emmanuel Doelsch^{4,5}, Benoit Cournoyer^{1‡} and Wessam Galia^{1*‡}

OPEN ACCESS

Edited by:

Dimitrios Georgios Karpouzas,
University of Thessaly, Greece

Reviewed by:

Heribert Insam,
University of Innsbruck, Austria
Panagiotis A. Karas,
University of Thessaly, Greece

*Correspondence:

Wessam Galia
wessam.galia@vetagro-sup.fr

† Present address:

Axel Aigle,
ILM, Villeurbanne, France

‡ These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Microbiotechnology,
a section of the journal
Frontiers in Microbiology

Received: 11 February 2021

Accepted: 06 April 2021

Published: 14 May 2021

Citation:

Aigle A, Bourgeois E, Marjolet L,
Houot S, Patureau D, Doelsch E,
Cournoyer B and Galia W (2021)
Relative Weight of Organic Waste
Origin on Compost and Digestate
16S rRNA Gene Bacterial Profilings
and Related Functional Inferences.
Front. Microbiol. 12:667043.
doi: 10.3389/fmicb.2021.667043

¹ Univ Lyon, UMR Ecologie Microbienne (LEM), Université Claude Bernard Lyon 1, CNRS 5557, INRAE 1418, VetAgro Sup, Marcy L'Etoile, France, ² UMR ECOSYS, INRAE, AgroParisTech, Thiverval-Grignon, France, ³ INRAE, Univ Montpellier, LBE, Narbonne, France, ⁴ CIRAD, UPR Recyclage et risque, Montpellier, France, ⁵ Recyclage et Risque, Univ Montpellier, CIRAD, Montpellier, France

Even though organic waste (OW) recycling via anaerobic digestion (AD) and composting are increasingly used, little is known about the impact of OW origin (fecal matters and food and vegetable wastes) on the end products' bacterial contents. The hypothesis of a predictable bacterial community structure in the end products according to the OW origin was tested. Nine OW treatment plants were selected to assess the genetic structure of bacterial communities found in raw OW according to their content in agricultural and urban wastes and to estimate their modifications through AD and composting. Two main bacterial community structures among raw OWs were observed and matched a differentiation according to the occurrences of urban chemical pollutants. Composting led to similar 16S rRNA gene OTU profiles whatever the OW origin. With a significant shift of about 140 genera (representing 50% of the bacteria), composting was confirmed to largely shape bacterial communities toward similar structures. The enriched taxa were found to be involved in detoxification and bioremediation activities. This process was found to be highly selective and favorable for bacterial specialists. Digestates showed that OTU profiles differentiated into two groups according to their relative content in agricultural (manure) and urban wastes (mainly activated sludge). About one third of the bacterial taxa was significantly affected by AD. In digestates of urban OW, this sorting led to an enrichment of 32 out of the 50 impacted genera, while for those produced from agricultural or mixed urban/agricultural OW (called central OW), a decay of 54 genera over 60 was observed. Bacteria from activated sludge appeared more fit for AD than those of other origins. Functional inferences showed AD enriched genera from all origins to share similar functional traits, e.g., chemoheterotrophy and fermentation, while being often taxonomically distinct. The main functional traits among the dominant genera in activated sludge supported a role in

AD. Raw OW content in activated sludge was found to be a critical factor for predicting digestate bacterial contents. Composting generated highly predictable and specialized community patterns whatever the OW origin. AD and composting bacterial changes were driven by functional traits selected by physicochemical factors such as temperature and chemical pollutants.

Keywords: organic waste, anaerobic digestion, composting, 16S rRNA gene meta-barcoding, functional traits

INTRODUCTION

Organic wastes (OWs) have increased 10-fold since the last century and are likely to double by 2025 as a result of the growth of the world population (Hoornweg et al., 2013). There is thus a strong need for sustainable OW management. From a waste management standpoint, aerobic (composting) and anaerobic digestions (AD) are the most obvious operational processes prior to soil applications. These recycling treatments can also be combined by performing a composting of digestates. A major constraint in these OW recycling scenarios is the capacity to assess the related OW health hazards, which are, in fact, dependent on their origin and contents in terms of chemical and microbiological contaminants, and the efficacy of the treatments at reducing these hazards.

AD implies a breakdown of organic materials naturally found in raw OW into biogas (50–75% CH₄, 25–50% CO₂, and 1–2% H₂S, H₂, and NH₃) and organic residues (called digestate) by a microbial consortium (Atelge et al., 2020). The success of this process relies on the synergies between microorganisms acting in a coordinated succession driven by the environmental changes occurring during the process (Rivière et al., 2009; Yu et al., 2010). To avoid deviation of microbial communities during the AD process, parameters such as volatile fatty acids (VFAs), ammonia and hydrogen sulfide concentrations, acetate:propionate ratios, and pH are regularly monitored (Kayhanian, 1994; Chen et al., 2008; González-Fernández and García-Encina, 2009; Wang et al., 2009; Franke-Whittle et al., 2014; Wagner et al., 2014). The microbial reactions of these processes can be performed by psychrophiles, mesophiles, and thermophiles (Moseet et al., 2015; Saady and Massé, 2015; Hupfauf et al., 2018). Mesophilic digestion is considered the most stable AD process (Gou et al., 2014) because diversity of bacterial and archaeal assemblages are greater at mesophilic (35–37°C) temperatures (Levén et al., 2007; Pycke et al., 2011). The initial hydrolysis step of OW is performed by many facultative anaerobes or strictly anaerobic bacteria. This hydrolysis can degrade complex molecules into simple sugar monomers, amino acids, and alcohols. Fermentative bacteria will then convert these monomers into short-chain fatty acids, primary alcohols, as well as H₂ and CO₂. The production of acetate from the fatty acids will be performed by homoacetogenic bacteria like *Butyrivibrio* and *Acetobacterium*, that of H₂ will be performed by the acetogenic microbiota such as *Syntrophomonas*, and that of CO₂ and H₂S will be performed by sulfate-reducing bacteria such as *Desulfovibrio* and *Desulfobacter*. Some of these compounds will then be used by methanogens to generate CH₄. Hydrogenotrophic methanogens such as *Methanobacterium* and *Methanogenium* sp. convert H₂ and CO₂

into CH₄, and acetotrophic ones such as *Methanosarcina* will convert acetate into CH₄ (Demirel and Scherer, 2008; Lykidis et al., 2011).

In the same way, composting is a natural process characterized by the conversion of OW into by-products that can be used as soil conditioners and/or organic fertilizers (Reinikainen and Herranen, 2001; Ahmad et al., 2007). This process is performed by an aerobic microbial consortium composed of *Proteobacteria*, *Actinobacteria*, and fungi. During the process of composting, microbial community successions are driven by environmental parameters such as moisture, temperature, and the aeration conditions (Blanc et al., 1999; Tian et al., 2013; Zhang et al., 2016). Composting in its initial phase can be carried out by several mesophiles including bacteria, which represent environmental health hazards such as *E. coli*, *Salmonella*, *Klebsiella*, and *Nocardia* (Insam and de Bertoldi, 2007). These organisms can break down soluble organic compounds by exothermic reactions. As the temperature rises above 40°C, these mesophiles will be outcompeted by more thermophilic bacteria such as *Bacillus stearothermophilus* and species from the *Deinococcus/Thermus* group (Insam and de Bertoldi, 2007). These will continue the breakdown processes and generate higher temperatures that will partly sanitize the end products by killing heat-sensitive hazardous microbial agents. A compost pile can reach 65–70°C. As the supply of high-energy compounds will get low, thermo-tolerant mesophiles will be favored and will perform the last transformation or “stabilization” reactions (Day and Shaw, 2001). The cool zones in the composting systems, particularly with those having no or little mixing, can favor a survival and (re)growth of some bacterial pathogens (Noble and Roberts, 2004).

Beyond the general microbial structures described previously for AD and composting, several questions remain, including how the high inputs of microorganisms and cells from the feeding substrates (OW) under conditions of continuous flow impact the microbial community structure in the end products. On one hand, some studies suggested that the microbial communities of digested OW are unique to a digester because of high differences in substrate and operating conditions (Werner et al., 2011; Sundberg et al., 2013; Zhang et al., 2014; De Vrieze et al., 2016). However, some of these analyses on full-scale AD plants fed with a mixture of substrates (De Vrieze et al., 2016) or with activated sludge (Wang et al., 2018) also indicated that the microbial communities could be stable over time. Nevertheless, in other studies, the microbial communities in AD were found variable over time (among a same digester), even when substrate and operational conditions remained constant (Supaphol et al., 2011; De Vrieze et al., 2013; Williams et al., 2013; Vanwonterghem et al., 2014). On the other hand, during composting, microbial

communities of the raw OW were found to trigger a rise and decrease in temperature (Palaniveloo et al., 2020). These changes are highly selective but their incidence on microbial community patterns of mature compost remain to be evaluated according to the origin of OW. A knowledge gap thus remains about the relative contribution of the raw OW microbiome on the digestate and compost structural and functional microbial community patterns.

Here, we made the hypothesis that the microbiomes found among digesters and composts will be (i) significantly structured by the communities of the raw OW inputs, but that (ii) functional redundancy among these communities would generate distinct community profiles between treatment plants. Similar biodegradation activities between distinct treatment plants would thus be achieved by different microbial organizations because of this functional redundancy. To test these hypotheses, nine full-scale treatment plants located in various rural and urban backgrounds were investigated. These treatment plants were fed with agricultural or urban wastes or a mixture of both. Bacterial genetic structures were inferred from 16S rRNA gene meta-barcoding approaches, and patterns were compared between process types and according to the origin and nature of the raw OW. Functional traits were inferred from the 16S rRNA gene taxonomic allocations and databases recording the metabolic processes of well-defined bacterial taxa (FAPROTAX). The theoretical metabolic pathways of the genera impacted by treatment were further inferred from sequenced genomes using the MACADAM software (Le Boulch et al., 2019).

MATERIALS AND METHODS

OW Recycling Sites and Samplings

Nine full-scale OW treatment plants (numbered as Urban-1-2-3, Agri-1-2-3 and Central-1-2-3) were considered in this study (Table 1). The sampling strategy implied the use of at least three independent treatment plants per sample categories. OWs were treated by AD (Urban-1-2, Agri-1-2, and Central-1-2-3) or composting (Urban-1-3 and Agri-1-3). Five of the AD plants applied a post-treatment to the digestate: solid/liquid separation (Central-1-2), solid/liquid separation followed by composting (Urban-1 and Agri-1), or solid/liquid separation followed by solid phase drying (Urban-2). Raw OWs were chosen to represent the most common origins of OW: urban [sewage sludge produced by wastewater treatment plants (WWTP): Urban-1-2-3], agricultural (exclusively or mainly composed of animal manure and/or slurry: Agri-1-2-3), and central (a mix of food industry waste, harvest residues, livestock effluents, and/or sewage sludge: Central-1-2-3) (see Table 1 for details of the OWs compositions). For all sites, samples were taken at each treatment step to capture treatment effects on the microbial community's composition and abundance. Thus, a raw OW sample was taken from each site, resulting in nine samples, respectively, defined as urban (Urban-1-2-3-OW), agricultural (Agri-1-2-3-OW), and central (Central-1-2-3-OW) (Table 1). Digestates were available and sampled for six sites (Urban-1-D, Agri-1-2-D, and Central-1-2-3-D), as well as solid and dried

digestates, respectively, from liquid/solid phase separation, and from solid phase drying (Urban-1-2-SD, Agri-1-SD, Central-1-2-SD, and Urban-2-DD) (Table 1). Liquid digestates were available for four sites (Urban-2-LD, Agri-1-LD, and Central-1-2-LD), and finally, compost was produced and sampled also at four sites (Urban-1-3-C and Agri-1-3-C) (Table 1). At least 1 kg of OW, composts, or digestates was collected per treatment. Samples were coded as indicated in Le Bars et al. (2018). For each sample, certain pollutants and physicochemical parameters were measured (Supplementary Tables 1, 2). These datasets have been presented in Le Bars et al. (2018) and Sertillanges et al. (2020) and further analyzed in order to verify the chemical differentiation of samples according to their origin (urban, agricultural, and central). Total nitrogen (ISO 13878) and total carbon (ISO 10694) were determined by dry combustion with an elemental NC 2100 Soil Analyzer (Thermo Electron Corp.). Inorganic carbon (ISO 10693) was determined with a Bernard Calcimeter by adding HCl and measuring released CO₂. Organic carbon (C_{org}) was deduced from the difference between total and inorganic carbon. Concentrations of trace elements (TE) were measured using an inductively coupled plasma mass spectrometer (ICP-MS iCAP Q Thermo Fisher Scientific). pH was directly measured on liquid samples, or according to the ISO 10390. More details for those procedures are described in Le Bars et al. (2018). The concentrations of all the elements are expressed on a dry matter (DM) basis. Nonylphenols (NPs) and polycyclic aromatic hydrocarbons (PAHs) were measured in 1 g subsample according to Trably et al. (2004) and Mailler et al. (2017). More details for these monitorings are given in Sertillanges et al. (2020).

DNA Extractions and MST

DNA was extracted from 0.5 g of solid samples or 500 µl of liquid samples using the FastDNA Spin kit for soil and the FastPrep-24 (MP Biomedicals) following the manufacturer's instructions. All the DNA extractions were triplicated for each sample. DNA was eluted in 50 µl of ultrapure water, and quantity and quality were measured using a nanodrop-One UV-Vis Spectrophotometer (Thermo Fisher Scientific). DNA extracts were visualized after electrophoresis at 6 V/cm using a TBE buffer [89 mM Tris-borate, 89 mM boric acid, and 2 mM EDTA (pH 8.0)] through a 0.8% (w/v) agarose gel, and DNA staining with 0.4 mg ml⁻¹ ethidium bromide. A Gel Doc XR+ System (Bio-Rad, France) was used to observe the stained DNA and confirm their relative quantities and qualities. DNA was kept at -80°C and shipped on ice within 24 h to the DNA sequencing services when appropriate.

MST methods were used to monitor host fecal contaminations by human, bovine, and pig (Supplementary Table 3). These tests were performed on all DNA extracts including all replicates from the same sample. MST qPCR amplifications of general and host-specific *Bacteroides* markers, as well as general 16S rRNA gene qPCR assays, were performed as described in Marti et al. (2017) and Voisin et al. (2020). The human-specific marker (HF183) and the 16S rRNA genes were quantified by the SYBR Green chemistry while the bovine-specific, pig-specific, and the total *Bacteroidetes* markers were quantified using the Taqman chemistry. Six qPCR assays were performed per sample using DNA obtained from three independent extractions

TABLE 1 | Organic waste treatment plants and sample features.

Sampling sites	Reactor size (m ³)/temperature for AD	Samples	Duration ^a /treatment ^b	Sample code	Organic waste composition and co-substrate used for composting
Urban-1 French wastewater treatment plant and composting platform	6,200 32–42°C	Organic waste Digestate Solid digestate Compost ^c	38d/AD 4w/C	Urban-1-OW Urban-1-D Urban-1-SD Urban-1-C	Drained activated sludge Co-substrate: green waste
Urban-2 French wastewater treatment plant	6,000 32–42°C	Organic waste Liquid digestate Solid digestate Dried digestate	34d/AD	Urban-2-OW Urban-2-LD Urban-2-SD Urban-2-DD	Drained sludge
Urban-3 French wastewater treatment plant and composting platform		Organic waste Compost ^d	6w/C	Urban-3-OW Urban-3-C	Dehydrated activated sludge Co-substrate: green waste
Agri-1 Anaerobic digestion plant and composting platform (France)	950 32–42°C	Organic waste Digestate Liquid digestate Solid digestate Compost ^c	38d/AD 12w/C	Agri-1-OW Agri-1-D Agri-1-LD Agri-1-SD Agri-1-C	Porcine slurry (for 6–7 T/day) and grass clippings, food and vegetable wastes, corn silage and straws Co-substrate: none
Agri-2 Anaerobic digestion plant (France)	740 45°C	Organic waste Digestate	42d/AD	Agri-2-OW Agri-2-D	80% of bovine manure and 20% of stercoral material, poultry litter, corn cob, cooking oils, lawn mowing and bran
Agri-3 Composting platform (France)		Manure Compost ^d	12w/C	Agri-3-OW Agri-3-C	Raw cattle, sheep and equine manures Co-substrate: none
Central-1 Anaerobic digestion plant (France)	1,600 32–42°C	Organic waste Digestate Liquid digestate Solid digestate	70d/AD	Central-1-OW Central-1-D Central-1-LD Central-1-SD	50% livestock manures (N ₂ manure + animal slurry), 28% wastewater treatment plant sludge, 22% food industry by-products
Central-2 Anaerobic digestion plant (France)	6000 32–42°C	Organic waste Digestate Liquid digestate Solid digestate	30d/AD	Central-2-OW Central-2-D Central-2-LD Central-2-SD	45% pig slurry, 45% wastes from food industries, 10% cereal mixture
Central-3 Anaerobic digestion plant (France)	8,000 32–42°C	Organic waste Digestate	90d/AD	Central-3-OW Central-3-D	50% biowastes (community, catering, mass distribution), 20% dairy cow slurry, 30% wastes from food industries and agricultural by-products

^aProcess duration ($d = \text{days}/w = \text{weeks}$). ^bTreatments (AD = anaerobic digestion/C = composting). ^cThe substrates used for composting were solid digestates from the matching site (Urban-1 or Agri-1). ^dRaw organic wastes were used for the composting at the Urban-3 and Agri-3 sites.

(**Supplementary Table 4**). Standard curves were generated using 10-fold serial dilutions (ranging from 10^5 to 10^0 gene copies) of cloned targeted sequences (Marti et al., 2017). MST qPCR datasets were expressed per copy number of DNA targets per gram of dry sample and normalized by dividing their numbers by the 16S rRNA gene copy numbers per gram of dry sample.

V5–V6 16S rRNA Gene Sequencing and Reads Processing

The V5–V6 region of the 16S rRNA genes were amplified using the 799F forward (barcode + ACCMGGATTAGATACCKG) and 1193R reverse (CRTCCMCACCTTCCTC) primers (Chelius and Triplett, 2001; Bodenhausen et al., 2013; **Supplementary Table 3**) by the Molecular Research DNA lab (MrDNA, Shallowater, United States) and were sequenced by the Illumina MiSeq V3 technology. PCR amplifications were performed on pooled DNA extracts obtained per sample (equimolar mixes

of the triplicates) in order to handle variability between DNA extracts. The HotStarTaq Plus Master Mix Kit (Qiagen, United States) using the following temperature cycles was used to generate the PCR products: 94°C for 3 min, followed by 28 cycles of 94°C for 30 s, 53°C for 40 s, and 72°C for 1 min, with a final elongation step at 72°C for 5 min. PCR products and blank control samples were verified using a 2% agarose gel and following the electrophoretic procedure described above. PCR products obtained from field samples showed sizes around 430 bp but blanks did not show detectable and quantifiable PCR products. Dual-index adapters were ligated to the PCR fragments using the TruSeq[®] DNA Library Prep Kit, which also involved quality controls of the ligation step (Illumina, Paris, France). Illumina MiSeq DNA sequencings of the PCR products were paired-end and set up to obtain around 40K reads per sample.

The generated sequences were processed with the Mothur package (Schloss et al., 2009) according to a standard operation

protocol (SOP) for MiSeq-based microbial community analysis (Schloss et al., 2009; Kozich et al., 2013), so-called MiSeq SOP, and available at http://www.mothur.org/wiki/MiSeq_SOP. Briefly, primers and barcodes were trimmed according to quality and size filtration, followed by sequence alignment. Size filtration was refined, and unique sequences were extracted. The `pre.cluster` command was applied in order to mask the remaining sequencing errors as well as the `split.abund` command where the cutoff was set to 1. Chimera removal was performed using the `chimera.uchime` command and chimeric sequences were removed. At this stage, sequences were processed following two normalization strategies: (i) subsampling according to the sample containing the lowest number of sequences, and (ii) no subsampling (see statistical analysis for normalization strategy). Finally, for both datasets, the SILVA 16S rRNA Bacterial reference library was used for taxonomic allocation of the OTUs (cutoff 80)¹ and OTU clustering was performed with a similarity threshold of 97% for subsampled reads (Supplementary Table 5) or using the phylotype command at the genus level, for non-subsampled reads (Supplementary Table 6). The Mothur package was also used to calculate diversity indices (Shannon, Simpson, evenness, and richness) on the 16S rRNA subsampled OTU table (Supplementary Table 7).

Statistical Analyses

Relations between sample groupings and chemical datasets were tested by principal component analysis using the `rda` function and plotted using the `biplot` function; confidence ellipses were drawn to show the variance observed per group. Kruskal–Wallis tests were performed using the `kruskal.test` function on 16S rRNA diversity indices, MST markers ratio, and physical–chemical data. When significant differences were found ($p < 0.05$), a Dunn test was applied using the `kwAllPairsDunnTest` function from the `PMCMRplus` package (Pohlert, 2020) for *post hoc* analysis.

Hellinger transformation was applied on the 16S rRNA subsampled OTU table using the `decostand` function (method = “total”), and Bray–Curtis dissimilarities were computed between pairs of samples using the `vegdist` function (“bray”). NMDS was generated using the `metaNMDS` function. `ordiellipse` option was used to display the variance observed per group, and PERMANOVA tests were performed using the `pairwise.adonis` and `vegdist` functions (Bray–Curtis dissimilarities) to confirm the significance of differences between groups.

The non-subsampled 16S rRNA gene reads genus table (Supplementary Table 6) was processed with the DESEQ2 package (Love et al., 2014) in order to reveal genera with significant change in abundance before and after treatment. Counts normalized by the median of ratios method (Anders and Huber, 2010) were recovered and relative abundances were calculated in order to present the percentage of significant differences in abundances at the genus level before and after treatment. Change in abundance has been considered significant if a genus shows a log₂ fold change ≤ -2 or $\geq +2$ with a minimum Base Mean value of 10 and adjusted p value of < 0.05 . To facilitate

global comparison with other studies, computations were also performed at the phylum level. All the functions used without package indication were from the `vegan` package (Oksanen et al., 2019) and all analyses were performed on RStudio (R v3.3.3). The statistical experimental design (Kruskal–Wallis tests on 16S rRNA diversity indices and MST markers, as well as NMDS and DESEQ2 analyses) implied first testing relations according to treatment (AD against composting), regardless of the OW origin and regardless of the post-treatments. Thus, all OW samples were compared to composted samples and digestates. Second, the OW origins were considered in the comparisons to infer their impact on the V5–V6 16S rRNA gene datasets.


The 16S rRNA gene sequences produced in this study are available from webin under the accession number PRJEB40193.

RESULTS

General Chemical and Microbiological Features of the Raw OWs, Digestates, and Composts

Principal component analyses were performed on the physicochemical datasets (organic carbon, total nitrogen, C:N ratio, pH, and phosphorus) and pollutants (iron and zinc concentrations, PAHs, NPs, and TE) measured on all samples (Supplementary Tables 1, 2 and Supplementary Figure 1). The first two axes of these PCAs explained a significant part of the variability observed among these datasets ($> 60\%$) (Supplementary Figure 1). The origin of the sample was found to explain the chemical dataset (Supplementary Figure 1). The urban samples (treated or not) were significantly differentiated from the agricultural and central (mixed) samples by a set of pollutants including several TE (Ag, Al, As, Cd, Ce, Co, Cr, Cu, Ga, K, Mo, Ni, P, Pb, and Ti), 4-nonylphenol (A2), phenanthrene (H5), anthracene (H6), and total nitrogen concentrations (KW–Dunn’s *post hoc* tests; $p < 0.05$) (Supplementary Figure 2). Similarly, global indicators of the microbiological content of the samples were analyzed by qPCR using microbial source tracking DNA targets (Supplementary Table 4 and Figure 1). These qPCR analyses showed that 16S rRNA gene numbers varied from 4.26×10^5 copies/g sample up to 6.41×10^{11} copies/g (Supplementary Table 4). Lowest 16S rRNA gene copy numbers/g sample were obtained for digestates. Monitorings of total *Bacteroides* cell numbers were correlated with the 16S rRNA gene dataset (Kendall’s rank correlation tau; $p < 0.01$) but the range of values was lower (2.44×10^2 copies/g sample up to 1.12×10^9 copies/g). The DNA indicators for pig and bovine fecal contaminations were most significant among central and agricultural raw OW (Figure 1 and Supplementary Table 4). These distributions were in agreement with the information given by the operators of the OW treatment sites. The human DNA fecal marker was most often detected among the raw OW but in significantly higher numbers (KW and Dunn’s *post hoc*; $p < 0.05$) among the urban and central groups of samples (Supplementary Table 4 and Figure 1). Digestates and composts showed lower occurrences of these markers but

¹<https://www.arb-silva.de/documentation/>



a significant number of HF183 DNA was detected among the urban digestates (**Figure 1**).

Bacterial Diversity of Raw OWs, Digestates, and Composts Inferred From V5 to V6 16S rRNA Gene Sequences

About 5 million DNA sequences were obtained for the samples investigated in this work, and 45% were validated as high-quality 16S rRNA gene sequences. The number of high-quality V5–V6 16S rRNA gene sequences per sample ranged from 26,465 to 119,793 (**Supplementary Table 7**). The number of reads per sample were negatively correlated to the 16S rRNA gene qPCR dataset (Spearman Rank Correlation; $\rho = -0.45$, $p < 0.05$). This highlighted a variability in the dataset that required a normalization of the 16S rRNA gene dataset prior to performing statistical analyses. The global representation of the dataset and the computing of diversity indices were thus performed after a subsampling of the gene reads at the lowest value for the number of reads on a sample, i.e., 26,465 reads. This subsampled dataset led to a grouping of reads into 7755 OTUs, which varied in their distribution from 318 to 1404 per sample (**Supplementary Table 5**). Accordingly, Shannon diversity, Shannon-evenness, and the Simpson diversity indices were computed (**Supplementary Table 7** and **Supplementary Figure 3**). Significant differences in these indices were observed between those computed for the urban digestates and those of the agricultural or central samples (KW test; $p < 0.05$). More diverse OTU contents and even distributions of the reads between OTUs were observed among the urban digestates than the other ones. However, the initial OTU contents of raw OW were found similar between sites, and diversity indices remained similar between raw OW and digestates of a same site (**Supplementary Figure 3**).

To better resolve differences in the genetic diversity among samples, Bray–Curtis dissimilarities between OTU profiles of

each pair of samples were computed. NMDS plots were then performed to visualize the differences (**Figure 2A**). All OTU profiles could be differentiated according to their site category (urban, central, or agricultural) (PERMANOVA $p < 0.05$). A significant effect on the profiles of the OW treatment by composting or AD could also be resolved (PERMANOVA $p < 0.05$). Digestates' OTU profiles could not be differentiated into three categories (PERMANOVA $p > 0.05$) but into two entities with (i) grouping the agricultural/mixed central samples and (ii) the urban ones (PERMANOVA $p < 0.01$) (**Figure 2B**). A further differentiation of OTU profiles according to post-treatments of the digestates was tested. The OTU patterns of liquid and solid digestates could not be differentiated (**Supplementary Figure 4**). Composting was found to drive bacterial community changes toward similar structures even though the nature of the OWs was variable (WWTP activated sludge, manure, green wastes, etc.) (PERMANOVA $p < 0.05$) (**Figure 2A**).

Incidence of AD or Composting on Bacterial Taxa as Inferred From 16S rRNA Gene Taxonomic Allocations

The chemical and V5–V6 16S rRNA gene OTU pattern analyses indicated in the previous sections showed (i) a significant proximity between the agricultural and central samples (raw OW and digestates), (ii) a significant differentiation of the urban samples (raw OW and digestates) from the agricultural/central samples, and (iii) a significant differentiation of the compost samples from all the other samples. Accordingly, in the following analyses, the datasets were grouped into these latter three entities. Informative groupings of V5–V6 DNA reads matching significant changes (higher or lower) in relative abundances between these entities were analyzed.




FIGURE 2 | Non-metric multidimensional scaling (NMDS) plot of Bray-Curtis dissimilarities computed from the 16S rRNA V5-V6 OTU profiles (subsamped, 97% identity) of the organic samples before (raw OW) and after treatment by anaerobic digestion or composting. **(A)** NMDS ordinations were built from the full dataset shown in **Table 1**; stress value was 0.1227. **(B)** NMDS ordinations were built from a dataset limited to raw OW and digestates. The ellipses indicate the variance observed within each group of samples. PERMANOVA tests are indicated in the text and confirmed the significance ($p < 0.05$) of several groupings.

Variations at the Scale of Phyla

Raw OWs showed a high variability in their OTU contents (**Figure 2A**). However, these OTUs can be grouped into larger taxonomic entities, which can be conserved between raw OWs, and contribute reproducibly to the buildup of the microbial communities among composts or digestates. These core taxa were found to be distributed among 6 major phyla: *Firmicutes* (31.2%), *Proteobacteria* (26.4%), *Bacteroidetes* (26.2%), *Actinobacteria* (6.1%), *Tenericutes* (4.4%), and *Spirochaetae* (0.7%) but discrepancies were observed in terms of proportion of reads per phylum according to the site category (urban, central, and agricultural) (**Supplementary Table 8**). Raw urban OW showed lower relative numbers (7.1%) of *Firmicutes* [converted into 2.05×10^9 equivalent 16S rRNA gene copy numbers (16S) per gram of OW] than the raw OWs from the central and agricultural sites (29.9%; equivalent to 1.01×10^{10} 16S per gram) (**Supplementary Table 9** and **Figure 3**). These urban OWs were found to have higher relative counts of *Proteobacteria* (46.6%;

equivalent to 3.97×10^{10} 16S per gram) and *Actinobacteria* (11.7%; equivalent to 6.51×10^9 16S per gram) than the central and agricultural samples (respectively, 29.3%; equivalent to 8.74×10^9 16S per gram and 2.9%; equivalent to 1.2×10^9 16S per gram) (**Supplementary Table 9** and **Figure 3**). It is to be noted that 16S rRNA gene reads allocated to the *Nitrospirae*, *Chloroflexi*, and *Acidobacteria* represented more than 1% (equivalent to 7.41×10^9 16S rRNA gene copies per gram) of the urban OW dataset, but these phyla occurred at a much lower score (equivalent to 4.32×10^6 16S rRNA gene copies per gram) among the OW of the other sites (**Supplementary Table 9** and **Figure 3**).

As observed among raw OW, *Firmicutes* and *Proteobacteria* were among the most abundant phyla in digestates and composts (**Supplementary Table 9** and **Figure 3**). However, a decrease in the relative abundance of the *Proteobacteria* was observed in all treated samples (**Supplementary Table 9** and **Figure 3**) and was supported by a significant statistical test for the urban




FIGURE 3 | Relative averaged abundances of V5-V6 16S rRNA gene reads allocated to phyla before and after treatment of raw organic wastes (OWs) by anaerobic digestion or composting performed by treatment plants located in rural areas (agricultural/central sites) or urban ones. Only phyla with a relative abundance above 0.5% are displayed (see **Supplementary Table 9** for the complete dataset).

samples (KW $p \leq 0.05$). These changes in proportion were not correlated to a significant reduction in bacterial cell numbers (between 1.77×10^9 and 3.19×10^{10} 16S rRNA gene copies per gram of digestates or composts). The observed change in relative abundance thus implies a significant re-shuffling of bacterial phyla with some of these taking advantage of the treatment conditions or becoming negligible. For example, increases in the relative proportion of *Firmicutes* from the raw OW to the digestates were observed among both categories of samples (agricultural/central, and urban). *Firmicutes* showed the highest occurrence among the central/agricultural digestates (52.2% of the bacterial taxa). These relative changes matched an averaged one log increase in *Firmicutes* 16S rRNA gene copies between the raw OW and the digestates of the urban sites or composts (Supplementary Table 9 and Figure 3). Similarly, relative increases in *Actinobacteria*, *Deferribacteres*, and *Thermotogae* among composts matched significant increases in 16S rRNA gene copies estimated from the qPCR datasets (Supplementary Table 9 and Figure 3; KW $p \leq 0.05$). Similar trends were also observed between raw OW and urban digestates for the *Cloacimonetes*, which showed up to a five-log increase in 16S rRNA gene copies per gram. Significant decreases in the relative abundance of some phyla were also recorded such as the raw OW *Fusobacteria*, which became negligible among the digestates and composts (Supplementary Table 9 and Figure 3; KW $p \leq 0.05$).



Variations at the Scale of Genera Among Digestates

To go deeper into the understanding of community changes, variations of V5–V6 16S rRNA gene reads allocated at the genus level were investigated. A DESEQ2 analysis was performed to highlight these changes (Figures 4, 5 and Supplementary Tables 10, 11). These analyses were performed with the non-resampled sequenced gene reads dataset (Supplementary Table 6) because a normalization scheme is directly run by DESEQ2. This approach revealed that the digestates of the central/agricultural category of samples led to a significant increase in the relative proportion of six genera: *Gelria* (*Firmicutes*), *Sporosarcina* (*Firmicutes*), *Dethiobacter* (*Firmicutes*), and three unclassified groups belonging to the *Spirochaetales*, *Cloacimonetes*, and *Firmicutes*. Four (*Gelria*, *Sporosarcina* and two others unclassified at genus level) of these genera were part of the core bacterial taxa found among digestates (Figure 4 and Supplementary Table 10). Four of these were also found to be part of the *Firmicutes* and confirmed the trends observed at the scale of phyla, which indicated an increase in *Firmicutes*. These six genera were two to five times (Log2 fold change) more abundant in the digestates than the non-treated samples (raw OWs). They represented from 0.1 to 6.1% (according to the mean normalized counts) of the total digestate bacterial community (Supplementary Table 10). Conversion of relative read numbers into 16S rRNA gene copies from the subsampled contingency table supported these trends and indicated an abundance of about 3.0×10^7 *Gelria* 16S rRNA gene copies per gram of dry weight in the raw OW while 5.0×10^8 *Gelria* 16S rRNA gene copies were estimated in average in the digestates (Supplementary Table 9). Similarly, *Sporosarcina* and *Dethiobacter* 16S rRNA

gene copies per gram of dry weight indicated a gain of one log between the raw OW and digestates. Regarding genera that showed a relative decrease, 54 (of which 15 are parts of the core bacterial taxa found among digestates) (Supplementary Table 8) were subjected to a significant decrease after AD of OW of the agricultural/central sites (Figure 4 and Supplementary Table 10). This relative reduction varied from 2.6- to 9.9-fold when compared to raw OW (Figure 4 and Supplementary Table 10). *Lactobacillus* (4.8–0.1%), *Bacteroides* (4.1–0.4%), *Prevotella* 9 (1.8–0.0%), and *Methanobrevibacter* (0.5–0.1%) were found to have undergone the most significant reductions (Figure 4 and Supplementary Table 10). The decrease in bacterial numbers was confirmed through a conversion of the dataset into 16S rRNA gene copies. *Lactobacillus* 16S rRNA gene copies went down by about two logs, and those of *Bacteroides* went down by about one log (supported by Wilcoxon rank sum tests; $p < 0.05$). Similar trends were observed for the other genera (Supplementary Table 9).

Several bacterial genera of the urban OWs were shown to be significantly enriched in the digestates. A total of 32 genera (of which 25 were part of the core microbiome) showed a significant increase (Figure 5 and Supplementary Table 11). These genera were enriched by three to sevenfold while comparing with the concentrations observed among OW. These genera represented from 0.03 to 3.6% of the total digestate microbial community. The most significant increases in the digestates were observed for V5–V6 16S rRNA gene reads allocated to the *Smithella* (0.03–3.0%), part of the *vadinBC27_wastewater-sludge_group* (0.1–2.1%), *Candidatus_Caldatribacterium* (0.1–1.9%), *Gracilibacter* (0.0–1.8%), *Proteiniphilum* (0.02–1.6%), *Sedimentibacter* (0.01–1.5%), *Syntrophorhabdus* (0.01–1.0%), and *Petrimonas* genus (0.01–0.5%). Conversion of these values into 16S rRNA gene copies showed the *Smithella* to have undergone an increase from 6.15×10^6 copies per gram of OW to 2.64×10^9 copies per gram of digestates, and the *Gracilibacter* 16S rRNA gene copies went up from about 3.0×10^6 to 1.61×10^9 (Supplementary Table 9). Similar trends were observed for the other above-cited genera (Supplementary Table 9) but a significant number ($n = 18$) of genera also underwent a relative reduction in V5–V6 gene reads from the OW to the digestates (Figure 5 and Supplementary Table 11). These decreases were about four- to ninefold in comparison to their relative numbers among the raw OW samples. Hence, the *Uliginosibacterium* (1.3–0.1%), *Simidiuia* (0.3–0.0%), *Cellvibrio* (0.1–0.01%), *Fonticella* (0.1–0.0%) *Polyangium* (0.04–0.0%), and *Clostridium_sensu_stricto_12* (0.03–0.0%) genera showed such a decrease (Figure 5 and Supplementary Table 11). These shifts were confirmed while converting these datasets into 16S rRNA gene copy numbers (Supplementary Table 9).

Global changes in the relative abundance of digestates' bacterial genera significantly affected by ADs over the dataset were investigated. DESEQ2 analyses revealed that 30.8 and 35.5% of the total relative abundance of the digestate genera were significantly affected by the AD treatment, respectively, at the urban and agricultural/central sites (Supplementary Tables 10, 11). The relative abundances of 50 and 60




genera were significantly impacted by AD in, respectively, the urban and central/agricultural digestates. Despite this similar effect, a remarkable difference depending on OW origins was highlighted. In digestates of urban origin, this sorting mainly involved an enrichment of genera (32 out of the 50 impacted genera) while those produced from agricultural/central OW

led essentially to a reduction of several taxa (54 out of 60 impacted genera) (**Supplementary Tables 10, 11**). These comparisons suggested a full re-shuffling of agricultural/central OW bacterial taxa while those of the urban OW appeared to be well adapted for the living conditions prevailing in the anaerobic digester.

Variations at the Scale of Genera Among Composts

Composting was found to drive microbial communities toward similar V5–V6 genetic structures whatever the origin of the OW including those that had undergone AD (Figure 2A). These conserved compost microbiomes appeared to have been selected by strong environmental constraints that led to an increase in the relative abundance of, among others, *Actinobacteria*, *Deferribacteres*, *Firmicutes*, and *Thermotogae*. These trends were confirmed while considering the numbers of 16S rRNA gene copies between OW and composts for these phyla. In fact, a significant one log increase was observed for the *Actinobacteria* and *Firmicutes* per gram of composts (Supplementary Table 9). Furthermore, the 16S rRNA gene copies allocated to the *Thermotogae* and *Deferribacteres* went from a negligible score up to several millions. Simultaneously, V5–V6 reads of six phyla (*Nitrospirae*, *Elusimicrobia*, *Acidobacteria*, *Saccharibacteria*, TM6, and GOUTA4) became negligible and were associated to a two-log reduction in 16S rRNA gene copies (Supplementary Table 9). These shifts among these phyla were confirmed at the scale of genera (Figure 6 and Supplementary Table 12). The compost microbiome was enriched in 36 genera of which only 6 were versatile and not found among all composts (Figure 6 and Supplementary Table 12). Relative distribution patterns of these genera changed by 4.1- to 8.1-Log2 fold when compared to the non-treated OW samples (going from 0.04 to 5.7% of the total compost microbial community). The most significant relative distribution increases were observed for reads allocated to the *Actinomadura*, *Longispora*, *Parapedobacter*, *Saccharomonospora*,

Brachybacterium, *Pedobacter*, *Thermobifida*, *Planifilum*, *Oceanobacillus*, *Olivibacter*, and *Paenalcaligenes*. These genera are part of the *Actinobacteria* (50%), *Bacteroidetes*, or *Firmicutes*. In most cases, about two log increases in 16S rRNA gene copy numbers were associated with these changes (all statistical tests were significant; Wilcoxon $p < 0.05$) (Supplementary Table 9). For example, the number of 16S rRNA copies for the *Olivibacter* went up from 2.0×10^8 to 1.0×10^{10} . Most impressive was the shift from negligible numbers of *Longispora* among the OW going up to about 2.0×10^9 16S rRNA gene copies (Supplementary Table 9). These changes were associated with the relative decreases of V5–V6 16S rRNA gene reads allocated to 105 genera including 29 found in the raw OWs core microbiome (Figure 6 and Supplementary Table 12). This reduction went from 3.8- to 8.1-Log2 fold change compared to OW where these genera originally represented 0.02–4.7% of the total raw OW microbial community (Figure 6 and Supplementary Table 12). Reads from *Tetrasphaera* (1.7–0.04%), *Microthrinx* (0.6–0.03%), *Intestinibacter_9* (0.4–0.04%), *Treponema_2* (0.4% to 0.0%), *Proteiniphilum* (0.4–0.04%), *Simplicispira* (0.3–0.0%), *Christensenellaceae_R7_group* (0.3–0.03%), *Lautropia* (0.2–0.0%), *Macellibacteroides* (0.2–0.01%), BD17_clade (0.1–0.0%), *Propionivibrio* (0.1–0.0%), *Proteocatella* (0.1–0.0%), and *Streptococcus* (0.1–0.0%) in the OW were the most significantly impacted (Figure 6 and Supplementary Table 12). These decreases among the community were confirmed by an analysis of the equivalent 16S rRNA gene copies for these genera in the OW and composts (Supplementary Table 9). DESEQ2 analyses



indicated that a re-shuffling of 52.6% of the read distribution patterns occurred during composting of OW. Composting was thus highly selective with a decline of 28.3% of the total relative abundance of the compost's bacterial taxa and an enrichment of 24.4% of the other taxa in the composts (**Supplementary Table 12**).

Incidence of OW Digestion Processes on Bacterial Functional Groups

Functional traits selected by AD and composting were inferred from the taxonomic allocations performed at the genus level. At first, the sum of relative abundances of enriched genera potentially covering a functional trait extracted from the FAPROTAX database using MACADAM was calculated (**Supplementary Table 13**). The contribution of each genus to differences in hypothetical functional profiles is based on relative abundances before and after treatment (**Supplementary Table 13**). Based on the most abundant taxa in digestates from urban origin, the five main hierarchical functional traits were chemoheterotrophy (6.3%), fermentation (6.2%), human and mammal gut colonizers (each feature at 2.1%), and animal parasites or symbionts (2.1%) (**Supplementary Table 13D**). Similarly, these functional traits were also associated with the most abundant taxa in digestates from agricultural and central origin (**Supplementary Table 13F**). In composts, the functional traits of the enriched genera (as suggested from the number of equivalent 16S rRNA gene copies) were related to aliphatic non-methane hydrocarbon degradation, arsenate detoxification, dissimilatory arsenate reduction, hydrocarbon degradation, nitrate and nitrite ammonification, nitrate and nitrite respiration, and oil bioremediation. These taxa had increased by at least 8.5 log after composting of raw OWs (**Supplementary Table 13B**).

Functional traits related to health hazards were also investigated based on changes in relative abundances of bacterial genera before and after OW treatment (**Supplementary Table 13**). AD and composting processes contributed significantly at reducing genera potentially harboring infectious agents. The number of 16S rRNA gene copies of genera potentially causing diseases in human, plants, or fishes decreased by 9.3, 7.4, and 7.9 log, respectively, after composting of OW (**Supplementary Table 13A**). Similar trends could be observed during AD for genera harboring potential human pathogens. The number of 16S rRNA gene copies of genera potentially associated with human pathogens went down by about 7.4 and 1.7 log, respectively, in digestates from urban and agricultural/central categories (**Supplementary Tables 13C,E**). Interestingly, the number of 16S rRNA gene copies of genera possibly including lung pathogens or plant pathogens went down by about 7.5 log in digestates from the agricultural/central category (**Supplementary Table 13E**).

The theoretical metabolic pathways of the genera identified in this study were further inferred from sequenced genomes using the MACADAM software. Metabolic pathways (with a $PS \geq 0.5$) potentially present in genera significantly impacted by the investigated digestion processes were counted, and

their relative distribution patterns were compared (**Table 2** and **Supplementary Table 14**). Distribution patterns between these functional categories observed among OWs from urban sites and from central/agricultural ones were found positively correlated (Spearman Rank Correlation; $\rho = 0.69$, $p < 0.05$). Such correlations were not observed between the anaerobic and aerobic digestion processes (digestates and composts).

After AD treatments, biosynthesis pathways for cofactors, nucleotides, amino acids, lipids, and aminoacyl-tRNAs charging were more abundant in the enriched genera than in the decaying ones whatever the raw OW origin (**Table 2** and **Supplementary Table 14**). Whereas biosynthesis pathways for secondary metabolites, cell structure, metabolic regulators, and alcohol and hormone biosynthesis were less abundant in the enriched genera than in the decaying ones (**Table 2** and **Supplementary Table 14**). Similarly, degradation pathways for fatty acids and lipids, carbohydrates, secondary metabolites, aromatic compounds, and sulfur-containing compounds metabolism degradation were less abundant in the enriched genera than the decaying ones (**Table 2** and **Supplementary Table 14**). Interestingly, in digestates from agricultural and central origin, nitrogen cycle-related processes (nitrate-reduction, ammonia-assimilation, ammonia-oxidation, denitrification, taurine-degradation, and nitrogen-fixation pathways) were more abundant in the enriched genera than the decaying ones, but the opposite situation occurred among digestates of urban origin (**Table 2** and **Supplementary Table 14**). After composting treatment, excepting the biosynthesis pathways involved in cell structure and the aromatic compounds degradation pathways, the inferred pathways ($n = 29$) had a similar distribution among the impacted genera (**Table 2** and **Supplementary Table 14**). Cell structure biosynthesis pathways were less abundant in the enriched genera after both types of treatment (AD and composting) than in the decaying ones. Aromatic compounds degradation pathways were more abundant in the enriched genera found among composts but not among those of digestates (**Table 2** and **Supplementary Table 14**).

DISCUSSION

OW recycling of sewage sludge and livestock effluents represent a major challenge because not only the bioenergy production but also environmental and sanitary issues must be considered. A key criterion in the assessment of the microbiological risks of OW recycling that remain poorly defined is the nature and structure of the various microbiomes getting into these processes and their potential health and ecological hazards. Here, the core and flexible components of the end product microbiomes were inferred, and the likelihood of these at being reproducibly found among highly diverse OW treatment plants was evaluated. Nine full-scale treatment plants distributed over France were selected to obtain a global view of microbiome bacterial taxa in the French OW treatment sector. These inferred bacterial taxa (from metabarcoding 16S rRNA gene datasets) are discussed below according to process type (AD, composting) and OW origins (fecal wastes from WWTP or agricultural sites, and food

products). Key functional traits of the bacterial genera impacted by the digestion processes were investigated to go further with the identification of key structuring bacterial processes.

According to NMDS ordinations of 16S rRNA gene metabarcoding datasets, two main bacterial community structures could be identified and found to match a segregation

TABLE 2 | Metabolic pathways recorded among the bacterial genera significantly impacted by the anaerobic digestion or composting processes as observed by the MACADAM approach.

Main hierarchical groups of pathways		Percentage of theoretical metabolic pathways ^a based on available sequenced genomes of isolates correlated to significantly impacted genera ^b in								
		Digestates from urban origin			Digestates from agricultural/central origin			Compost		
		TMP-EG (n = 252)	TMP-DG (n = 645)	R	TMP-EG (n = 321)	TMP-DG (n = 746)	R	TMP-EG (n = 753)	TMP-DG (n = 800)	R
Biosynthesis pathways	Cofactor ^c	23.0	15.0	1.5	21.8	11.7	1.9	13.8	13.5	1.0
	Amino acid	13.9	8.7	1.6	12.8	6.4	2.0	8.6	8.6	1.0
	Nucleotide	9.9	6.1	1.6	8.4	3.6	2.3	5.1	5.3	1.0
	Lipid	7.5	4.3	1.7	4.7	4.0	1.2	4.3	4.5	1.0
	Carbohydrates	4.8	5.1	0.9	2.2	4.8	0.5	4.5	5.4	0.8
	Aminoacyl-tRNAs-charging	1.2	0.5	2.4	0.9	0.1	9.0	0.4	0.4	1.0
	Secondary metabolite	1.2	2.2	0.5	0.9	3.0	0.3	2.7	2.4	1.1
	Cell structure	0.8	1.6	0.5	0.6	1.3	0.5	1.1	1.8	0.6
	Metabolic regulators	0.8	1.4	0.6	0.3	1.6	0.2	1.7	1.6	1.1
	Aromatic compounds	0.8	1.1	0.7	1.3	1.1	1.2	1.3	1.1	1.2
Degradation pathways	Alcohol	0.4	1.7	0.2	0.6	1.3	0.5	1.2	1.3	0.9
	Hormone	0.0	0.6	0	0.0	0.4	0	0.5	0.5	1.0
	Amino acid	6.4	7.4	0.9	8.1	8.3	0.9	7.6	7.8	1.0
	Nucleotide	4.8	3.6	1.3	4.4	2.7	1.6	2.8	3.0	0.9
	Fatty acid and lipid	0.0	1.4	0	0.6	1.2	0.5	1.2	1.4	0.9
	Carbohydrates	4.8	7.1	0.7	2.5	7.6	0.3	5.4	6.4	0.8
	Secondary metabolite	2.0	3.3	0.6	1.6	4.2	0.4	4.1	3.3	1.2
	Aromatic compounds	0.0	2.0	0	0.9	7.4	0.1	7.7	5.3	1.5
	Alcohol	1.2	1.7	0.7	1.3	1.3	1.0	1.3	1.5	0.9
	Carboxylates	3.2	5.0	0.6	4.1	5.6	0.7	5.1	5.1	1.0
C1 compounds ^d	C1 compounds ^d	2.0	2.3	0.9	2.8	2.7	1.0	2.1	2.4	0.9
	Non-carbon nutrients	1.2	2.3	0.5	2.2	4.0	0.6	2.4	2.9	0.8
	Sulfur metabolism									
Energy-metabolism	Phosphorus compounds	0.4	0.8	0.5	1.3	1.2	1.1	1.2	1.0	1.2
	Anaerobic respiration	1.6	1.9	0.8	1.9	3.2	0.6	2.8	2.9	1.0
	Aerobic respiration	1.2	1.6	0.8	2.8	1.6	1.8	1.6	1.6	1.0
	Fermentation	1.2	4.7	0.3	3.1	3.5	0.9	3.3	3.3	1.0
	Hydrogen-production	1.2	0.8	1.5	0.6	0.7	0.9	0.7	0.6	1.2
	Hydrogen-oxidation	0.4	0.5	0.8	0.3	0.4	0.8	0.4	0.4	1.0
Others	Nitrogen cycling processes ^e	1.6	3.1	0.5	4.1	2.7	1.5	2.8	2.8	1.0
	Detoxification ^f	1.2	1.9	0.6	2.2	2.0	1.1	1.9	1.9	1.0
	Nucleic acid processing	1.6	0.6	2.7	0.9	0.3	3.0	0.5	0.5	1.0

^aA minimum pathway score (PS) and pathway frequency score (PSF) of 0.5 was used for hypothetical metabolic pathways filtering.

^bTo see the list of organisms with the targeted metabolic pathway, see **Supplementary Tables 14A–F**. ^cCofactor biosynthesis are cobyrinate-diamide biosynthesis, vitamin biosynthesis, CoA biosynthesis, reductants, NAD metabolism, porphyrin compounds biosynthesis, quinone biosynthesis, polyprenyl biosynthesis, molybdenum cofactor, lipoate biosynthesis, and tetrapyrrole biosynthesis. ^dC1 compounds are CO₂ fixation, formaldehyde oxidation, methanol oxidation, formaldehyde assimilation, and methane oxidation. ^eNitrogen cycling processes consist of nitrate reduction, ammonia assimilation, ammonia oxidation, denitrification, taurine degradation, and nitrogen fixation. ^fDetoxification involves acid resistance, arsenate detoxification, methylglyoxal detoxification, antibiotic resistance, and reactive oxygen species degradation. Abbreviations: TMP-EG, theoretical metabolic pathways in the enriched genera after treatments; TMP-DG, theoretical metabolic pathways in the decaying genera; R, relative proportion of pathways per category between the enriched genera and the decaying ones.

between bacterial taxa from OW containing agricultural wastes or not (restricted to urban WWTP activated sludge). This segregation was also in line with a differentiation between their contents in micropollutants particularly the TEs. The urban samples (treated or not) had significantly more micropollutants including 14 TEs (Ag, Al, As, Cd, Ce, Co, Cr, Cu, Ga, Mo, Ni, P, Pb, and Ti), 4-nonylphenol, and 2 PAHs (phenanthrene and anthracene) than the agricultural and mixed (agricultural/urban; termed “central” in the text) samples. The 16S rRNA gene microbial community structures of the investigated OW urban samples were in agreement with previous studies (Juretschko et al., 2002; Wong et al., 2005; Hu et al., 2012; Díaz et al., 2018). Inferred 16S rRNA gene taxonomic allocations were predominantly among the *Proteobacteria*, *Bacteroidetes*, *Actinobacteria*, *Firmicutes*, and *Nitrospirae* phyla. *Proteobacteria* were reported to be recurrent among WWTP activated sludge and have been shown to represent almost 50% of the bacterial taxa (Juretschko et al., 2002; Wong et al., 2005; Hu et al., 2012). *Actinobacteria* were also previously mentioned as highly abundant organisms in such wastes and being favored by the high phosphorus content of activated sludge (Wagner et al., 1994). *Nitrospirae* known to be nitrite-oxidizing bacteria were previously considered as common in activated sludge but were not mentioned as being abundant core organisms (Saunders et al., 2016).

After AD of both categories of raw OWs (urban and central/agricultural categories), 16S rRNA gene inferred phyla with similar relative changes were observed. These shifts occurred whatever the raw OW origin. Both categories of digestates (urban and central/agricultural categories) were impacted by selective environmental changes, which led to a decrease in the relative abundance of *Proteobacteria* but an enrichment in *Firmicutes* (which nearly doubled in relative abundance). *Firmicutes* had been reported by several authors as one of the most abundant phyla in AD processes of activated sludge (Hu et al., 2012; Zhang et al., 2012; Abendroth et al., 2015; Díaz et al., 2018). Sundberg et al. (2013) also found that *Firmicutes* were the most prevalent in samples taken from reactors fed with various combinations of wastes (from slaughterhouses, restaurants, households, etc.). In this work, with 52.2% of V5–V6 rRNA gene reads, *Firmicutes* were dominant in the digestates from central (mixed)/agricultural sites. Furthermore, they were enriched by 2.6-fold in digestates from urban sites to reach an 18.4% relative abundance. These results seem to indicate that the AD conditions favored their growth whatever the OW origins. To go deeper into the understanding of these community changes, the impact of AD on the structuration of end product microbiomes was assessed by grouping 16S rRNA gene reads according to their allocations at the genus level, and making comparisons by DESEQ2. These analyses revealed that about one third of the total relative abundance (according to the mean normalized counts) of the digestate bacterial taxa were significantly affected by the AD treatment whatever the raw OW origin. The relative abundances of 50 and 60 genera were significantly impacted by AD as observed from, respectively, urban and central/agricultural digestates. However, differences in the shifts of the 16S rRNA gene reads of these taxa were found

related to the origin of the OW. In digestates of urban origin, this sorting mainly implied an increase in the relative abundance of OW urban taxa while those generated from agricultural/central OW led essentially to a reduction in the relative abundance of most of the initial OW taxa. This indicates that the coalescence of bacterial taxa of raw OW with the microbial community of the digesters will be largely dependent on their origin (and nature). In fact, urban OWs (mainly composed of activated sludge) were found to contain better fit communities for the AD process that could interact with the digesters' microbial communities. Bacterial taxa present in agricultural raw OW were outcompeted by the digesters' communities. This could be explained by the pre-treatment conditions performed in municipal wastewater facilities, leading to an increase of specialized bacterial taxa in the activated sludge. These taxa can perform multiple biodegradation processes (Shchegolkova et al., 2016) and can form heterogeneous structures termed flocs (Ju and Zhang, 2015). Functional inferences through FAPROTAX (Louca et al., 2016) were undertaken to highlight their potential contributions to the AD process.

The functional traits of the dominant microbial taxa inferred from the 16S rRNA gene datasets in the AD end products were investigated. Based on FAPROTAX analyses, functional traits such as chemoheterotrophy and fermentation were highly common among the taxa found in digestates of the urban and agricultural/central categories. Chemoheterotrophic bacteria are usually acting as primary decomposers (hydrolysis step), which are responsible for OW recycling in diverse ecosystems (Kämpfer et al., 1993). Fermentation, a widespread anaerobic pathway, is an essential step in AD process where fermentative bacteria convert simple monomers such as simple sugar, amino acids, and alcohols into short-chain fatty acids, primary alcohols, as well as H₂ and CO₂ (Cardinali-Rezende et al., 2009; Bajpai, 2017; Anukam et al., 2019). Interestingly, these functions (chemoheterotrophy and fermentation) were also previously reported by Wei et al. (2018) as the most dominant functions in activated sludge. This elucidates why bacterial genera from activated sludge could directly contribute to the observed diversity in the digestates. This observation is in line with the hypothesis of a selection of the most fit K-strategists (specialists) with ecologically relevant functions during the AD process. Also, these results confirmed that the changes in the AD community composition were driven by functional traits and less constrained by taxonomic relatedness.

To corroborate the above conclusion, closer analyses of the metabolic potential of bacterial communities were performed. Pathways inferred to be present in the enriched genera after treatment against those of the decreased ones were compared according to the origin/natures of the OWs. Interestingly, AD treatment was confirmed to impact similarly numerous metabolic pathways in both categories of OWs (urban and central/agricultural categories). Ten biosynthesis pathways (10/12) involved in the production of secondary metabolites, cell structure, metabolic regulators, biosynthesis, etc., and nine degradation pathways (9/11) (e.g., aromatic compounds and sulfur-containing compounds degradation) were affected similarly in both digestates from urban and agricultural/central categories. Nevertheless, some categories of pathways were

not impacted in the same way according to the OW origin. For example, in digestates from agricultural and central origin, nitrogen cycling processes (e.g., nitrate-reduction, ammonia-assimilation, denitrification, and nitrogen-fixation pathways) were more abundant in the enriched genera than in the decaying ones, but the opposite situation was observed for digestates of urban origin. These pathways (related to nitrogen cycle) have been reported by Wei et al. (2018) as central key functions in activated sludge. Thus, these pathways could be critical in the ecology of activated sludge but less important during AD.

The impact of composting per OW type on the bacterial communities was also investigated. After composting, a NMDS plot grouped together the 16S rRNA community profiles whatever the OW origin. Composting was found to drive bacterial communities toward similar genetic structures in all samples without differentiating OW inputs that had previously undergone AD. This conserved microbiome appeared to have been selected by highly selective constraints leading to a significant decrease in the relative abundance of *Bacteroidetes*, *Chlorobi*, *Chloroflexi*, *Lentisphaerae*, *Planctomycetes*, *Proteobacteria*, *Verrucomicrobia*, and *Spirochaetae* and the loss of six phyla (*Nitrospirae*, *Elusimicrobia*, *Acidobacteria*, *Saccharibacteria*, *TM6*, and *GOUTA4*) from the inputs OW. Other studies reported similar findings (Steger et al., 2007a; Song et al., 2014). In a compost pile, changes in physicochemical factors such as pH, temperature, moisture, and organic carbon were found to be correlated with bacterial growth. Microbial community structures were found to be affected by these evolving parameters (Cahyani et al., 2003; Liang et al., 2003; Tang et al., 2007; Zhang et al., 2011). Among these, temperature was shown to be highly selective (Palmisano and Barlaz, 1996; Steger et al., 2007a). Interestingly, with their large shift in relative abundance (4.3-fold) observed between OW and composts, *Actinobacteria* showed a major re-shuffling with an up to 20% increase. These are clearly key players in the composting process.

To go deeper into the understanding of these effects, changes in the composition of the compost bacterial taxa were investigated at genus level. A total of 141 genera were found to be affected by composting, with 105 genera showing a significant decrease (equivalent to 53.7% of the total relative abundance of the compost's bacterial taxa). Only three of these decaying genera were allocated to the *Actinobacteria*. However, 16 of the 36 enriched genera were part of this phylum. Previous studies have reported the persistence of *Actinobacteria* over the composting process from the thermophilic to the curing stage, regardless of the nature of treated OW (Danon et al., 2008). This bacterial group also constituted a significant part of the microbial community in the late composting stages (Herrmann and Shann, 1997; Tiquia et al., 2002; Hiraishi et al., 2003; Steger et al., 2007b). *Actinobacteria* are Gram-positive bacteria with high G + C contents in DNA, widely distributed in a diversity of ecological environments, including extreme ecosystem (Qin et al., 2011, 2019; Dhakal et al., 2017; Goodfellow et al., 2018). They can perform several important functions. Some *Actinobacteria* can synthesize a wide diversity of secondary

metabolites, which may promote survival in highly selective environments (Fang et al., 2017; Puttaswamygowda et al., 2019; Qin et al., 2019). Several studies have highlighted the importance of temperature as a selective factor of thermo-tolerant *Actinobacteria*. In fact, *Saccharomonospora viridis*, *Thermobifida fusca*, and *Thermobispora bispora* have been reported in the later composting stages. In our study, many genera of thermo-tolerant *Actinobacteria*, e.g., *Thermobifida*, *Actinomadura*, *Saccharomonospora*, *Arthrobacter*, *Thermomonospora*, and *Thermocrispum* were enriched in compost irrespective of the origin of the treated OW. *Actinobacteria* are capable of breaking down a wide range of animal wastes and plant debris particularly complex compounds (e.g., lignocelluloses), playing an important role in the later stages of composting (Herrmann and Shann, 1997; Wang et al., 2016). It is to be noted that thanks to the occurrences of bacterial genera degrading xenobiotics, composting offers the advantage of decreasing the concentration of chemical contaminants (Semple et al., 2001; Ghaly, 2011; Cascant et al., 2016; Ren et al., 2018). In line with these reports, genera of xenobiotic degrading *Actinobacteria*, e.g., *Arthrobacter*, were found to be enriched in the 16S rRNA gene datasets generated from composts DNA. To go deeper, closer analyses of functional potentials *per taxa* enriched after composting were performed. The functional profiles of the enriched genera suggested that the number of 16S rRNA copies of genera, e.g., *Arthrobacter*, *Alcanivorax*, *Rhodococcus*, *Brevibacterium*, etc., known to consist of many strains having functional traits involved in arsenate detoxification, dissimilatory arsenate reduction, plastic degradation, and oil bioremediation have significantly increased through composting. This further supports the idea that composting is an efficient bioremediation process.

CONCLUSION

In conclusion, AD and composting induced bacterial changes that were found driven by pH, temperature, but also the C- and energy sources which require specialized metabolic properties. These factors were more selective in composting than AD. Consequently, composting generated highly predictable community patterns whatever the OW origin. In AD, the relative contribution of the raw OW microbiomes on the digestate structural and functional bacterial community patterns was dependent on its origin. Bacterial genera from activated sludge were found to be well-suited for the AD process and contributed directly to the observed diversity observed in the digestates. Content in activated sludge among the OW was thus considered to be a critical factor for predicting the digestate bacterial contents. Our findings provided new insights into our understanding of bacterial coalescence phenomena in aerobic and anaerobic digesters. However, knowledge gap remains regarding potential health hazards associated with the emerging communities observed in the end products. These organic matter biodegradation processes being built from a large panel of exogenous communities, they might offer conditions

favorable for horizontal gene transfer (HGT) events such as the HGT of antibiotic resistance genes (ARG). The next step in these analyses will be to evaluate the frequency of these HGT.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: European Nucleotide Archive (<https://www.ebi.ac.uk/research>) under the accession number PRJEB40193.

AUTHOR CONTRIBUTIONS

AA, EB, LM, BC, and WG conceived and designed experiments. AA, EB, LM, SH, DP, ED, BC, and WG performed the experiments. AA, EB, BC, and WG performed the data curation and statistical analyses. AA, BC, and WG wrote the manuscript. All authors read and approved the final manuscript.

REFERENCES

- Abendroth, C., Vilanova, C., Günther, T., Luschign, O., and Porcar, M. (2015). Eubacteria and archaea communities in seven mesophile anaerobic digester plants in Germany. *Biotechnol. Biofuels* 8:87. doi: 10.1186/s13068-015-0271-6
- Ahmad, R., Jilani, G., Arshad, M., Zahir, Z. A., and Khalid, A. (2007). Bio-conversion of organic wastes for their recycling in agriculture: an overview of perspectives and prospects. *Ann. Microbiol.* 57, 471–479. doi: 10.1007/BF03175343
- Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biol.* 11:R106.
- Anukam, A., Mohammadi, A., Naqvi, M., and Granström, K. (2019). A review of the chemistry of anaerobic digestion: methods of accelerating and optimizing process efficiency. *Processes* 7:504. doi: 10.3390/pr7080504
- Atelg, M. R., Krisa, D., Kumar, G., Eskicioglu, C., Nguyen, D. D., Chang, S. W., et al. (2020). Biogas production from organic waste: recent progress and perspectives. *Waste Biomass Valorization* 11, 1019–1040. doi: 10.1007/s12649-018-00546-0
- Bajpai, P. (2017). “Basics of anaerobic digestion process,” in *Anaerobic Technology in Pulp and Paper Industry SpringerBriefs in Applied Sciences and Technology*, ed. P. Bajpai, (Singapore: Springer), 7–12. doi: 10.1007/978-981-10-4130-3_2
- Blanc, M., Marilley, L., Beffa, T., and Aragno, M. (1999). Thermophilic bacterial communities in hot composts as revealed by most probable number counts and molecular (16S rDNA) methods. *FEMS Microbiol. Ecol.* 28, 141–149. doi: 10.1111/j.1574-6941.1999.tb00569.x
- Bodenhausen, N., Horton, M. W., and Bergelson, J. (2013). Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS One* 8:e56329. doi: 10.1371/journal.pone.0056329
- Cahyani, V. R., Matsuya, K., Asakawa, S., and Kimura, M. (2003). Succession and phylogenetic composition of bacterial communities responsible for the composting process of rice straw estimated by PCR-DGGE analysis. *Soil Sci. Plant Nutr.* 49, 619–630. doi: 10.1080/00380768.2003.10410052
- Cardinali-Rezende, J., Debarry, R. B., Colturato, L. F. D. B., Carneiro, E. V., Chartone-Souza, E., and Nascimento, A. M. A. (2009). Molecular identification and dynamics of microbial communities in reactor treating organic household waste. *Appl. Microbiol. Biotechnol.* 84, 777–789. doi: 10.1007/s00253-009-2071-z
- Cascant, M. M., Sisouane, M., Tahiri, S., Krati, M. E., Cervera, M. L., Garrigues, S., et al. (2016). Determination of total phenolic compounds in compost by infrared spectroscopy. *Talanta* 153, 360–365. doi: 10.1016/j.talanta.2016.03.020

FUNDING

This study was partly funded by the ANR (Agence Nationale de la Recherche, France) DIGESTATE project ANR-15-CE340003-01, Bacter’eau project ANR-16-CE04-0014-01, and the “Auvergne-Rhône-Alpes” Region (France) “pack ambition recherche” project ITAS-MZ2M.

ACKNOWLEDGMENTS

We wish to thank J. P. Delgenes, N. Sertillanges, D. Montenach, A. Savoie (INRAe), M. Montels, and C. Chevassut-Rosset (CIRAD) who provided us the samples analyzed in this project.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.667043/full#supplementary-material>

- Chelius, M. K., and Triplett, E. W. (2001). The diversity of archaea and bacteria in association with the roots of *Zea mays* L. *Microb. Ecol.* 41, 252–263. doi: 10.1007/s002480000087
- Chen, Y., Cheng, J. J., and Creamer, K. S. (2008). Inhibition of anaerobic digestion process: a review. *Bioresour. Technol.* 99, 4044–4064. doi: 10.1016/j.biortech.2007.01.057
- Danon, M., Franke-Whittle, I. H., Insam, H., Chen, Y., and Hadar, Y. (2008). Molecular analysis of bacterial community succession during prolonged compost curing. *FEMS Microbiol. Ecol.* 65, 133–144. doi: 10.1111/j.1574-6941.2008.00506.x
- Day, M., and Shaw, K. (2001). “Biological, chemical and physical processes of composting,” in *Compost Utilization In Horticultural Cropping Systems*, eds P. J. Stoffella, and B. A. Kahn, (Boca Raton FL: Lewis Publication), 17–50.
- De Vrieze, J., Raport, L., Roume, H., Vilchez-Vargas, R., Jáuregui, R., Pieper, D. H., et al. (2016). The full-scale anaerobic digestion microbiome is represented by specific marker populations. *Water Res.* 104, 101–110. doi: 10.1016/j.watres.2016.08.008
- De Vrieze, J., Verstraete, W., and Boon, N. (2013). Repeated pulse feeding induces functional stability in anaerobic digestion. *Microb. Biotechnol.* 6, 414–424. doi: 10.1111/1751-7915.12025
- Demirel, B., and Scherer, P. (2008). The roles of acetotrophic and hydrogenotrophic methanogens during anaerobic conversion of biomass to methane: a review. *Rev. Environ. Sci. Biotechnol.* 7, 173–190. doi: 10.1007/s11157-008-9131-1
- Dhokal, D., Pokhrel, A. R., Shrestha, B., and Sohng, J. K. (2017). Marine rare actinobacteria: isolation, characterization, and strategies for harnessing bioactive compounds. *Front. Microbiol.* 8:1106. doi: 10.3389/fmicb.2017.01106
- Díaz, A. I., Oulego, P., Collado, S., Laca, A., González, J. M., and Díaz, M. (2018). Impact of anaerobic digestion and centrifugation/decanting processes in bacterial communities fractions. *J. Biosci. Bioeng.* 126, 742–749. doi: 10.1016/j.jbiosc.2018.05.024
- Fang, B.-Z., Salam, N., Han, M.-X., Jiao, J.-Y., Cheng, J., Wei, D.-Q., et al. (2017). Insights on the effects of heat pretreatment, pH, and calcium salts on isolation of rare actinobacteria from Karstic caves. *Front. Microbiol.* 8:1535. doi: 10.3389/fmicb.2017.01535
- Franke-Whittle, I. H., Walter, A., Ebner, C., and Insam, H. (2014). Investigation into the effect of high concentrations of volatile fatty acids in anaerobic digestion on methanogenic communities. *Waste Manag.* 34, 2080–2089. doi: 10.1016/j.wasman.2014.07.020
- Ghaly, (2011). Biodegradation of phenolic compounds in creosote treated wood waste by a composting microbial culture augmented with the fungus

- Thermoascus aurantiacus*. *Am. J. Biochem. Biotechnol.* 7, 90–103. doi: 10.3844/ajbbsp.2011.90.103
- González-Fernández, C., and García-Encina, P. A. (2009). Impact of substrate to inoculum ratio in anaerobic digestion of swine slurry. *Biomass Bioenergy* 33, 1065–1069. doi: 10.1016/j.biombioe.2009.03.008
- Goodfellow, M., Nouioui, I., Sanderson, R., Xie, F., and Bull, A. T. (2018). Rare taxa and dark microbial matter: novel bioactive actinobacteria abundant in Atacama Desert soils. *Antonie Van Leeuwenhoek* 111, 1315–1332. doi: 10.1007/s10482-018-1088-7
- Gou, C., Yang, Z., Huang, J., Wang, H., Xu, H., and Wang, L. (2014). Effects of temperature and organic loading rate on the performance and microbial community of anaerobic co-digestion of waste activated sludge and food waste. *Chemosphere* 105, 146–151. doi: 10.1016/j.chemosphere.2014.01.018
- Herrmann, R. F., and Shann, J. F. (1997). Microbial community changes during the composting of municipal solid waste. *Microb. Ecol.* 33, 78–85. doi: 10.1007/s002489900010
- Hiraishi, A., Narihira, T., and Yamanaka, Y. (2003). Microbial community dynamics during start-up operation of flowerpot-using fed-batch reactors for composting of household biowaste. *Environ. Microbiol.* 5, 765–776. doi: 10.1046/j.1462-2920.2003.00473.x
- Hoornweg, D., Bhada-Tata, P., and Kennedy, C. (2013). Environment: waste production must peak this century. *Nature* 502, 615–617. doi: 10.1038/502615a
- Hu, M., Wang, X., Wen, X., and Xia, Y. (2012). Microbial community structures in different wastewater treatment plants as revealed by 454-pyrosequencing analysis. *Bioresour. Technol.* 117, 72–79. doi: 10.1016/j.biortech.2012.04.061
- Hupfaut, S., Plattner, P., Wagner, A. O., Kaufmann, R., Insam, H., and Podmirseg, S. M. (2018). Temperature shapes the microbiota in anaerobic digestion and drives efficiency to a maximum at 45°C. *Bioresour. Technol.* 269, 309–318. doi: 10.1016/j.biortech.2018.08.106
- Insam, H., and de Bertoldi, M. (2007). “Chapter 3 microbiology of the composting process,” in *Waste Management Series Compost Science and Technology*, eds L. F. Diaz, M. de Bertoldi, W. Bidlingmaier, and E. Stentiford, (Amsterdam: Elsevier), 25–48. doi: 10.1016/S1478-7482(07)80006-6
- Ju, F., and Zhang, T. (2015). Bacterial assembly and temporal dynamics in activated sludge of a full-scale municipal wastewater treatment plant. *ISME J.* 9, 683–695. doi: 10.1038/ismej.2014.162
- Juretschko, S., Loy, A., Lehner, A., and Wagner, M. (2002). The microbial community composition of a nitrifying-denitrifying activated sludge from an industrial sewage treatment plant analyzed by the full-cycle rRNA approach. *Syst. Appl. Microbiol.* 25, 84–99. doi: 10.1078/0723-2020-00093
- Kämpfer, P., Steiof, M., Becker, P. M., and Dott, W. (1993). Characterization of chemoheterotrophic bacteria associated with the in situ bioremediation of a waste-oil contaminated site. *Microb. Ecol.* 26, 161–188.
- Kayhanian, M. (1994). Performance of a high-solids anaerobic digestion process under various ammonia concentrations. *J. Chem. Technol. Biotechnol.* 59, 349–352. doi: 10.1002/jctb.280590406
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D. (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl. Environ. Microbiol.* 79, 5112–5120. doi: 10.1128/AEM.01043-13
- Le Bars, M., Legros, S., Levard, C., Chaurand, P., Tella, M., Rovezzi, M., et al. (2018). Drastic change in zinc speciation during anaerobic digestion and composting: instability of nanosized Zinc Sulfide. *Environ. Sci. Technol.* 52, 12987–12996. doi: 10.1021/acs.est.8b02697
- Le Boulch, M., Déhais, P., Combes, S., and Pascal, G. (2019). The MACADAM database: a metabolic pathways database for microbial taxonomic groups for mining potential metabolic capacities of archaeal and bacterial taxonomic groups. *Database* 2019:baz049. doi: 10.1093/database/baz049
- Levén, L., Eriksson, A. R. B., and Schnürer, A. (2007). Effect of process temperature on bacterial and archaeal communities in two methanogenic bioreactors treating organic household waste. *FEMS Microbiol. Ecol.* 59, 683–693. doi: 10.1111/j.1574-6941.2006.00263.x
- Liang, C., Das, K. C., and McClendon, R. W. (2003). The influence of temperature and moisture contents regimes on the aerobic microbial activity of a biosolids composting blend. *Bioresour. Technol.* 86, 131–137. doi: 10.1016/S0960-8524(02)00153-0
- Louca, S., Parfrey, L. W., and Doebeli, M. (2016). Decoupling function and taxonomy in the global ocean microbiome. *Science* 353, 1272–1277. doi: 10.1126/science.aaf4507
- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *bioRxiv* [Preprint] doi: 10.1101/002832
- Lykidis, A., Chen, C.-L., Tringe, S. G., McHardy, A. C., Copeland, A., Kyrpides, N. C., et al. (2011). Multiple syntrophic interactions in a terephthalate-degrading methanogenic consortium. *ISME J.* 5, 122–130. doi: 10.1038/ismej.2010.125
- Mailler, R., Gasperi, J., Patureau, D., Vulliet, E., Delgenes, N., Danel, A., et al. (2017). Fate of emerging and priority micropollutants during the sewage sludge treatment: Case study of Paris conurbation. Part 1: contamination of the different types of sewage sludge. *Waste Manag.* 59, 379–393. doi: 10.1016/j.wasman.2016.11.010
- Marti, R., Bécouze-Lareure, C., Ribun, S., Marjolet, L., Souibgui, C. B., Aubin, J.-B., et al. (2017). Bacteriome genetic structures of urban deposits are indicative of their origin and impacted by chemical pollutants. *Sci. Rep.* 7:13219. doi: 10.1038/s41598-017-13594-8
- Moset, V., Poulsen, M., Wahid, R., Højberg, O., and Møller, H. B. (2015). Mesophilic versus thermophilic anaerobic digestion of cattle manure: methane productivity and microbial ecology. *Microb. Biotechnol.* 8, 787–800. doi: 10.1111/1751-7915.12271
- Noble, R., and Roberts, S. J. (2004). Eradication of plant pathogens and nematodes during composting: a review. *Plant Pathol.* 53, 548–568. doi: 10.1111/j.0032-0862.2004.01059.x
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2019). *vegan: Community Ecology Package*. Available online at: <https://CRAN.R-project.org/package=vegan> (accessed November 3, 2020).
- Palaniveloo, K., Amran, M. A., Norhashim, N. A., Mohamad-Fauzi, N., Peng-Hui, F., Hui-Wen, L., et al. (2020). Food waste composting and microbial community structure profiling. *Processes* 8:723. doi: 10.3390/pr8060723
- Palmaniano, A. C., and Barlaz, M. A. (1996). *Microbiology of Solid Waste*. Boca Raton, FL: CRC Press.
- Pohlert, T. (2020). *PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended*. Available online at: <https://CRAN.R-project.org/package=PMCMRplus> (accessed November 3, 2020).
- Puttaswamygowda, G. H., Olakkaran, S., Antony, A., and Kizhakke Purayil, A. (2019). “Chapter 22 – present status and future perspectives of marine actinobacterial metabolites,” in *Recent Developments in Applied Microbiology and Biochemistry*, ed V. Buddolla, (Cambridge, MA: Academic Press), 307–319. doi: 10.1016/B978-0-12-816328-3.00022-2
- Pycke, B., Etchebehere, C., Van de Caveye, P., Negroni, A., Verstraete, W., and Boon, N. (2011). A time-course analysis of four full-scale anaerobic digesters in relation to the dynamics of change of their microbial communities. *Water Sci. Technol.* 63, 769–775. doi: 10.2166/wst.2011.307
- Qin, S., Li, W.-J., Klenk, H.-P., Hozzein, W. N., and Ahmed, I. (2019). Editorial: actinobacteria in special and extreme habitats: diversity, function roles and environmental adaptations, second edition. *Front. Microbiol.* 10:944. doi: 10.3389/fmicb.2019.00944
- Qin, S., Xing, K., Jiang, J.-H., Xu, L.-H., and Li, W.-J. (2011). Biodiversity, bioactive natural products and biotechnological potential of plant-associated endophytic actinobacteria. *Appl. Microbiol. Biotechnol.* 89, 457–473. doi: 10.1007/s00253-010-2923-6
- Reinikainen, O., and Herranen, M. (2001). Different methods for measuring compost stability and maturity. *Acta Hort.* 549, 99–104. doi: 10.17660/ActaHortic.2001.549.10
- Ren, X., Zeng, G., Tang, L., Wang, J., Wan, J., Wang, J., et al. (2018). The potential impact on the biodegradation of organic pollutants from composting technology for soil remediation. *Waste Manag.* 72, 138–149. doi: 10.1016/j.wasman.2017.11.032
- Rivière, D., Desvignes, V., Pelletier, E., Chaussonnerie, S., Guermazi, S., Weissenbach, J., et al. (2009). Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. *ISME J.* 3, 700–714. doi: 10.1038/ismej.2009.2
- Saady, N. M. C., and Massé, D. I. (2015). A start-up of psychrophilic anaerobic sequence batch reactor digesting a 35 % total solids feed of

- dairy manure and wheat straw. *AMB Express* 5:144. doi: 10.1186/s13568-015-0144-1
- Saunders, A. M., Albertsen, M., Vollertsen, J., and Nielsen, P. H. (2016). The activated sludge ecosystem contains a core community of abundant organisms. *ISME J.* 10, 11–20. doi: 10.1038/ismej.2015.117
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541. doi: 10.1128/AEM.01541-09
- Semple, K. T., Reid, B. J., and Fermor, T. R. (2001). Impact of composting strategies on the treatment of soils contaminated with organic pollutants. *Environ. Pollut.* 112, 269–283. doi: 10.1016/s0269-7491(00)00099-3
- Sertillanges, N., Haudin, C.-S., Bourdat-Deschamps, M., Bernet, N., Serre, V., Danel, A., et al. (2020). Process type is the key driver of the fate of organic micropollutants during industrial scale treatment of organic wastes. *Sci. Total Environ.* 734:139108. doi: 10.1016/j.scitotenv.2020.139108
- Shchegolkova, N. M., Krasnov, G. S., Belova, A. A., Dmitriev, A. A., Kharitonov, S. L., Klimina, K. M., et al. (2016). Microbial community structure of activated sludge in treatment plants with different wastewater compositions. *Front. Microbiol.* 7:90. doi: 10.3389/fmicb.2016.00090
- Song, C., Li, M., Jia, X., Wei, Z., Zhao, Y., Xi, B., et al. (2014). Comparison of bacterial community structure and dynamics during the thermophilic composting of different types of solid wastes: anaerobic digestion residue, pig manure and chicken manure. *Microb. Biotechnol.* 7, 424–433. doi: 10.1111/1751-7915.12131
- Steger, K., Jarvis, Å., Vasara, T., Romantschuk, M., and Sundh, I. (2007a). Effects of differing temperature management on development of Actinobacteria populations during composting. *Res. Microbiol.* 158, 617–624. doi: 10.1016/j.resmic.2007.05.006
- Steger, K., Sjögren, Å.M., Jarvis, Å., Jansson, J. K., and Sundh, I. (2007b). Development of compost maturity and Actinobacteria populations during full-scale composting of organic household waste. *J. Appl. Microbiol.* 103, 487–498. doi: 10.1111/j.1365-2672.2006.03271.x
- Sundberg, C., Al-Soud, W. A., Larsson, M., Alm, E., Yekta, S. S., Svensson, B. H., et al. (2013). 454 pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas digesters. *FEMS Microbiol. Ecol.* 85, 612–626. doi: 10.1111/1574-6941.12148
- Supaphol, S., Jenkins, S. N., Intomo, P., Waite, I. S., and O'Donnell, A. G. (2011). Microbial community dynamics in mesophilic anaerobic co-digestion of mixed waste. *Bioresour. Technol.* 102, 4021–4027. doi: 10.1016/j.biortech.2010.11.124
- Tang, J.-C., Shibata, A., Zhou, Q., and Katayama, A. (2007). Effect of temperature on reaction rate and microbial community in composting of cattle manure with rice straw. *J. Biosci. Bioeng.* 104, 321–328. doi: 10.1263/jbb.104.321
- Tian, W., Sun, Q., Xu, D., Zhang, Z., Chen, D., Li, C., et al. (2013). Succession of bacterial communities during composting process as detected by 16S rRNA clone libraries analysis. *Int. Biodeterior. Biodegrad.* 78, 58–66. doi: 10.1016/j.ibiod.2012.12.008
- Tiquia, S. M., Wan, H. C., and Tam, N. F. Y. (2002). Microbial population dynamics and enzyme activities during composting. *Compost Sci. Util.* 10, 150–161. doi: 10.1080/1065657X.2002.10702075
- Trably, E., Delgènes, N., Patureau, D., and Delgènes, J. P. (2004). Statistical tools for the optimization of a highly reproducible method for the analysis of polycyclic aromatic hydrocarbons in sludge samples. *Int. J. Environ. Anal. Chem.* 84, 995–1008. doi: 10.1080/03067310412331298412
- Vanwongerghem, I., Jensen, P. D., Dennis, P. G., Hugenholtz, P., Rabaey, K., and Tyson, G. W. (2014). Deterministic processes guide long-term synchronised population dynamics in replicate anaerobic digesters. *ISME J.* 8, 2015–2028. doi: 10.1038/ismej.2014.50
- Voisin, J., Cournoyer, B., Marjolet, L., Vienney, A., and Mermillod-Blondin, F. (2020). Ecological assessment of groundwater ecosystems disturbed by recharge systems using organic matter quality, biofilm characteristics, and bacterial diversity. *Environ. Sci. Pollut. Res.* 27, 3295–3308. doi: 10.1007/s11356-019-06971-5
- Wagner, A. O., Reitschuler, C., and Illmer, P. (2014). Effect of different acetate:propionate ratios on the methanogenic community during thermophilic anaerobic digestion in batch experiments. *Biochem. Eng. J.* 90, 154–161. doi: 10.1016/j.bej.2014.05.014
- Wagner, M., Erhart, R., Manz, W., Amann, R., Lemmer, H., Wedi, D., et al. (1994). Development of an rRNA-targeted oligonucleotide probe specific for the genus *Acinetobacter* and its application for in situ monitoring in activated sludge. *Appl. Environ. Microbiol.* 60, 792–800.
- Wang, C., Dong, D., Wang, H., Müller, K., Qin, Y., Wang, H., et al. (2016). Metagenomic analysis of microbial consortia enriched from compost: new insights into the role of Actinobacteria in lignocellulose decomposition. *Biotechnol. Biofuels* 9:22. doi: 10.1186/s13068-016-0440-2
- Wang, P., Yu, Z., Zhao, J., and Zhang, H. (2018). Do microbial communities in an anaerobic bioreactor change with continuous feeding sludge into a full-scale anaerobic digestion system? *Bioresour. Technol.* 249, 89–98. doi: 10.1016/j.biortech.2017.09.191
- Wang, Y., Zhang, Y., Wang, J., and Meng, L. (2009). Effects of volatile fatty acid concentrations on methane yield and methanogenic bacteria. *Biomass Bioenergy* 33, 848–853. doi: 10.1016/j.biombioe.2009.01.007
- Wei, Z., Liu, Y., Feng, K., Li, S., Wang, S., Jin, D., et al. (2018). The divergence between fungal and bacterial communities in seasonal and spatial variations of wastewater treatment plants. *Sci. Total Environ.* 628–629, 969–978. doi: 10.1016/j.scitotenv.2018.02.003
- Werner, J. J., Knights, D., Garcia, M. L., Scalfone, N. B., Smith, S., Yarasheski, K., et al. (2011). Bacterial community structures are unique and resilient in full-scale bioenergy systems. *Proc. Natl. Acad. Sci. U.S.A.* 108, 4158–4163. doi: 10.1073/pnas.1015676108
- Williams, J., Williams, H., Dinsdale, R., Guwy, A., and Esteves, S. (2013). Monitoring methanogenic population dynamics in a full-scale anaerobic digester to facilitate operational management. *Bioresour. Technol.* 140, 234–242. doi: 10.1016/j.biortech.2013.04.089
- Wong, M.-T., Mino, T., Seviour, R. J., Onuki, M., and Liu, W.-T. (2005). In situ identification and characterization of the microbial community structure of full-scale enhanced biological phosphorous removal plants in Japan. *Water Res.* 39, 2901–2914. doi: 10.1016/j.watres.2005.05.015
- Yu, Z., Morrison, M., and Schanbacher, F. L. (2010). “Production and utilization of methane biogas as renewable fuel,” in *Biomass to Biofuels*, eds A. A. Vertès, N. Qureshi, H. P. Blaschek, and H. Yukawa, (Hoboken, NJ: Blackwell Publishing Ltd), 403–433. doi: 10.1002/9780470750025.ch20
- Zhang, J., Zeng, G., Chen, Y., Yu, M., Yu, Z., Li, H., et al. (2011). Effects of physico-chemical parameters on the bacterial and fungal communities during agricultural waste composting. *Bioresour. Technol.* 102, 2950–2956. doi: 10.1016/j.biortech.2010.11.089
- Zhang, L., Zhang, H., Wang, Z., Chen, G., and Wang, L. (2016). Dynamic changes of the dominant functioning microbial community in the compost of a 90-m³ aerobic solid state fermentor revealed by integrated meta-omics. *Bioresour. Technol.* 203, 1–10. doi: 10.1016/j.biortech.2015.12.040
- Zhang, T., Shao, M.-F., and Ye, L. (2012). 454 Pyrosequencing reveals bacterial diversity of activated sludge from 14 sewage treatment plants. *ISME J.* 6, 1137–1147. doi: 10.1038/ismej.2011.188
- Zhang, W., Werner, J. J., Agler, M. T., and Angenent, L. T. (2014). Substrate type drives variation in reactor microbiomes of anaerobic digesters. *Bioresour. Technol.* 151, 397–401. doi: 10.1016/j.biortech.2013.10.004

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Aigle, Bourgeois, Marjolet, Houot, Patureau, Doelsch, Cournoyer and Galia. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.