



HAL
open science

Assessing sensitivity and persistence of updated initial conditions through Particle filter and EnKF for streamflow forecasting

Gaia Piazzzi, Guillaume Thirel, Charles Perrin

► **To cite this version:**

Gaia Piazzzi, Guillaume Thirel, Charles Perrin. Assessing sensitivity and persistence of updated initial conditions through Particle filter and EnKF for streamflow forecasting. EGU General Assembly 2020, May 2020, Vienna, Austria. 10.5194/egusphere-egu2020-18694 . hal-03266313

HAL Id: hal-03266313

<https://hal.inrae.fr/hal-03266313>

Submitted on 21 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU2020-18694

<https://doi.org/10.5194/egusphere-egu2020-18694>

EGU General Assembly 2020

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Assessing sensitivity and persistence of updated initial conditions through Particle filter and EnKF for streamflow forecasting

Gaia Piazza, Guillaume Thirel, and Charles Perrin

Université Paris-Saclay, INRAE, HYCAR Research Unit, Antony, France (gaia.piazza@inrae.fr)

Skillful streamflow forecasts provide a key support to several water-related applications. Ensemble forecasting systems are gaining a widespread interest, since they allow accounting for different sources of uncertainty. Because of the critical impact of the initial conditions (ICs) on the forecast accuracy, it is essential to improve their estimates via data assimilation (DA). This study aims at assessing the sensitivity of the DA-based estimation of forecast ICs to several sources of uncertainty and to the update of different model states and parameters of a conceptual rainfall-runoff model. The performance of two sequential ensemble-based techniques are compared, namely Ensemble Kalman filter and Particle filter, in terms of both efficiency and temporal persistence of the updating effect through the assimilation of observed discharges at the forecast time. Several experiments specifically address the impact of the meteorological, model state and parameter uncertainties over 232 catchments in France. Results show that the benefit of the DA-based estimation of ICs for forecasting is the largest when focusing on the level of the model routing store, which is the internal state the most correlated to streamflow. While the EnKF-based forecasts outperform the PF-based ones when accounting for the meteorological uncertainty, the representation of the model state uncertainty allows greatly improving the accuracy of the PF-based predictions, with a longer-lasting updating effect (up to 10 days). Conversely, the forecasting skill is undermined when accounting for the parameter uncertainty, due to the change in the hydrological responsiveness through the update of both the production and routing store levels. A further effort is focused on assessing the impact of the spatial resolution of the hydrological model on the predictive accuracy of DA-based streamflow forecasts.