

Tree diversity effects on forest productivity: disentangling the effects of tree species addition vs. substitution

Maude Toïgo, Bastien Castagneyrol, Hervé Jactel, Xavier Morin, Céline Meredieu

► To cite this version:

Maude Toïgo, Bastien Castagneyrol, Hervé Jactel, Xavier Morin, Céline Meredieu. Tree diversity effects on forest productivity: disentangling the effects of tree species addition vs. substitution. 2021. hal-03276166

HAL Id: hal-03276166 https://hal.inrae.fr/hal-03276166

Preprint submitted on 1 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 TREE DIVERSITY EFFECTS ON FOREST PRODUCTIVITY: 2 DISENTANGLING THE EFFECTS OF TREE SPECIES ADDITION VS. 3 SUBSTITUTION

4 AUTHORS

- 5 Toïgo Maude^{a,b}, Castagneyrol Bastien^b, Jactel Hervé^b, Morin Xavier^a, Meredieu Celine^b
- ^aCEFE UMR 5175, CNRS Université de Montpellier Université Paul-Valéry Montpellier EPHE– IRD,
 1919 Route de Mende, F-34293 Montpellier, France
- 8 ^b INRAE, Univ. Bordeaux, BIOGECO, F-33612 Cestas, France
- 9
- 10 Correspondence author
- 11 Toïgo Maude
- 12 °CEFE UMR 5175, CNRS Université de Montpellier Université Paul-Valéry Montpellier EPHE–
- 13 IRD, 1919 Route de Mende, F-34293 Montpellier, France
- 14 maude.toigo@gmail.com

15 ABSTRACT

- 16 1. Mixture effect on stand productivity is usually apprehended through a substitutive approach, 17 whereby productivity in mixed stands is compared to productivity in monocultures, at equivalent stand 18 density. This approach has proved that in many cases mixed stands perform better than monospecific 19 forests, however, we do not yet have a solid theory about species behaviour in the mixture or even 20 guidelines for combining species. The addition of a second tree species to an existing mono-specific 21 stand has received much less consideration. Yet, this approach has the potential to separate the 22 facilitation effect from the complementarity effect.
- 23 2. We compared the effect of tree species substitution vs. addition on the productivity of maritime
 24 pine and silver birch in a young tree diversity experiment implemented in 2008 in SW France.
- 3. Substituting pines with birches to create two-species mixtures resulted in an increase of tree productivity at stand level beyond what was expected from monocultures (i.e., overyielding). In contrast, creating mixture through the addition of birches to pine stands had no effect on the maritime pine stand productivity (transgressive mixture effect not significant). This absence of effect is produced by two distinct density-dependence responses at an individual level.

4. Our results allow clarifying the cases in which a mixed stand can be considered as an alternative to
a monoculture of a productive species. In particular, the addition of a pioneer and soil low-demanding
species during young developmental stages is a possibility to diversify the stand and potentially to
increase ecosystem services without altering the productivity of the target species.

34 KEYWORDS

35 Betula pendula; Pinus pinaster; biodiversity; ecosystem functioning; overyielding; transgressive 36 overyielding; forest

37 INTRODUCTION

38 1§ CHALLENGES AND DETERMINANTS OF MIXED PLANTATIONS.

39 Despite ample evidence that mixed stands provide more ecosystem services than monospecific forests 40 under various ecological conditions (Baeten et al., 2019), most planted forests are still managed as 41 monocultures. Moving towards ecologically intensive and sustainable forest management requires a 42 sound understanding of the drivers that could improve or hamper the benefits of mixed forests (Felton 43 et al., 2010). Tree species diversity has well documented positive effects on tree productivity (Gamfeldt 44 et al., 2013). Such positive effects are driven by both complementarity and selection effects (Loreau & 45 Hector, 2001). Complementarity mostly refers to niche partitioning processes whereby mixed stands 46 are better able to capture resources than monospecific stands (Jucker et al., 2015), and to facilitation, where one species in the mixture benefit to the others, e.g. via improved resource quality (N-fixing 47 species) acquisition (water uplifting), or protection against herbivores (Caspersen et al., 2018; Kunz et 48 49 al., 2019). Selection effect refers to situations where a highly productive species recruited in the mixed 50 stand drives positive mixture effect (Fox, 2005; Loreau & Hector, 2001). However, recent studies have 51 highlighted that positive diversity-productivity relationship is strongly context-dependent. For 52 instance, species functional characteristics or stand structure can modify the shape and strength of the 53 diversity-productivity relationship (Forrester, 2014; Grossman et al., 2017). Disentangling drivers of 54 the mixture effect require innovative conceptual framework supported by novel experimental approaches based on stand density, a major component of stand structure that can be controlled by 55 56 thinning operations.

57 2§ STAND DENSITY A KEY DETERMINANT OF MIXTURE EFFECT ON STAND PRODUCTIVITY

58 Stand density influences the degree of canopy closure, which in turn participate in the regulation of 59 light transmittance, the interception of water precipitations, belowground competition for water and 60 can modify understory microclimate, understory vegetation, and soil biodiversity (Baeten et al., 2019; 61 Gaudio et al., 2011; Henneron et al., 2017; Ligot et al., 2014; Perot et al., 2017). Stand density is also a 62 major driver of tree-tree competition, being used to calculate several competition indices in forest 63 (Biging & Dobbertin, 1992). Due to the considerable effects of tree density on canopy packing and abiotic factors in forest stands, it is surprising that only a few studies addressing the effect of tree 64 diversity on productivity in temperate forest explicitly questioned the importance of stand density 65 (Forrester, 2014; Jucker et al., 2016). Yet, complementarity among species and intra-specific 66 67 competition both intensify with stand density. This was documented in mixed stands of latesuccessional species (Amoroso & Turnblom, 2006; Forrester et al., 2013) of slow- with fast-growing 68 69 tree species (Condés et al., 2013) and of species with contrasted shade tolerance (del Rio & Sterba, 70 2009). However, it remains unclear how the mixture effect can possibly be modified by stand density, 71 especially in young plantations of fast-growing tree species.

72 3§ CONTROLLING STAND DENSITY TO COMPARE MONOCULTURES TO MIXED STANDS: OVERYIELDING,

73 THE CLASSICAL INDEX BASED ON SPECIES SUBSTITUTION

74 Our understanding of the diversity-productivity relationship in mixed forests is further hampered by 75 unresolved methodological issues. The net biodiversity effect generally simply compares the observed 76 productivity of a mixture to a theoretical mixture assembled with the same proportion of trees drawn 77 from the component monocultures (Loreau, 1998; Loreau & Hector, 2001). As such, overyielding can 78 be seen as a measure of changes in stand productivity due to the substitution of a species by others. 79 Estimating the effect of species mixture on productivity through overyielding has several advantages. 80 First, it provides a quantitative estimate of the net biodiversity effect on stand productivity (Tobner et 81 al., 2016). Second, because it compares the productivity of the mixture to the weighted productivity 82 of the component monocultures, it allows addressing whether the mixture performs better than the average of monocultures (overyielding) or the most productive monoculture (transgressive 83 84 overyielding).

4§ LIMITATIONS LINKED TO SPECIES SUBSTITUTION AND ITS RELATED OVERYIELDING

The use of overyielding estimate has also several shortcomings. First, because it is inherently defined at the stand level, overyielding does not account for species-specific responses to tree diversity and knowing which species benefits or not from the mixture is of primary importance, particularly when it comes to harvest species at different times because of differences in growth patterns. Still the effects of tree diversity may not be symmetrical (del Rio & Sterba, 2009), which is a major concern to understand the functioning of mixed species forests. As a consequence, considering the mixture effect on species productivity and on individual tree productivity is a first step to disentangle the mechanisms underlying the diversity-productivity relationship (Nadrowski et al., 2010). Moreover, from a practical
point of view, the conversion of monocultures to mixed stands through species substitution is not
without management problems. On the one hand, the silviculture of mixed stands, particularly in cases
of intimate mixing, is complicated by the difference in growth rates of the different species and the
lack of knowledge about the optimum growing space for trees of each species. On the other hand,
wood product processing chains are often specialized in a limited number of species and may not be
able to offer a market for substitute species.

100 5§ SPECIES ADDITION AS AN ALTERNATIVE TO SPECIES SUBSTITUTION

101 An alternative to species substitution is the addition of a new species within an existing stand. A species 102 addition could be less constraining than species substitution by making it possible to keep the same 103 harvesting rate for the target tree species, for example in the case of alternate-row mixing. Second if 104 resource complementarity cannot be distinguished from facilitation through species substitution, any 105 gain of productivity observed through species addition should be the signature of facilitative processes 106 (i.e. comparison of one tree species productivity with vs. without any heterospecific neighbours). So, 107 either species addition or substitution should be considered to design and manage mixed-species 108 forests and dedicated experiments are needed to disentangle their specific effects on productivity.

109 5§ OBJECTIVES AND TESTED HYPOTHESES

Using a long-term tree diversity experiment, we experimentally uncoupled the effect of species 110 111 addition vs. substitution on forest stand productivity while controlling for stand and species-specific 112 density to gain further insight into the mechanisms underlying the effect of species addition and 113 substitution. We focused on two-species mixtures of maritime pine (Pinus pinaster Ait.) and silver birch 114 (Betula pendula Roth) at two stand densities. Although the two species studied are fast-growing 115 species, they are nevertheless distinct in terms of growth dynamics and tree sizes. In the case of species 116 substitution, we expected a positive global mixture effect (ME) with a positive specific effect for both 117 pine and birch. By contrast we anticipated a negative transgressive effect (TME) as birch is notably less 118 productive than maritime pine in the local conditions of the experiment. In the case of species addition, we hypothesized opposite patterns of response: a negative ME due to increased competition between 119 120 trees (due to higher tree density) but a positive TME due to a tree packing effect and a weak 121 competition from silver birch in pine stands. Lastly, we expected that all mixture effects would intensify 122 with stand density.

123 METHODS

Both maritime pine and silver birch are light demanding, fast-growing tree species and native to the site. The area of distribution of maritime pine is mainly restricted to Spain, the south-west of France and the north-west of Italy. Maritime pine is a highly drought tolerant species and a major species of production in France yielded exclusively in monoculture. Conversely silver birch is widely distributed across Europe from the Atlantic to eastern Siberia. Silver birch is yielded in the Northern and Eastern Europe and despite the interest shown by these countries, in the Atlantic this tree species is depreciated (Hynynen et al., 2010).

131 EXPERIMENTAL DESIGN

The ORPHEE experiment is located 40km south of Bordeaux (44°440 N, 00° 460 W) and belongs to the 132 133 worldwide Tree Diversity Network (TreeDivNet). The experimental plantation was established in 2008 134 on a clear cut of former maritime pine stands on a sandy podzol. Stumps were removed and the site 135 was ploughed and fertilized with phosphorus and potassium before planting. In total, 25 600 trees of 136 five native species (Silver birch, Betula pendula; pedunculate oak, Quercus robur; Pyrenean oak, 137 Quercus pyrenaica; holm Oak, Quercus ilex and maritime pine, Pinus pinaster) were planted within a 12ha area. Eight blocks were established, with 32 plots in every block, corresponding to the 31 possible 138 139 combinations of one to five species, with an additional replicate of the combination of five species. 140 Each plot contained 10 rows of 10 trees planted 2 m apart, resulting in 100 trees per plot, with a plot 141 area of 400 m². The total initial stand density was therefore 2500 tree per hectare in each plot. Tree 142 species mixtures were established according to a substitutive design, keeping the total tree number 143 (n=100) equal in all plots. Within plots, individual trees from different species were planted in a regular 144 alternate pattern, such that a tree from a given species had at least one neighbour from each of the 145 other species within a 2 m radius. Plots were three meters apart and were randomly distributed within 146 blocks. The entire experimental site was fenced to prevent grazing by mammalian herbivores.

147 PLOT SELECTION

148 The present study analyses growth data collected in 2014 (7 years-old) on the target trees at the center 149 of the plots in order to avoid edge effects (measured planting locations = 36). Importantly, at this time 150 of plot development, oak trees were on average 112 cm high and have a negligible growth in diameter, 151 whereas pines and birches were on average five times taller than oaks (563 and 510 cm high, 152 respectively). As a consequence, oak trees were confounded with the understorey vegetation. By 153 considering oak seedlings as part of the understorey vegetation we therefore focused solely on birch 154 and pine growth. However we do not deny the existence of belowground interactions as the 155 understorey can represent a large part of the fine root biomass in maritime pine stands (Bakker et al.,

2006). But the three oak species represent only a few individuals among the 25 species found in the 156 157 understorey (i.e. the most common are Molinia caerulae, Ulex minor and Pteridium aquilinum Corcket 158 et al., 2020), which allows us to reasonably assume that the impact of these relatively few oak individuals on the productivity of pine and birch at these developmental stages is negligible. We tested 159 160 the effect of species addition and substitution on tree and stand volume by selecting monocultures and mixed stands of birch and pine at equal stand density: the "high-density plots" (2500 t/ha) had 161 162 100 pines or 100 birches for monoculture plots or a mixture of 50 pines and 50 birches. The "medium-163 density plots" (1250t/ha) had 50 pines or 50 birches for monoculture plots, or a mixture of 25 pines 164 and 25 birches. We completed the sampling by selecting "low-density plots" as monocultures (625 165 t/ha) with 25 pines or 25 birches (figure 1). To avoid bias when comparing volume of mixed stands and monocultures we eventually selected plots with less than 15% of dead trees as an optimal balance 166 167 between the number of plots per treatments and the number of trees per plot (Supporting Information 168 Table S1).



170

Figure 1 Schematic representation of the experimental treatments consisting in three levels of stand density (low: 625t/ha, medium: 1250t/ha and high: 2500t/ha) and stand compositions, from left to right: *Pinus pinaster* in monocultures, mixed *Betula pentula - P. pinaster* stands (proportion of 50% of each species) and monoculture stands of *B. pendula*. Arrows indicate the between-treatment comparisons disentangling heterospecific addition (solid arrows) and species substitution (black arrows). Arrows are numbered according to the different experimental treatments compared in the result section.

178 DENDROMETRIC DATA

179 We measured the height of every 36 innermost planted trees at the center of every plot using a 180 graduate pole, each year from 2008 to 2014. We therefore measured 36, 18 or 9 pines or birches in 181 the high, medium and low-density plots, respectively. We also measured circumferences at 1.30m 182 from 2012 to 2014 on 7 randomly chosen pines and 7 randomly chosen birches per plot, irrespective 183 of plot composition. We used height-circumference relationships to estimate the circumferences of trees that have not been measured in 2014 (Supporting Information Figure S1), then we estimated 184 185 tree volume following the generic model developed by (Deleuze et al., 2014). We assigned a minimum 186 volume of $0.01m^3$ /ha to the few trees below 1.30m in height (corresponding to the minimum volume 187 found in the data set). Eventually we estimated dimensions of missing trees by averaging diameter, 188 height and volume of trees in the plot. Given the negligible tree dimensions at the time of plantation (2008) volumes of trees in 2014 were used as a measure of tree productivity (i.e. volume increment: 189 190 VI). Stand dendrometric characteristics are summarized in table 1.

191

192 Table 1 Mean (minimal-maximal) values of tree height, tree circumference and plot basal area of

193 maritime pine (*Ppin*) and silver birch (*Bpen*) in monocultures (*mo*) and mixed stands (*mx*) at the three

194 stand densities studied: low (625 t/ha), medium (1250 t/ha) and high (2500 t/ha).

		Stand density				
		Low	Medium		High	
		Мо	Мо	Mx	Мо	Mx
Plot basal area (m²/ha)	Bpen	1.41	2.39		3.55	
		0.84-2.25	0.80-4.62		1.71-5.10	
	Ppin	7.22	11.4		17.2	
		6.17-8.30	4.69-14.7		11.3-21.3	
	Bpen + Ppin			7.42		11.2
				4.54-9.10		8.24-14.7
Tree height (cm)	Bpen	526	537	513	517	508
		85-764	120-900	106-795	95-791	148-729
	Ppin	570	575	581	571	567
		232-706	133-801	250-780	200-810	161-795
Tree circumference at	Bpen	16.6	14.9	13.3	13.0	12.0
breast height (cm)		2.4-26.7	1.52-28.2	1.5-28.5	2.3-24.2	1.1-22.5
	Ppin	37.6	33.4	36.1	29.1	30.6
		9.5-51	0.7-50.2	10.1-49.9	5.3-45.2	2.6-47

195

196 TRANSGRESSIVE MIXTURE EFFECT AND MIXTURE EFFECTS FOR SPECIES SUBSTITUTION AND ADDITION

197 We calculated two integrated indices of mixture effect for heterospecific addition and species 198 substitution: mixture effect (ME) and transgressive mixture effect (TME). These indices were adapted 199 from the classic calculation of transgressive overyielding and overyielding. Transgressive overyielding 200 and overyielding are two standardized indices of mixture effect on stand productivity calculated by 201 comparing monocultures to mixed stands at a same stand density (i.e. in case of species substitution). 202 Yet, the major difference between species substitution and species addition is that total stand density 203 increases from monoculture to mixed stands in an additive design, while it is kept constant in a 204 substitutive design. It follows that the reference monoculture used to calculate ME and TME differed 205 between additive and substitutive designs. We calculated TME the same way for species substitution 206 and species addition at medium and high stand density (n = 2500 t/ha and n = 1250 t/ha) by comparing 207 the mean total stand volume increment (SVI) of stands between mixed stands (mx) and monoculture 208 stands (mo) of the most productive species, i.e. maritime pine (Fig. 2):

209 TME =
$$(SVI_{mx} - SVI_{mo.pine}) / SVI_{mo.pine}$$
 Eq. 1

- Where SVI_{mx} was the stand volume increment since plantation of mixed stands averaged per block;
 SVI_{mo} the stand volume increment in monoculture of birch or pine averaged per block.
- The mixture effect (ME) was calculated for each block separately at medium and high levels of stand
 density (n = 2500 t/ha and n = 1250 t/ha) as:

214
$$ME = (SVI_{mx} - SVI_{exp})/SVI_{exp}$$
 Eq. 2

- where SVI_{mx} was the observed volume increment of mixed pine-birch stands and SVI_{exp} was the expected volume increment of mixed stands. While SVI_{mx} was the same for both species substitution and species addition, SVI_{exp} differed between the additive and substitutive scenarios.
- 218 For species substitution, the calculation of SVI_{exp.sub} was:

219
$$SVI_{exp.sub} = 0.5 \times SVI_{mo.pine} + 0.5 \times SVI_{mo.birch}$$
 Eq. 3

- where SVI_{mo. pine} and SVI_{mo. birch} were the stand volume increment of pine and birch monocultures averaged per block and 0.5 corresponds to the species proportion.
- 222 For species addition, we compared SVI_{mx} to SVI_{exp.add} at an equal number of trees per species. Thus, for
- $\label{eq:solution} 223 \qquad a \ SVI_{mx} \ at \ a \ density \ of \ n \ trees, \ we \ derived \ SVI_{exp.add} \ by \ summing \ volume \ increments \ in \ monocultures \ of$
- n/2 trees (see Figure 2):
- SVIexp.add.n = SVI_{mo} . pine. $n_{/2}$ + SVI_{mo} . birch. $n_{/2}$

- 226 Figure 2 Schematic representation of the calculations of transgressive mixture effect (TME) and
- 227 mixture effect (ME) of stand volume increment (SVI) for species substitution and species addition
- 228 based on observed (obs.) values and expected (exp.) values.



229

230 STATISTICAL ANALYSES

All the analyses were performed with R 3.2.5 and functions gam, lme and glht in packages mgcv, nlme

and multcomp.

We conducted separate analyses at the stand and tree levels by fitting a set of linear mixed effect models. At the plot level, we analysed four response variables: (i) total stand volume increment (SVI) estimated by summing tree volumes at plot level, (ii) mixture effect (ME) resulting from a species substitution (ME_{sub}) and from a species addition (ME_{add}) and (iv) transgressive mixture effect (TME) resulting from a species substitution (TME_{sub}) and from a species addition (TME_{add}). We completed the analyses at the plot level by considering also tree volume increment (TVI) of maritime pine and silver birch individual trees in monocultures and mixed plots.

Models of SVI and TVI included the effects of stand density (low, medium and high) and tree diversity (monoculture vs. two-species mixture) as fixed effect factors. Models of ME and TME included the effects of stand density (high and medium) and mixture scenario (substitution vs. addition) as fixed effect factors. We added Block as a random effect estimating between-block variability, except for analyses conducted at the level of individual tree where we nested plot within block to account for the non-independence of multiple trees sampled within the same plots and blocks. To consider the residual heteroscedasticity, analyses of SVI and TVI were carried out by introducing a variance model into the linear mixed models allowing unequal variance among experimental treatments (Pinheiro & Bates, 2006).

249 RESULTS

250 TREE SPECIES SUBSTITUTION

The substitution of silver birch by maritime pine multiplied significantly SVI by 3.6 at medium stand density (Figure 3, 38.8±5.67 m³/ha, n=18; S1 in Figure 1) as well as at high stand density (Figure 3, 56.6±10.6 m³/ha, n=8; S2 in Figure 1). Conversely, species substitution of maritime pine by silver birch decreased significantly SVI by 35% (Figure 3, S3 in Figure 1) and 36% (Figure 3, S4 in Figure 1) at medium and high stand density.

Birch-pine mixtures obtained through the substitution scenario were significantly less productive than the most productive (pine) monoculture (TME_{sub} < 0, Figure 4) at both medium (-0.35 \pm 0.08, n= 8) and high (-0.35 \pm 1.45, n=8) species density. Mixture effect (ME_{sub}, figure 4) indicated that pine-birch mixtures was marginally significantly more productive (overyielding) than their component monocultures at medium (0.10 \pm 0.14, n=8) and at high stand density (0.10 \pm 0.20, n=8, Figure 4).

Species substitution had opposite effects on TVI (tree volume increment) of the two studied species at medium stand density: a substitution of maritime pine by silver birch caused a significant increase of 15% in pine TVI (Figure 5), but a significant reduction of 23% in birch TVI (Figure 5). At high stand density, species substitution did not have any significant effect on TVI of silver birch or maritime pine (Figure 5).

266 HETEROSPECIFIC TREE ADDITION

Heterospecific species addition of maritime pine in silver birch stands multiplied significantly SVI by 5.8
at medium stand density (Figure 3, H1 in Figure 1) and by 5.2 at high stand density (H3 in Figure 1).
Heterospecific species addition of silver birch in maritime pine stands did not have any significant
effect on SVI neither at medium stand density (Figure 3, H3 in Figure 1) nor at high stand density (H4
in Figure 1).

ME_{add} indicated that pine-birch mixtures were significantly less productive (underyielding) than their component monocultures at intermediate (-0.14±0.07, n=8) and at high stand density (-0.19±0.16, n=8, Figure 4). TME_{add} at medium (0.01±0.10, n=6) and high (-0.05±0.20, n=8) stand density were not

- significantly different from zero indicating that SVI of mixed stands did not differ from SVI of pine in
- 276 monoculture, i.e. no transgressive overyielding (Figure 4).
- 277 Heterospecific addition of silver birch in monoculture of maritime pine did not cause any significant
- change in TVI of maritime pine at medium stand density (Figure 5) but a significant reduction of 17%
- 279 (Figure 5) at high stand density. Heterospecific addition of maritime pine in in monoculture of silver
- birch caused a significant reduction in TVI of silver birch of 42% (Figure 5) and 36% (Figure 5) at medium
- and high stand density, respectively.

bioRxiv preprint doi: https://doi.org/10.1101/2021.05.07.443133; this version posted May 9, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.



283 Figure 3 Stand volume increment (SVI) for different plot composition: in monoculture (mo) the SVI is 284 the sum of tree volume increment of the target species; in mixed plots (Ppin + Bpen) the SVI cumulates the tree volume increment of silver birch (Bpen) and maritime pine (Ppin). Effect of heterospecific 285 286 addition (H, see solid arrows in Figure 1) and species substitution (S, see black arrows in Figure 1) on 287 stand volume increment (SVI) of silver birch (a) and maritime pine (b) at low, medium and high stand 288 density. Black dots indicate mean values. Significance at a level of 5% were indicated by stars ((.) 0.1 >p-values > 0.05; * 0.05 > p-values > 0.01; ** 0.01 > p-values > 0.001; *** p-values > 0.001; ns no 289 290 significant effect). Note that SVI Ppin + Bpen of the medium and high density mixed stands are the same in 291 a) and b).

bioRxiv preprint doi: https://doi.org/10.1101/2021.05.07.443133; this version posted May 9, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.



293

Figure 4 Transgressive mixture effect (TME) and mixture effect (ME) at the stand level for species substitution and species addition at medium and high stand density. Black dot indicates the mean effects. Significance at a level of 5% were indicated by stars ((.) 0.1 > p-values > 0.05; * 0.05 > p-values > 0.01; ** 0.01 > p-values > 0.001; *** p-values > 0.001; ns no significant effect)





299

Figure 5 Effect of heterospecific species addition (H) and species substitution (S) on tree volume increment (TVI) of silver birch (*Betula pendula*) (a) and maritime pine (*Pinus pinaster*) (b) at low, medium and high stand density and in monoculture (Mo) or mixed (Mx) plots. Black dots indicate mean values. Significance at a level of 5% were indicated by stars ((.) 0.1 > p-values > 0.05; * 0.05 > p-values > 0.01; ** 0.01 > p-values > 0.001; *** p-values > 0.001; ns no significant effect). Note that the scales of the two figures (m³/ha) are different.

306 DISCUSSION

Our study assessed the role of tree species addition and species substitution in mixture effect in stands at an early age. We highlighted that when controlling for stand density, overyielding in young silver birch – maritime pine stands was due to a relaxation of intra-individual competition for pine. Conversely addition of silver birch (the least productive species) in a maritime pine stand (the most productive species) did not have a negative impact on stand productivity, which implies a nonsignificant transgressive mixture effect. Eventually stand density had little impact on the mixture effects tested and rather contribute to the species responses at an individual scale.

314 1§ SPECIES SUBSTITUTION TRIGGED OVERYIELDING IN MIXTURE OF TWO PIONEER SPECIES

315 Respective growth rates of tree species are crucial for interactions between species in the early stages 316 of development of mixed forests and our results confirm that positive effects of biodiversity on 317 productivity are mainly due to selection effect (Tobner et al., 2016) i.e. a fast-growth and productive species driving ecosystem functioning. Competitive advantage is common in young forests and positive 318 319 diversity-productivity relationship at this stage are often attributed in a lesser extent to 320 complementarity, particularly in harsher conditions (Van de Peer et al., 2018). Such positive effects are 321 commonly attributed to differences in shade tolerance as species with rapid growth rates benefit from 322 a relaxation of intraspecific competition, which may or may not be accompanied by niche separation 323 favouring shade-tolerant species rapidly overtopped due to their lower height growth rate (Boyden et 324 al., 2009; Tobner et al., 2016). However, we evidenced that overyielding can be triggered by species 325 similarities in their shade tolerance. Mixture effect was not conditioned by different light acquisition 326 strategies but more probably by their unequal ability to both tolerate drought. The experimental 327 plantation was on sandy heathlands that experience intense drought episodes in summer, water availability is an important limiting factor for tree growth, especially in silver birch, which has the 328 329 lowest drought tolerance. Maritime pine is able to maintain its stem growth during a longer period and 330 even to restart height grow in autumn (fast-growth evergreen species, (Heuret et al., 2006)). Silver 331 birch remains a species sensitive to interspecific competition at a young age, even in the Nordic 332 countries where temperature is a more limiting factor for growth than water (Jucker et al., 2020), but it is likely that dry conditions further accentuate its competitive disadvantage. 333

Nonetheless effects that we observed 8 (7?) years after planting will change very quickly, the growing gap in height between maritime pine and silver birch being detrimental to birch under current climatic conditions, tree mortality will intensity (Morin et al., 2020). Long term simulations of pine and birch stand showed a lasting overyielding due to the relaxation of intraspecific competition for pine over time (Morin et al., 2020). Oaks species with their slower growth rates and varying drought and shade tolerances will gradually establish into stands, leading to a stand stratification possibly suitable to mixed stands, even though at this point, there is no consensus on the stage of forest development for which the positive effect of diversity peak (Jucker et al., 2020; Taylor et al., 2020).

342 2§ TRANSGRESSIVE MIXTURE EFFECT WAS NOT SIGNIFICANTLY DIFFERENT FROM ZERO IN THE343 ADDITION SCENARIO

We did not find any transgressive mixture effect in mixed birch-pine stand created by addition of the two species. Conversely the substitutive approach caused a loss of mixed stand productivity compares to pine monocultures due to the substitution of a high productive species (maritime pine) by a low productive species (silver birch). These findings mirror what has been found in colder and more humid sites for the same species that we studied (Frivold & Frank, 2002), more generally in mixed-forest (Jactel et al., 2018) and in plant community where positive transgressive overyielding were rarely reported (Cardinale et al., 2007).

351 At a medium stand density in the additive scenario the absence of any competition effect of silver birch 352 on maritime pine can be explained by two inseparable mechanisms: either, a purely neutral effect of 353 the addition of the least productive species due to an insufficient proximity of stems, or a facilitating 354 effect of birch on soil resource that compensates for a weak competitive constraint due to species 355 addition. Leaves of silver birch have a rate of decomposition higher than needles of Pinus species 356 (Palviainen et al., 2004) and depending on the stand structure, nutrient cycling can be higher in birch 357 regeneration than in pine regeneration (De Schrijver et al., 2009). Hence in the studied site, carbon 358 and nitrogen at an intermediate soil depth have been found higher in mixed stands than on monospecific stands (Maxwell et al., 2020), even though there is no evidence on belowground 359 360 complementarity of fine roots (Altinalmazis-Kondylis et al., 2020).

These findings are also of great ecological relevance because they demonstrate that it is possible to diversify pine monocultures with addition of birch at an early age and then benefit from ecosystem services such as pest protection (Damien et al., 2016; Jactel et al., 2019) and increased diversity of predatory insects (Jouveau et al., 2020) without compromising the wood production of the target species (here maritime pine). Long-term simulations of pine and birch growth on the study sites supported our results (Morin et al., 2020) showing that the ecosystem services associated with pine monoculture diversification can persist as the stand ages. 368 3§ MIXTURE EFFECTS AND TRANSGRESSIVE MIXTURE EFFECTS DO NOT CHANGE WITH STAND DENSITY,

369 BUT TREE PRODUCTIVITY DOES.

370 In young stands, high stand densities usually speed up mixture effects (Tobner et al., 2016; Van de Peer 371 et al., 2018). In this study, we did not observe any intensification with stand density neither of the 372 mixture effect nor of transgressive mixture effect. However, an intensification of interactions with stands density was observed at the tree level: at medium density, heterospecific addition did not affect 373 374 maritime pine trees (the most productive species) but silver birch trees (the least productive species). 375 At high density, the intensification of interspecific competition led to a reduction of productivity for 376 both species. Regarding species substitution, at medium density maritime pine (the most productive 377 species) benefit from mixture at the expense of silver birch (the least productive species). When stand 378 density increased, the difference between inter and intraspecific competition becomes narrower, 379 which might explain the absence of any effect of species substitution. This illustrates on the one hand 380 that the same response pattern in term of mixture effect can emerge from different mechanisms at 381 the individual levels and. On the other hand, it is consistent with what has been observed elsewhere 382 in young and dynamic stages with an intensification of competitive interactions (Boyden et al., 2009) 383 or at least a decrease in overyielding with density (Kweon & Comeau, 2019). Finally, these results 384 contrast with the intensification of the positive diversity productivity relationship observed as forest 385 stands become older (Huang et al., 2018) particularly when shade tolerant and shade intolerant species 386 are mixed (Brunner, 2020; del Rio & Sterba, 2009).

387 CONCLUSION

388 By controlling stand density and species identity, we have shown that selection effect is the main driver 389 of positive diversity – productivity relationships in the early stages of mixed species forests. This calls 390 for a careful choice of which tree species to associate when designing plantations of mixed species, 391 especially of fast-growing species. In addition, our results also showed that the addition of a pioneer 392 and low-demand species to a monoculture of a high productive species in young developmental stages 393 offers the opportunity to benefit from ecosystem services associated to mixed stands without affecting 394 the productivity of the target species. The addition of tree species is a promising way to promote 395 multifunctionality in mixed-species plantations and to preserve the harvest of a particular species for 396 timber production, thus circumventing two major obstacles in the implementation of mixed-species 397 forestry.

398 ACKNOWLEDGEMENTS

- We thank the Forest experimental Facility (UEFP-https://doi.org/10.15454/1.5483264699193726E12)
- and especially Bernard Issenhuth for the maintenance and the measurements of the ORPHEEexperiment

402 AUTHORS' CONTRIBUTIONS

All authors contributed critically to conceived the ideas and designed methodology; HJ designed the
 ORPHEE experiment; MT analysed the data; MT and CM led the writing of the manuscript. All authors

405 contributed critically to the drafts and gave final approval for publication.

406 DATA AVAILABILITY STATEMENT

407 Data will be available from the INRAE Digital Repository <u>https://data.inrae.fr/dataverse/biogeco</u>.

408 REFERENCES

- 409 Altinalmazis-Kondylis, A., Muessig, K., Meredieu, C., Jactel, H., Augusto, L., Fanin, N., & Bakker, M. R.
- 410 (2020). Effect of tree mixtures and water availability on belowground complementarity of fine
 411 roots of birch and pine planted on sandy podzol. *Plant and Soil*, 457(1), 437–455.
- 412 Amoroso, M. M., & Turnblom, E. C. (2006). Comparing productivity of pure and mixed Douglas-fir and
- 413 western hemlock plantations in the Pacific Northwest. *Canadian Journal of Forest Research*,
- 414 *36*(6), 1484–1496. https://doi.org/10.1139/x06-042
- 415 Baeten, L., Bruelheide, H., van der Plas, F., Kambach, S., Ratcliffe, S., Jucker, T., Allan, E., Ampoorter,
- 416 E., Barbaro, L., & Bastias, C. C. (2019). Identifying the tree species compositions that maximize 417 ecosystem functioning in European forests. *Journal of Applied Ecology*, *56*(3), 733–744.
- 418 Bakker, M. R., Augusto, L., & Achat, D. L. (2006). Fine root distribution of trees and understory in
- 419 mature stands of maritime pine (Pinus pinaster) on dry and humid sites. *Plant and Soil, 286*(1–
- 420 2), 37–51. https://doi.org/10.1007/s11104-006-9024-4
- 421 Biging, G. S., & Dobbertin, M. (1992). A comparison of distance-dependent competition measures for
- 422 height and basal area growth of individual conifer trees. *Forest Science*, *38*(3), 695–720.

- 423 Boyden, S. B., Reich, P. B., Puettmann, K. J., & Baker, T. R. (2009). Effects of density and ontogeny on 424 size and growth ranks of three competing tree species. *Journal of Ecology*, *97*(2), 277–288.
- 425 https://doi.org/10.1111/j.1365-2745.2008.01477.x
- Brunner, A. (2020). Tree species mixture effects on stem growth vary with stand density An analysis
 based on individual tree responses. *Forest Ecology and Management*, 15.
- 428 Cardinale, B. J., Wright, J. P., Cadotte, M. W., Carroll, I. T., Hector, A., Srivastava, D. S., Loreau, M., &
- 429 Weis, J. J. (2007). Impacts of plant diversity on biomass production increase through time 430 because of species complementarity. *Proceedings of the National Academy of Sciences*,
- 431 *104*(46), 18123–18128. https://doi.org/10.1073/pnas.0709069104
- Caspersen, J. P., Thürig, E., Rigling, A., & Zimmermann, N. E. (2018). Complementarity of gymnosperms
 and angiosperms along an altitudinal temperature gradient. *Oikos*, *127*(12), 1787–1799.
- Condés, S., Del Rio, M., & Sterba, H. (2013). Mixing effect on volume growth of Fagus sylvatica and
 Pinus sylvestris is modulated by stand density. *Forest Ecology and Management, 292*, 86–95.
 https://doi.org/10.1016/j.foreco.2012.12.013
- 437 Corcket, E., Alard, D., Halder, I., Jactel, H., Garrido Diaz, B., Reuzeau, E., & Castagneyrol, B. (2020).
 438 Canopy composition and drought shape understorey plant assemblages in a young tree
 439 diversity experiment. *Journal of Vegetation Science*, *31*(5), 803–816.
 440 https://doi.org/10.1111/jvs.12903
- Damien, M., Jactel, H., Meredieu, C., Régolini, M., van Halder, I., & Castagneyrol, B. (2016). Pest
 damage in mixed forests: Disentangling the effects of neighbor identity, host density and host
 apparency at different spatial scales. *Forest Ecology and Management*, *378*, 103–110.
 https://doi.org/10.1016/j.foreco.2016.07.025
- De Schrijver, A., Geudens, G., Wuyts, K., Staelens, J., Gielis, L., & Verheyen, K. (2009). Nutrient cycling
 in two continuous cover scenarios for forest conversion of pine plantations on sandy soil. I.
 Nutrient cycling via aboveground tree biomass. *Canadian Journal of Forest Research*, *39*(2),
 441–452. https://doi.org/10.1139/X08-176

bioRxiv preprint doi: https://doi.org/10.1101/2021.05.07.443133; this version posted May 9, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

del Rio, M., & Sterba, H. (2009). Comparing volume growth in pure and mixed stands of Pinus sylvestris

- 450 and Quercus pyrenaica. Annals of Forest Science, 66(5), 502.
 451 https://doi.org/10.1051/forest/2009035
- Deleuze, C., Morneau, F., renaud, J.-P., vivien, Y., Rivoire, M., santenoise, philippe, Longuetaud, F.,
 mothe, frédéric, Hervé, J.-C., & Vallet, P. (2014). Estimer le volume total d'un arbre, quelles
 que soient l'essence, la taille, la sylviculture, la station. *Rendez-Vous Techniques*, 44(44), 22–
 32.
- Felton, A., Lindbladh, M., Brunet, J., & Fritz, Ö. (2010). Replacing coniferous monocultures with mixedspecies production stands: An assessment of the potential benefits for forest biodiversity in
 northern Europe. *Forest Ecology and Management, 260*(6), 939–947.
 https://doi.org/10.1016/j.foreco.2010.06.011
- Forrester, D. I. (2014). The spatial and temporal dynamics of species interactions in mixed-species
 forests: From pattern to process. *Forest Ecology and Management*, *312*, 282–292.
 https://doi.org/10.1016/j.foreco.2013.10.003
- Forrester, D. I., Kohnle, U., Albrecht, A. T., & Bauhus, J. (2013). Complementarity in mixed-species
 stands of Abies alba and Picea abies varies with climate, site quality and stand density. *Forest Ecology and Management*, *304*, 233–242. https://doi.org/10.1016/j.foreco.2013.04.038
- 466 Fox, J. W. (2005). Interpreting the 'selection effect' of biodiversity on ecosystem function. *Ecology*467 *Letters*, 8(8), 846–856. https://doi.org/10.1111/j.1461-0248.2005.00795.x
- Frivold, L. H., & Frank, J. (2002). Growth of Mixed Birch-Coniferous Stands in Relation to Pure
 Coniferous Stands at Similar Sites in South-eastern Norway. *Scandinavian Journal of Forest Research*, *17*(2), 139–149. https://doi.org/10.1080/028275802753626782
- 471 Gamfeldt, L., Snall, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M. C., Froberg,
- 472 M., Stendahl, J., Philipson, C. D., Mikusinski, G., Andersson, E., Westerlund, B., Andren, H.,
 473 Moberg, F., Moen, J., & Bengtsson, J. (2013). Higher levels of multiple ecosystem services are

474 found in forests with more tree species. *Nature Communications*, *4*, 1340.
475 https://doi.org/10.1038/ncomms2328

- Gaudio, N., Balandier, P., Dumas, Y., & Ginisty, C. (2011). Growth and morphology of three forest
 understorey species (Calluna vulgaris, Molinia caerulea and Pteridium aquilinum) according to
 light availability. *Forest Ecology and Management*, 261(3), 489–498.
- Grossman, J. J., Cavender-Bares, J., Hobbie, S. E., Reich, P. B., & Montgomery, R. A. (2017). Species
 richness and traits predict overyielding in stem growth in an early-successional tree diversity
 experiment. *Ecology*, *98*(10), 2601–2614.
- Henneron, L., Aubert, M., Archaux, F., Bureau, F., Dumas, Y., Ningre, F., Richter, C., Balandier, P., &
 Chauvat, M. (2017). Forest plant community as a driver of soil biodiversity: experimental
 evidence from collembolan assemblages through large-scale and long-term removal of oak

485 canopy trees Quercus petraea. *Oikos*, *126*(3), 420–434.

- Heuret, P., Meredieu, C., Coudurier, T., Courdier, F., & Barthélémy, D. (2006). Ontogenetic trends in
 the morphological features of main stem annual shoots of Pinus pinaster (Pinaceae). *American Journal of Botany*, *93*(11), 1577–1587.
- 489 Huang, Y., Chen, Y., Castro-Izaguirre, N., Baruffol, M., Brezzi, M., Lang, A., Li, Y., Härdtle, W., von
- 490 Oheimb, G., Yang, X., Liu, X., Pei, K., Both, S., Yang, B., Eichenberg, D., Assmann, T., Bauhus, J.,
- 491 Behrens, T., Buscot, F., ... Schmid, B. (2018). Impacts of species richness on productivity in a
- 492 large-scale subtropical forest experiment. *Science*, *362*(6410), 80–83.
 493 https://doi.org/10.1126/science.aat6405
- Hynynen, J., Niemisto, P., Vihera-Aarnio, A., Brunner, A., Hein, S., & Velling, P. (2010). Silviculture of
 birch (Betula pendula Roth and Betula pubescens Ehrh.) in northern Europe. *Forestry*, *83*(1),
- 496 103–119. https://doi.org/10.1093/forestry/cpp035
- 497 Jactel, H., Gritti, E. S., Drössler, L., Forrester, D. I., Mason, W. L., Morin, X., Pretzsch, H., & Castagneyrol,
- 498 B. (2018). Positive biodiversity–productivity relationships in forests: climate matters. *Biology*
- 499 *Letters*, 14(4), 20170747. https://doi.org/10.1098/rsbl.2017.0747

500	Jactel, H., Poeydebat, C., van Halder, I., & Castagneyrol, B. (2019). Interactive Effects of Tree Mixing
501	and Drought on a Primary Forest Pest. Frontiers in Forests and Global Change, 2, 77.
502	https://doi.org/10.3389/ffgc.2019.00077

- Jouveau, S., Toïgo, M., Giffard, B., Castagneyrol, B., Halder, I., Vétillard, F., & Jactel, H. (2020). Carabid
 activity-density increases with forest vegetation diversity at different spatial scales. *Insect Conservation and Diversity*, *13*(1), 36–46. https://doi.org/10.1111/icad.12372
- Jucker, T., Avăcăriței, D., Bărnoaiea, I., Duduman, G., Bouriaud, O., & Coomes, D. A. (2016). Climate
 modulates the effects of tree diversity on forest productivity. *Journal of Ecology*, *104*(2), 388–
 398. https://doi.org/10.1111/1365-2745.12522
- Jucker, T., Bouriaud, O., & Coomes, D. A. (2015). Crown plasticity enables trees to optimize canopy
 packing in mixed-species forests. *Functional Ecology*, *29*(8), 1078–1086.
 https://doi.org/10.1111/1365-2435.12428
- 512 Jucker, T., Koricheva, J., Finér, L., Bouriaud, O., Iacopetti, G., & Coomes, D. A. (2020). Good things take 513 time—Diversity effects on tree growth shift from negative to positive during stand 514 development in boreal forests. Journal Ecology, 108(6), 2198-2211. of https://doi.org/10.1111/1365-2745.13464 515
- 516 Kunz, M., Fichtner, A., Härdtle, W., Raumonen, P., Bruelheide, H., & Oheimb, G. (2019). Neighbour 517 species richness and local structural variability modulate aboveground allocation patterns and 518 crown morphology of individual trees. Ecology Letters, 22(12), 2130-2140. 519 https://doi.org/10.1111/ele.13400
- 520 Kweon, D., & Comeau, P. G. (2019). Factors influencing overyielding in young boreal mixedwood stands
 521 in western Canada. *Forest Ecology and Management*, *432*, 546–557.
- Ligot, G., Balandier, P., Courbaud, B., Jonard, M., Kneeshaw, D., & Claessens, H. (2014). Managing
 understory light to maintain a mixture of species with different shade tolerance. *Forest Ecology and Management*, *327*, 189–200. https://doi.org/10.1016/j.foreco.2014.05.010

bioRxiv preprint doi: https://doi.org/10.1101/2021.05.07.443133; this version posted May 9, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

- 525 Loreau, M. (1998). Separating Sampling and Other Effects in Biodiversity Experiments. Oikos, 82(3),
- 526 600. https://doi.org/10.2307/3546381
- Loreau, M., & Hector, A. (2001). Partitioning selection and complementarity in biodiversity
 experiments. *Nature*, *412*(6842), 72–76. https://doi.org/10.1038/35083573
- 529 Maxwell, T. L., Augusto, L., Bon, L., Courbineau, A., Altinalmazis-Kondylis, A., Milin, S., Bakker, M. R.,
- 530 Jactel, H., & Fanin, N. (2020). Effect of a tree mixture and water availability on soil nutrients
- and extracellular enzyme activities along the soil profile in an experimental forest. *Soil Biology and Biochemistry*, *148*, 107864.
- 533 Morin, X., Damestoy, T., Toigo, M., Castagneyrol, B., Jactel, H., de Coligny, F., & Meredieu, C. (2020).
- 534 Using forest gap models and experimental data to explore long-term effects of tree diversity 535 on the productivity of mixed planted forests. *Annals of Forest Science*, *77*, 1–19.
- Nadrowski, K., Wirth, C., & Scherer-Lorenzen, M. (2010). Is forest diversity driving ecosystem function
 and service? *Current Opinion in Environmental Sustainability*, 2(1–2), 75–79.
 https://doi.org/10.1016/j.cosust.2010.02.003
- 539 Palviainen, M., Finér, L., Kurka, A.-M., Mannerkoski, H., Piirainen, S., & Starr, M. (2004). Decomposition
- 540 and nutrient release from logging residues after clear-cutting of mixed boreal forest. *Plant and*
- 541 Soil, 263(1), 53–67. https://doi.org/10.1023/B:PLSO.0000047718.34805.fb
- 542 Perot, T., Mårell, A., Korboulewsky, N., Seigner, V., & Balandier, P. (2017). Modeling and predicting
 543 solar radiation transmittance in mixed forests at a within-stand scale from tree species basal
 544 area. *Forest Ecology and Management, 390*, 127–136.
- Pinheiro, J., & Bates, D. (2006). *Mixed-effects models in S and S-PLUS*. Springer Science & Business
 Media.
- Taylor, A. R., Gao, B., & Chen, H. Y. H. (2020). The effect of species diversity on tree growth varies
 during forest succession in the boreal forest of central Canada. *Forest Ecology and Management*, 455, 117641. https://doi.org/10.1016/j.foreco.2019.117641

- 550 Tobner, C. M., Paquette, A., Gravel, D., Reich, P. B., Williams, L. J., & Messier, C. (2016). Functional
- identity is the main driver of diversity effects in young tree communities. *Ecology Letters*, 19(6),
- 552 638–647. https://doi.org/10.1111/ele.12600
- 553 Van de Peer, T., Verheyen, K., Ponette, Q., Setiawan, N. N., & Muys, B. (2018). Overyielding in young
- 554 tree plantations is driven by local complementarity and selection effects related to shade
- tolerance. *Journal of Ecology*, *106*(3), 1096–1105. https://doi.org/10.1111/1365-2745.12839

556