



HAL
open science

Sperm Morphology Assessment in the Era of Intracytoplasmic Sperm Injection: Reliable Results Require Focus on Standardization, Quality Control, and Training

Ashok Agarwal, Rakesh Sharma, Sajal Gupta, Renata Finelli, Neel Parekh, Manesh Kumar Panner Selvam, Ralf Henkel, Damayanthi Durairajanayagam, Camila Pompeu, Sarah Madani, et al.

► **To cite this version:**

Ashok Agarwal, Rakesh Sharma, Sajal Gupta, Renata Finelli, Neel Parekh, et al.. Sperm Morphology Assessment in the Era of Intracytoplasmic Sperm Injection: Reliable Results Require Focus on Standardization, Quality Control, and Training. *The World Journal of Men's Health*, 2021, 39, 10.5534/wjmh.210054 . hal-03278904

HAL Id: hal-03278904

<https://hal.inrae.fr/hal-03278904>

Submitted on 1 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Sperm Morphology Assessment in the Era of Intracytoplasmic Sperm Injection: Reliable Results Require Focus on Standardization, Quality Control, and Training

Ashok Agarwal¹, Rakesh Sharma¹, Sajal Gupta¹, Renata Finelli¹, Neel Parekh¹, Manesh Kumar Panner Selvam^{1,2}, Ralf Henkel^{1,3,4,5}, Damayanthi Durairajanayagam⁶, Camila Pompeu⁷, Sarah Madani⁸, Andrea Belo⁹, Neha Singh¹⁰, Simryn Covarrubias¹¹, Sara Darbandi¹², Raha Sadeghi¹¹, Mahsa Darbandi¹², Paraskevi Vogiatzi¹³, Florence Boitrelle^{14,15}, Mara Simopoulou¹⁶, Ramadan Saleh¹⁷, Mohamed Arafa^{1,18,19}, Ahmad Majzoub^{1,19}, Hussein Kandil²⁰, Armand Zini²¹, Edmund Ko²², Juan G. Alvarez²³, Marlon Martinez²⁴, Jonathan Ramsay²⁵, Sunil Jindal²⁶, Gian Maria Busetto²⁷, Hassan Sallam²⁸, Israel Maldonado²⁹, Christina Anagnostopoulou³⁰, Marco G. Alves³¹, Pallav Sengupta³², Kambiz Gilany³³, Donald P. Evenson³⁴, Sheena E.M. Lewis^{35,36}, Jaime Gosalvez³⁷, Rafael F. Ambar³⁸, Rupin Shah³⁹

¹American Center for Reproductive Medicine, Cleveland Clinic, Cleveland, OH, ²Department of Urology, Tulane University Health Sciences Center, New Orleans, LA, USA, ³Department of Metabolism, Digestion and Reproduction, Imperial College London, London, UK, ⁴Department of Medical Bioscience, University of the Western Cape, Bellville, South Africa, ⁵LogixX Pharma, Theale, Reading, Berkshire, UK, ⁶Department of Physiology, Faculty of Medicine, Universiti Teknologi MARA, Sungai Buloh Campus, Selangor, Malaysia, ⁷Instituto Ideia Fértil, Santo André, Brazil, ⁸Department of Biology and Physiology of Organisms, Faculty of Biological Sciences, University of Science and Technology, Houari Boumediene, Algiers, Algeria, ⁹Huntington Centro de Medicina Reprodutiva S/A, Sao Paulo, São Paulo, Brazil, ¹⁰NOVA IVF, Gorakhpur, India, ¹¹Department of Physiology, University of San Francisco, CA, USA, ¹²Fetal Health Research Center, Hope Generation Foundation, Tehran, Iran, ¹³Andromed Health & Reproduction, Fertility Diagnostics Laboratory, Maroussi, Athens, Greece, ¹⁴Reproductive Biology, Fertility Preservation, Andrology, CECOS, Poissy Hospital, Poissy, ¹⁵Paris Saclay University, UVSQ, INRAE, BREED, Jouy-en-Josas, France, ¹⁶Department of Experimental Physiology, School of Health Sciences, Faculty of Medicine, National and Kapodistrian University of Athens, Athens, Greece, ¹⁷Department of Dermatology, Venereology and Andrology, Faculty of Medicine, Sohag University, Sohag, Egypt, ¹⁸Department of Urology, Hamad Medical Corporation, ¹⁹Department of Urology, Weill Cornell Medicine-Qatar, Doha, Qatar, ²⁰Fakih IVF Fertility Center, Abu Dhabi, UAE, ²¹Department of Surgery, McGill University, Montreal, QC, Canada, ²²Department of Urology, Loma Linda University Health, Loma Linda, CA, USA, ²³Centro ANDROGEN, La Coruña, Spain, ²⁴Section of Urology, University of Santo Tomas Hospital, Manila, Philippines, ²⁵Department of Andrology, Hammersmith Hospital, London, UK, ²⁶Department of Andrology and Reproductive Medicine, Jindal Hospital, Meerut, India, ²⁷Department of Urology and Renal Transplantation, University of Foggia Policlinico Riuniti of Foggia, Foggia, Italy, ²⁸Department of Obstetrics and Gynaecology, Alexandria University Faculty of Medicine, Alexandria, Egypt, ²⁹Citmer Reproductive Medicine, IVF LAB, Mexico City, Mexico, ³⁰IVF Clinic "Akeso-Embryo ART", Athens, Greece, ³¹Department of Anatomy, Unit for Multidisciplinary Research in Biomedicine (UMIB), Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, Porto, Portugal, ³²Department of Physiology, Faculty of Medicine, MAHSA University, Selangor, Malaysia, ³³Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran, ³⁴SCSA Diagnostics, Brookings, SD, USA, ³⁵Queens University Belfast, ³⁶Examenlab Ltd., Weavers Court, Belfast, Northern Ireland, UK, ³⁷Genetic Unit, Department of Biology, Universidad Autónoma de Madrid, Madrid, Spain, ³⁸Department of Urology, Centro Universitario em Saude do ABC/Andrology Group at Ideia Fértil Institute of Human Reproduction, Santo André, Brazil, ³⁹Department of Urology, Lilavati Hospital and Research Centre, Mumbai, India

Received: Mar 29, 2021 Revised: Apr 21, 2021 Accepted: May 9, 2021 Published online Jun 17, 2021

Correspondence to: Ashok Agarwal  <https://orcid.org/0000-0003-0585-1026>

Andrology Center and American Center for Reproductive Medicine, Cleveland Clinic, Mail Code X-11, 10681 Carnegie Avenue, Cleveland, OH 44195, USA.

Tel: +1-216-444-9485, Fax: +1-216-445-6049, E-mail: agarwaa@ccf.org, Website: www.Clevelandclinic.org/ReproductiveResearchCenter

Semen analysis is the first, and frequently, the only step in the evaluation of male fertility. Although the laboratory procedures are conducted according to the World Health Organization (WHO) guidelines, semen analysis and especially sperm morphology assessment is very difficult to standardize and obtain reproducible results. This is mainly due to the highly subjective nature of their evaluation. ICSI is the choice of treatment when sperm morphology is severely abnormal (teratozoospermic). Hence, the standardization of laboratory protocols for sperm morphology evaluation represents a fundamental step to ensure reliable, accurate and consistent laboratory results that avoid misdiagnoses and inadequate treatment of the infertile patient. This article aims to promote standardized laboratory procedures for an accurate evaluation of sperm morphology, including the establishment of quality control and quality assurance policies. Additionally, the clinical importance of sperm morphology results in assisted reproductive outcomes is discussed, along with the clinical management of teratozoospermic patients.

Keywords: Abnormality, teratozoospermia; Morphology, stain; Sperm

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Infertility is defined as the inability to achieve a successful pregnancy after 12 months or more of regular, unprotected sexual intercourse [1]. The World Health Organization (WHO) demographics estimate that about 200 million people are affected by infertility globally [2,3], with male factor contributing to 50% of infertility cases [4]. A meta-analysis published in 1992 reported a decline in sperm quality over a period of 50 years (1938 to 1991) [5] while, more recently, Levine et al [6] reported a 50% to 60% decline in sperm count from 1973 to 2011 across North America, Europe, New Zealand, and Australia. Furthermore, other researchers have reported a significant decline in sperm concentration, total count, morphology, and motility between 2000 to 2017 [7]. A negative association is also reported between poor morphology and gonadotropins [8]. As the analysis of sperm parameters is the primary approach for identifying and diagnosing male infertility, semen analysis is of particular importance since it defines fertility status and potential, as well as the course of natural or assisted reproduction [9]. In this framework, sperm morphology is established as the most prominent component, as this parameter cannot be surpassed even by the most invasive interventions such as intracytoplasmic sperm injection (ICSI).

Interestingly, there have been significant changes in the classification of sperm morphology over the years. Normal sperm morphology reference values have been revised dramatically from $\geq 80.5\%$ reported in the 1st

edition of the WHO manual [10,11] to $\geq 14\%$ in the 4th edition [12], and even lower to $\geq 4\%$ in the 5th and most recent edition [13]. The 'strict criteria' for assessing sperm morphology classify semen as normal if the percentage of normal sperm is $\geq 14\%$ [14,15]. Based on this classification, abnormal sperm morphology has been associated with poor fertilization and clinical outcomes after assisted reproductive technology (ART), thereby establishing sperm morphology as a predictor of ART outcomes [16-18]. The cut-off for sperm morphology based on strict criteria was further revised in the 5th edition of the WHO manual [13] which uses the 5th centiles (and their 95% confidence intervals [CIs]) as the lowest reference limit ($\geq 4\%$ cutoff for normal sperm morphology). The lowest limit for morphologically normal forms of spermatozoa is 4% (95% CI, 3%–4%) and all borderline forms are considered abnormal. This classification serves as a surrogate tool to choose the most appropriate type of ART procedure for infertile couples [19]; if the percentage of normal forms is $\geq 4\%$, the probability of fertilization with ART is high and procedures such as intrauterine insemination (IUI) or *in vitro* fertilization (IVF) may be selected. In specimens with otherwise normal count and motility, poor morphology may itself be a determinant in deciding whether or not to proceed with ICSI. When the percentage of sperm with normal morphology is $< 4\%$, fertilization with IUI and IVF is poor, and ICSI should be preferred [20].

The predictive value of sperm morphology in ART continues to be a matter of debate. Recent studies sug-

gest that morphology may not be a good predictor of fertilization or pregnancy outcome in ICSI [21-25]. Abnormal fertilization following ICSI has been associated with abnormal parameters in semen analysis. Oligoasthenozoospermia and oligoasthenoteratozoospermia have been suggested to be further associated with lower cleavage and blastocyst formation rates [26]. A careful assessment is critical to identify the underlying cause of infertility, as this can serve as a tool to choose the most appropriate type of ART method [27]. However, heterogeneity of study groups, differences in staining methods, intra- and inter-laboratory variations, differences in the scoring classifications, and manual *versus* computer assisted semen analyzer (CASA) scoring may be among the many contributory factors for the lack of robust predictive power [19,28,29].

Diagnostic and predictive value of sperm morphology can be improved by uniform and consistent application of the defined strict criteria for scoring sperm morphology, standardization of the staining methods, internal and external quality controls (QCs), and training of the laboratory personnel [11,19,30].

In this review, we aim to: a) summarize standardized laboratory procedures for a proper evaluation of sperm morphology; b) highlight the importance of QC and quality assurance (QA) in laboratory assessment of sperm morphology; c) discuss the association between abnormal sperm morphology (teratozoospermia) and ART outcomes; and d) review the clinical management of men with abnormal sperm morphology.

MORPHOLOGICAL ASSESSMENT OF SPERM BY IMPLEMENTATION OF STANDARDIZATION AND QUALITY CONTROL

1. Preparation of semen sample

Semen sample is collected in a sterile collection container according to the recommendations provided by the WHO (2010) and incubated at 37°C for 30 minutes to allow liquefaction [13]. If the ejaculate is viscous, proteolytic enzymes such as α -chymotrypsin or bromelain can be added to the sample and incubated at 37°C for an additional 10 minutes [13,31]. The liquefied sample is vortexed for 10 seconds, and a 10 μ L aliquot is quickly removed. If the sperm concentration is $<2 \times 10^6$ /mL, the sample is centrifuged at 600 g for 10 minutes to remove most of the supernatant, leaving about 100 μ L

of seminal plasma in the underlying concentrate. The pellet is resuspended by gentle pipetting to redilute the sample, not exceeding 50×10^6 /mL. Centrifugation may affect sperm morphology and should be recorded in the patient's worksheet [32].

2. Preparing a smear for sperm morphology

On a clean frosted slide with the patient identifiers, 10 μ L of well-mixed semen aliquot is added on one end of the slide. A second slide is used with an angle of 45° to quickly spread the drop of semen along the frosted slide, forming a smooth and even smear. The slides are prepared in duplicate and then air-dried before staining.

The smear is stained using a rapid stain such as Diff-Quik, which consists of a fixative (triarylmethane dye, methanol), solution I (xanthene dye, sodium azide, pH buffer), and solution II (thiazine dye, pH buffer) (Fig. 1). According to the Diff-Quik staining protocol, once the smear slide is dry, it is immersed in the fixative five times and allowed to dry completely for 15 minutes. Once dried, the slide is immersed three times in solution I for 10 seconds, following which the excess stain is drained, and the slide is immersed five times in solution II for 10 seconds. The excess stain is quickly rinsed in sterile water, and the slide is placed vertically on absorbent paper to dry. Finally, a few drops of the mounting medium Cytoseal are placed on the slide, and the slide is covered with a coverslip. The slide is allowed to dry completely before it is examined under a bright field microscope [33]. The stained smear is exam-




Fig. 1. Main components of the Diff-Quik staining method: fixative, Solution I, Solution II, and water to rinse excess stain.





Fig. 2. (A) A magnified image showing sperm morphology staining as seen on an external monitor. (B) Microscope equipped with 100× oil-immersion brightfield objective used for the analysis of sperm morphology.

Cleveland Clinic
©2021



Bright-field microscope

Ocular micrometer

Lint-free wipes

Differential counter

Immersion oil

Lens cleaner solution

Cleveland Clinic
©2021

Fig. 3. Supplies and instruments used for the evaluation of sperm morphology.

ined using a bright field microscope with 100× objective and 10× eyepiece (Fig. 2). For optimum sharpness, the immersion oil used at 100× magnification should have a refractive index (RI) of 1.52, similar to that of glass and the cells being examined. An ocular micrometer is placed in one of the microscope's eyepieces to accurately measure the sperm dimensions (Fig. 3) [13]. Based on the strict criteria, a spermatozoon must conform to all normal morphological criteria as outlined above. Without the aid of an ocular micrometer, a precise evaluation of morphology cannot be performed [13].

3. Evaluation of sperm morphology

Evaluation of sperm morphology is highly subjective and depends largely on the perception of the observer scoring the slide. To obtain reliable and reproducible

results, the andrology laboratory must develop a detailed step-by-step protocol. In addition, the use of an ocular micrometer is essential to measure the sperm dimensions. As reported in the WHO guidelines, the sperm head should be oval in shape, smooth, and regularly contoured, 5 to 6 μm long and 2.5 to 3.5 μm wide [13]. The acrosome must be well-defined, occupy between 40% to 70% of the total area of the head, and should not contain more than two small vacuoles. The vacuoles must not occupy more than 20% of the area of the sperm head. The acrosomal region stains light blue, while the post-acrosomal region stains dark blue. The post-acrosomal region must not contain any vacuoles [14,33,34]. The mid-piece must be regular, slender, about the same length as the sperm head and stained a purple-red color. The mid-piece must also be aligned

with the axis of the head of the sperm. If a residual cytoplasm larger than one-third of the area of the head is present, the sperm should be considered as abnormal [35]. If present, excess residual cytoplasm may be seen around the midpiece and is stained pink/red or reddish orange depending on the type of stain used. The tail should be approximately 45 μm long, uniform along its length, appear thinner than the mid-piece, and stained a blue or reddish color. At least 2 replicates of 100 spermatozoa must be scored, with all borderline forms being considered as abnormal. When this strict classification is followed, the reference threshold is $\geq 4\%$ for morphologically normal forms [13].

QUALITY CONTROL AND QUALITY ASSURANCE IN SPERM MORPHOLOGY ASSESSMENT


Sperm morphology remains one of the most controversial semen parameters, as an incorrect categorization can lead to lack of predictive value for ART outcomes, as reported in several recent publications, although sperm morphology has shown strong predictive values in the early nineties [21-25].

The WHO 5th edition recommends Papanicolaou (gold standard), Shorr and Diff-Quik staining as methods of choice [13]. These methods are adequate for spermatozoa staining and observation under a bright field microscope [36-38]. Any other non-validated staining method can alter the morphological appearance, by causing swelling of the sperm due to osmotic changes [39]. When assessing morphology under bright field optics, spermatozoa in a well-produced and stained smear should show well-differentiated stained regions, such as the acrosomal region, the midpiece, the tail, and the excess of residual cytoplasm, when present. An example of a poor and a good seminal smear staining is shown in Fig. 4 (A and B respectively). Accurate results can be achieved by following a standard operating procedure

(SOP) and when the test is performed by a higher-level trained technologist.

QC is crucial in sperm morphology assessment and involves not only the preparation of a good smear, but also a good staining technique. It includes checking the storage conditions, *i.e.* shelf-life, storage temperature, bacterial growth or oily surface film, and changing the stains frequently, depending on the volume/number of slides that have been stained. Reference slides prepared using semen samples of fertile donors or pre-assessed morphology slides can be used to check the quality of the freshly prepared stains. The quality of the stains should be monitored regularly either bi-weekly or monthly, according to usage and recorded on the QC forms.

QA is an ongoing process to monitor QC, scoring of morphology, reporting of results, and ensuring low inter- and intraobserver variability. A Bland Altman plot generated with morphology results is used to assess interobserver variability. Validation of sperm morphology results is achieved by testing the samples provided by an external testing agency such as the College of American Pathologists (CAP) in the United States. The laboratory results for sperm morphology analyte need to be within ± 2 standard deviation of the mean of the participating laboratories as reported by the CAP. The analytes are tested routinely like other patient samples in the laboratory, and the results are reported using the WHO 5th edition criteria. Corrective action must be taken if laboratory technician's results are out of the range [40]. Among the appropriate measures that could be taken are staff retraining/counseling, recalibration of the instruments, review of the current protocol to identify any potential causes of the discrepancy, and if needed, to implement a change in procedures or protocols, or to institute new laboratory checks. Ongoing monitoring should take place to minimize any recurrence [40,41]. All QA activities, including any problems identified, and corrective actions taken should be docu-



Cleveland
Clinic
©2021

Fig. 4. Example of (A) poor (left side) and (B) good (right side) staining quality.

mented [42].

Proper training of the andrology laboratory staff that follows the guidelines for the strict criteria is the best way to ensure correct evaluation of sperm morphology, uniformity across the laboratory and standardized results. To accomplish this, the Technical Supervisor should perform and document competency assessments through direct observation (side-by-side scoring), by monitoring the recording and reporting of test results, review of intermediate test results or worksheets, direct observation of performing instrument function checks, evaluation of previously tested samples, and assessment of problem-solving skills [43].

CHALLENGES IN MORPHOLOGICAL ASSESSMENT

1. Subjectivity in sperm morphology assessment

Even though sperm morphology is conducted routinely in the general and Andrology laboratories, unlike sperm motility or sperm concentration, it remains a challenge because of the large variability across the staining methods used, or the WHO guidelines followed (1999 *vs.* 2010) by practitioners. In addition, the subjectivity in scoring the sperm can also result from merely observing the sperm and categorizing it as normal or abnormal, without using an ocular micrometer, or confirming potential borderline normal forms with an experienced supervisor to avoid subjectivity. In addition to QC and QA, laboratories need to do an internal audit looking at their morphology scores in their various patient groups and correlate with pregnancies - either natural, IUI, IVF, or ICSI. If they find that a large proportion of their fertile patients are misdiagnosed with teratozoospermia then they would need to look at their

criteria for morphology assessment and perhaps revise their scoring accordingly.

2. Differences in technique/protocols in morphology assessment

Staining methods are very important for observing different regions of the spermatozoa under the microscope. Although the Papanicolaou stain is considered the gold standard for sperm morphology, many ART laboratories require a fast turn-around-time for diagnostic reporting of morphology results, and the Papanicolaou protocol can be time consuming. Rapid staining such as pre-stained Testsimplets and Diff-Quik have also been used. The staining procedure for Testsimplets does not require a fixation step and uses a wet mount on special pre-stained slides. Testsimplets however is not recommended by WHO as a standard staining method for the evaluation of sperm morphology [38]. Diff-Quik is a good alternative, which ascertains that results can be reported in a short time.

3. Inter- and intra-observer and laboratory variability

Inter- and intra-observer as well as laboratory variability are some of the challenges in morphological assessment, which can lead to inaccurate results. This can lead to misdiagnosis and mismanagement of the patient. It is important to keep a low inter-observer variability among technicians within the laboratory [44]. With the emphasis on strict morphological criteria, many laboratories appear to have become overly strict in their morphological assessment. As a result, many sperm that are potentially fertile, and therefore “normal”, may be labeled as “abnormal”, and when the majority of samples are diagnosed as teratozoospermic, the test loses its discriminatory ability. This may be the




Fig. 5. Schematic representation of how the micrometer is used to measure the dimensions of the sperm head.

reason why several recent studies [25] report that IUI pregnancy rates were unaffected by teratozoospermia. This is indeed a very important issue that needs to be addressed if morphology is to remain a useful predictor of fertility.

The improved predictive power of sperm morphology for ART outcome can be achieved by standardizing the methodology, using an optimal staining method such as Diff-Quik, and an ocular micrometer to measure the sperm dimensions correctly (Fig. 5) and following the WHO 5th edition guidelines for reporting results. The external QC for sperm morphology assessment is met by participating in proficiency testing and performing competency assessments of laboratory personnel. The lab needs to meet the quality standards for internal and external QC and eliminate the inter- and intra-observer variability.

CLINICAL IMPORTANCE OF SPERM MORPHOLOGY ASSESSMENT

In a prospective study with 45 couples undergoing IVF, Kruger et al [14] demonstrated that a threshold of >14% normal sperm morphology was associated with higher fertilization and pregnancy rates. The authors introduced a classification model where they referred to the >14% normal sperm morphology as normal or N-pattern. Patients showing normal morphology rates between 5% and 14% were referred to as good prognosis or G-pattern. In these patients, sperm morphology described by Kruger et al [14] was associated with a significantly higher fertilization rates (%) per oocyte and the average number of embryos per patient ($p=0.0001$). Sperm morphology of <5% was referred to as poor prognosis or P-pattern. Obara et al [18] examined the relationship between the percentage of normal sperm morphology and fertilization rate in IVF-embryo transfer (IVF-ET) cycles. The fertilization rate (80.5%) in 110 "normal" samples (>14% normal forms) was significantly higher than that (55.4%) in 27 samples with "poor prognosis" (those with 5%–14% normal forms) ($p<0.01$). The study concluded that ICSI may be required for achieving comparable fertilization rates in samples with a low percentage of normal forms [18].

Sperm morphology according to strict criteria has a strong association with fertilization. Numerous studies have also shown that sperm morphology and total progressive motile sperm count can help in the selection

of patients for either IVF or ICSI. In a meta-analysis, which included four retrospective cohort studies covering 2,853 IVF/ICSI cycles, patients with and without severe teratozoospermia (<2% normal forms) ($n=673$ and $2,183$, respectively) were analyzed [22]. Results showed that sperm morphology assessed by strict criteria had little or no prognostic value in ICSI cycle outcomes. However, it should be highlighted that a marked heterogeneity, with a relative risk ranging from 0.40 to 3.36 characterized the ICSI group [22].

A large observational study with 8,846 subjects undergoing 3,676 IVF/ICSI cycles concluded that sperm morphology has poor or no prognostic value in pregnancy outcomes [23]. However, the staining technique employed was not in accordance with the WHO 5th edition standards. Overall, studies assessing the predictive ability of sperm morphology have their own limitations, such as heterogeneity of study groups, staining methods used, scoring and classifications. The importance of morphology is during the initial assessment when consulting with the couple regarding natural conception, IUI or IVF due to the swimming patterns of normal forms. The interaction of the sperm with corona radiata and the sperm head with the oocyte receptors is important to allow acrosome reaction. However, in ICSI, the operator injects a sperm that appears as morphologically normal, unless using the intracytoplasmic morphologically selected sperm injection (IMSI method), where the subjectivity is a huge issue. In ICSI, a spermatozoon is forced into the oocyte independently of its strict morphology (apart from rough head/neck shape and vacuoles), as DNA content/chromatin quality are more important to determine the fate of the embryo. Teratozoospermia should be an independent indication for further clinical investigations as the assumption that poor morphology only requires ICSI misses the opportunity for the diagnosis of male partner. Furthermore, the use of a single sperm in ICSI bypasses natural selection and therefore sperm morphology may contribute to the lack of predictive ability due to this intervention.

Current studies that show the poor predictive ability of sperm morphology have many shortcomings including lack of standardization, deviation from WHO recommended protocols and different outcome measures. Strict criteria classification is based on the concept of natural selection. It can predict sperm fertilizing ability, indicate underlying pathology and be used as a

tool to choose the type of ART method employed. If the percentage of normal forms is $\geq 4\%$, then IUI or IVF can be performed. Alternatively, ICSI should be performed when the percentage of sperm with morphologically normal forms is $< 4\%$. Standardized morphology assessment by 'strict criteria' remains relevant in making clinical decisions for infertile couples. Future studies should examine the association between sperm morphology, ICSI and birth outcomes.

CLINICAL MANAGEMENT OF PATIENTS WITH ABNORMAL SPERM MORPHOLOGY

Kovac et al [24] reported that natural conception can still occur in up to 30% of men with absolute teratozoospermia (0% normal forms), suggesting that sperm morphology assessment may not be a robust predictor of infertility. On the other hand, various reports have shown significantly higher pregnancy rates in males with normal sperm morphology than that of those with abnormal sperm morphology [27,45,46]. Several studies also established a significant relationship between normal sperm morphology and vital sperm functions including acrosome reaction, binding to the zona pellucida, sperm DNA and chromatin integrity [47-51]. Therefore, treatment directed at improving the percentage of normal sperm morphology is undoubtedly warranted. Furthermore, morphology profiles are characteristic for specific clinical conditions and can therefore provide the clinician with valuable indications for the treatment [52]. Morphological profiles, as seen below, require specific treatments, without which fertilization and conception may not be feasible.

MORPHOLOGICAL PROFILES PROVIDING CLINICAL IMPLICATIONS

1. Globozoospermia

Globozoospermia is a unique sperm morphological defect characterized by round sperm head with the absence of acrosome, and identified in less than 0.1% of male infertility cases [53]. It may be associated with cytoskeletal defects in the sperm as well as sperm DNA fragmentation (SDF) [54-56]. Globozoospermia may be classified as total, with 100% of sperm showing the same defect (Type I), or partial, with the defect found

in up to 90% of sperm, and the presence of morphologically normal sperm in semen (Type II). Globozoospermia is frequently found in clusters in certain families denoting its genetic origin [57]. The semen analysis in such cases is usually characterized by a reduction in motility and normal morphology, with or without decrease in sperm concentration [53]. Furthermore, men with globozoospermia may carry a DPY19L2 mutation, which is common in North African men. Globozoospermic spermatozoa are not able to fertilize oocytes, as the acrosomal factor which initiates the oocyte activation, oscillin, is not present in globozoospermic sperm [53]. Moreover, the increased SDF and aneuploidy rates found in globozoospermic sperm decrease their fertilizing ability [53]. This can hinder natural pregnancy as well as the success of ART. ICSI is the only effective treatment in cases of complete globozoospermia, and it should be combined with oocyte activation using calcium ionophore to induce oocyte activation and fertilization [56].

2. Tapered sperm head

Sperm with elongated or pyriform-shaped heads are commonly seen in patients with testicular hyperthermia, commonly secondary to varicocele [58]. This morphological profile may be associated with an increased incidence of tail defects, sperm chromatin abnormalities and SDF, decreasing the fertilizing potential of the sperm [59,60]. These patients may benefit from varicocelectomy or, alternatively, selection of morphologically normal sperm during ICSI [61].

3. Macrozoospermia

Large-headed spermatozoa are defined as those with a length $> 4.7 \mu\text{m}$ and a width $> 3.2 \mu\text{m}$ [13]. A number of different terms have been used to refer to these spermatozoa, including "macrozoospermia" "macrocephalic sperm", "megalohead" spermatozoa, "enlarged-head spermatozoa" or "enlarged forms" "macronuclear spermatozoa" and "large head spermatozoa".

A high proportion of irregularly shaped, multi-tailed spermatozoa is also associated with severe male factor infertility. High rates of polyploidy and aneuploidy have been described in these cases, in association with Aurora kinase C mutation, frequently seen among North African men [57]. This specific sperm morphology carries a poor prognosis for ICSI, and therefore it is important to document in the semen analysis. Even

in cases of a moderate percentage of large-head sperm, genetic analysis such as sperm FISH could be useful to determine the ploidy of spermatozoa [62].

4. Pin head

A pin head is a spermatozoon lacking chromatin with only a tail [63]. This defect results from errors in the formation of the connecting piece during spermiogenesis. Recent genetic studies related the defect to a mutation in the *Sad1* and *UNC84* domain containing 5 (*SUN5*) gene, which is normally responsible for the formation of complexes linking the nucleus to the cytoskeleton [64]. Using pin head sperm in ICSI may result in fertilization but with no progression to cleavage-stage embryo [65,66].

5. Residual cytoplasmic droplet

Normal mature spermatozoa do not have a cytoplasmic droplets around the midpiece [67]. In contrast, immature, abnormal sperm have excess cytoplasm that is not removed during spermiogenesis. These are referred to as cytoplasmic droplets and reflect defective maturation processes, and epididymal function associated with infertility [68]. Cytoplasmic droplets are considered an anomaly when they exceed one third of the sperm head size, and they are prevalent in the semen of infertile patients with varicocele [35,69]. Smoking is reported to be associated with increased incidence of cytoplasmic droplets [70]. Antioxidant therapy, varicocelectomy along with lifestyle modification may be beneficial in restoring fertility in such types of sperm morphological abnormalities [71].

6. Tail defects

Tail defects are usually associated with asthenozoospermia. Stump tails and coiled tails are linked to toxic and chemicals substances' exposure [72]. In cases of primary ciliary dyskinesia, sperm tails may appear normal despite total sperm immotility. This is caused by the dynein arm defect in the sperm tail [73]. It is usually associated with chronic respiratory problems and with situs inversus in some cases (Kartagener's syndrome). ICSI can be performed in these cases using ejaculated sperm or testicular sperm. Additional testing such as the hypoosmotic swelling test should be performed to identify viable sperm. Studies have shown that immotile testicular sperm show a better prognosis in ICSI than in the immotile ejaculated sperm [74-76].

GENERAL TREATMENT MODALITIES FOR ABNORMAL SPERM MORPHOLOGY

1. Lifestyle modification and limitation of occupational exposure

Overall, certain lifestyle habits and occupational exposures are known to affect semen quality and fertility potential. However, few studies have investigated the impact of these risk factors on sperm morphology, revealing contradictory results. Pacey et al [77] indicated that modifiable lifestyle factors such as obesity, smoking or alcohol intake/ consumption have little effect on the percentage of sperm morphology. Similarly, Cherry et al [78] examined the occupational exposures of 2,011 men attending infertility clinics across the UK and failed to find any significant relation with sperm morphology. Several other studies reported a significant negative impact of alcohol and smoking on various semen parameters including sperm morphology [79-81]. Similarly, other studies reported a negative impact of exposure to heavy metals (lead, cadmium), bisphenol A, pesticides, radiation, and excessive heat on sperm morphology [82-85]. Regardless of whether lifestyle modification can affect semen quality, adopting healthy habits and avoiding environmental pollutants can positively influence overall health and longevity.

2. Antioxidant supplementation

Antioxidant supplementation is commonly used in the treatment of male infertility to alleviate the detrimental effects of oxidative stress on sperm production [86]. Although of low quality, recent evidence has shown a significant benefit for antioxidant use on conventional semen parameters [86,87]. A systematic review identified the antioxidants vitamin E, N-acetyl cysteine, lycopene, selenium, and zinc to be particularly useful when attempting to address abnormal sperm morphology [88].

3. Varicocelectomy

Varicocele is the most common correctable cause of male infertility. Its detrimental effects on various semen parameters have been established. Treatment is indicated in men with clinically palpable disease in the presence of abnormal semen parameters, including sperm morphology. The current evidence suggests that varicocele ligation significantly improves semen

parameters including morphology. It is also associated with an improvement in fertility potential [89].

4. Assisted reproduction

The use of morphologically normal sperm during ICSI has been shown to overcome most of the sperm abnormalities resulting in better reproductive outcomes. Certain morphological abnormalities such as globozoospermia and primary ciliary dyskinesia require the use of ICSI. Sperm selected for ICSI using ultra-high magnification are reportedly associated with better fertilization and pregnancy rates in couples with male factor infertility or those with a history of ART failure [90].

CONCLUSIONS

In conclusion, evaluation of sperm morphology requires laboratory skills with consistent training and monitoring and the use of an ocular micrometer for accurate measurement of sperm dimensions. Sources of error should be promptly identified, and rectified. QC and QA for sperm morphology are crucial for laboratory accreditation and practices for dependable results. Further, proficiency testing is an external QC tool for reporting accurate morphology results. Although the oocyte and the embryo are key players in ART, spermatozoa equally contribute to a successful outcome. Therefore, the interpretation of sperm morphology is important in the clinical management of infertile couples and in their choice of ART.

ACKNOWLEDGEMENTS

Authors are thankful to the artists from the Cleveland Clinic's Center for Medical Art & Photography for their help with the illustrations. The study was supported by the American Center for Reproductive Medicine.

Conflict of Interest

The authors have nothing to disclose.

Author Contribution

Conceptualization: AA, RSharma, SG. Writing – original draft: All the authors. Writing – review & editing: All the authors.

REFERENCES

1. Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, et al. The international glossary on infertility and fertility care, 2017. *Hum Reprod* 2017;32:1786-801.
2. Kumar N, Singh AK. Trends of male factor infertility, an important cause of infertility: a review of literature. *J Hum Reprod Sci* 2015;8:191-6.
3. Rutstein SO, Iqbal HS. Infecundity, infertility, and childlessness in developing countries. DHS comparative reports no. 9. Calverton (MD): ORC Macro; 2004. Co-published by the World Health Organization.
4. Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol Endocrinol* 2015;13:37.
5. Carlsen E, Giwercman A, Keiding N, Skakkebaek NE. Evidence for decreasing quality of semen during past 50 years. *BMJ* 1992;305:609-13.
6. Levine H, Jørgensen N, Martino-Andrade A, Mendiola J, Weksler-Derri D, Mindlis I, et al. Temporal trends in sperm count: a systematic review and meta-regression analysis. *Hum Reprod Update* 2017;23:646-59.
7. Mínguez-Alarcón L, Williams PL, Chiu YH, Gaskins AJ, Nassan FL, Dadd R, et al.; Earth Study Team. Secular trends in semen parameters among men attending a fertility center between 2000 and 2017: identifying potential predictors. *Environ Int* 2018;121(Pt 2):1297-303.
8. Trussell JC, Coward RM, Santoro N, Stetter C, Kunselman A, Diamond MP, et al.; Reproductive Medicine Network. Association between testosterone, semen parameters, and live birth in men with unexplained infertility in an intrauterine insemination population. *Fertil Steril* 2019;111:1129-34.
9. Agarwal A, Baskaran S, Parekh N, Cho CL, Henkel R, Vij S, et al. Male infertility. *Lancet* 2021;397:319-33.
10. Comhaire FH. Towards more objectivity in diagnosis and management of male infertility: results of a World Health Organization multicentre study. Oxford; Blackwell Scientific; 1987.
11. Menkveld R. Clinical significance of the low normal sperm morphology value as proposed in the fifth edition of the WHO laboratory manual for the examination and processing of human semen. *Asian J Androl* 2010;12:47-58.
12. World Health Organization (WHO). WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction. 4th ed. Cambridge: Cambridge University Press; 1999.
13. World Health Organization (WHO). WHO laboratory manual for the examination and processing of human semen. 5th

- ed. Geneva: World Health Organization; 2010.
14. Kruger TF, Menkveld R, Stander FS, Lombard CJ, Van der Merwe JP, van Zyl JA, et al. Sperm morphologic features as a prognostic factor in in vitro fertilization. *Fertil Steril* 1986;46:1118-23.
 15. Menkveld R, Stander FS, Kotze TJ, Kruger TF, van Zyl JA. The evaluation of morphological characteristics of human spermatozoa according to stricter criteria. *Hum Reprod* 1990;5:586-92.
 16. Kruger TF, Acosta AA, Simmons KF, Swanson RJ, Matta JF, Oehninger S. Predictive value of abnormal sperm morphology in in vitro fertilization. *Fertil Steril* 1988;49:112-7.
 17. Enginsu ME, Pieters MH, Dumoulin JC, Evers JL, Geraedts JP. Male factor as determinant of in-vitro fertilization outcome. *Hum Reprod* 1992;7:1136-40.
 18. Obara H, Shibahara H, Tsunoda H, Taneichi A, Fujiwara H, Takamizawa S, et al. Prediction of unexpectedly poor fertilization and pregnancy outcome using the strict criteria for sperm morphology before and after sperm separation in IVF-ET. *Int J Androl* 2001;24:102-8.
 19. Gatimel N, Moreau J, Parinaud J, Léandri RD. Sperm morphology: assessment, pathophysiology, clinical relevance, and state of the art in 2017. *Andrology* 2017;5:845-62.
 20. Zhu DL, Zhang HG, Wang RX, Jiang YT, Liu RZ. Re-evaluation of the value of sperm morphology in classical in vitro fertilization in a Northeastern Chinese population. *J Int Med Res* 2019;47:4134-42.
 21. French DB, Sabanegh ES Jr, Goldfarb J, Desai N. Does severe teratozoospermia affect blastocyst formation, live birth rate, and other clinical outcome parameters in ICSI cycles? *Fertil Steril* 2010;93:1097-103.
 22. Hotaling JM, Smith JF, Rosen M, Muller CH, Walsh TJ. The relationship between isolated teratozoospermia and clinical pregnancy after in vitro fertilization with or without intracytoplasmic sperm injection: a systematic review and meta-analysis. *Fertil Steril* 2011;95:1141-5.
 23. van den Hoven L, Hendriks JC, Verbeet JG, Westphal JR, Wetzels AM. Status of sperm morphology assessment: an evaluation of methodology and clinical value. *Fertil Steril* 2015;103:53-8.
 24. Kovac JR, Smith RP, Cajipe M, Lamb DJ, Lipshultz LI. Men with a complete absence of normal sperm morphology exhibit high rates of success without assisted reproduction. *Asian J Androl* 2017;19:39-42.
 25. Kohn TP, Kohn JR, Ramasamy R. Effect of sperm morphology on pregnancy success via intrauterine insemination: a systematic review and meta-analysis. *J Urol* 2018;199:812-22.
 26. Pantos K, Sfakianoudis K, Maziotis E, Rapani A, Karantzali E, Gounari-Papaioannou A, et al. Abnormal fertilization in ICSI and its association with abnormal semen parameters: a retrospective observational study on 1855 cases. *Asian J Androl* 2021. doi: 10.4103/aja.aja_84_20 [Epub].
 27. Jedrzejczak P, Taszarek-Hauke G, Hauke J, Pawelczyk L, Duleba AJ. Prediction of spontaneous conception based on semen parameters. *Int J Androl* 2008;31:499-507.
 28. Danis RB, Samplaski MK. Sperm morphology: history, challenges, and impact on natural and assisted fertility. *Curr Urol Rep* 2019;20:43.
 29. Talarczyk-Desole J, Berger A, Taszarek-Hauke G, Hauke J, Pawelczyk L, Jedrzejczak P. Manual vs. computer-assisted sperm analysis: can CASA replace manual assessment of human semen in clinical practice? *Ginekol Pol* 2017;88:56-60.
 30. Carrell DT, De Jonge CJ. The troubling state of the semen analysis. *Andrology* 2016;4:761-2.
 31. Schallmoser A, Bakjaji F, Königsberger S, John J, Färber C, Schmidt E, et al. Effect of mild α -chymotrypsin treatment of highly viscous semen samples on fertilization rates. *Transl Androl Urol* 2021;10:448-54.
 32. Zhu WJ. Preparation and observation methods can produce misleading artefacts in human sperm ultrastructural morphology. *Andrologia* 2018;50:e13043.
 33. Agarwal A, Gupta S, Sharma R. Sperm morphology stain (Diff-Quik®). In: Agarwal A, Gupta S, Sharma R, editors. *Andrological evaluation of male infertility: a laboratory guide*. Cham: Springer; 2016;79-82.
 34. Menkveld R, Wong WY, Lombard CJ, Wetzels AM, Thomas CM, Merkus HM, et al. Semen parameters, including WHO and strict criteria morphology, in a fertile and subfertile population: an effort towards standardization of in-vivo thresholds. *Hum Reprod* 2001;16:1165-71.
 35. Mortimer D, Menkveld R. Sperm morphology assessment--historical perspectives and current opinions. *J Androl* 2001; 22:192-205.
 36. Singh S, Sharma S, Jain M, Chauhan R. Importance of papanicolaou staining for sperm morphologic analysis: comparison with an automated sperm quality analyzer. *Am J Clin Pathol* 2011;136:247-51.
 37. Oral E, Yetis O, Elibol F, Senol H, Irez T, Aksu FM. Assessment of human sperm morphology by strict criteria: comparison of wet preparation versus stained with the modified Diff-Quik method. *Arch Androl* 2002;48:307-14.
 38. Henkel R, Schreiber G, Sturmhoefel A, Hipler UC, Zermann DH, Menkveld R. Comparison of three staining methods for the morphological evaluation of human spermatozoa. *Fertil Steril* 2008;89:449-55.
 39. Björndahl L, Mortimer D, Barratt CLR, Castilla JA, Menkveld

- R, Kvist U, et al. A practical guide to basic laboratory andrology. Cambridge: Cambridge University Press; 2010.
40. College of American Pathologists. Commission on laboratory accreditation reproductive laboratory checklist: laboratory accreditation program. Northfield (IL): College of American Pathologists; 2007;1-70.
 41. Brazil C, Swan SH, Tollner CR, Treece C, Drobnis EZ, Wang C, et al.; Study for Future Families Research Group. Quality control of laboratory methods for semen evaluation in a multicenter research study. *J Androl* 2004;25:645-56.
 42. Keel BA. Quality control, quality assurance, and proficiency testing in the andrology laboratory. *Arch Androl* 2002;48:417-31.
 43. U.S. Departments and Agencies of the Federal Government. 42 CFR Ch. IV. Part 493 - laboratory requirements. In: U.S. Government, editor. Code of Federal Regulations. Washington, D.C.: U.S. Government; 2011;516-637.
 44. Comhaire F, Schoonjans F, Vermeulen L, De Clercq N. Methodological aspects of sperm morphology evaluation: comparison between strict and liberal criteria. *Fertil Steril* 1994;62:857-61.
 45. Sripada S, Townend J, Campbell D, Murdoch L, Mathers E, Bhattacharya S. Relationship between semen parameters and spontaneous pregnancy. *Fertil Steril* 2010;94:624-30.
 46. Kruger TF, Lacquet FA, Sarmiento CA, Menkveld R, Ozgür K, Lombard CJ, et al. A prospective study on the predictive value of normal sperm morphology as evaluated by computer (IVOS). *Fertil Steril* 1996;66:285-91.
 47. Majzoub A, Arafa M, Mahdi M, Agarwal A, Al Said S, Al-Emadi I, et al. Oxidation-reduction potential and sperm DNA fragmentation, and their associations with sperm morphological anomalies amongst fertile and infertile men. *Arab J Urol* 2018;16:87-95.
 48. Liu DY, Baker HW. Disordered acrosome reaction of spermatozoa bound to the zona pellucida: a newly discovered sperm defect causing infertility with reduced sperm-zona pellucida penetration and reduced fertilization in vitro. *Hum Reprod* 1994;9:1694-700.
 49. Liu DY, Baker HW. Calcium ionophore-induced acrosome reaction correlates with fertilization rates in vitro in patients with teratozoospermic semen. *Hum Reprod* 1998;13:905-10.
 50. Abu Hassan Abu D, Franken DR, Hoffman B, Henkel R. Accurate sperm morphology assessment predicts sperm function. *Andrologia* 2012;44 Suppl 1:571-7.
 51. Brahem S, Mehdi M, Elghezal H, Saad A. Detection of DNA fragmentation and meiotic segregation in human with isolated teratozoospermia. *J Assist Reprod Genet* 2011;28:41-8.
 52. Auger J, Jouannet P, Eustache F. Another look at human sperm morphology. *Hum Reprod* 2016;31:10-23.
 53. Fesahat F, Henkel R, Agarwal A. Globozoospermia syndrome: an update. *Andrologia* 2020;52:e13459.
 54. Vozdova M, Rybar R, Kloudova S, Prinosilova P, Textl P, Rubes J. Total globozoospermia associated with increased frequency of immature spermatozoa with chromatin defects and aneuploidy: a case report. *Andrologia* 2014;46:831-6.
 55. Han F, Liu C, Zhang L, Chen M, Zhou Y, Qin Y, et al. Globozoospermia and lack of acrosome formation in GM130-deficient mice. *Cell Death Dis* 2017;8:e2532.
 56. Eskandari N, Tavalae M, Zohrabi D, Nasr-Esfahani MH. Association between total globozoospermia and sperm chromatin defects. *Andrologia* 2018;50:e12843.
 57. Ounis L, Zoghmar A, Coutton C, Rouabah L, Hachemi M, Martinez D, et al. Mutations of the aurora kinase C gene causing macrozoospermia are the most frequent genetic cause of male infertility in Algerian men. *Asian J Androl* 2015;17:68-73.
 58. Parikh FR, Kamat SA, Kodwaney GG, Balaiah D. Computer-assisted semen analysis parameters in men with varicocele: is surgery helpful? *Fertil Steril* 1996;66:440-5.
 59. Lee JD, Kamiguchi Y, Yanagimachi R. Analysis of chromosome constitution of human spermatozoa with normal and aberrant head morphologies after injection into mouse oocytes. *Hum Reprod* 1996;11:1942-6.
 60. Prisant N, Escalier D, Soufir JC, Morillon M, Schoevaert D, Misrahi M, et al. Ultrastructural nuclear defects and increased chromosome aneuploidies in spermatozoa with elongated heads. *Hum Reprod* 2007;22:1052-9.
 61. Prasivoravong J, Marcelli F, Lemaître L, Pigny P, Ramdane N, Peers MC, et al. Beneficial effects of varicocele embolization on semen parameters. *Basic Clin Androl* 2014;24:9.
 62. Guthauser B, Pollet-Villard X, Boitrelle F, Vialard F. Is intracouple assisted reproductive technology an option for men with large-headed spermatozoa? A literature review and a decision guide proposal. *Basic Clin Androl* 2016;26:8.
 63. Ray PF, Toure A, Metzler-Guillemain C, Mitchell MJ, Arnoult C, Coutton C. Genetic abnormalities leading to qualitative defects of sperm morphology or function. *Clin Genet* 2017;91:217-32.
 64. Yassine S, Escoffier J, Abi Nahed R, Pierre V, Karaouzene T, Ray PF, et al. Dynamics of Sun5 localization during spermatogenesis in wild type and Dpy19L2 knock-out mice indicates that Sun5 is not involved in acrosome attachment to the nuclear envelope. *PLoS One* 2015;10:e0118698.
 65. Chemes HE, Puigdomenech ET, Carizza C, Olmedo SB, Zanchetti F, Hermes R. Acephalic spermatozoa and abnormal development of the head-neck attachment: a human syn-

- drome of genetic origin. *Hum Reprod* 1999;14:1811-8.
66. Saïas-Magnan J, Metzler-Guillemain C, Mercier G, Carles-Marcorelles F, Grillo JM, Guichaoua MR. Failure of pregnancy after intracytoplasmic sperm injection with decapitated spermatozoa: case report. *Hum Reprod* 1999;14:1989-92.
 67. Huszar G, Vigue L. Spermatogenesis-related change in the synthesis of the creatine kinase B-type and M-type isoforms in human spermatozoa. *Mol Reprod Dev* 1990;25:258-62.
 68. Gomez E, Buckingham DW, Brindle J, Lanzafame F, Irvine DS, Aitken RJ. Development of an image analysis system to monitor the retention of residual cytoplasm by human spermatozoa: correlation with biochemical markers of the cytoplasmic space, oxidative stress, and sperm function. *J Androl* 1996;17:276-87.
 69. Zini A, Defreitas G, Freeman M, Hechter S, Jarvi K. Varicocele is associated with abnormal retention of cytoplasmic droplets by human spermatozoa. *Fertil Steril* 2000;74:461-4.
 70. Mak V, Jarvi K, Buckspan M, Freeman M, Hechter S, Zini A. Smoking is associated with the retention of cytoplasm by human spermatozoa. *Urology* 2000;56:463-6.
 71. Rengan AK, Agarwal A, van der Linde M, du Plessis SS. An investigation of excess residual cytoplasm in human spermatozoa and its distinction from the cytoplasmic droplet. *Reprod Biol Endocrinol* 2012;10:92.
 72. Dávila Garza SA, Patrizio P. Reproductive outcomes in patients with male infertility because of Klinefelter's syndrome, Kartagener's syndrome, round-head sperm, dysplasia fibrous sheath, and 'stump' tail sperm: an updated literature review. *Curr Opin Obstet Gynecol* 2013;25:229-46.
 73. Sha YW, Ding L, Li P. Management of primary ciliary dyskinesia/Kartagener's syndrome in infertile male patients and current progress in defining the underlying genetic mechanism. *Asian J Androl* 2014;16:101-6.
 74. Nijs M, Vanderzwalmen P, Vandamme B, Segal-Bertin G, Lejeune B, Segal L, et al. Fertilizing ability of immotile spermatozoa after intracytoplasmic sperm injection. *Hum Reprod* 1996;11:2180-5.
 75. Westlander G, Barry M, Petrucco O, Norman R. Different fertilization rates between immotile testicular spermatozoa and immotile ejaculated spermatozoa for ICSI in men with Kartagener's syndrome: case reports. *Hum Reprod* 2003;18:1286-8.
 76. Esteves SC, Sánchez-Martín F, Sánchez-Martín P, Schneider DT, Gosálvez J. Comparison of reproductive outcome in oligozoospermic men with high sperm DNA fragmentation undergoing intracytoplasmic sperm injection with ejaculated and testicular sperm. *Fertil Steril* 2015;104:1398-405.
 77. Pacey AA, Povey AC, Clyma JA, McNamee R, Moore HD, Baillie H, et al.; Participating Centres of Chaps-UK. Modifiable and non-modifiable risk factors for poor sperm morphology. *Hum Reprod* 2014;29:1629-36.
 78. Cherry N, Povey AC, McNamee R, Moore H, Baillie H, Clyma JA, et al.; participating centres of CHAPS-UK. Occupation exposures and sperm morphology: a case-referent analysis of a multi-centre study. *Occup Environ Med* 2014;71:598-604.
 79. Boeri L, Capogrosso P, Ventimiglia E, Pederzoli F, Cazzaniga W, Chierigo F, et al. Heavy cigarette smoking and alcohol consumption are associated with impaired sperm parameters in primary infertile men. *Asian J Androl* 2019;21:478-85.
 80. Gaur DS, Talekar M, Pathak VP. Effect of cigarette smoking on semen quality of infertile men. *Singapore Med J* 2007;48:119-23.
 81. Gaur DS, Talekar MS, Pathak VP. Alcohol intake and cigarette smoking: impact of two major lifestyle factors on male fertility. *Indian J Pathol Microbiol* 2010;53:35-40.
 82. Zamkowska D, Karwacka A, Jurewicz J, Radwan M. Environmental exposure to non-persistent endocrine disrupting chemicals and semen quality: an overview of the current epidemiological evidence. *Int J Occup Med Environ Health* 2018;31:377-414.
 83. Wijsekera GU, Fernando DM, Wijerathna S, Bandara N. Environmental and occupational exposures as a cause of male infertility. *Ceylon Med J* 2015;60:52-6.
 84. Pollard SH, Cox KJ, Blackburn BE, Wilkins DG, Carrell DT, Stanford JB, et al. Male exposure to bisphenol A (BPA) and semen quality in the home observation of periconceptual exposures (HOPE) cohort. *Reprod Toxicol* 2019;90:82-7.
 85. Meeker JD, Rossano MG, Protas B, Diamond MP, Puscheck E, Daly D, et al. Cadmium, lead, and other metals in relation to semen quality: human evidence for molybdenum as a male reproductive toxicant. *Environ Health Perspect* 2008;116:1473-9.
 86. Agarwal A, Leisegang K, Majzoub A, Henkel R, Finelli R, Panner Selvam MK, et al. Utility of antioxidants in the treatment of male infertility: clinical guidelines based on a systematic review and analysis of evidence. *World J Mens Health* 2021;39:233-90.
 87. Arafa M, Agarwal A, Majzoub A, Panner Selvam MK, Basakaran S, Henkel R, et al. Efficacy of antioxidant supplementation on conventional and advanced sperm function tests in patients with idiopathic male infertility. *Antioxidants (Basel)* 2020;9:219.
 88. Majzoub A, Agarwal A. Systematic review of antioxidant types and doses in male infertility: benefits on semen parameters, advanced sperm function, assisted reproduction and live-birth rate. *Arab J Urol* 2018;16:113-24.
 89. Agarwal A, Deepinder F, Cocuzza M, Agarwal R, Short RA,

Sabanegh E, et al. Efficacy of varicoelectomy in improving semen parameters: new meta-analytical approach. *Urology* 2007;70:532-8.

90. Karabulut S, Aksunger O, Korkmaz O, Eren Gozel H, Keskin I. Intracytoplasmic morphologically selected sperm injection, but for whom? *Zygote* 2019;27:299-304.