



**HAL**  
open science

## Lactoferrin/ $\beta$ -lactoglobulin complex coacervates: rheological properties

Hachfi Soussi Rima, Marie-Hélène Famelart, Florence Rousseau, Pascaline  
Hamon, Said Bouhallab

### ► To cite this version:

Hachfi Soussi Rima, Marie-Hélène Famelart, Florence Rousseau, Pascaline Hamon, Said Bouhallab. Lactoferrin/ $\beta$ -lactoglobulin complex coacervates: rheological properties. *Genova Colloids 2021*, Apr 2021, Edition virtuelle, Switzerland. . hal-03279352

**HAL Id: hal-03279352**

**<https://hal.inrae.fr/hal-03279352v1>**

Submitted on 6 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0  
International License

# Lactoferrin/ $\beta$ -lactoglobulin complex coacervates: rheological properties

Soussi, H.R., Famelart M.H., Rousseau F., Hamon, P., Bouhallab, S.

*INRAE, Institut Agro, STLO, F-35042 Rennes, France*

[rma.hachfi-soussi@inrae.fr](mailto:rma.hachfi-soussi@inrae.fr)

Abstract: Complex coacervation is a liquid-liquid phase separation that leads to the formation of highly concentrated phase (coacervates). Despite its multiple use, especially in the field of food science, little has been reported on hetero-protein complex coacervation. In our previous work, we reported on the ability of two globular proteins Lactoferrin (LF) and  $\beta$ -lactoglobulin ( $\beta$ -LG) to form coacervates under specific physicochemical conditions [1]. Here, different experiments were conducted in order to characterize the rheological properties of the formed LF/ $\beta$ -LG coacervates. We showed that unlike some polysaccharide/protein systems, LF/ $\beta$ -LG coacervates has a liquid like behavior; the loss modulus  $G''$  was 100 times higher than the storage modulus  $G'$ . This behavior was confirmed under creep-recovery tests. In addition, the coacervates exhibited a Newtonian viscous flow under low shear rate and a shear thinning behavior above  $10\text{ s}^{-1}$ . These results suggest a structural change probably due to the weaknesses of electrostatic interactions inside the protein network at high shear rates. This structural change was proved to be reversible and the rearrangement of the structure was time-dependent. These results allow better understanding of the involved interactions in concentrated protein solutions and open new avenues for the use of coacervates as texturizing agents in food products.

Key words: Complex coacervation, Co-assembly, Rheology,  $\beta$ -Lactoglobulin, Lactoferrin.

## References:

- [1]. Tavares, G. M., Croguennec, T., Hamon, P., Carvalho, A. F. & Bouhallab, S. Selective coacervation between lactoferrin and the two isoforms of  $\beta$ -lactoglobulin. (2015) *Food Hydrocolloids* 48, 238–247.