



## Rheological properties of lactoferrin / $\beta$ -lactoglobulin complex coacervates

Hachfi Soussi Rima, Marie-Hélène Famelart, Florence Rousseau, Pascaline Hamon, Said Bouhallab

### ► To cite this version:

Hachfi Soussi Rima, Marie-Hélène Famelart, Florence Rousseau, Pascaline Hamon, Said Bouhallab. Rheological properties of lactoferrin / $\beta$ -lactoglobulin complex coacervates. Journées scientifiques Ecole doctorale EGAAL, Yannick OUTREMAN, Professeur Agrocampus Ouest, Alain VIAN, Professeur Université d'Angers; Hervé PREVOST, Professeur Oniris Nantes; Annick CHAUVIN, Professeure Université Rennes 1, Jun 2021, Rennes édition virtuelle, France. hal-03279366

**HAL Id: hal-03279366**

**<https://hal.inrae.fr/hal-03279366>**

Submitted on 6 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

✓ Oral presentation



## **RHEOLOGICAL PROPERTIES OF LACTOFERRIN/ $\beta$ -LACTOGLOBULIN COMPLEX COACERVATES**

Soussi, H.R.\*, Famelart M.H., Rousseau F., Hamon, P., Bouhallab S.

*INRAE, Institut Agro, STLO, F-35042 Rennes, France*

**Key words:** Alimentation, Complex coacervation, Rheology

Complex coacervation is a liquid-liquid phase separation that leads to the formation of a highly concentrated phase or coacervates. The formed coacervates are proposed as carriers for bioactives because of their high encapsulation efficiency under mild processing conditions. Despite its multiple use, especially in the fields of food science, little has been reported on hetero-protein complex coacervation. In our previous work, we reported on the ability of two globular proteins, Lactoferrin (LF) and  $\beta$ -lactoglobulin ( $\beta$ -LG) to form coacervates under specific physicochemical conditions <sup>[1]</sup>. In the present work, we aimed to characterize the rheological properties of the formed LF/ $\beta$ -LG coacervates. We show that unlike some polysaccharide/protein systems, LF/ $\beta$ -LG coacervates has a liquid-like behavior; the loss modulus  $G''$  was 100 times higher than the storage modulus  $G'$ . This behavior was confirmed under creep-recovery tests. In addition, the coacervates exhibited a Newtonian viscous flow under low shear rate and a shear thinning behavior above  $10 \text{ s}^{-1}$ . These results suggest a structural change probably due to the weaknesses of electrostatic interactions inside the coacervates at high shear rates. This structural change was proved to be reversible and the rearrangement of the structure was time-dependent. These results allow better understanding of the involved interactions in concentrated protein solutions and open new avenues for the use of coacervates as texturizing agents in food matrices.

### **References:**

- [1]. Tavares, G. M., Croguennec, T., Hamon, P., Carvalho, A. F. & Bouhallab, S. Selective coacervation between lactoferrin and the two isoforms of  $\beta$ -lactoglobulin. (2015) *Food Hydrocolloids* 48, 238–247.