

# Determination of the Regulon of CcpA, a pleiotropic regulator, in the pathogen Streptococcus agalactiae

Anne-Emmanuelle Roux, Laurent Mereghetti, Emilie Camiade

#### ▶ To cite this version:

Anne-Emmanuelle Roux, Laurent Mereghetti, Emilie Camiade. Determination of the Regulon of CcpA, a pleiotropic regulator, in the pathogen Streptococcus agalactiae. World Microbe Forum, Jun 2021, Online, France. hal-03280053

## HAL Id: hal-03280053 https://hal.inrae.fr/hal-03280053

Submitted on 7 Jul2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### Determination of the Regulon of CcpA, a Pleiotropic Regulator, in the Pathogen Streptococcus agalactiae

<u>Anne-Emmanuelle Roux</u> <sup>(a)</sup>, Laurent Mereghetti <sup>(a,b)</sup> and Emilie Camiade <sup>(a)</sup> (a) ISP, Université de Tours, INRAE, Tours, F-37000, France

(b) CHRU de Tours, Service de Bactériologie-Virologie, F-37044 Tours, France

## Abstract

The commensal bacterium *Streptococcus agalactiae* is responsible for various infections in a wide variety of hosts including humans. Its broad spectrum of hosts shows its ability to acquire nutrients in variable conditions. The carbon catabolite repression allows bacteria to prioritize the uptake and the catabolism of the environmental sugars. In Gram-positive bacteria, CcpA (catabolite control protein A), a pleiotropic transcriptional regulator, plays a key role in catabolite repression. Complexed to its phosphorylated co-effector HPr (histidine-containing phosphocarrier protein), CcpA binds to a specific DNA consensus sequence named « *cre* » (for catabolite response element) in the regulatory region of the target genes, thereby causing their activation or repression. Studies have shown the involvement of carbon catabolite repression in the adaptation and stress resistance of pathogenic bacteria. The goal of this study is to determine the regulon and the role(s) of CcpA in the physiology and adaptation of *S. agalactiae*.

We studied the effect of CcpA deletion in *Streptococcus agalactiae* A909 by comparing, with an RNAseq analysis, the global transcription of the WT and  $\Delta ccpA$  strains. We showed that 13.5% of the *S. agalactiae* genome was regulated by CcpA and that it acts mostly as a repressor.

In order to determine the direct targets of CcpA, we searched the genes harbouring a putative cre site in the *S. agalactiae* genome with an in-silico approach (RegPrecise, Virtual Footprint). Among the 274 genes significantly regulated by CcpA, 79 have at least one *cre* site. We are validating the consensus cre sequence with gelshift experiments on approximatively 20 putative direct targets.

In-silico and RNAseq analysis highlight the CcpA direct regulation of genes coding for proteins putatively involved in acidic and oxidative stresses. As an example, one of these proteins coded by the SAK\_1689 gene that is highly overexpressed in the  $\Delta ccpA$  strain, has more than 50% identity with *lmo\_1580*, an UspA protein of *Listeria monocytogenes* involved in the oxidative and stress responses. We are constructing transcriptional fusion vectors in order to confirm the implication of CcpA in their regulation and their expression in stress responses, and therefore the role of this regulator in the adaptation of *S. agalactiae*.