

# On synergistic co-infection in crop vector-borne diseases

Americo Matusse, Michael Chapwanya, Yves Dumont

### ▶ To cite this version:

Americo Matusse, Michael Chapwanya, Yves Dumont. On synergistic co-infection in crop vector-borne diseases. BIOMATH 2021 - International Conference on Mathematical Methods and Models in Biosciences, Jun 2021, Pretoria, South Africa. 2 p. hal-03282739

HAL Id: hal-03282739 https://hal.inrae.fr/hal-03282739

Submitted on 9 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## BIOMATH 2021

International Conference on Mathematical Methods and Models in Biosciences

20 - 25 June 2021 Protea Hotel, Pretoria Loftus Park South Africa

 $\begin{array}{c} \textbf{Hosted by Department of} \\ \textbf{Mathematics \& Applied Mathematics} \\ \textbf{and DSI/NRF SARChI Chair } M^3B^2 \\ \textbf{University of Pretoria} \end{array}$ 

## On synergistic co-infection in crop vector-borne diseases

<u>Americo Matusse</u><sup>1</sup> & Michael Chapwanya<sup>1</sup> & Yves Dumont<sup>1,2</sup>
University of Pretoria, Department of Mathematics and Applied Mathematics, South Africa

<sup>1</sup>CIRAD, UMR AMAP, St Pierre, F-97410, Reunion Island, France ajmatusse@gmail.com

Major constraint to crop productivity are pests and diseases that cause major food insecurity throughout the world, especially in Southern Countries. The Food and Agriculture Organization estimates that pests and diseases are responsible for about 25% of crop loss. A new deadly disease of maize was reported in many Southern Countries, the Maize Lethal Necrosis Disease (MLND). For instance, in Kenya alone, the MLND affected around 77 000 ha, translating into an estimated loss of US \$ 52 million. The MLND results from a synergistic interaction between the Maize Chlorotic Mottle Virus (MCMV) and one of several viruses from the Potyviridae family, like the Sugarcane Mosaic Virus (SCMV). In this talk, we focus on modelling and analysis of MLND. However, our model is sufficiently generic to be applied to different co-infection vector borne diseases. A theoretical analysis shows that different equilibria and thresholds exist: the basic reproduction numbers and the invasion reproduction numbers. We show that these thresholds drive the dynamics of the MLND system. In particular, the invasion reproduction numbers are essential for the emergence or not of the MLND. We illustrate our results through numerical simulations and discuss potential control methods. This work has been published in [1].

**Key words:** Crop disease, synergistic interaction, basic reproduction number, invasion reproduction number, numerical simulations.

#### References

[1] Michael Chapwanya, Americo Matusse, and Yves Dumont. On synergistic coinfection in crop diseases. The case of the maize lethal necrosis disease. Applied Mathematical Modelling, 90:912-942, 2021.