



HAL
open science

Nitric oxide production is involved in maintaining energy state in Alfalfa (*Medicago sativa* L.) nodulated roots under both salinity and flooding

Fatma Aridhi, Hajer Sghaier, Allyzée Gaitanaros, Ayda Khadri, Samira Aschi-Smiti, Renaud Brouquisse

► To cite this version:

Fatma Aridhi, Hajer Sghaier, Allyzée Gaitanaros, Ayda Khadri, Samira Aschi-Smiti, et al.. Nitric oxide production is involved in maintaining energy state in Alfalfa (*Medicago sativa* L.) nodulated roots under both salinity and flooding. *Planta*, 2020, 252 (2), 10.1007/s00425-020-03422-1 . hal-03285554

HAL Id: hal-03285554

<https://hal.inrae.fr/hal-03285554v1>

Submitted on 28 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nitric oxide production is involved in maintaining energy state in Alfalfa (*Medicago sativa* L.) nodulated roots under both salinity and flooding

Fatma Aridhi^{1,2} · Hajer Sghaier¹ · Allyzée Gaitanaros² · Ayda Khadri¹ · Samira Aschi-Smiti¹ · Renaud Brouquisse² 

Accepted: 8 July 2020 / Published online: 16 July 2020

Abstract

Main conclusion In *Medicago sativa* nodulated roots, NR-dependent NO production is involved in maintaining energy state, presumably through phytohemoglobin NO respiration, under both salinity and hypoxia stress.

Abstract The response to low and average salinity stress and to a 5 day-long flooding period was analyzed in *M. sativa* nodulated roots. The two treatments result in a decrease in the biological nitrogen fixation capacity and the energy state (evaluated by the ATP/ADP ratio), and conversely in an increase nitric oxide (NO) production. Under salinity and hypoxia treatments, the use of either sodium tungstate, an inhibitor of nitrate reductase (NR), or carboxy-PTIO, a NO scavenger, results in a decrease in NO production and ATP/ADP ratio, meaning that NR-dependent NO production participates to the maintenance of the nodulated roots energy state.

Keywords Hypoxia · Legume · Nitrogen-fixing symbiosis · Phytohemoglobin NO respiration · Salt stress

✉ Renaud Brouquisse
renaud.brouquisse@inrae.fr

Fatma Aridhi
aridhifatma@gmail.com

Hajer Sghaier
sghaier.hajer@hotmail.com

Allyzée Gaitanaros
allyzee.gaitanaros@hotmail.fr

Ayda Khadri
khadriayda@yahoo.fr

Samira Aschi-Smiti
samira.smiti@planet.tn

¹ Unité de Recherche d'Ecologie Végétale, Faculté des Sciences de Tunis, Université de Tunis El Manar, Campus Universitaire Farhat Hached, Tunis, Tunisia

² UMR INRAE 1355, CNRS 7254, Université Côte d'Azur, Institut Sophia Agrobiotech, Sophia Antipolis, France

Abbreviations

ARA	Acetylene reducing activity
BNF	Biological nitrogen fixation
cPTIO	2-[4-carboxyphenyl]-4,4,5,5-tetramethylimidazole-1-oxyl-3-oxide
DAF-2	4,5-diaminofluorescein
NR	Nitrate reductase
Phytogb	Phytohemoglobin

Introduction

Nitric oxide (NO) is a reactive gaseous molecule with a broad spectrum of regulatory functions in plant growth and development (Besson-Bard et al. 2008), and response to biotic (Hichri et al. 2015; Thalineau et al. 2016) and abiotic stresses, including salinity and hypoxia (Blokina and Fagerstedt 2010; Simontacchi et al. 2015). Under salt stress, enhancement of NO production is accompanied by the exclusion of Na⁺ and retention of K⁺, through increased membrane H⁺-ATPase and H⁺-PPase activities (Zhang et al. 2006; Wang et al. 2009; Zhao et al. 2018). NO induces the expression of genes related to osmotic adjustment, heat-shock protein, and ROS-scavenging processes (Uchida et al. 2002; Kopyra and Gwózdź 2003). It contributes to the

maintenance of cellular redox homeostasis by controlling the NADPH level (Liu et al. 2007) and by inducing antioxidant enzyme activities (Tanou et al. 2009). Overall speaking, NO protects plants by helping them to control water status, maintain ionic homeostasis and reduce oxidative damage imposed during salt stress (Molassiotis et al. 2010; Simon-tacchi et al. 2015).

On the other hand, increased NO production is a hallmark of plant response to flooding and hypoxia (Blokhina and Fagerstedt 2010; Gupta and Igamberdiev 2011). Hypoxia compromises mitochondrial respiration and leads to an insufficiency in ATP for energy-demanding processes (Bailey-Serres and Voesenek 2008). In these conditions, NO contributes to the recycling of NADH and the synthesis of ATP through the setting up of an alternative respiration called “phytoglobin (Phytogb)-NO respiration” (Stoimenova et al. 2007; Igamberdiev and Hill 2009; Gupta and Igamberdiev 2011). NO is also essential for the development of lysigenous aerenchyma that enhance O₂ diffusion along with the roots (Wany et al. 2017). Furthermore, the signaling role of NO in the perception and the response to hypoxia was recently elucidated in *Arabidopsis thaliana* (Hartman et al. 2019). An increase in NO production, therefore, appears to be a common response to both salt stress and hypoxia.

Exposure of nitrogen-fixing legumes to high salt concentrations resulted in a rapid decrease in nitrogenase activity and nodule respiration, and this decrease was compensated by raising pO₂ (Serraj et al. 1994; Serraj and Drevon 1998). Similarly, by comparing nodule respiration of various legumes under NaCl treatments, Delgado et al. (1994) concluded that under low salinity the supply of O₂ to nodules is the limiting factor of the biological nitrogen fixation (BNF). These observations suggest that under salt stress, the nodulated roots of legumes could also face hypoxia and that this could limit BNF.

To test this hypothesis, using the *Medicago sativa* legume model, we analyzed BNF, NO production and energy state in the nodulated roots of plants grown in the presence of low to average salinity and subjected to short-term flooding. Our data show that, under moderate salt stress as under flooding, a nitrate reductase (NR)-dependent NO production is induced which participates to the maintenance of the energy state of the root system.

Materials and methods

Biological material and growth conditions

Seeds of *Medicago sativa* (var. Siriver; obtained from Espave Vert Tunisie, Tunis, Tunisia) were sterilized and germinated as previously described (del Giudice et al. 2011). Five-day-old germinated seeds were transplanted

in pots (2 plants/pot) filled with 250 cm³ of B5 sand (0.6–1.6 mm diameter). Plants growth conditions and nutrient solution are described in Sghaier et al. (2020). To set salinity conditions, nutrient solution also contained either 0.2 mM KNO₃ (Control, Ctrl), 0.2 mM KNO₃ and 20 mM NaCl (Treatment 20, T20), or 0.9 mM KNO₃ and 50 mM NaCl (Treatment 50, T50). Plants were inoculated 7 days after transplanting with *Ensifer meliloti* 2011 bacteria (Sghaier et al. 2020).

In the first set of experiments (experiment 1), at 28 days post-inoculation (dpi), half of the Ctrl, T20 and T50 plants were subjected to flooding for 5 days by submerging pots in their respective nutrient solutions to sand level (Fig. S1a). Non-flooded plants were watered with nutrient solutions at 28 and 31 dpi (Fig. S1b). At 32 dpi, nodulated roots were harvested and either used for growth parameters analysis (9 plants/assay), N₂-fixing capacity (2 plants/assay), NO production (2 plants/assay), or immediately frozen in liquid N₂, ground in powder and stored at – 80 °C until analysis of protein and adenine nucleotide contents (2 plants/assay).

In the second set of experiments (experiment 2), at 28 dpi, the plants were transferred either into syringes filled with 35 ml nutrient solution and traversed by a humidified airflow (normoxia, Fig. S2a and b) or into glass tubes filled with 40 ml nutrient solution previously flushed with a mixture of 4.5–95.5%: O₂-N₂ (hypoxia, Fig. S2c and d). The plants (2 plants/syringe or tube) were cultured either in normoxia or hypoxia for 5 days. To compensate for the evaporation, syringes and tubes were supplemented daily with nutrient solution. One day before harvest, nutrient solutions were supplemented with either 0.1 mM 2-[4-carboxyphenyl]-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (cPTIO), a scavenger of NO, or 1 mM sodium tungstate (Tg), an inhibitor of NR, or water (control). At 32 dpi, nodulated root was promptly excised, and used for the analysis of NO production (3 × 2 plants/assay), or immediately frozen in liquid N₂, ground in powder and stored at – 80 °C until adenine nucleotide measurements (3 × 2 plants/assay).

Vegetative growth analysis

Root systems were washed in distilled water, blotted on filter paper and weighted. Dry weights were measured after drying of samples for 48 h at 70 °C.

Biological nitrogen-fixing capacity

BNF capacity of nodulated roots was determined in vivo by measuring the acetylene reducing activity (ARA) as previously described (Pierre et al. 2014).

Measurement of NO production

Segments (2 cm-long) of nodulated roots were incubated in the dark, at 23 °C, in 5 mL tubes containing 2 mL of detection medium (10 mM Tris–HCl pH 7.5, 10 mM KCl) in the presence of 10 µM 4,5-diaminofluorescein (DAF-2, Coger, with excitation at 495 nm and emission at 515 nm) fluorescent probe. NO production and controls were carried out as in Horchani et al. (2011). In addition, to test the specificity of the DAF-2 probe, NO production was also analysed with the Cu(II) fluorescein (CuFL) fluorescent probe (Strem Chemicals, with excitation at 495 nm and emission at 515 nm), which is known to react rapidly and specifically with NO itself (Lim et al. 2006).

Extraction and measurement of adenine nucleotides and proteins

Adenine nucleotides were extracted essentially as in Horchani et al. (2011). All extraction steps were carried out at 4 °C. Frozen material (40–60 mg) was crushed in liquid nitrogen with 300 mL of perchloric acid solution, containing 7% (v/v) HClO₄ and 25 mM Na₂EDTA, with a mortar and pestle. After thawing, the extract was taken and the mortar was rinsed with 200 mL of perchloric acid solution, which was then pooled with the extract. The sample was centrifuged for 5 min at 13,000g. The supernatant was quickly and carefully neutralized at pH 5.6–6.0 using a 2 M KOH–0.3 M MOPS solution. KClO₄ precipitate was discarded by centrifugation (5 min, 13,000g). Adenine nucleotides of the supernatant were measured in a Xenius spectrofluorimeter-luminometer (Safas, Monaco) using the ATPlite one-step assay system (ATPLT1STP-0509; Perkin-Elmer) according to the manufacturer's instructions. Proteins were extracted and quantified on clarified extracts as in Horchani et al. (2010).

Results

Salinity and flooding affect nodulated roots nitrogen-fixing capacity and energy state, and increase NO production

Salt and flooding treated *M. sativa* roots were first analysed for their biomass and nodule production. As reported in Table S1, root system dry weight is not affected by the two treatments. However, T20 and T50 salt treatments decrease by 38 and 52%, respectively, the number of nodules per plant, indicating that salt stress affects nodulation. On its side, flooding treatment has no effect on the number of nodules per plant. It can be noted that, regardless of the treatment, the protein content of the root system remains

unchanged (Fig. S3), suggesting that, as a whole, the root system is not senescent and is still functional.

The effects of salinity and flooding on *M. sativa* BNF capacity were assessed by measuring the ARA of nodules (Fig. 1a). As compared with control nodules in normoxia, T20 and T50 treatments result in an 18 and 75% decrease in ARA, respectively. When compared to normoxia, flooding also results in a 33 and 52% decrease in ARA for control and T20 nodules, respectively, but is without additional effect for T50 nodules.

To test the effects of salinity and flooding on the energy state, we analyzed the ATP/ADP ratio (experiment 1). ATP/ADP ratio of control roots is 7.4 (Fig. 1b), which is in agreement with previous analysis in either roots or nodules (Brouquisse et al. 1991; Horchani et al. 2011). ATP/ADP ratio is not modified in T20 (7.0), but significantly decreased in T50 roots (6.5), indicating that average but not low salt stress impairs the energy status of the nodulated roots. After a 5-day flooding period, ATP/ADP ratios decrease close to 5.0, 4.5 and 4.2 in Ctrl, T20 and T50 treated roots, respectively (Fig. 1b). These results indicate that the saline and flooding stresses, independently of each other, affect the energy state of the nodulated roots and that their combined effect is partly cumulative.

In roots and nodules under hypoxia, NO production has been associated with the establishment of a Phyto gb -NO respiration, whose function is to maintain the energy state of the tissues when O₂ concentration decreases (Igamberdiev and Hill 2009; Horchani et al. 2011). We checked whether NO production could be involved in the maintenance of the root system energy state under each stress. As compared with control, T20 and T50 treatments caused a 1.8 and 3.8-fold increase in NO production (Fig. 1c). Similarly, flooding induced a 1.3 to 2.7-fold increase in NO production compared to normoxic conditions. As a control, NO production was also measured with Cu(II) fluorescein probe with similar results (Fig. S3), indicating that measurements with DAF-2 are reliable and relevant in this context. Thus, NO production was increased by both saline and flooding treatments, suggesting (1) that the effects of salinity and flooding on NO production and energy state could result from hypoxic environment, and (2) that Phyto gb -NO respiration could be activated in response to both saline and flooding/hypoxia treatments.

NO production contributes to the maintenance of energy state under both salinity and hypoxia

To test the above hypothesis, we analyzed the effects of cPTIO (a NO scavenger) and Tg (a NR and Phyto gb -NO respiration inhibitor) on ATP/ADP ratio and NO production in the root system of control (0) and T50 (50) treated plants submitted (H) or not (N) to hypoxia. To this end, we




Fig. 1 **a** Nitrogen-fixing activity, **b** energy state and **c** NO production of *Medicago sativa* nodulated roots submitted to saline and hypoxic treatments. T20 and T50 refer to 20 and 50 mM NaCl treatments, respectively. Nitrogen-fixing activity is measured through acetylene reducing activity (ARA). Energy state is evaluated through ATP/ADP ratio. The values are mean \pm SE ($n=3$ for ATP/ADP ratio; $n=4$ for ARA and NO production) of 4 independent experiments. Values followed by different letters are significantly different according to one-way ANOVA analysis followed by a Tukey test ($P < 0.05$)

used a specific hydroponic experimental setup where the root system was incubated in nutrient solution supplemented or not with the inhibitors (experiment 2, Fig. S2). Compared to the control (0-N), NR inhibition by Tg results in a 43, 33 and 41% decrease of ATP/ADP ratio in 0-H, 50-N and 50-H roots, respectively (Table 1). Similarly, NO scavenging by cPTIO results in a 34, 37 and 37% decrease of ATP/ADP ratio in 0-H, 50-N, and 50-H roots, respectively, compared to control conditions. cPTIO and Tg has no significant effect on the ATP/ADP ratio in 0-N roots. Treatment with Tg triggers a strong decrease in NO production in 0-H, 50-N and 50-H treated root systems, but not in 0-N treated root system (Table 1). In the presence of cPTIO, NO production is totally abolished. Taken together, these data indicate that both the functioning of NR and the production of NO are necessary to maintain root and nodule energy state under salinity and hypoxia, alone or combined together.

Discussion

The present data show that nodulation, but not root growth, of *M. sativa* plants is affected by average salinity (Table S1). This is in line with literature data reporting that alfalfa is a moderately salt-tolerant legume (Bruning and Rozema 2013) and that its growth is significantly affected only from NaCl concentrations above 80 mM (Rogers et al. 2008, 2009, 2011). However, salinity is known to inhibit BNF more rapidly than growth in legumes (Bruning and Rozema 2013). In our study, ARA is the more inhibited as the salt concentration increases (Fig. 1). Under low salt stress, the supply of O₂ to nodules was suggested to be the limiting factor for BNF (Bergersen 1982; Delgado et al. 1994; Serraj et al. 1994). This inhibition would be due to the fact that the salt stress disrupts the symplastic connections between nodule cells, decreases the permeability of the nodules to O₂ and increases its critical O₂ pressure, which would have the effect of inhibiting the energy metabolism and the regeneration of ATP. These interpretations were supported by Serraj et al. (1998) and Serraj and Drevon (1998) who showed that the BNF of common bean, soybean and alfalfa submitted to 50–100 mM NaCl concentrations is partly inhibited compared to that of control plants when measured at 21% O₂, but that the inhibition is reversed by increasing pO₂. Considered together, these studies suggest that the decrease in ARA under salt stress is due to an O₂ limitation within the nodules leading to a reduction in energy metabolism.

Short-term effects of flooding on BNF have been little studied. However, in *Trifolium repens* (Pugh et al. 1995) and soybean (Sánchez et al. 2010) nodulated roots, a short period of flooding (6–7 days) resulted in a complete or half inhibition of ARA, respectively. Similarly, our data show that in *M. sativa* nodulated roots, ARA is inhibited after

Table 1 Effect of Tg and cPTIO on energy state (ATP/ADP ratio) and NO production in *Medicago sativa* nodulated roots submitted to saline and hypoxic treatments (Experiment 2)

	0-N			0-H			50-N			50-H		
	Ctr	Tg	cPTIO	Ctr	Tg	cPTIO	Ctr	Tg	cPTIO	Ctr	Tg	cPTIO
ATP/ADP ratio	6.2±0.6 a	5.0±0.7 ab	5.1±0.6 ab	4.4±0.4 b	2.5±0.6 d	2.9±0.3 cd	5.4±0.8 ab	3.6±0.9 bc	3.4±0.1 c	4.1±0.4 bc	2.4±0.4 d	2.6±0.4 d
NO production (FU h ⁻¹ gFW ⁻¹)	18.0±9.7 a	11.5±3.3 a	5.5±2.2 b	54.1±11.4 c	18.4±6.1 a	4.8±3.0 b	91.8±27.5 d	19.7±12.3 ab	5.2±2.5 b	88.5±16.4 d	12.8±10.6 ab	5.0±2.2 b

Plants were grown in the absence (0) or in the presence of 50 mM NaCl (50) and submitted to either normoxia (N) or hypoxia (H) as indicated in the Materials and methods. Ctr, control; Tg, 1 mM sodium tungstate; cPTIO, 0.1 mM cPTIO. Data are mean±SE ($n=3$) of 3 independent experiments. Values followed by different letters are significantly different according to one-way ANOVA analysis followed by a Tukey test ($P<0.05$)

5 days of flooding (Fig. 1), confirming that flooding inhibits ARA presumably through flooding-associated O₂ shortage.

In the root system of many plant species, including legumes, submitted to flooding or hypoxia, the induction of a Phyto**g**-NO respiration occurs, allowing cell energy status retention (Dordas et al. 2003; Gupta and Igamberdiev 2011; Berger et al. 2018). The Phyto**g**-NO respiration cycle involves (1) the reduction of NO₃⁻ by NR (2) the translocation of NO₂⁻ from the cytosol to the mitochondrial matrix, (3) the reduction of NO₂⁻ to NO via the mitochondrial electron transfer chain (ETC) allowing ATP regeneration, and finally (4) the passive diffusion of NO to the cytosol where it is oxidized back to NO₃⁻ by Phyto**g** (Gupta and Igamberdiev 2011). Functional legume nodules are naturally characterized by a microoxic environment, and the existence of such a Phyto**g**-NO respiration was also evidenced in *M. truncatula* nodules (Horchani et al. 2011). In these nodules, a hypoxic treatment triggers an increase in NO production, which is abolished by the addition of either cPTIO or Tg, and the energy state (ATP/ADP) is significantly inhibited when nodules are incubated in the presence of Tg (Horchani et al. 2011). In our study, under either salt or hypoxic treatment, or both, the use of Tg and cPTIO leads to a reduction of the ATP/ADP ratio and to an inhibition of NO production (Table 1) indicating that both NR activity and NO production are necessary to maintain the energy state. Taken together our data first confirm that in *M. sativa* nodulated roots under flooding/hypoxia, NO production is involved in maintaining energy state presumably through the establishment of a Phyto**g**-NO respiration. Second, our study shows that NR-dependent NO production is also involved in maintaining energy state under salt stress, which argues that salt treatment triggers a hypoxic stress in the nodulated roots. Third, this study strongly suggests that the functioning of Phyto**g**-NO respiration is induced in nodulated roots under salt stress.

Author contributions statement RB and SA-S planned and designed the research. FA, HS, AG and RB performed experiments and conducted fieldwork. AK, SA-S and RB interpreted the data. RB wrote the manuscript.

Acknowledgements This work was supported by INRAE and by the Comité Mixte Franco-Tunisien pour la Coopération Universitaire (PHC Utique, Grant no. 17G0904).

References

Bailey-Serres J, Voesenek LACJ (2008) Flooding stress: acclimations and genetic diversity. *Annu Rev Plant Biol* 59:313–339. <https://doi.org/10.1146/annurev.arplant.59.032607.092752>

- Berger A, Brouquisse R, Pathak PK, Hichri I, Singh I et al (2018) Pathways of nitric oxide metabolism and operation of phytooglobins in legume nodules: missing links and future directions. *Plant Cell Env* 41:2057–2068. <https://doi.org/10.1111/pce.13151>
- Bergersen FJ (1982) Root nodules of legumes: Structure and functions. Research Studies Press/Wiley, Chichester
- Besson-Bard A, Pugin A, Wendehenne D (2008) New insights into nitric oxide signaling in plants. *Annu Rev Plant Biol* 59:21–39. <https://doi.org/10.1146/annurev.arplant.59.032607.092830>
- Blokhina O, Fagerstedt KV (2010) Oxidative metabolism, ROS and NO under oxygen deprivation. *Plant Physiol Biochem* 48:359–373. <https://doi.org/10.1016/j.plaphy.2010.01.007>
- Brouquisse R, James F, Raymond P, Pradet A (1991) Study of glucose starvation in excised maize root tips. *Plant Physiol* 96:619–626. <https://doi.org/10.1104/pp.96.2.619>
- Bruning B, Rozema J (2013) Symbiotic nitrogen fixation in legumes: perspectives for saline agriculture. *Env Exp Bot* 92:134–143. <https://doi.org/10.1016/j.envexpbot.2012.09.001>
- del Giudice J, Cam Y, Damiani I, Fung-Chat F, Meilhoc E, Bruand C, Brouquisse R, Puppo A, Boscari A (2011) Nitric oxide is required for an optimal establishment of the *Medicago truncatula*-*Sinorhizobium meliloti* symbiosis. *New Phytol* 191:405–417. <https://doi.org/10.1111/j.1469-8137.2011.03693.x>
- Delgado MJ, Ligeró F, Lluch C (1994) Effects of salt stress on growth and nitrogen fixation by pea, faba-bean, common bean and soybean plants. *Soil Biol Biochem* 26:371–376. [https://doi.org/10.1016/0038-0717\(94\)90286-0](https://doi.org/10.1016/0038-0717(94)90286-0)
- Dordas C, Hasinoff BB, Igamberdiev AU, Manac'h N, Rivoal J, Hill RD (2003) Expression of a stress-induced hemoglobin affects NO levels produced by alfalfa root cultures under hypoxic stress. *Plant J* 35:763–770. <https://doi.org/10.1046/j.1365-313x.2003.01846.x>
- Gupta KJ, Igamberdiev AU (2011) The anoxic plant mitochondrion as a nitrite: NO reductase. *Mitochondrion* 11:537–543. <https://doi.org/10.1016/j.mito.2011.03.005>
- Hartman S, Liu Z, van Veen H, Vicente J, Reinen E et al (2019) Ethylene-mediated nitric oxide depletion pre-adapts plants to hypoxia stress. *Nat Commun* 10:4020. <https://doi.org/10.1038/s41467-019-12045-4>
- Hichri I, Boscari A, Castella C, Rovere M, Puppo A, Brouquisse R (2015) Nitric oxide: a multifaceted regulator of the nitrogen-fixing symbiosis. *J Exp Bot* 66:2877–2887. <https://doi.org/10.1093/jxb/erv051>
- Horchani F, Hajri R, Khayati H, Brouquisse R, Aschi-Smiti S (2010) Does the source of nitrogen affect the response of subterranean clover to prolonged root hypoxia? *J Plant Nutr Soil Sci* 173:275–283. <https://doi.org/10.1002/jpln.200900280>
- Horchani F, Prevot M, Boscari A, Evangelisti E, Meilhoc E et al (2011) Both plant and bacterial nitrate reductases contribute to nitric oxide production in *Medicago truncatula* nitrogen-fixing nodules. *Plant Physiol* 155:1023–1036. <https://doi.org/10.1104/pp.110.166140>
- Igamberdiev AU, Hill RD (2009) Plant mitochondrial function during anaerobiosis. *Ann Bot* 103:259–268. <https://doi.org/10.1093/aob/mcn100>
- Kopyra M, Gwózdź EA (2003) Nitric oxide stimulates seed germination and counteracts the inhibitory effect of heavy metals and salinity on root growth of *Lupinus luteus*. *Plant Physiol Biochem* 41:1011–1017. <https://doi.org/10.1016/j.plaphy.2003.09.003>
- Lim MH, Xu D, Lippard SJ (2006) Visualization of nitric oxide in living cells by a copper-based fluorescent probe. *Nat Chem Biol* 2:375–380. <https://doi.org/10.1038/nchembio794>
- Liu Y, Wu R, Wan Q, Xie G, Bi Y (2007) Glucose-6-phosphate dehydrogenase plays a pivotal role in nitric oxide-involved defense against oxidative stress under salt stress in red kidney bean roots. *Plant Cell Physiol* 48:511–522. <https://doi.org/10.1093/pcp/pcm020>
- Molassiotis A, Tanou G, Diamantidis G (2010) NO says more than ‘YES’ to salt tolerance salt priming and systemic nitric oxide signaling in plants. *Plant Signal Behav* 5:209–212. <https://doi.org/10.4161/psb.5.3.10738>
- Pierre O, Hopkins J, Combier M, Baldacci F, Engler G et al (2014) Involvement of papain and legumain proteinase in the senescence process of *Medicago truncatula* nodules. *New Phytol* 202:849–863. <https://doi.org/10.1111/nph.12717>
- Pugh R, Witty JF, Mytton LR, Minchin FR (1995) The effect of waterlogging on nitrogen fixation and nodule morphology in soil-grown white clover (*Trifolium repens* L.). *J Exp Bot* 46:285–290. <https://doi.org/10.1093/jxb/46.3.285>
- Rogers ME, Colmer TD, Frost K, Henry D, Cornwall D et al (2008) Diversity in the genus *Melilotus* for tolerance to salinity and waterlogging. *Plant Soil* 304:89–101. <https://doi.org/10.1007/s11104-007-9523-y>
- Rogers ME, Colmer TD, Frost K, Henry D, Cornwall D et al (2009) The influence of NaCl salinity and hypoxia on aspects of growth in *Trifolium* species. *Crop Past Sci* 60:71–82. <https://doi.org/10.1071/CP08123>
- Rogers ME, Colmer TD, Nichols PGH, Hughes SJ, Frost K et al (2011) Salinity and waterlogging tolerance amongst accessions of messina (*Melilotus siculus*). *Crop Past Sci* 62:225–235. <https://doi.org/10.1071/CP10270>
- Sánchez C, Gates AJ, Meakin GE, Uchiumi T, Girard L, Richardson DJ, Bedmar EJ, Delgado MJ (2010) Production of nitric oxide and nitrosylhemoglobin complexes in soybean nodules in response to flooding. *Mol Plant-Mi Inter* 23:702–711. <https://doi.org/10.1094/MPMI-23-5-0702>
- Serraj R, Drevon JJ (1998) Effects of salinity and nitrogen source on growth and nitrogen fixation in alfalfa. *J Plant Nutr* 21:1805–1818. <https://doi.org/10.1080/01904169809365525>
- Serraj R, Roy G, Drevon JJ (1994) Salt stress induces a decrease in the oxygen uptake of soybean nodules and their permeability to oxygen diffusion. *Physiol Plant* 91:161–168. <https://doi.org/10.1111/j.1399-3054.1994.tb00414.x>
- Serraj R, Vasquez-Diaz H, Drevon JJ (1998) Effects of salt stress on nitrogen, fixation, oxygen diffusion, and ion distribution in soybean, common bean and alfalfa. *J Plant Nutr* 21:475–488. <https://doi.org/10.1080/01904169809365418>
- Sghaier H, Aloui A, Khadri A, Aschi-Smiti S, Brouquisse R (2020) Effects of combined salt and flooding stresses on the growth and physiological behavior of alfalfa (*Medicago sativa* L.). *Afric J Agro Res* 16:758–764. <https://doi.org/10.5897/AJAR2020.14831>
- Simontacchi M, Galatro A, Ramos-Artuso F, Santa-María G (2015) Plant survival in a changing environment: the role of nitric oxide in plant responses to abiotic stress. *Front Plant Sci* 6:977. <https://doi.org/10.3389/fpls.2015.00977>
- Stoimenova M, Igamberdiev AU, Gupta KJ, Hill RD (2007) Nitrite-driven anaerobic ATP synthesis in barley and rice root mitochondria. *Planta* 226:465–474. <https://doi.org/10.1007/s00425-007-0496-0>
- Tanou G, Molassiotis A, Diamantidis G (2009) Hydrogen peroxide- and nitric oxide-induced systemic antioxidant prime-like activity under NaCl-stress and stress free conditions in citrus plants. *J Plant Physiol* 166:1904–1913. <https://doi.org/10.1016/j.jplph.2009.06.012>
- Thalineau E, Truong H-N, Berger A, Fournier C, Boscari A et al (2016) Cross-regulation between N metabolism and nitric oxide (NO) signaling during plant immunity. *Front Plant Sci* 7:472. <https://doi.org/10.3389/fpls.2016.00472>
- Uchida A, Jagendorf AT, Hibino T, Takabe T, Takabe T (2002) Effects of hydrogen peroxide and nitric oxide on both salt and heat stress tolerance in rice. *Plant Sci* 63:515–523. [https://doi.org/10.1016/S0168-9452\(02\)00159-0](https://doi.org/10.1016/S0168-9452(02)00159-0)

- Wang H, Liang X, Wan Q, Wang X, Bi Y (2009) Ethylene and nitric oxide are involved in maintaining ion homeostasis in *Arabidopsis* callus under salt stress. *Planta* 230:293–307. <https://doi.org/10.1007/s00425-009-0946-y>
- Wany A, Kumari A, Gupta KJ (2017) Nitric oxide is essential for the development of aerenchyma in wheat roots under hypoxic stress. *Plant Cell Environ* 40:3002–3017. <https://doi.org/10.1111/pce.13061>
- Zhang Y, Wang L, Liu Y, Zhang Q, Wei Q, Zhang W (2006) Nitric oxide enhances salt tolerance in maize seedlings through increasing activities of proton-pump and Na^+/H^+ antiport in the tonoplast. *Planta* 224:545–555. <https://doi.org/10.1007/s00425-006-0242-z>
- Zhao G, Zhao Y, Yu X, Kiprotich F, Han H, Guan R, Wang R, Shen W (2018) Nitric oxide is required for melatonin-enhanced tolerance against salinity stress in rapeseed (*Brassica napus L*) seedlings. *Int J Mol Sci* 19:1912. <https://doi.org/10.3390/ijms19071912>