

Consideration of mass effect processes in bioindication allows more accurate bioassessment of water quality

Thibault Leboucher, Lucas Mignien, Marie Wach, Sébastien Boutry, Aurélien

Jamoneau, Sophia I Passy, Juliette Tison-Rosebery

▶ To cite this version:

Thibault Leboucher, Lucas Mignien, Marie Wach, Sébastien Boutry, Aurélien Jamoneau, et al.. Consideration of mass effect processes in bioindication allows more accurate bioassessment of water quality. Ecological Indicators, 2021, 127, pp.107791. 10.1016/j.ecolind.2021.107791. hal-03290177

HAL Id: hal-03290177 https://hal.inrae.fr/hal-03290177

Submitted on 19 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Contents lists available at ScienceDirect

Ecological Indicators

ELSEVIER



journal homepage: www.elsevier.com/locate/ecolind

Consideration of mass effect processes in bioindication allows more accurate bioassessment of water quality

any biotic index.

Thibault Leboucher^{a,*}, Lucas Mignien^a, Marie Wach^a, Sébastien Boutry^a, Aurélien Jamoneau^a, Sophia I. Passy^b, Juliette Tison-Rosebery^a

^a INRAE, EABX, Cestas Cedex, France

^b Department of Biology, University of Texas at Arlington, Box 19498, Arlington, TX 76019, USA

ARTICLE INFO	A B S T R A C T				
Keywords: Biological Diatom Index Bio-assessment Community assembly Diatoms Dispersal Mass effect Streams	Bioassessment is widely used to measure ecological integrity of natural habitats following anthropogenic dis- turbances and modifications. Traditionally, bioassessment has been based exclusively on species-environment interactions, <i>i.e.</i> niche processes. However, dispersal processes, and in particular mass effect, could mask the influence of niche processes and lead to erroneous conclusions about ecosystem health. To circumvent this problem, we identified 40 diatom species with distributions driven primarily by mass effect and propose an alternative version of the Biological Diatom Index (IBD ₂₀₀₇) excluding these species. We tested the environmental responses of both the original IBD (IBD ₂₀₀₇) and the modified IBD (IBD _{mod}) with a benthic diatom dataset from France, collected between 2007 and 2013 and including 9487 samples from 3913 spatially distinct localities. Our results indicate a better relationship between the IBD _{mod} scores and environmental conditions, compared to the IBD ₂₀₀₇ scores, leading to a more accurate determination of river ecological status, especially in conditions of moderate nutrient enrichment. This study supports the idea that mass effect may result in biased evaluation of water quality. It is advocated that this process is considered in other diatom-based indices, and by extension, in				

1. Introduction

Bioassessment is broadly implemented by state and government agencies in an effort to measure how species communities respond to anthropogenic stressors, *e.g.* EU Water Framework Directive (European Parliament, 2000). A key instrument of bioassessment are biotic indices, which have been traditionally derived from species tolerance and sensitivity to anthropogenic perturbations (Moog et al., 2018). In freshwater biomonitoring, diatom-based indices are commonly used to capture species and community responses to environmental conditions, such as eutrophication, acidification, and organic pollution (Stevenson and Bahls, 1999; Passy and Bode, 2004) or even to specific stressors (Vilmi et al., 2016) in order to determine the level of habitat impairment. The Biological Diatom Index (IBD₂₀₀₇) (Lenoir and Coste, 1996; Coste et al., 2009) is a standardized method (AFNOR, 2007) created and widely used in France for the monitoring of water quality in accordance with the Water Framework Directive requirements.

Environmental bioassessment is typically performed using

information on species tolerance and sensitivity to disturbance and measures how community composition changes under anthropogenic influences. However, community composition is driven not only by environmental factors but also by dispersal (Leibold et al., 2004). Therefore, to accurately determine ecological status, dispersal processes must be integrated in bioassessment metrics (Brown et al., 2011; Heino et al., 2013). Depending on species traits and environmental settings, dispersal can be limited or unlimited, the latter leading to mass effect. Dispersal limitation, preventing species from reaching suitable habitats, is now considered in new environmental assessment methods, focusing on smaller regions and appropriate scales (Pont et al., 2006; Smucker and Vis, 2011). Mass effect, on the other hand, allowing species persistence in unfavorable conditions due to dispersal from source habitats, has not been considered in bioassessment, despite evidence for its importance in freshwaters (Bottin et al., 2016; Goldenberg Vilar et al., 2014; Jamoneau et al., 2018; Leboucher et al., 2020). There is thus a need to evaluate the potential of mass effect to distort bioassessment results.

* Corresponding author. *E-mail address*: thibault.leboucher@inrae.fr (T. Leboucher).

https://doi.org/10.1016/j.ecolind.2021.107791

Received 15 January 2021; Received in revised form 3 May 2021; Accepted 5 May 2021 Available online 17 May 2021 1470-160X/© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-ad/4.0/). A combination of spatial and trait analyses can measure the importance of mass effect in structuring active disperser communities, *e.g.*, benthic macroinvertebrates (Brown and Swan, 2010). However, this type of approach cannot be used for passive dispersers such as microalgae, due to the absence of functional traits clearly related to the dispersal capabilities of these organisms. This hampers our ability to evaluate the potential influence of mass effect in diatom-based bioassessment methods. Here we propose to circumvent this problem by implementing an innovative method, based on a species distribution modeling, with environmental and spatial factors computed from asymmetric eigenvector maps, coupled with negative co-occurrence patterns (Leboucher et al., 2020). Using this method, we identified 40 species as being influenced by mass effect, referred to as "mass effect species" henceforth.

Our objective was to evaluate whether mass effect species can weaken bioassessment and generate erroneous stream classification in terms of water quality. To achieve this goal, we examined the sensitivity to eutrophication of IBD_{2007} and modified IBD_{2007} (IBD_{mod}) after the removal of mass effect species.

2. Materials & methods

2.1. Diatom and environmental condition datasets

We analyzed a benthic diatom dataset from France, including 9487 samples collected between 2007 and 2013 from 3913 spatially distinct localities (Fig. 1), following a standardized protocol (NFT 90-354, (AFNOR, 2007). Diatoms were sampled on hard substrates between May and October, in order to reduce seasonal variability in species assemblages. Cells were identified at 1000× magnification by examining 400 cleaned diatom frustules in permanent slides using among others Krammer and Lange-Bertalot (1986-1991, 1995-2015, 2000-2013) as identification references. A taxonomic homogenization at the species level was performed with Omnidia 5.3 software (Lecointe et al., 1993).

Corresponding physico-chemical data were gathered by the French Water Agencies. Physico-chemical variables included water pH, specific conductance (at 25 °C, mS.cm⁻¹), biological oxygen demand (mg of O₂ consumed per liter during 5 days of incubation at 20 °C), concentrations of total phosphorous (mg.L⁻¹ of $_{15}$ P), orthophosphate (mg.L⁻¹ of $_{15}$ P), ammonium (mg.L⁻¹ of $_{7}$ N), nitrate (mg.L⁻¹ of $_{7}$ N) and nitrite (mg.L⁻¹ of $_{7}$ N). We used the median values of all environmental variable



Fig. 1. Map of the studied area, including 3913 spatially distinct localities.

measurements obtained during the 30 days before and the 15 days after the diatom sampling date.

2.2. Biological Diatom Index (IBD₂₀₀₇)

The IBD₂₀₀₇ (Coste et al., 2009) is calculated using the sum of the presence probability of indicator taxa along seven water quality classes following the equation

$$BDI = \sum_{j=1}^{7} j \times \frac{\sum_{i=1}^{n} A_i \times V_i \times P_{ij}}{\sum_{i=1}^{n} A_i \times V_i}$$

where A_i is the abundance of taxon *i*, V_i is the degree of stenoecy (*i.e.* tolerance) of the taxon *i*, P_{ij} is the presence probability of the taxon *i* for the quality class *j*.

 IBD_{2007} scores are related to reference values for the type of river considered, to calculate corresponding Ecological Quality Ratios (EQR) and to assign an ecological status class to the considered site (JORF, 2015). These 5 ecological status classes are, from the best to the worst, high, good, moderate, poor and bad. A modified IBD (IBD_{mod}) was calculated after removing from the list of indicator taxa 40 species (Table 1) previously identified as influenced by mass effect (Leboucher et al., 2020). To ensure that the removal of the 40 species does not generate a mathematical artifact, we also calculated 999 alternative versions of the IBD₂₀₀₇ (referred to as randomized IBD₂₀₀₇) after removing 40 species randomly drawn among the indicator species not subjected to mass effect.

Table 1

List of the 40 indicator taxa identified as influenced by mass effect in Leboucher et al. (2020).

ID	Species
DVUL	Diatoma vulgaris Bory
ADEU	Achnanthidium eutrophilum (Lange-Bertalot) Lange-Bertalot
ACOP	Amphora copulata (Kützing) Schoeman & Archibald
AOVA	Amphora ovalis (Kützing) Kützing
APED	Amphora pediculus (Kützing) Grunow
CNTH	Cocconeis neothumensis Krammer
CPED	Cocconeis pediculus Ehrenberg
CPLA	Cocconeis placentula Ehrenberg
COPL	Cocconeis pseudolineata (Geitler) Lange-Bertalot
DMON	Diatoma moniliformis Kützing
DOCU	Diploneis oculata (Brebisson) Cleve
ENLB	Encyonema lange-bertalotii Krammer
ECPM	Encyonopsis minuta Krammer & Reichardt
EOMI	Eolimna minima (Grunow) Lange-Bertalot
FPYG	Fallacia pygmaea (Kützing) Stickle & Mann
FSLU	Fallacia sublucidula (Hustedt) Mann
GMIN	Gomphonema minutum (Agardh)
GOLI	Gomphonema olivaceum (Hornemann) Brubisson
GPUM	Gomphonema pumilum Reichardt & Lange-Bertalot
GYAT	Gyrosigma attenuatum (Kützing) Rabenhorst
KAPL	Karayevia ploenensis (Hustedt) Bukhtiyarova
LGOE	Luticola goeppertiana (Bleisch in Rabenhorst) Mann
NANT	Navicula antonii Lange-Bertalot
NCPR	Navicula capitatoradiata Germain
NCTE	Navicula cryptotenella Lange-Bertalot
NGER	Navicula germainii Wallace
NAMP	Nitzschia amphibia Grunow
NCPL	Nitzschia capitellata Hustedt in Schmidt
NFON	Nitzschia fonticola Grunow
NREC	Nitzschia recta Hantzsch in Rabenhorst
NSOC	Nitzschia sociabilis Hustedt
NSUA	Nitzschia subacicularis Hustedt in Schmidt
NTUB	Nitzschia tubicola Grunow
PGRN	Planothidium granum (Hohn & Hellerman) Lange-Bertalot
PRST	Planothidium rostratum (Oestrup) Lange-Bertalot
PTCO	Platessa conspicua (Mayer) Lange-Bertalot
PBTG	Pseudostaurosira brevistriata (Grunow in Van Heurck) Williams & Round
RUNI	Reimeria uniseriata Sala Guerrero & Ferrario
RABB	Rhoicosphenia abbreviata (Agardh) Lange-Bertalot
SIDE	Simonsenia delognei Lange-Bertalot

2.3. Relationships between IBD₂₀₀₇ indices and environmental conditions

We performed a principal component analysis (PCA) on powertransformed physico-chemical variables. We used Box-Cox maximum likelihood estimation (Box and Cox, 1964) to select the powertransformation best approximating normality. We simplified the analysis by retaining only the components with eigenvalues greater than those randomly generated by a broken-stick model (Jackson, 1993; Legendre and Legendre, 2012). We used sample coordinates on the principal component axes (PC) selected by the broken-stick model to evaluate the sample site position on the environmental gradient of the dataset.

Akaike information criterion (minimum theoretical information criterion, AIC) (Akaike, 1974) was implemented to compare linear models between the different versions of IBD (IBD₂₀₀₇, IBD_{mod}, randomized IBD₂₀₀₇) and the PCA coordinates. The smallest value of AIC indicates the best model, and therefore, the IBD version with the best response to environmental conditions (original, modified and the randomized IBD₂₀₀₇). Difference between the AICs of randomized IBD₂₀₀₇ and the AICs of the original and modified IBD were tested using z-scores with the criterion -2 < Z < 2.

2.4. Comparison of ecological status assessment

We compared the number of samples per water quality category among the original and the modified IBD with a χ^2 test. We also analyzed the differences in ecological status assessment among the different index versions, and whether these differences were correlated with the nutrient gradient (derived from the PCA coordinates) with ANOVA, followed by Tukey post-hoc tests. Only categories with at least 30 values were tested.

All data analyses were performed with R 3.4.4 (R Core Team, 2018), using the 'vegan' package (Oksanen et al., 2018).

3. Results

Following the broken-stick model, only the first principal component (PC1) axis was significant explaining 40% of the total variance (Fig. 2).

PC1 represented a gradient of nutrient enrichment mainly driven by phosphorous (orthophosphates and total phosphorous), nitrogen (nitrites and ammonia) and biological oxygen demand. Only PC1 scores were then used to compare the response of the different IBD indices to environmental conditions. AIC of the linear models between IBD indices and PC1 scores was smaller for the IBD_{mod} (-45477) compared to the IBD₂₀₀₇ (-45235) indicating a better relationship.

AIC of the IBD₂₀₀₇ was not significantly different from those of the randomized IBD₂₀₀₇ (z-score = -0.45). Conversely, AIC of the IBD_{mod} was significantly smaller than AIC of the randomized IBD₂₀₀₇ (z-score = -2.30).

Original and modified IBD based ecological status classifications of sites were significantly different (χ^2 test, p < 0.001) largely because a significant number of sites originally belonging to average classes (good and moderate) were placed in the extreme classes (high, poor and bad) (Table 2). The overall number of samples originally belonging to high and bad quality status did not significantly change (Table 2).

The IBD_{mod} allowed for a better discrimination of ecological status assessments (Fig. 3), given that the new classes obtained with the IBD_{mod} were better correlated with the nutrient gradient from the PCA. Samples upgraded with the IBD_{mod} to higher ecological status class displayed low nutrient concentrations (low PC1 scores) whereas samples downgraded

Table 2

Number of samples across ecological status classes obtained with the original and the modified version of IBD. The sum of each row represents the number of samples placed in the respective class by the IBD_{2007} , and the sum of each column represents the number of samples placed in the respective class by the IBD_{mod} .

		Modifi					
		High	Good	Moderate	Poor	Bad	Total
Original IBD ₂₀₀₇	High Good Moderate Poor Bad Total	2061 745 12 0 0 2818	9 2353 297 0 0 2659	0 483 2379 12 0 2874	0 2 536 443 0 981	0 0 2 100 53 155	2070 3583 3226 555 53 9487



Fig. 2. Principal component analysis biplot showing site distribution along the first two principal component axes. Environmental variables are represented by arrows, inclding CONDUC.25: specific conductance at 25 °C; BDO5: biological oxygen demand; NO2: nitrite concentration; NH4: ammonium concentration; P. TOTAL: total phosphorous concentration; ORTHOP: orthophosphate concentrations.



Fig. 3. Boxplot of the site coordinates on PC1 in correspondence with 1) ecological status classes from the original IBD₂₀₀₇ version (grey boxes on the left) and 2) ecological status classes from the modified IBD₂₀₀₇ version (box and whisker plots on the right). Only categories with at least 30 values were tested with ANOVA followed up by Tukey post-hoc tests when multiple classes were compared (n.s.: not significant; ***: p-value < 0.001).

showed high nutrient concentrations (high PC1 scores). These differences were, for the most part, significant (see ANOVA and Tukey tests on Fig. 3).

4. Discussion

Mass effect is an important driver of diatom community structure and thus has the potential to mask the effect of the local environment, and lead to an inaccurate evaluation of water quality (Heino et al., 2015; Jamoneau et al., 2018). However, until recently, there were no statistical methods for identifying species driven by mass effect (Leboucher et al., 2020). In this study, we implemented such method and determined the extent to which mass effect altered water quality assessment. We compared the performance of a broadly used diatom index in its original form and after the removal of species influenced by mass effect. We showed for the first time that accounting for mass effect leads to more reliable estimates of ecological status. We discuss the implications of this finding below. Research on bioindication is just beginning to recognize the role of spatial effects (Heino, 2013), which may even exceed this of the environment. For example, Vilmi et al. (2016) simultaneously considered environmental and spatial effects on different diatom indices designed to measure water quality. Variance partitioning revealed that score variation across indices was largely associated with pure effects of spatial variables, or shared fractions of spatial variables and physico-chemical variables. In contrast, water chemistry (*i.e.* water quality or the eutrophication) did not capture much of the variation in the diatom indices. Although the results of Vilmi et al. (2016) clearly showed the importance of spatial factors, traditional methods, such as variance partitioning, were unable distinguish between the forms of dispersal, *i.e.* limited or unlimited (Cottenie, 2005), which makes them less useful for bioassessment.

In this study, we specifically targeted species identified as being influenced by mass effect by a newly developed spatial method (Leboucher et al., 2020). Using a large-scale database of diatom samples collected along a eutrophication gradient in rivers throughout France, we demonstrated a significantly improved capability of a common diatom index, IBD_{2007} , to measure water quality after the exclusion of mass effect species. Specifically, the relationship between IBD_{mod} scores and the eutrophication gradient was stronger than for the original index, which was created to respond to this type of gradient (Coste et al., 2009; Prygiel et al., 2002; Karthikeyan et al., 2018; Torrisi and Dell'Uomo, 2006).

IBD_{mod} also better discriminated ecological status of the study streams compared to IBD₂₀₀₇ and resulted in reclassification of 2198 out of the 9487 study sites (23.2%). We, therefore, recommend reevaluation of ecological status generated by IBD₂₀₀₇. The risk of assigning an incorrect ecological status to a site with IBD₂₀₀₇ was recently studied by Wach et al. (2019). Authors pointed out the role of the inter-operator variability (human error) in potential misclassification, but we recommend here to investigate the role of mass effect, and we expect a lower variability in ecological status assignment with IBDmod scores. For example, using IBD₂₀₀₇, Szczepocka et al. (2018) reported an unexpected ecological status of good for highly impacted urban sites. Communities from these sites were dominated by one or two taxa, including Diatoma vulgaris a species identified here as influenced by mass effect. In such contexts, when communities show both strong abundance inequalities and dominance of mass effect species, we consider IBD_{mod} more appropriate than IBD₂₀₀₇.

In summary, the results presented here from a national database, indicated that mass effect processes can bias ecological status evaluation. Therefore, identification of mass effect species and their removal from biotic indices will provide more accurate bioassessment.

5. Authors' contributions

T.L., supported by J.T.R. and S.P., was responsible for the development of the study. All authors participated in designing the study and developing aims and research questions. T.L. designed methodology, extracted data and made the analyses, supported by L.M., S.B. and M.W. T.L. led the writing of the manuscript supported by J.T.R. and S.P.. All authors contributed critically to the drafts, contributed to the final version of the manuscript, and gave final approval for publication.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank all French Water Agencies for data sharing, and the French Agency of Biodiversity for its financial support.

References

- AFNOR, 2007. Norme NFT 90-354 Qualité de l'eau Détermination de l'Indice Biologique Diatomées (IBD).
- Akaike, H., 1974. A new look at the statistical model identification. IEEE Trans. Autom. Control 19, 716–723. https://doi.org/10.1109/TAC.1974.1100705.
- Bottin, M., Soininen, J., Alard, D., Rosebery, J., 2016. Diatom cooccurrence shows less segregation than predicted from niche modeling. PLoS ONE 11, e0154581.
- Box, G.E.P., Cox, D.R., 1964. An analysis of transformations. J. R. Statist. Soc. Series B (Methodological) 26, 211–252. https://doi.org/10.1111/j.2517-6161.1964. tb00553.x.
- Brown, B.L., Swan, C.M., 2010. Dendritic network structure constrains metacommunity properties in riverine ecosystems. J. Anim. Ecol. 79, 571–580. https://doi.org/ 10.1111/j.1365-2656.2010.01668.x.
- Brown, B.L., Swan, C.M., Auerbach, D.A., Campbell Grant, E.H., Hitt, N.P., Maloney, K. O., Patrick, C., 2011. Metacommunity theory as a multispecies, multiscale framework for studying the influence of river network structure on riverine communities and ecosystems. J. North Am. Benthol. Soc. 30, 310–327. https://doi. org/10.1899/10-129.1.

- Coste, M., Boutry, S., Tison-Rosebery, J., Delmas, F., 2009. Improvements of the Biological Diatom Index (IBD): description and efficiency of the new version (IBD-2006). Ecol. Ind. 9, 621–650. https://doi.org/10.1016/j.ecolind.2008.06.003.
- Cottenie, K., 2005. Integrating environmental and spatial processes in ecological community dynamics. Ecol. Lett. 8, 1175–1182. https://doi.org/10.1111/j.1461-0248.2005.00820.x.
- European Parliament, 2000. Directive 2000/60/EC of the European Parliament and of the Council establishing a framework for Community action in the field of water policy. O.J.L327, 72pp.
- Goldenberg Vilar, A., van Dam, H., van Loon, E.E., Vonk, J.A., van Der Geest, H.G., Admiraal, W., 2014. Eutrophication decreases distance decay of similarity in diatom communities. Freshw Biol 59, 1522–1531. https://doi.org/10.1111/fwb.12363.
- Heino, J., 2013. The importance of metacommunity ecology for environmental assessment research in the freshwater realm. Biol. Rev. 88, 166–178. https://doi. org/10.1111/j.1469-185X.2012.00244.x.
- Heino, J., Grönroos, M., Ilmonen, J., Karhu, T., Niva, M., Paasivirta, L., 2013. Environmental heterogeneity and beta diversity of stream macroinvertebrate communities at intermediate spatial scales. Freshwater Sci. 32, 142–154. https:// doi.org/10.1899/12-083.1.
- Heino, J., Melo, A.S., Siqueira, T., Soininen, J., Valanko, S., Bini, L.M., 2015. Metacommunity organisation, spatial extent and dispersal in aquatic systems: patterns, processes and prospects. Freshw. Biol. 60, 845–869. https://doi.org/ 10.1111/fwb.12533.
- Jackson, D.A., 1993. Stopping rules in principal components analysis: a comparison of heuristical and statistical approaches. Ecology 74, 2204–2214. https://doi.org/ 10.2307/1939574.
- Jamoneau, A., Passy, S.I., Soininen, J., Leboucher, T., Tison-Rosebery, J., 2018. Beta diversity of diatom species and ecological guilds: response to environmental and spatial mechanisms along the stream watercourse. Freshw Biol 63, 62–73. https:// doi.org/10.1111/fwb.12980.
- JORF, 2015. Arrêté du 27 juillet 2015 modifiant l'arrêté du 25 janvier 2010 relatif aux méthodes et critères d'évaluation de l'état écologique, de l'état chimique et du potentiel écologique des eaux de surface pris en application des articles R. 212-10, R. 212-11 et R. 212-18 du code de l'environnement. Journal Officiel de la République Française.
- Karthikeyan, P., Venkatachalapathy, R., Vennila, G., 2018. Using of biological diatom Indices (IBD) for ecological status of the Cauvery River, India. Indian J. Geo Marine Sci. 47, 426–436. http://nopr.niscair.res.in/handle/123456789/43576.
- Krammer, K., and H. Lange-Bertalot (1986-1991, 1995-2015, 2000-2013). Süßwasserflora von Mitteleuropa. Pages Bacillariophyceae 1. Teil: Naviculaceae. 876 p.; 2. Teil: Bacillariaceae, Epithemiaceae, Surirellaceae, 596 p.; 3. Teil: Centrales, Fragilariaceae, Eunotiaceae, 576 p.; 4. Teil: Achnanthaceae. Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema. 437 p. G. Fischer Verlag.
- Leboucher, T., Tison-Rosebery, J., Budnick, W.R., Jamoneau, A., Vyverman, W., Soininen, J., Boutry, S., Passy, S.I., 2020. A metacommunity approach for detecting species influenced by mass effect. J Appl Ecol 57, 2031–2040. https://doi.org/ 10.1111/1365-2664.13701.
- Lecointe, C., Coste, M., Prygiel, J., 1993. "Omnidia": software for taxonomy, calculation of diatom indices and inventories management. Hydrobiologia 269–270, 509–513. https://doi.org/10.1007/BF00028048.

Legendre, P., Legendre, L.F.J., 2012. Numerical Ecology. Elsevier.

- Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., Holt, R.D., Shurin, J.B., Law, R., Tilman, D., Loreau, M., Gonzalez, A., 2004. The metacommunity concept: a framework for multi-scale community ecology. Ecol. Lett. 7, 601–613. https://doi.org/10.1111/j.1461-0248.2004.00608.x.
- Lenoir, A., Coste, M., 1996. Development of a practical diatom index of overall water quality applicable to the French National Water Board Network. In International symposium, Volksbildungsheim Grilhof Vill, AUT, 17-19 September 1995 (pp. 29-43). Universität Innsbruck.
- Moog, O., Schmutz, S., Schwarzinger, I., 2018. Biomonitoring and Bioassessment. In: Schmutz, S., Sendzimir, J. (Eds.), Riverine Ecosystem Management. Aquatic Ecology Series 8. Springer, Cham. https://doi.org/10.1007/978-3-319-73250-3 19.
- Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Wagner, H., 2018. vegan : Community ecology package. R package version 117–118.
- Passy, S.I., Bode, R.W., 2004. Diatom model affinity (DMA), a new index for water quality assessment. Hydrobiologia 524, 241–252. https://doi.org/10.1023/B: HYDR.0000036143.60578.e0.
- Pont, D., Hugueny, B., Beier, U., Goffaux, D., Melcher, A., Noble, R., Rogers, C., Roset, N., Schmutz, S., 2006. Assessing river biotic condition at a continental scale: a European approach using functional metrics and fish assemblages. J. Appl. Ecol. 43, 70–80. https://doi.org/10.1111/j.1365-2664.2005.01126.x.
- Prygiel, J., Carpentier, P., Almeida, S., et al., 2002. Determination of the biological diatom index (IBD NF T 90–354): results of an intercomparison exercise. J. Appl. Phycol. 14, 27–39. https://doi.org/10.1023/A:1015277207328.
- R Core Team, 2018. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Smucker, N.J., Vis, M.L., 2011. Diatom biomonitoring of streams: reliability of reference sites and the response of metrics to environmental variations across temporal scales. Ecol. Ind. 11, 1647–1657. https://doi.org/10.1016/j.ecolind.2011.04.011.
- Stevenson, R. J., Bahls, L. L., 1999. Periphyton protocols. Rapid bioassessment protocols for use in wadeable streams and rivers: periphyton, benthic macroinvertebrates, and fish. EPA.
- Szczepocka, E., Nowicka-Krawczyk, P., Kruk, A., 2018. Deceptive ecological status of urban streams and rivers—evidence from diatom indices. Ecosphere 9, e02310.

T. Leboucher et al.

- Torrisi, M., Dell'Uomo, A., 2006. Biological monitoring of some Apennine rivers (central Italy) using the diatom-based eutrophication/pollution index (EPI-D) compared to other European diatom indices. Diatom Res. 21, 159–174. https://doi.org/10.1080/ 0269249X.2006.9705657.
- Vilmi, A., Karjalainen, S.M., Hellsten, S., Heino, J., 2016. Bioassessment in a metacommunity context: Are diatom communities structured solely by species sorting? Ecol. Ind. 62, 86–94. https://doi.org/10.1016/j.ecolind.2015.11.043.
- Wach, M., Guéguen, J., Chauvin, C., Delmas, F., Dagens, N., Feret, T., Loriot, S., Tison-Rosebery, J., 2019. Probability of misclassifying river ecological status: a large-scale approach to assign uncertainty in macrophyte and diatom-based biomonitoring. Ecol. Ind. 101, 285–295. https://doi.org/10.1016/j.ecolind.2019.01.028.