



HAL
open science

Predicting the retinal content in omega-3 fatty acids for age-related macular-degeneration

Niyazi Acar, Bénédicte M. J. Merle, Soufiane Ajana, Zhiguo He, Stéphane Grégoire, Boris P. Hejblum, Lucy Martine, Benjamin Buaud, Alain Marie Bron, Catherine Creuzot-garcher, et al.

► To cite this version:

Niyazi Acar, Bénédicte M. J. Merle, Soufiane Ajana, Zhiguo He, Stéphane Grégoire, et al.. Predicting the retinal content in omega-3 fatty acids for age-related macular-degeneration. *Clinical and Translational Medicine*, 2021, 11 (7), pp.e404. 10.1002/ctm2.404 . hal-03293612

HAL Id: hal-03293612

<https://hal.inrae.fr/hal-03293612>

Submitted on 21 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LETTER TO EDITOR

Predicting the retinal content in omega-3 fatty acids for age-related macular-degeneration

Dear Editor,

Current treatments for age-related macular degeneration (AMD)—the leading cause of blindness in industrialized countries—are restricted to a single form of the disease and cannot always avoid severe visual loss.¹ Among emerging preventive strategies, ω -3 polyunsaturated fatty acids (PUFAs) are enjoying interest as they promote normal retinal structure and function, reduce incidence, and slow progression of AMD, consistently with their abundance in retinal neurons. However, direct assessment of retinal ω -3 PUFAs content in humans is impossible, making the use of a systemic biomarker a mandatory surrogate for tissue composition. Here, we determine which circulating lipids have the highest predictive performance for retinal content in ω -3 PUFAs.

As major components of the human retina, ω -3 PUFAs are essential for retinal physiology. The highest content in ω -3 PUFAs is found in retinal photoreceptors, where they inhibit several cellular and molecular mechanisms involved in AMD pathogenesis and the related visual loss. While more than 20 epidemiological studies have consistently shown a 40%-reduction in risk of AMD in subjects with high dietary intake of ω -3 PUFAs,² and human donor studies have measured lower ω -3 PUFAs concentration in AMD eyes compared to age-matched controls,³ supplementation trials failed to impact disease progression.^{4,5} This controversy lies partly in the impossibility of directly assess the retinal ω -3 PUFAs content, making the use of a systemic biomarker a mandatory surrogate for tissue composition.

In the Biomarkers of Lipid Status and Metabolism in Retinal ageing project, we determined which circulating lipids have the highest predictive performance for retinal content in ω -3 PUFAs. In addition to plasma total lipids, we focused our attention on the longer- and medium-term markers of dietary fat, namely red blood cell phospholipids and plasma phosphatidylcholines and cholesteryl esters. By using human donor samples (Table S1) and advanced lipidomics,⁶ we generated an extremely

exhaustive database, corresponding to 304 different lipid species (Tables 1 and S2). Selecting the molecular species relevant for the prediction of retinal ω -3 PUFAs in this high-dimensional, low sample size context is not trivial. This is why we tested several methods combining dimension reduction, variable selection, and prior group structure.⁷ The model with the lowest error of prediction was obtained using sparse group partial least squares and was characterized by a cross-validated r of 0.62 between observed and predicted values of retinal ω -3 PUFAs content (Figure 1A, see Ajana et al⁷ for details on statistical methods). It was based on an algorithm combining the levels of seven plasma cholesteryl esters.⁸ Three of them were characterized by the presence of PUFAs from ω -3 family, the four remaining species consisting in a cholesterol molecule esterified to ω -6 PUFAs (Figure 1B). Interestingly, cholesteryl esters species with ω -3 PUFAs contributed positively to the estimation of retinal ω -3 PUFAs content whereas those with ω -6 PUFAs lowered the prediction index, which is consistent with the well-established competitive metabolism of ω -3 and ω -6 PUFAs. Previous attempts (including ours) on RBC or total plasma lipids have failed in finding strong associations between circulating lipids and absolute levels of retinal ω -3 PUFAs.^{6,9} Here, we confirm that our predictor is more robust (cross-validated $r = 0.62$, Figure 1A) than the use of the ω -3 PUFAs content of RBC ($r = 0.40$) and total plasma ($r = 0.14$).

In the second phase, we further explored the associations of AMD with ω -3 PUFAs status. First, in a post-mortem case-control study, we confirmed previous data showing a lower ω -3 PUFAs content in retinas affected by AMD when compared to those from healthy donors (Figure 2).^{3,9} After statistical adjustments (Table S3), the difference between AMD and control retinas was of -2.41% of total fatty acids. Then, using the algorithm based on seven plasma cholesteryl esters developed at phase 1, we compared the predicted retinal ω -3 PUFAs content between 31 subjects affected by advanced AMD and 31 controls (Table 2) and found that it was lower in cases compared

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Clinical and Translational Medicine* published by John Wiley & Sons Australia, Ltd on behalf of Shanghai Institute of Clinical Bioinformatics

TABLE 1 Fatty acid composition (%) of blood and eye tissues of human donors

	Retina		Red blood cells		Plasma					
					Total		Phosphatidylcholine		Cholesteryl esters	
	Median	(IQR)	Median	(IQR)	Median	(IQR)	Median	(IQR)	Median	(IQR)
C14:0	0.35	(0.13–0.46)	0.68	(0.32–1.39)	0.58	(0.43–0.83)	0.20	(0.11–0.25)	0.36	(0.16–0.77)
C15:0	0.10	(0.07–0.13)	0.23	(0.17–0.29)	0.22	(0.18–0.27)	0.23	(0.15–0.31)	0.20	(0.015–0.27)
C16:0	19.65	(14.76–21.00)	24.34	(21.99–26.74)	23.42	(22.58–25.91)	27.15	(21.83–32.03)	12.67	(11.57–14.44)
C16:1ω-9	0.75	(0.57–0.87)	0.30	(0.23–0.39)	0.30	(0.25–0.34)	0.16	(0.11–0.24)	0.33	(0.24–0.43)
C16:1ω-7	0.55	(0.42–0.77)	2.30	(1.71–2.76)	2.72	(1.87–3.18)	0.48	(0.31–0.77)	3.27	(2.09–4.34)
C17:0	0.15	(0.12–0.18)	0.36	(0.32–0.44)	0.33	(0.30–0.36)	0.47	(0.42–0.57)	0.13	(0.11–0.17)
C18:0	20.52	(19.40–21.33)	9.21	(7.55–10.81)	6.39	(5.73–6.94)	13.33	(10.92–16.07)	0.87	(0.73–1.35)
C18:1trans	0.12	(0.08–0.17)	0.44	(0.35–0.54)	0.47	(0.31–0.64)	0.42	(0.27–0.57)	0.58	(0.20–0.98)
C18:1ω-9	15.46	(14.12–16.18)	29.13	(24.93–34.84)	29.00	(26.52–31.94)	15.42	(13.07–17.31)	25.97	(24.06–28.62)
C18:1ω-7	3.13	(2.88–3.45)	2.28	(1.96–2.55)	2.50	(2.22–2.83)	2.37	(1.99–2.87)	2.14	(1.83–2.52)
C18:2ω-6	1.61	(1.35–1.95)	12.29	(10.30–14.86)	19.95	(17.40–21.97)	16.91	(14.68–20.62)	41.84	(38.03–45.98)
C20:0	0.53	(0.47–0.63)	0.17	(0.13–0.21)	0.14	(0.09–0.15)	0.17	(0.09–0.29)	n.d.	–
C18:3ω-6	0.13	(0.11–0.16)	n.d.	–	0.13	(0.08–0.22)	0.21	(0.00–0.26)	0.37	(0.24–0.61)
C20:1ω-9	0.56	(0.48–0.68)	0.39	(0.33–0.49)	0.27	(0.21–0.32)	0.25	(0.20–0.45)	n.d.	–
C18:3ω-3	0.15	(0.13–0.19)	0.30	(0.20–0.41)	0.37	(0.31–0.54)	0.18	(0.14–0.26)	0.45	(0.31–0.56)
C20:2ω-6	0.16	(0.13–0.20)	0.22	(0.18–0.25)	0.19	(0.16–0.24)	0.38	(0.30–0.52)	n.d.	–
C20:3ω-9	0.31	(0.21–0.40)	n.d.	–	0.14	(0.11–0.19)	0.23	(0.19–0.33)	n.d.	–
C22:0	0.36	(0.30–0.49)	0.23	(0.16–0.32)	0.17	(0.12–0.21)	n.d.	–	n.d.	–
C20:3ω-6	1.57	(1.29–2.05)	0.67	(0.42–0.93)	0.78	(0.59–1.07)	2.34	(1.78–2.95)	0.60	(0.45–0.67)
C22:1ω-9	0.10	(0.07–0.13)	n.d.	–	n.d.	–	n.d.	–	n.d.	–
C20:4ω-6	10.92	(10.31–12.15)	6.18	(3.24–8.12)	5.43	(4.33–6.79)	9.14	(6.79–11.44)	6.30	(4.99–8.34)
C24:0	0.30	(0.25–0.41)	0.39	(0.27–0.52)	0.10	(0.00–0.16)	n.d.	–	n.d.	–
C20:5ω-3	0.18	(0.14–0.24)	0.28	(0.19–0.40)	0.43	(0.22–0.59)	0.63	(0.37–0.86)	0.62	(0.42–0.83)
C24:1ω-9	0.11	(0.08–0.17)	0.64	(0.48–0.92)	0.42	(0.26–0.57)	n.d.	–	n.d.	–
C22:4ω-6	1.37	(1.26–1.62)	0.83	(0.48–1.12)	0.26	(0.20–0.30)	0.40	(0.31–0.45)	n.d.	–
C22:5ω-6	0.53	(0.43–0.71)	0.17	(0.11–0.27)	0.15	(0.12–0.19)	0.27	(0.21–0.37)	n.d.	–
C22:5ω-3	1.04	(0.86–1.26)	0.83	(0.40–1.05)	0.45	(0.38–0.49)	1.02	(0.70–1.15)	n.d.	–
C22:6ω-3	15.20	(13.09–16.87)	1.84	(0.70–2.31)	1.54	(1.27–1.89)	2.99	(2.21–4.20)	0.53	(0.41–0.67)
total SFAs	45.34	(41.90–46.87)	38.41	(35.52–40.58)	32.40	(31.07–34.21)	42.63	(39.17–45.63)	14.80	(13.24–17.03)
total MUFAs	21.49	(19.94–22.81)	36.48	(31.50–29.26)	36.42	(32.56–39.02)	19.48	(17.15–21.38)	33.14	(31.10–35.54)
total PUFAs	33.41	(30.87–36.60)	25.25	(16.59–29.26)	30.70	(27.04–34.26)	36.86	(32.49–40.10)	51.7	(47.95–54.87)
total ω-6	16.42	(15.55–19.07)	21.32	(14.80–24.91)	27.48	(24.18–30.89)	10.75	(7.82–12.66)	6.90	(5.23–8.72)
total ω-3	16.79	(14.46–18.43)	3.41	(1.72–4.09)	2.89	(2.39–3.66)	4.95	(3.96–6.08)	1.63	(1.40–2.05)
ω-6 / ω-3 ratio	1.04	(0.91–1.20)	7.45	(5.36–9.28)	9.49	(7.77–11.33)	2.12	(1.63–2.76)	4.34	(3.39–5.51)

Abbreviations: IQR, interquartile range; MUFAs, monounsaturated fatty acids; n.d., not detected; PUFAs, polyunsaturated fatty acids; SFAs, saturated fatty acids.

to controls. The difference in predicted retinal ω -3 PUFAs content between AMD cases and controls was of -1.39% . These observations are consistent with the pathophysiology of AMD that is characterized by the loss of the ω -3 PUFA-rich photoreceptor cells.

Considering the crucial functions played by ω -3 PUFAs in retinal structure and function and its depletion in AMD eyes, the idea that maintaining high retinal lev-

els of ω -3 PUFAs would protect against the development and/or the progression of the disease makes sense. This hypothesis is consolidated by a large number of studies showing a significantly reduced risk for developing AMD in subjects with high dietary consumption of ω -3 PUFAs.² However, in the two randomized trials AREDS2 and NAT2,^{4,5} ω -3 PUFAs supplementation did not modify disease progression. One possible explanation




FIGURE 1 Plasma cholesteryl esters as markers of retinal content in ω -3 PUFAs. (A) Output from the prediction model resulting from sgPLS analysis showing a higher robustness of predictor when compared to total plasma and RBC ω -3 PUFAs in estimating retinal ω -3 PUFA content. (B) Molecular species of cholesteryl esters whose plasma concentration values are used to predict retinal ω -3 PUFA content. The plasma concentration values of cholesteryl esters with ω -3 PUFAs contributed positively to the calculated index whereas those of ω -6 PUFA cholesteryl esters lowered it




FIGURE 2 Predicted levels of retinal ω -3 PUFAs are discriminant for AMD and equivalent to a direct assessment: case-control studies. Adjusted differences black squares and 95% confidence intervals (CIs) (horizontal lines) of observed and predicted retinal ω -3 PUFAs contents for age-related macular degeneration in donor eyes and humans patients. For postmortem study, adjusted difference was assessed by linear regression adjusted for age, gender, and time after death. For clinical study, adjusted difference was assessed by mixed linear regression adjusted for age, body mass index, smoking, ω -3 supplement use, HDL-cholesterol, and LDL-cholesterol

TABLE 2 Characteristics of the participants to the case-control and LIMPIA studies

Case-control study	cases (n = 31)	controls (n = 31)	p value
Age, mean (SD) (range), y	84.5 (4.2) (76.7–92.5)	84.2 (4.3) (76.8–90.7)	0.4
Female, no (%)	22 (71.0)	22 (71.0)	–
BMI, mean (SD), kg/m ²	24.8 (3.0) (18.9–31.6)	24.8 (3.6) (18.1–34.7)	0.98
Plasma total cholesterol, mean (SD) (range), mmol/L	5.89 (0.98) (3.32–8.62)	5.59 (1.00) (3.00–7.56)	0.24
Plasma HDL cholesterol, mean (SD) (range), mmol/L	1.62 (0.34) (1.01–2.18)	1.45 (0.38) (0.82–2.22)	0.05
Plasma LDL cholesterol, mean (SD) (range), mmol/L	3.73 (0.84) (1.67–6.01)	3.57 (0.94) (1.14–5.55)	
Plasma triglycerides, mean (SD) (range), mmol/L	1.19 (0.53) (0.69–3.22)	1.26 (0.56) (0.40–2.37)	0.49
ω -3 Supplement use, yes (%)	6 (19.4)	1 (3.2)	0.10
Smoking, no (%)			0.55
Never smoker	19 (61.3)	21 (67.7)	
<20 pack-year	5 (16.1)	6 (19.3)	
\geq 20 pack-year	7 (22.6)	4 (13.0)	
LIMPIA study	Supplemented (n = 29)	Placebo (n = 26)	p value
Age, mean (SD) (range), y	57.7 (6.3) (44.6–70.2)	55.6 (6.9) (44.3–66.1)	0.22
Female, no (%)	22 (75.9)	20 (76.9)	0.93
BMI, mean (SD), kg/m ²	24.2 (3.5) (19.2–31.3)	23.8 (3.8) (18.1–32.8)	0.74
Plasma total cholesterol, mean (SD) (range), mmol/L	5.85 (0.81) (4.32–7.62)	5.84 (0.72) (4.28–7.53)	0.95
Plasma HDL cholesterol, mean (SD) (range), mmol/L	1.73 (0.37) (1.14–2.54)	1.74 (0.56) (0.98–3.44)	0.94
Plasma LDL cholesterol, mean (SD) (range), mmol/L	3.58 (0.77) (1.74–4.98)	3.58 (0.73) (1.37–4.82)	0.99
Plasma triglycerides, mean (SD) (range), mmol/L	1.19 (0.44) (0.61–2.16)	1.14 (0.42) (0.50–1.98)	0.68
Smoking, no (%)			0.85
Never smoker	16 (61.5)	16 (55.2)	
<20 pack-year	6 (23.1)	7 (24.1)	
\geq 20 pack-year	4 (15.4)	6 (20.7)	

Abbreviations: BMI, body mass index; HDL, high density lipoprotein; LDL, low density lipoprotein; SD, standard deviation.

to these contradictory results lies on subjects' sensitivity to nutritional supplementation. Indeed, whereas ω -3 PUFAs supplementation had no effect on AMD progression when considering total population, we have shown in NAT2 study that subjects who maintained steadily high blood levels of ω -3 PUFAs had a significantly lower risk for developing advanced AMD.¹⁰ This finding confirms the need of monitoring the metabolic status of subjects participating in nutritional interventions. This is why we evaluated the sensitivity of the predicted retinal ω -3 PUFAs content to a dietary supplementation with ω -3 PUFAs. We analyzed blood samples collected within the LIMPIA randomized clinical trial in which participants received ω -3 PUFAs for 6 months (Table 2). Whereas the median values

of predicted retinal ω -3 PUFAs content at baseline were similar in both groups, they were significantly increased in supplemented subjects at 3 months, this difference being maintained after 6 months of supplementation (Figure 3).

In summary, we identified a blood biomarker of retinal ω -3 PUFAs status, which is inversely related to the risk for advanced AMD and increased with supplementation in ω -3 PUFAs. If confirmed in larger studies, it may serve as a practical tool for scientists to set-up and conduct clinical trials, but also to help doctors to diagnose the condition earlier and single out individuals who are more likely to benefit from nutritional treatment. Finally, an independent model validation dataset and samples from




FIGURE 3 Predicted levels of retinal ω -3 PUFAs are sensitive to a dietary supplementation with ω -3 PUFAs. Medians (horizontal lines) and interquartile ranges (boxes) of predicted retinal ω -3 PUFAs content among participants to the LIMPIA study. Linear regression model adjusted for age, gender, body mass index, smoking, plasma LDL-cholesterol, and plasma HDL-cholesterol

intermediate-AMD subjects would also strengthen our findings.

CONFLICT OF INTEREST

The authors declare no conflict of interest related to this study.

ACKNOWLEDGMENT

This work was supported by grants from Agence Nationale de la Recherche (ANR-14-CE12-0020-01 BLISAR; ANR-11-LABX-0021-01), the Conseil Régional Bourgogne, Franche-Comté (PARI grant), the FEDER (European Funding for Regional Economical Development), and the Fondation de France/Fondation de l'oeil.

AUTHOR CONTRIBUTIONS

Designed the studies: Niyazi Acar, Benjamin Buaud, Lionel Bretillon, and Cécile Delcourt. Conducted experiments: Zhiguo He, Stéphane Grégoire, Lucy Martine, and Benjamin Buaud. Acquired data: Stéphane Grégoire, Lucy Martine, Benjamin Buaud, Olivier Berdeaux, Jean-François Korobelnik, Catherine P. Creuzot-Garcher, and Alain M. Bron. Analyzed data: Niyazi Acar, Bénédicte M. J. Merle, Soufiane Ajana, Boris P. Hejblum, Benjamin Buaud, Olivier Berdeaux, Héléne Jacqmin-Gadda, Lionel Bretillon, and Cécile Delcourt. Wrote manuscript: Niyazi Acar, Bénédicte M. J. Merle, Soufiane Ajana, Zhiguo He, Stéphane Grégoire, Boris P. Hejblum, Lucy Martine, Benjamin Buaud, Alain M. Bron, Catherine P. Creuzot-Garcher, Jean-François Korobelnik, Olivier Berdeaux, Héléne Jacqmin-Gadda, Lionel Bretillon, and Cécile Delcourt.

AVAILABILITY OF DATA AND MATERIALS

The datasets generated and analyzed during the current study are neither publicly available due to personal information involved nor from the corresponding author due to a pending patent.

Niyazi Acar¹ 
 Bénédicte M. J. Merle^{3,8}
 Soufiane Ajana^{3,8}
 Zhiguo He⁴
 Stéphane Grégoire¹
 Boris P. Hejblum⁵
 Lucy Martine¹
 Benjamin Buaud⁶
 Alain M. Bron^{1,7}
 Catherine P. Creuzot-Garcher^{1,7}
 Jean-François Korobelnik^{2,8}
 Olivier Berdeaux²
 Héléne Jacqmin-Gadda^{3,8}
 Lionel Bretillon^{1,8}
 Cécile Delcourt^{3,8}

for the Biomarkers of Lipid Status And metabolism in Retinal ageing (BLISAR) Study Group[#]

¹ Eye and Nutrition Research Group, AgroSup Dijon, CNRS, INRAE, Centre des Sciences du Goût et de l'Alimentation Université Bourgogne Franche-Comté, Dijon, France

² ChemoSens Platform, Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE, Université Bourgogne Franche-Comté, Dijon, France

³ Team LEHA, UMR 1219 Bordeaux Population Health Research Center, Univ. Bordeaux, INSERM, F-33000 Bordeaux, France

⁴ Biology, Imaging, and Engineering of Corneal Graft, Faculty of Medicine, Jean Monnet University, Saint-Etienne, France

⁵ Team SISTM, UMR 1219 Bordeaux Population Health Research Center, Univ. Bordeaux, INSERM, INRIA BSO, F-33000 Bordeaux, France

⁶ ITERG - Equipe Nutrition – Santé and Biochimie des Lipides, Canéjan, France

⁷ Department of Ophthalmology, University Hospital, Dijon, France

⁸ CHU Bordeaux, Service d'ophtalmologie, Bordeaux, France

Correspondence

Niyazi Acar, Eye and Nutrition Research Group, Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE, Université Bourgogne Franche-Comté, Dijon, F-21000, France.
Email: niyazi.acar@inrae.fr

[§]These authors contributed equally to this work.

[#]BLISAR STUDY GROUP: Niyazi Acar¹, Soufiane Ajana², Olivier Berdeaux¹, Lionel Bretillon¹, Alain Bron^{1,3}, Benjamin Buaud⁴, Stéphanie Cabaret¹, Philippe Gain⁵, Audrey Cougnard-Grégoire², Catherine Creuzot-Garcher^{1,3}, Cécile Delcourt², Marie-Noelle Delyfer^{2,6}, Catherine Féart², Valérie Febvret¹, Stéphane Grégoire¹, Zhiguo He⁵, Jean-François Korobelnik^{2,6}, Lucy Martine¹, Bénédicte MJ Merle², Gilles Thuret⁵, Carole Vaysse⁴

¹Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE, Université Bourgogne Franche-Comté, Dijon, France ²Univ. Bordeaux, INSERM, Bordeaux Population Health Research Center, team LEHA, UMR 1219, F-33000 Bordeaux, France

³Department of Ophthalmology, University Hospital, Dijon, France ⁴ITERG - Equipe Nutrition Santé & Biochimie des Lipides, Canéjan, France ⁵Laboratory for Biology, Imaging, and Engineering of Corneal Grafts, EA2521, Faculty of Medicine, University Jean Monnet, Saint-Etienne, France ⁶CHU de Bordeaux, Service d'Ophtalmologie, Bordeaux, F-33000, France

ORCID

Niyazi Acar  <https://orcid.org/0000-0002-3401-8079>

REFERENCES

- Mitchell P, Liew G, Gopinath B, Wong TY. Age-related macular degeneration. *Lancet*. 2018;392(10153):1147–1159.
- van Leeuwen EM, Emri E, Merle BMJ, et al. A new perspective on lipid research in age-related macular degeneration. *Prog Retin Eye Res*. 2018;67:56–86.
- Liu A, Chang J, Lin Y, et al. Long-chain and very long-chain polyunsaturated fatty acids in ocular aging and age-related macular degeneration. *J Lipid Res*. 2010;51(11):3217–3229.
- Age-Related Eye Disease Study 2 Research Group. Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *JAMA*. 2013;309(19):2005–2015.
- Souied EH, Delcourt C, Querques G, et al. Oral docosahexaenoic acid in the prevention of exudative age-related macular degeneration: the nutritional AMD treatment 2 study. *Ophthalmology*. 2013;120(8):1619–1631.
- Acar N, Berdeaux O, Gregoire S, et al. Lipid composition of the human eye: are red blood cells a good mirror of retinal and optic nerve fatty acids?. *PLoS One*. 2012;7(4):e35102.
- Ajana S, Acar N, Bretillon L, et al. Benefits of dimension reduction in penalized regression methods for high dimensional grouped data: a case study in low sample size. *Bioinformatics*. 2019;35(19):3628–3634.
- Delcourt C, Acar N, Bretillon L, et al. Prédiction du contenu de la rétine en acides gras polyinsaturés oméga-3 à travers le dosage de 7 molécules d'esters de cholestérol. Vol. FR 1910515 (Patent FR 1910515; France, 2019).
- Gorusupudi A, Liu A, Hageman GS, Bernstein PS. Associations of human retinal very long-chain polyunsaturated fatty acids with dietary lipid biomarkers. *J Lipid Res*. 2016;57(3):499–508.
- Merle BM, Benlian P, Puche N, et al. Circulating omega-3 fatty acids and neovascular age-related macular degeneration. *Invest Ophthalmol Vis Sci*. 2014;55(3):2010–2019.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.