



**HAL**  
open science

## Confrontation of the “Dual Tracer” Indirect Method With Direct Ileal Sampling for Indispensable Amino Acid Digestibility of Sunflower Isolate in Humans

Romain Tessier, Juliane Calvez, Nadezda Khodorova, Alain Quinsac, Romain Kapel, Olivier Galet, Daniel Tomé, Gheorghe Airinei, Robert Benamouzig, Claire C. Gaudichon

### ► To cite this version:

Romain Tessier, Juliane Calvez, Nadezda Khodorova, Alain Quinsac, Romain Kapel, et al.. Confrontation of the “Dual Tracer” Indirect Method With Direct Ileal Sampling for Indispensable Amino Acid Digestibility of Sunflower Isolate in Humans. *Nutrition* 2021, Jun 2021, online, United States. 10.1093/cdn/nzab048\_019 . hal-03296232

**HAL Id: hal-03296232**

**<https://hal.inrae.fr/hal-03296232>**

Submitted on 29 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

### Confrontation of the “Dual Tracer” Indirect Method With Direct Ileal Sampling for Indispensable Amino Acid Digestibility of Sunflower Isolate in Humans

Romain Tessier<sup>1</sup>, Juliane Calvez<sup>1</sup>, Nadezda Khodorova<sup>1</sup>, Alain Quinsac<sup>2</sup>, Romain Kapel<sup>3</sup>, Olivier Galet<sup>4</sup>, Daniel Tomé<sup>1</sup>, Gheorghe Airinei<sup>1</sup>, Robert Benamouzig<sup>1</sup>, and Claire Gaudichon<sup>1</sup>

<sup>1</sup>Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, France; <sup>2</sup>Terres Inovia; <sup>3</sup>LRGP Unité Mixte de Recherche CNRS 7274, Université de Lorraine, Plateforme SVS; and <sup>4</sup>Avril

**Objectives:** The direct assessment of ileal samples of amino acid (AA) digestibility is invasive in humans. A less invasive but indirect method, namely « dual tracer » was recently developed. It relies on the plasma isotopic enrichment ratio of two labeled protein, a reference protein or alternately AAs labeled with <sup>13</sup>C and the test protein labeled with <sup>15</sup>N. This recent method has not yet been challenged against direct measurement of ileal digestibility.

**Methods:** Seven healthy volunteers were intubated with naso-ileal tube. Every 30min for 4h, they ingested sunflower biscuits containing a total of 25g of <sup>15</sup>N intrinsically labeled sunflower protein isolate. They also ingested 60g chocolate containing a total of 400mg of a mix of <sup>13</sup>C algal individual AAs. Ileal contents were collected continuously for 8h following the first meal and plasma was sampled every 30min for 4h and hourly between 4 and 8h. <sup>15</sup>N and <sup>13</sup>C indispensable amino

acid (IAA) ileal digestibility were determined by measuring <sup>15</sup>N and <sup>13</sup>C enrichment in AAs by GC-C-IRMS and AA content by UHPLC in ileal effluent. Plasma and meal <sup>15</sup>N and <sup>13</sup>C IAA enrichment were measured by GC-C-IRMS. Isotopic <sup>15</sup>N/<sup>13</sup>C ratio were determined using area under the curve value for each isotope.

**Results:** Using direct ileal sampling, average IAA ileal digestibility was: (i)  $88.5 \pm 5.0\%$  for sunflower isolate (<sup>15</sup>N) with values ranged from  $85.8 \pm 5.1\%$  for threonine to  $91.1 \pm 5.8\%$  for methionine, and (ii)  $97.6 \pm 1.7\%$  for free AAs (<sup>13</sup>C) with values ranged from  $95.9 \pm 2.3\%$  for lysine to  $98.8 \pm 0.8\%$  for phenylalanine. With the “dual tracer” method, digestibility of isoleucine, leucine, threonine and valine was significantly lower than with ileal determination (from 7.9% for threonine to 24.3% for leucine), Methionine and phenylalanine values were aberrant (over 100%) For lysine, the difference between the two methods was not statistically different (4.7%,  $p = 0.49$ ).

**Conclusions:** With our methodological conditions, the “dual tracer” method provides physiological values for most IAA except methionine and phenylalanine. However, values were low compared to ileal digestibility (about 10%) and interindividual variability was high. This less invasive method is promising but requires methodological improvements.

**Funding Sources:** French Research National Agency (ANR), financial support of SOFIPROTEOL under the FASO Project PRODIAL.