

Plasmodesmal Components Involved in Cell-to-cell Transport of Potyviruses—Focus on HVA22a candidate

Mingshuo Xue, Luc Sofer, Vincent Simon, Roxane Lion, Jens Tilsner², Sylvie German-Retana

▶ To cite this version:

Mingshuo Xue, Luc Sofer, Vincent Simon, Roxane Lion, Jens Tilsner², et al.. Plasmodesmal Components Involved in Cell-to-cell Transport of Potyviruses—Focus on HVA22a candidate. EMBO Workshop- Intercellular communication and plasmodesmata in plant development and disease, Jul 2021, Virtual, France. hal-03305277

HAL Id: hal-03305277 https://hal.inrae.fr/hal-03305277v1

Submitted on 28 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EMBO Workshop- Intercellular communication and plasmodesmata in plant development and disease-12-16 July 2021/Virtual

Plasmodesmal Components Involved in Cell-to-cell Transport of Potyviruses---Focus on HVA22a candidate

Mingshuo Xue¹, Luc Sofer¹, Vincent Simon¹, Roxane Lion¹, Jens Tilsner^{2,3} and Sylvie German-Retana¹

- ¹ UMR 1332, Biologie du Fruit et Pathologie, INRAe, Univ. Bordeaux, Equipe de Virologie, 71 Av. E. Bourlaux, CS 20032, 33882 Villenave d'Ornon CEDEX, France
- ² Biomedical Sciences Research Complex, Univ. St. Andrews, St. Andrews, United Kingdom
- ³ Cell and Molecular Sciences, James Hutton Institute, Dundee, United Kingdom

Potyvirus is one of the largest genera of plant viruses responsible for serious diseases in vegetable and fruit crops worldwide (Scholthof *et al*, 2011). The potyviral species *Turnip mosaic virus* (TuMV) represents one of the rare examples of plant viruses that utilize the host endomembrane system to produce membranous vesicles mobile between cells. The 6K2 protein of TuMV, a small transmembrane protein, induces the formation of endoplasmic reticulum (ER)-derived viral vesicles, important not only for replication but also for intracellular and intercellular movement (Cotton *et al*, 2009; Laliberté & Zheng, 2014).

In *Arabidopsis thaliana*, AtHVA22a (*Hordeum vulgare abscisic* acid responsive gene 22) belongs to a multigenic family of transmembrane proteins, homologous of reticulons and DP1/Yop1 family proteins (Chen *et al*, 2002), which are responsible for the constriction of ER tubules (Shibata *et al*, 2008). Reticulons play a role in positive strand RNA virus replication in animals and plants (Diaz *et al*, 2010). Moreover, semi-quantitative proteomics analysis of plasmodesmata (PD) fractions purified from *A. thaliana* suspension cells showed that AtHVA22c and AtHVA22a are highly enriched in PD proteome (Brault *et al*, 2019).

In our study, we showed that TuMV-6K2 interacts with AtHVA22a by split-ubiquitin yeast two hybrid assay (SuY2H) in yeast and further confirmed this interaction in planta by Bimolecular Fluorescence Complementation (BiFC). Overexpression of AtHVA22a increases TuMV propagation in *Nicotiana benthamiana*. Furthermore, AtHVA22a is partially re-localized at the level of the viral replication complex (VRC) during TuMV infection and the 6K2-induced vesicles at the PD. The interaction between AtHAV22a and 6K2 observed in BiFC is also localized at the VRC during viral infection.

Altogether, our results indicate that HVA22a is a candidate protein potentially involved in replication and cell to cell movement of TuMV.

References:

- Brault ML, Crowet J, et al (2019) Multiple C2 domains and transmembrane region proteins (MCTP s) tether membranes at plasmodesmata. EMBO reports 20.
- Chen C-N, David T-H (2002) AtHVA22 gene family in Arabidopsis: phylogenetic relationship, ABA and stress regulation, and tissue-specific expression. 12
- Cotton S, Laliberté J-F (2009) Turnip Mosaic Virus RNA Replication Complex Vesicles Are Mobile, Align with Microfilaments, and Are Each Derived from a Single Viral Genome. *Journal of Virology* 83: 10460–10471
- Diaz A, Ahlquist P (2010) Membrane-shaping host reticulon proteins play crucial roles in viral RNA replication compartment formation and function. *Proceedings of the National Academy of Sciences* 107: 16291–16296
- Grangeon R, Laliberté J-F (2013) 6K2-induced vesicles can move cell to cell during turnip mosaic virus infection. Frontiers in Microbiology 4
- Laliberté J-F & Zheng H (2014) Viral Manipulation of Plant Host Membranes. *Annual Review of Virology* 1: 237–259 Scholthof K-BG, Ahlquist P, *et al* (2011) Top 10 plant viruses in molecular plant pathology: Top 10 plant viruses. *Molecular Plant Pathology* 12: 938–954
- Shibata Y, Voeltz GK (2008) The Reticulon and Dp1/Yop1p Proteins Form Immobile Oligomers in the Tubular Endoplasmic Reticulum. *Journal of Biological Chemistry* 283: 18892–18904