



**HAL**  
open science

## **HVA22a and HVA22c, two candidate proteins involved in potyvirus replication and movement**

Mingshuo Xue, Luc Sofer, Simon Vincent, Roxane Lion, Jens Tilsner, Sylvie  
German-Retana

► **To cite this version:**

Mingshuo Xue, Luc Sofer, Simon Vincent, Roxane Lion, Jens Tilsner, et al.. HVA22a and HVA22c, two candidate proteins involved in potyvirus replication and movement. Journée de l'École Doctorale des Sciences de la Vie et de la Santé de Bordeaux-, May 2021, Bordeaux, France. hal-03305383

**HAL Id: hal-03305383**

**<https://hal.inrae.fr/hal-03305383v1>**

Submitted on 28 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## HVA22a and HVA22c, two candidate proteins involved in *potyvirus* replication and movement

Mingshuo Xue<sup>1</sup>, Luc Sofer<sup>1</sup>, Simon Vincent<sup>1</sup>, Roxane Lion<sup>1</sup>, Jens Tilsner<sup>2,3</sup> and Sylvie German-Retana<sup>1</sup>.

<sup>1</sup> UMR 1332, Biologie du Fruit et Pathologie, INRAe, Univ. Bordeaux, Equipe de Virologie, 71 Av. E. Bourlaux, CS 20032, 33882 Villenave d'Ornon CEDEX, France

<sup>2</sup> Biomedical Sciences Research Complex, Univ. St. Andrews, St. Andrews, United Kingdom

<sup>3</sup> Cell and Molecular Sciences, James Hutton Institute, Dundee, United Kingdom

*Potyvirus* is one of the largest genera of plant viruses responsible for serious diseases in vegetable and fruit crops worldwide. The completion of the viral cycle results from a complex interplay between virus- and plant-encoded factors also called susceptibility factors. In this scheme, absence or non-adequacy of a single susceptibility factor leads to full or partial resistance to viruses. In this context, this work aims at identifying plant factors involved in the potyvirus cell-to-cell movement, a key step of the viral cycle. Indeed, to invade the whole plant after replication, plant viruses move intracellularly to reach the plasmodesmata (PD), the symplasmic tunnels between plant cells that are the gateway for this movement, cross them to enter in the neighboring cells, and enter into sieve elements.

The potyviral species *Turnip mosaic virus* (TuMV) represents one of the rare examples of plant viruses that utilizes the host endomembrane system to produce membranous vesicles mobile between cells, reminding of animal viruses that utilize membrane-derived vesicles for exit from infected cells and entry into healthy cells. This is the 6K2 protein of TuMV, a small transmembrane protein that is involved in this rearrangement of the endoplasmic reticulum (ER) for the generation of those viral vesicles important not only for replication but also for intracellular and intercellular movement. Viral RNA and proteins, together with host proteins essential for potyvirus replication are recruited in these small vesicles, which are transported through PD to neighboring plant cells to achieve the systematic infection.

In *Arabidopsis thaliana*, AtHVA22a and c (*Hordeum vulgare* abscisic acid responsive gene 22) belong to a multigenic family of transmembrane proteins, homologous of reticulons and DP1/Yop1 family, which are responsible for the constriction of ER tubules and that also play a role in positive strand RNA virus replication in animals and plants. Moreover, semi-quantitative proteomics analysis of PD fractions purified from *A. thaliana* suspension cells showed that AtHVA22c and AtHVA22a are highly enriched in PD proteome.

In our study, we showed that TuMV-6K2 interacts with AtHVA22a and AtHVA22c, by split-ubiquitin yeast two hybrid assay (SuY2H) in yeast and further confirmed those interactions in planta by Bimolecular Fluorescence Complementation (BiFC). Overexpression of AtHVA22a increases TuMV propagation in *Nicotiana benthamiana*. However, lack of AtHVA22c expression has a negative effect of TuMV propagation in *A. thaliana*. Furthermore, both AtHVA22a/c are partially localized at the viral replication complex (VRCs) during TuMV infection and have partial localization with 6K2-induced vesicles. The interactions between AtHVA22a/c and 6K2 observed in BiFC are also localized at the VRCs during viral infection.

Altogether, our results indicate that HVA22a and HVA22c, are candidate proteins potentially involved in replication and cell to cell movement of TuMV.