



HAL
open science

Microbial diversity and activity assessment in a 100-year-old lead mine

Sara Gallego, José María Esbrí, Juan Antonio Campos, Jesús Daniel Peco,
Fabrice Martin-Laurent, Pablo Higuera

► **To cite this version:**

Sara Gallego, José María Esbrí, Juan Antonio Campos, Jesús Daniel Peco, Fabrice Martin-Laurent, et al.. Microbial diversity and activity assessment in a 100-year-old lead mine. *Journal of Hazardous Materials*, 2021, 410, pp.124618. 10.1016/j.jhazmat.2020.124618 . hal-03310791

HAL Id: hal-03310791

<https://hal.inrae.fr/hal-03310791v1>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1

2 **Microbial diversity and activity assessment in a 100-year-old lead mine**

3

4 Sara Gallego^{1*}, José María Esbrí^{2,3}, Juan Antonio Campos^{2,4}, Jesús Daniel Peco^{2,4}, Fabrice
5 Martin-Laurent¹ and Pablo Higuera^{2,3}

6 ¹AgroSup Dijon, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, Agroécologie,
7 Dijon, France

8 ²Instituto de Geología Aplicada, IGeA. Universidad de Castilla-La Mancha. Plaza de Manuel
9 Meca, 1, 13400, Almadén, Ciudad Real, Spain

10 ³Escuela de Ingeniería Minera e Industrial de Almadén. Plaza de Manuel Meca, 1, 13400,
11 Almadén, Ciudad Real, Spain

12 ⁴Escuela de Ingenieros Agrónomos, Universidad de Castilla-La Mancha. Ronda de Calatrava,
13 7, 13071, Ciudad Real, Spain

14

15

16 *Corresponding author: Sara Gallego

17 Agroécologie, INRAE, 17 rue Sully, BP86510, 21065 Dijon Cedex, France. Tel: 33 3 80 69

18 34 06. Fax: 33 3 80 69 32 24. Email: sara.gallego.blanco@gmail.com

19 **Abstract**

20 Mining activities frequently leave a legacy of residues that remain in the area for long periods
21 causing the pollution of surroundings. We studied on a 100 year-old mine, the behaviour of
22 potentially toxic elements (PTEs) and their ecotoxicological impact on activity and diversity
23 of microorganisms. The PTEs contamination assessment allowed the classification of the
24 materials as highly (reference- and contaminated-samples) and very highly polluted (illegal
25 spill of olive mill wastes (OMW), tailings and dumps). OMW presented the lowest enzymatic
26 activities while tailings and dumps had low dehydrogenase and arylsulfatase activities. All the
27 alpha diversity indices studied were negatively impacted in dumps. Tailings had lower Chao1
28 and PD whole tree values as compared to those of reference-samples. Beta diversity analysis
29 showed similar bacterial community composition for reference- and contaminated-samples,
30 significantly differing from that of tailings and dumps. The relative abundance of
31 Gemmatimonadetes, Bacteroidetes and Verrucomicrobia was lower in OMW, tailings and
32 dumps as compared to reference-samples. Fifty-seven OTUs were selected as responsible of
33 the changes observed between samples. This study highlights that assessing the relationship
34 between physicochemical properties and microbial diversity and activity gives clues about
35 ongoing regulating processes that can be helpful for stakeholders to define an appropriate
36 management strategy.

37 **Keywords:** Potentially toxic elements, mining soil, enzyme activities, high-throughput
38 sequencing, microbial ecotoxicology,

39

40 **1. Introduction**

41 Mining activities are found all over the world because they provide access to mineral
42 resources that fuel various industrial activities in both developed and in developing countries.
43 Although recognized of great importance for the world gross domestic product, mining
44 operations are often viewed as an important source of pollution with negative impact on the
45 environment. During the processes of mineral extraction and preparation, large amounts of
46 ore wastes and debris are commonly accumulated in the proximity of the mining operation
47 site. These materials are essentially fractured rocks and soil devoid of vegetation,
48 characterized by high concentrations of heavy metals and metalloids. Consequently, the
49 environment is drastically transformed in highly polluted barren areas (Martín Duque et al.,
50 2015; Sánchez-Donoso et al., 2019), which can be toxic to human health and other life,
51 including plants and microorganisms (Giller et al., 1998; Nagajyoti et al., 2010; Tchounwou
52 et al., 2012). Furthermore, potentially toxic elements (PTEs) from these polluted areas can
53 transfer to surrounding aquatic and terrestrial compartments via leaching or runoff (Elmayel
54 et al., 2020; Fernández-Martínez et al., 2019; Jung and Thornton, 1996; Kisková et al., 2018),
55 disperse in the atmosphere (Esbrí et al., 2020; Naharro et al., 2020, 2018), and indirectly
56 contribute to pollutant dissemination. The recent interest in the reclamation of abandoned
57 mining sites in arid and semiarid regions for agricultural purposes highlights the need to
58 understand the biogeochemical processes contributing to soil health and fertility (Favas et al.,
59 2018; Higuera et al., 2019b; Krzaklewski and Pietrzykowski, 2002; Mendez et al., 2008)

60 Microorganisms such as bacteria and fungi are key players in soil ecosystem services.
61 They are involved in multiple geochemical cycles, influence plant growth and contribute to
62 climate regulation and soil restoration, among others (Ayangbenro et al., 2018; Cavicchioli et
63 al., 2019; Domeignoz-Horta et al., 2018, 2015; Van Der Heijden et al., 2008). Toxicity from
64 heavy metals and metalloids can compromise their growth and survival, with enormous

65 consequences on ecosystem functioning (Bell et al., 2005; Delgado-Baquerizo et al., 2020,
66 2017, 2016b, 2016a; Domeignoz-Horta et al., 2020; Griffiths and Philippot, 2013; Philippot et
67 al., 2013; Wagg et al., 2014). However, some microorganisms have developed several
68 resistance mechanisms to cope with their toxic effects such as extra and intracellular
69 sequestration, exclusion by permeability barriers, enzymatic detoxification, efflux-pumps and
70 specific reduction of cellular targets' sensibility (Hobman et al., 2007; Nies, 2003; Rouch et
71 al., 1995). In this frame, soil microorganisms constitute useful bio indicators of soil quality
72 (Alvarenga et al., 2018; Thiele-Bruhn et al., 2020). Measurement of microbial activity
73 through analysis of soil extracellular enzymatic activities has long been used as indicator of
74 soil disturbance and can allow the assessment of the impact of environmental contaminates on
75 soil microbial processes (Campos et al., 2018a, 2018b; Chu et al., 2007; Elmayel et al., 2020;
76 Higuera et al., 2019a; Hinojosa et al., 2004). Additionally, the implementation of state of the
77 art high throughput sequencing approaches could provide a high-resolution analysis of the
78 ecotoxicological effects of PTEs on the composition and diversity of soil microbial
79 populations (Fernandes et al., 2018; Gallego et al., 2019; Jacquiod et al., 2018; Kisková et al.,
80 2018).

81 The province of Ciudad Real, in the center of the Iberian Peninsula, is scattered with
82 abandoned and decommissioned mines (Bravo et al., 2019; García-Lorenzo et al., 2019;
83 Higuera et al., 2017, 2012; Martín-Crespo et al., 2015; Rodríguez et al., 2009; Ruiz et al.,
84 2009). In this regard, the San Quintín mine site (Cabezarados, Ciudad Real, Spain) offers an
85 unprecedented opportunity to shed light on this topic. This area covers about 100 Ha, with a
86 complex long-lasting history of mining and mineralurgical operations (Fig. 1). Between the
87 years 1888 and 1923, three mine shafts and a rudimentary froth flotation plant operated to
88 obtain Ag-rich galena (PbS) concentrates, leaving a first legacy of dumps, very heterogeneous
89 in grain size and containing high concentrations of Pb, and Zn (García-Lorenzo et al., 2019).

90 In 1973, a second froth flotation plant aimed to recuperate the sphalerite (ZnS) as well as
91 residual Ag-rich galena was established. During this period, the froth flotation process
92 favored the conversion of the dumps in wastes dams, which accumulated the residues from
93 this reprocessing. Additionally, cinnabar (HgS) from the adjacent Hg mines of Almadén
94 (Higuera et al., 2006; Tejero et al., 2015) was accumulated in the area during the last years of
95 activity of the froth flotation plant (1988-1990). In 2000, olive-oil milling wastes (OMW)
96 were illegally applied over the soils without any remediation interest. Currently, soil and
97 waste materials from the area are used in local road pavements, resulting in a redistribution of
98 the soil. Consequently, the area is deeply polluted with Pb, Zn and associated elements
99 (García-Lorenzo et al., 2019; Higuera et al., 2017; Martín-Crespo et al., 2015; Rodríguez et
100 al., 2009; Ruiz et al., 2009).

101 The characterization of the behavior of PTEs from heavily degraded mining soils and their
102 impact on the environment is the first step for planning restoration strategies in contaminated
103 areas. We hypothesized that decades of exposure to different levels of pollutants have exerted
104 a selective pressure on microbial communities in the San Quintín mine area. Within this
105 context, this study aims to investigate the link between geochemical characteristics and
106 microbial communities in a long-term polluted mining site and assess the effects of PTEs on
107 the activity, composition and diversity of microbial communities through enzymatic analysis
108 and high throughput sequencing.

109

110 **2. Materials and Methods**

111 **2.1 Samples collection**

112 Twenty samples of circa 800 g corresponding to the first 10-15 cm and relative to five
113 materials were collected (four independent replicates per material, each replicate was a

114 composite made out of three subsamples collected and homogenized *in situ*). The five
115 different materials included: non-directly polluted soil taken from the surrounding area and
116 used as local background (reference-samples); PTEs polluted soil from mine operations (via
117 leaching and runoff) and collected within the precincts of the mining site (contaminated-
118 samples); olive-oil mill wastes residues spilled in the area (OMW); processed waste
119 accumulated in the dams (tailings); and mine wastes without any treatment (dumps). Samples
120 were immediately transported to the laboratory and sieved (<2 mm) within 24 hours. Samples
121 were divided in three aliquots and stored at -4°C (physicochemical and enzyme analysis)
122 (Peoples and Koide, 2012) or -20°C (DNA extraction) until use. Location and main
123 characteristics of sampling points are described in Fig.1 and Table 1.

124

125 **2.2 Physicochemical analysis**

126 Physicochemical analyses were conducted in the Laboratory of Soil Biogeochemistry at
127 the EIMIA-UCLM, Almadén (Spain). Electric conductivity and pH were determined using
128 ISO-10390 (ISO-10390, 2005) and ISO-7888 (ISO-11265, 1994) protocols, respectively.
129 Analyses were performed on 2 g dwt of sieved samples. Texture was determined on dried
130 samples treated with a solution of 3% (v/v) H₂O₂ for 48 h to remove organic matter. Samples
131 were then wet and sieved at 2 mm and analyzed with a Fritsch Analysette MicroTec Plus 22
132 (Fritsch, Germany). A textural classification triangle plot was used to identify the
133 granulometry of samples (Gerakis and Baer, 1999). Organic matter (OM) was determined
134 using the Walkley-Black method (Walkley and Black, 1934).

135 Elemental concentration analyses were performed from ground dried samples with
136 particle length minor to 100 µm determined by means of X-Ray Fluorescence spectroscopy,

137 using an Epsilon 1 device (Malvern Panalytical, England). Pollution Load Index (PLI) (Jorfi
138 et al., 2017) was calculated according to the equation:

$$139 \quad PI = C_n/B_n$$

$$140 \quad PLI = \sqrt{PI_1 \times PI_2 \dots PI_n}$$

141 Where PI is the single factor pollution index of each metal, C_n and B_n is the concentration of
142 metal in the sample and background, respectively (mg Kg^{-1}). Reference average values from
143 this study were used as local background level. The values of PLI allow the qualification of
144 samples as follows: $PLI < 2$: moderately to unpolluted; $2 \leq PLI < 4$: moderately polluted; $4 \leq$
145 $PLI < 6$: highly polluted; $PLI > 6$: very highly polluted.

146

147 **2.3 Enzyme analysis**

148 Enzyme activities were measured through colorimetric techniques using a Biochrom Libra
149 S60 spectrophotometer (Biochrom, United Kingdom). Dehydrogenase (DHA) activity was
150 measured according to the method described by Casida (1977), with slight modifications.
151 Acid and alkaline phosphomonoesterase (PhA 6.5 and PhA 11, respectively) activities were
152 determined following Tabatabai and Bremner (1969) method. Arylsulfatase (ARS) activity
153 was measured as described by Tabatabai and Bremner, (1970) and β -galactosidase (β -Gal)
154 activity was assayed according to Eivazi and Tabatabai (1988).

155 Functional diversity from selected enzymes was assessed with the geometric mean index
156 (GMean) (Lessard et al., 2014) calculated as follow:

$$157 \quad GMean = \left(\prod_{i=1}^n y_i \right)^{\frac{1}{n}}$$

158 Where y_i is the mean value for each enzyme activity, n is the total number of enzymes.

159

160 **2.4 Microbiological analysis**

161 DNA was extracted using the Power soil DNA isolation kit (Qiagen). The extracted DNA
162 was quantified by using fluorescent dye of a Quant-iT™ PicoGreen® dsDNA assay kit
163 (Invitrogen). The diversity of the bacterial community was determined by high throughput
164 sequencing of 16S rRNA amplicons generated in a two-step PCR. First PCR reaction was
165 performed using the universal bacterial primers U341_F-805_R with overhang adapters
166 (forward adapter: TCGTCGGCAGCGTC AGATGTGTATAAGAGACAG, reverse adapter:
167 GTCTCGTGGGCTCGGAGATGTGTATA AGAGACAG). The reaction mixture contained 2
168 ng of DNA as template, 7.5 µL of 2X Phusion High Fidelity PCR Mastermix (Thermo
169 Scientific, Waltham, MA, USA), 250 ng of T4 gene 32 protein (MP Biomedicals, Santa Ana,
170 CA, USA), 0.375 µL of each primer (10 µM) and ultrapure sterile water to a total volume of
171 15 µL. Thermal conditions were 3 min at 98°C, 25 cycles at 98° C for 30 sec, 55° C for 30 sec
172 and 72° C for 30 sec followed by a final extension of 10 min at 72°C. Duplicates of each
173 amplicon were pooled and then 6 µL aliquot was used as template in a second PCR carried
174 out with multiplexed primers containing the universal overhang adaptors and specific
175 barcodes and using a 384 Nextera XT index kit (Illumina, San Diego, CA, USA). The reaction
176 mixture was carried out in 30 µL reaction volumes containing 2.5 µL sterile water, 15 µL 2X
177 Phusion HF master mix (Thermo Scientific, Waltham, MA, USA), 250 ng of T4 gene 32
178 protein (MP Biomedicals, Santa Ana, CA, USA), 3 µL of each primer (10 µM) and 6 µL of
179 the step-one PCR product. The thermal cycling was 98°C for 3 min, followed by eight cycles
180 of 98° C for 30 sec, 55°C for 30 sec and 72° C for 30 sec, with a final extension of 72° C for
181 10 min. The size of the amplicons was verified by electrophoresis on a 2 % agarose gel.
182 Amplicons were purified (amplicon library purification, PicoGreen® quantification and
183 pooling) and sequenced (Illumina MiSeq 2 x300bp) by Microsynth (Balgach, Switzerland).
184 Amplicons were normalized (SequalPrep™ kit), purified (Pippin prep) and sequenced by

185 Microsynth (Switzerland). The sequence data was analyzed using an in-house developed
186 Phyton notebook pipeline together with different bioinformatics tools: 16S rDNA sequences
187 were assembled using the PEAR software (Zhang et al., 2014) with the default settings;
188 further quality checks were conducted using the QIIME 1 pipeline (Caporaso et al., 2010a);
189 sequences shorter than 350 bp were removed; reference-based and *de novo* chimera detection,
190 as well as clustering in OTUs were performed with the VSEARCH software (Rognes et al.,
191 2016) using appropriate reference databases (Greengenes' representative set of sequences for
192 16S rDNA) with a threshold placed at 94 % identity; representative sequences for each OTU
193 were aligned using PyNAST (Caporaso et al., 2010b); phylogenetic trees were constructed
194 using FastTree (Price et al., 2009); taxonomy was assigned using UCLUST (Edgar, 2010) and
195 the latest released Greengenes database (v.05/2013, McDonald et al., 2012) for 16S rDNA
196 sequences; sequences were deposited in the GenBank to the sequence read archive (SRA)
197 under the accession number PRJNA646888. A range of bacterial α -diversity indices
198 pertaining to richness (Chao1, observed species), evenness (Simpson reciprocal, equitability,
199 dominance, Shannon) and relatedness (PD whole tree) were calculated based on rarefied
200 tables (13,000 sequences per sample). Bray Curtis distance matrices were also computed to
201 detect changes in the composition of microbial communities. Canonical Analysis of Principal
202 Coordinates (CAP) of OTU Bray Curtis distance matrices were also performed and plotted.
203 The relative abundance of the different bacterial phyla was also determined using a
204 comparative bar chart.

205

206 **2.5 Statistical analysis**

207 All statistical analyses were performed in R (<http://www.r-project.org>). The normality of
208 the data and residuals was checked (Shapiro Wilk's with $p > 0.05$) and the homogeneity of
209 variances was verified (Levene's test with $p > 0.05$). Root square, inverse and log-

210 transformations on the data were performed when necessary. For parametric distributions,
211 ANOVA followed by Tukey's test and t-student were used to determine differences. For non-
212 parametric distributions, data was compared using Kruskal Wallis test. To detect significant
213 differences in communities' structure among sample types, Permutational multivariate
214 analysis of variance (PerMANOVA) was used on Bray Curtis dissimilarity matrix using
215 Adonis function from R package "vegan" (Oksanen et al., 2018).

216 Integration and visualization of amplicon sequencing and physicochemical and enzymatic
217 analysis data sets were performed through the R package mixOmics (Rohart et al., 2017)
218 using DIABLO (Data Integration Analysis for Biomarker discovery using a Latent component
219 method for Omics studies) in order to identify correlated key omics variables in both datasets
220 (Singh et al., 2019).

221

222 **3. Results**

223 **3.1 Physicochemical and geochemical characterization of samples**

224 Table 2 shows the physicochemical parameters of the samples of the five selected
225 materials. According to their granulometry, samples were classified with some minor
226 variability in sandy loam (reference-, contaminated-samples and dumps), silt loam (tailings)
227 and loamy sand (OMW) (Table 2, Table S1 and Fig. S1). Reference-, contaminated-samples
228 and tailings presented a slightly acidic character while OMW and dumps had lower pH
229 values. EC varied from 62.4 ± 27.8 (reference-samples) to 2282.3 ± 1786.4 mV (dumps).
230 Overall, no significant differences were found in the pH and EC ($p > 0.055$), except for the
231 dumps, which presented significant lower pH ($p = 0.00371$) and significant higher EC values
232 ($p = 0.004$) as compared to the reference-samples. OM ranged from $1.57\% \pm 0.54$ (dumps) to
233 $60.60\% \pm 19.67$ (OMW). Similar organic content was found in reference-samples, tailings

234 and dumps, while higher values were found in contaminated-samples, although not significant
235 ($p > 0.5006$), probably due to their high heterogeneity. On the contrary, significant higher
236 organic matter was found in OMW as compared to the rest of the samples ($p = 0.0000271$).

237 The concentration of PTEs for the different samples is shown in Table 3. Among the
238 different PTEs studied, those which were part of the ore (Pb, Zn and Hg) were found at very
239 high concentrations in all samples except reference-samples, with significantly higher
240 concentrations for Pb ($p < 0.0043$) and Zn ($p < 0.00577$) in dumps and OMW as compared to
241 reference-samples. Similarly, Sb, Co and Cu concentrations were significantly higher in
242 dumps and OMW than in the three other samples ($p < 0.0407$). Regarding Mn, Ni and Th,
243 similar concentrations were recorded in all samples and no significant differences were
244 observed ($p > 0.05$). A high variability was found for all the PTEs studied. The pollution index
245 for each PTE was calculated (Table S2) and yielded values higher than 100 for Hg (in all
246 samples except reference-samples), Pb (dumps and OMW) and Zn (OMW). Pollution load
247 index (PLI) allowed to classify the samples as follows: reference- and contaminated-samples
248 as highly polluted ($4 \leq \text{PLI} < 6$) and OMW, dumps and tailings (from most polluted to less
249 polluted respectively) as very highly polluted ($\text{PLI} > 6$) (Table 3).

250

251 **3.2 Enzymatic analysis**

252 Five enzymatic activities including the dehydrogenase (DHA), arylsulfatase (ARS), the
253 acid and alkaline phosphomonoesterase (PhA 6.5 and PhA 11, respectively) and β -
254 galactosidase (β -Gal) were measured in the samples collected from the five sites studied in
255 mining area (Fig. 2 and Table S3). Reference- and contaminated-samples showed similar
256 enzyme activities ($p > 0.05$). The lowest enzymatic activities (ARS, PhA 6.5, PhA 11 and β -
257 Gal) were recorded in the OMW, which were significantly different from the values recorded

258 in reference- and contaminated-samples ($p < 0.0398351$) Tailings and dumps had significantly
259 lower DHA value (reduction of about 96 %) and ARS activities than reference-samples
260 ($p < 0.0018687$). Tailings had PhA 6.5 activities significantly lower than that of the reference-
261 samples ($p = 0.0223$). The geometric mean index was significantly lower in OMW samples
262 ($p = 0.0000002$) than that of all the other samples, which were similar ($p > 0.06922$) (Fig. S2).

263

264 **3.3 Bacterial composition and diversity**

265 The composition and diversity of the bacterial community was assessed by high
266 throughput sequencing of 16S rRNA amplicons amplified from DNA extracts. In total, after
267 de-multiplexing and removal of low-quality raw sequence reads, high throughput sequencing
268 generated 1,796,830 high quality sequences with an average sequence length of 461 ± 11 bp.
269 Using a threshold at 94% nucleotide sequence identity, these sequences were grouped in
270 16,299 different OTUs.

271 A range of α -diversity indices pertaining richness (Chao1), relatedness (PD whole tree)
272 and evenness (Simpson reciprocal) were calculated using rarefied data (Fig. 3). Overall,
273 Chao1, PD whole tree and Simpson reciprocal α -diversity indices were higher in reference-
274 samples than in contaminated-samples, OMW, tailings and dumps, ranked from the highest to
275 the lowest values. Chao1 values were significantly lower in OMW, tailings and dumps
276 ($p < 0.027$) while significant differences were found for tailings and dumps in PD whole tree
277 values as compared to reference-samples values ($p < 0.019$). Regarding Simpson reciprocal, no
278 significant differences were found between reference-, contaminated-samples, OMW and
279 tailings, and only significant lower values were found in dumps ($p = 0.024$).

280 Differences in the composition of bacterial communities measured in the different samples
281 were assessed by multivariate beta-diversity analysis using Canonical Analysis of Principal

282 Coordinates (CAP) based on Bray Curtis distances matrices (Fig. 4 & Fig. S3). A good
283 reproducibility between replicates of each sample type was observed regardless the dataset
284 used as explanatory factors. CAP ordinations using a range of PTEs (Co, Cu, Hg, Mn, Ni, Sb,
285 Th, Pb, Zn) as explanatory factors, explained up to 78 % of the variance observed, of which
286 29% is explained in the first two axes (17% and 12%, respectively) (Fig. 4a). The sample type
287 was the most explanatory variable (36.96%) followed by the Hg concentration (6.41%), both
288 associated with the first axis. All the other variables (Mn, Co, Ni, Cu, Zn, Sb, Pb and Th)
289 analyzed were not significant (Table S4). Likewise, CAP ordinations using physicochemical
290 parameters and enzymatic activities as variables explained 73% of the variance observed
291 (17% and 12% respectively for the first two axes) (Fig. 4b). None of the physicochemical
292 parameters nor the enzymatic activities were significant explanatory variables of the
293 composition of the bacterial community. For all CAP ordinations, reference- and
294 contaminated-samples showed similar bacterial community composition ($p=0.734$) that was
295 differing from those of tailings and dumps which were similar ($p=0.056$). The composition of
296 the bacterial community of OMW was markedly different from that of all the other samples
297 analyzed no matter the variables considered ($p<0.036$).

298 Taxonomic analysis of the 16S rRNA amplicon sequences at phylum level showed that
299 the relative abundance of the major bacterial phyla was very similar in reference- and
300 contaminated- samples. On the contrary, it markedly differed between reference-samples and
301 dumps. All samples were dominated by bacteria belonging to Proteobacteria, Actinobacteria,
302 Chloroflexi, Acidobacteria, Gemmatimonadetes, Bacteroidetes, Firmicutes, TM7,
303 Verrucomicrobia and Cyanobacteria (Fig. 5). Proteobacteria, Actinobacteria, Chloroflexi,
304 Acidobacteria, Firmicutes, TM7 and Cyanobacteria were observed in similar abundance in all
305 the sample types ($p>0.05$). Gemmatimonadetes and Bacteroidetes were found in significantly
306 lower amount in dumps than in reference-samples ($p<0.035$). Verrucomicrobia had

307 significantly lower relative abundances in dumps, tailings and OMW than in the reference-
308 samples ($p < 0.024$).

309 To further identify the OTUs responsible of differences in the composition of the bacterial
310 community between samples, a correlation analysis was performed with OTUs represented at
311 least once in all four replicates of a given sample type (a total of 76 different OTUs) using the
312 diablo R package. Among these 76 OTUs, 57 responsible of the differences observed between
313 substrate types (Pearson's correlation > 0.4) were plotted in a heatmap (Fig 6). All
314 discriminant OTUs affiliated to 7 of the 10 most abundant bacterial phyla detected
315 (Actinobacteria, Chloroflexi, Acidobacteria, Bacteroidetes, Cyanobacteria,
316 Gemmatimonadetes and Proteobacteria).

317 Similar abundances were found in discriminant OTUs for reference- and contaminated-
318 samples, confirming the results from α - and β -diversity analyses. Microorganisms belonging
319 to the orders Actinomycetales, Sphingomonadales, Burkholderiales, Rhodospirillales,
320 Stramenopiles, RB41, Solibacterales, Ellin 329 and some microorganisms belonging to the
321 order Rhizobiales were highly represented in reference- and contaminated-samples.
322 Microorganisms belonging to the families *Acidobacteriaceae*, *Sphingobacteriaceae* and
323 *Xanthomonadaceae*, the order Solirubrobacterales, N1423WL, some Actinomycetales and
324 some Rhizobiales; and the genera *Sphingomonas* and *Segetibacter* were highly represented in
325 OMW but scarcely detected in reference- and contaminated-samples. Regarding the dumps,
326 five OTUs belonging to Acidomicrobiales, B12-WMSP1, *Acidobacteriaceae* and
327 *Enterobacteriaceae* phylotypes were found in high abundance. Finally, a high heterogeneity
328 was observed for the discriminant OTUs in tailings.

329

330 **4. Discussion**

331 Mining soils are generally rich in PTEs that are known to have ecotoxicological effects on
332 soil microorganisms by disrupting their metabolism, affecting soil enzymatic activity and
333 having consequence on the abundance, composition and diversity of soil microbial
334 communities (Zhao et al., 2020). The analysis of the PTEs concentrations in the five different
335 sample types allowed the classification of the samples as highly polluted (reference- and
336 contaminated-samples) and very highly polluted (OMW, tailings and dumps). Although the
337 potential value of OMW as fertilizer or metal immobilization amendment in polluted soils
338 was shown in previous studies (Hmid et al., 2014; Paredes et al., 1999), here the application
339 of OMW showed no remediation activity. It is noteworthy that a number of recent studies
340 have shown the negative impact of OMW on aquatic and terrestrial ecosystems (Ntougias et
341 al., 2013). Soil enzymes are mainly produced by microorganisms, which play a crucial role in
342 nutrient cycling and consequently, in soil fertility (Adetunji et al., 2017). Microbial enzymes
343 are highly sensitive to metals, and their use as standard biochemical indicators has been
344 widely proposed to evaluate the quality of polluted soils (Pająk et al., 2018; Tang et al., 2019).
345 However, several studies have showed no effect of PTEs contamination on some soil
346 enzymatic activities, such as dehydrogenase, catalase, acid phosphomonoesterase and amylase
347 (Campos et al., 2018a; Liu et al., 2020; Yang et al., 2016). This phenomenon could be due to
348 three reasons: (I) the long-term presence of metals in the soil, which might promote the
349 development of heavy metal resistant microorganisms (Ciarkowska et al., 2014); (II)
350 variations in soil physicochemical properties, which may influence the activity of soil
351 microbial communities (Xian et al., 2015), the bioavailability and speciation of the pollutants
352 (Rieuwerts et al., 1998); and (III) antagonistic effects between various PTEs (Tang et al.,
353 2019). Our results showed that enzyme activities were similar in reference- and contaminated-
354 samples. This might be explained by the high organic matter content in these two sample
355 types, which may on the one hand, provide nutrient source for the development of

356 microorganisms and the production of enzymes and on the other hand, reduce the
357 bioavailability of metals and metalloids, limiting their toxicity towards microbial enzymatic
358 activities (De Santiago-Martín et al., 2013; Lair et al., 2007). It is important to note, that
359 organic matter can sometimes increase the mobility and bioavailability of pollutants (Lindsay,
360 1991; Meunier et al., 2011; Wang and Mulligan, 2006). Consequently, further research is
361 needed to better understand the processes governing the complex interactions between organic
362 matter and pollutants. Additionally, all enzymatic activities were very low in OMW as
363 confirmed by the geometric mean index. The important amount of phenols with powerful
364 antimicrobial properties and high content of PTEs, might explained the low enzymatic
365 activities recorded in OMW samples (Siles et al., 2014). DHA activities measured in tailings
366 and dumps were significantly lower than those of reference-samples. Keeping in mind the fact
367 that the inhibition of DHA activity in response to PTEs such as Pb, Cu, Cd, Zn has been
368 previously described (De Santiago-Martín et al., 2013; Pająk et al., 2018; Parelho et al., 2016;
369 Tang et al., 2019), one could hypothesize that the high concentration of PTEs in tailings and
370 dumps would explain the observed low enzyme activities. Similar results were found for ARS
371 activity, that is a key player in the sulfur cycle and known to be inhibited by PTEs (De
372 Santiago-Martín et al., 2013; Gülser and Erdoğan, 2008; Stefanowicz et al., 2020). No-matter
373 the sample type, acid PhA activities were higher than those of alkaline PhA probably due to
374 the acidic nature of all the samples (Adetunji et al., 2017). Acid PhA activity was lower in
375 tailings than in the reference-samples which is in accordance with known sensitivities of these
376 enzymes to heavy metals (Angelovičová et al., 2014; De Santiago-Martín et al., 2013; Gao et
377 al., 2010). Despite the fact that β -Gal has been reported as a sensitive bioindicator of metal
378 contamination of soils (De Santiago-Martín et al., 2013; Martínez-Iñigo et al., 2009), all
379 samples showed similar β -Gal activities except in dumps where slightly lower activities were
380 recorded.

381 As observed with microbial enzymatic activities, the exposure to PTEs can also affect key
382 microbial processes, which reflects on changes in microbial composition and diversity. On the
383 one hand, PTEs such as Pb, Zn or Hg can exert toxic effects compromising the survival and
384 growth of specific microbial guilds (Abdu et al., 2017; Doelman and Haanstra, 1984). On the
385 other hand, PTEs can promote the emergence of specific resistant microorganisms able to
386 cope with the ecotoxicity of these contaminants (Gillard et al., 2019; Martinez et al., 2006;
387 Sobolev and Begonia, 2008). Here, the lowest values of all the α -diversity indices (Chao1, PD
388 whole tree and Simpson reciprocal) were recorded in dumps, probably as consequence of the
389 very high concentrations of Pb, Zn, Sb, Co and Cu present in these samples and low organic
390 matter. Indeed, the reduction in soil bacterial α -diversity as consequence of Pb, Zn, Sb, Co or
391 Cu toxic effects has previously been reported (Luo et al., 2018; Nunes et al., 2016;
392 Stefanowicz et al., 2008; Xie et al., 2016). Similarly, Chao1 and PD whole tree values were
393 lower in tailings than in the reference-samples. It is noteworthy that PTEs concentrations in
394 tailings were not significantly different from that of reference-samples, suggesting that
395 differences observed in α -diversity may result from other physicochemical parameters
396 governing the bioavailability of PTEs. Chao1 diversity indices in OMW samples were
397 significantly lower than that of reference-samples. As observed for the enzymatic activities,
398 the high content of organic matter rich in phenols in OMW could have ecotoxicological
399 effects on survival and growth of phenol-sensitive microorganisms and therefore decreased
400 the microbial diversity. In this regard, changes in the structure of ammonium oxidizers
401 bacteria and Actinobacteria have been previously reported (Karpouzas et al., 2010; Mekki et
402 al., 2006). All alpha diversity indices were similar in both reference- and contaminated-
403 samples.

404 Multivariate β -diversity analysis based on Bray Curtis distance matrix confirmed the
405 results observed from α -diversity: reference- and contaminated-samples clustered together no

406 matter the explanatory factors (PTEs, enzymatic activities or texture). These two samples
407 were clearly separated from tailings and dumps. OMW was discriminated from all others
408 whatever the factors considered. Besides the sample type, which explained 37% of the
409 variance observed, concentration of Hg was the only significant explanatory factor (6% of the
410 variance). In line with our work, the influence of soil properties and heavy metals on the
411 structure of bacterial communities has been previously showed in contaminated soils (Li et
412 al., 2017; Liu et al., 2018; Nunes et al., 2016; Tipayno et al., 2018)

413 Phylogenetic analysis revealed that the majority of bacteria from all samples affiliated to
414 the Proteobacteria, Actinobacteria, Chloroflexi, Acidobacteria, Gemmatimonadetes,
415 Bacteroidetes, Firmicutes, TM7, Verrucomicrobia and Cyanobacteria phylotypes, in line with
416 previous studies performed in a range of PTEs polluted soils (Chen et al., 2018; Guo et al.,
417 2017; Pacwa-Płociniczak et al., 2018; Tipayno et al., 2018; Zhen et al., 2019). Overall, their
418 relative abundances were similar in all samples, with the exception of the bacteria belonging
419 to Gemmatimonadetes, Bacteroidetes and Verrucomicrobia, whose relative abundance were
420 significantly lower in dumps as compared to reference-samples. A significant reduction in the
421 relative abundance of Verrucomicrobia was also observed in tailings and OMW. The
422 significantly higher concentrations of PTEs found in in tailings and OMW may explain the
423 reduction of these phylotypes. This assumption is in line with the reduction of
424 Verrucomicrobia previously reported in a study performed on an agricultural paddy soil
425 polluted with cadmium (Luo et al., 2019) and Yellow river sediments heavily polluted with
426 cadmium, arsenic, lead and mercury (Chen et al., 2018). Similarly, the relative abundance of
427 members of several phyla including Verrucomicrobia and Bacteroidetes was found to
428 significantly decrease in a long-term Cu polluted site (Berg et al., 2012). In this regard, our
429 findings are consistent with a recent study that proposes that changes in the response of some
430 bacterial groups (the Verrucomicrobia/Chlamydiae ratio) can be used as bioindicators of

431 heavy metal pollution (Schneider et al., 2017). Contrary to our results, studies from a metal
432 polluted forest soil showed an increase in the Gemmatimonadetes (Azarbad et al., 2015) but a
433 negative correlation between the relative abundance of Verrucomicrobia and pollution levels
434 as observed in our study.

435 Further analysis at lower taxonomical level led to the identification of 57 OTUs
436 responsible of α - and β - diversity changes observed between different sample types.
437 Discriminant OTUs were found in similar abundances in reference- and contaminated-
438 samples. Interestingly, OTUs barely found in reference- and contaminated-samples were
439 significantly more abundant in OMW, suggesting that these OTUs may be able to grow in a
440 PTEs contaminated environment rich in organic matter. OTUs related to the B12-WMSP1
441 group described by Costello and Schmidt (Costello and Schmidt, 2006) in fumarolic soils
442 (Costello et al., 2009; Schmidt et al., 2018; Tebo et al., 2015) were found in higher amount in
443 dumps than in the reference-samples.

444

445 **5. Conclusions**

446 The importance of soil health and quality in ecosystem services underlines the need to
447 understand the processes governing the behavior and effects of PTEs on soil microorganisms
448 and supported functions. This assumption is especially relevant in highly polluted areas such
449 as mining areas. Here, we evaluated the degree of contamination in a long-term mining site
450 and assessed the ecotoxicological effects of PTEs on bacterial communities. We observed that
451 low PTEs concentrations can have a strong impact on the diversity and composition of
452 microbial communities while physicochemical parameters such as organic content can either
453 help counteract or enhance their negative effects. Understanding the relationship between
454 physicochemical parameters and diversity and composition of microorganisms can contribute

455 to the environmental risk assessment of long-term polluted sites and help create strategies to
456 restore contaminated sites and mitigate the pollution dissemination.

457 **Funding Sources**

458 Spanish Ministry of Economy and Competitiveness (MINECO) funded this work, Project
459 CGL2015-67644-R.

460

461 **ACKNOWLEDGEMENT**

462 The authors thank Sofía Rivera, Efrén García-Ordiales, Sandra Viso and Eva M^a García-
463 Noguero for their contribution in collecting samples and edaphological sample
464 characterization.

465 **ABBREVIATIONS**

466 ARS, Arylsulfatase enzyme activity

467 DHA, Dehydrogenase enzyme activity

468 EC, Electric conductivity

469 OMW, Olive-oil mill waste

470 OM, organic matter

471 PhA, Phosphomonoesterase enzyme activity

472 PTEs, Potentially toxic elements

473 β -Gal, β -galactosidase enzyme activity

474

475 **REFERENCES**

- 476 Abdu, N., Abdullahi, A.A., Abdulkadir, A., 2017. Heavy metals and soil microbes. *Environ.*
477 *Chem. Lett.* 15, 65–84. <https://doi.org/10.1007/s10311-016-0587-x>
- 478 Adetunji, A.T., Lewu, F.B., Mulidzi, R., Ncube, B., 2017. The biological activities of β -
479 glucosidase, phosphatase and urease as soil quality indicators: A review. *J. Soil Sci.*
480 *Plant Nutr.* 17, 794–807. <https://doi.org/10.4067/S0718-95162017000300018>
- 481 Alvarenga, P., Clemente, R., Garbisu, C., Becerril, J.M., 2018. Indicators for Monitoring
482 Mine Site Rehabilitation, in: *Bio-Geotechnologies for Mine Site Rehabilitation*. Elsevier,
483 pp. 49–66. <https://doi.org/10.1016/B978-0-12-812986-9.00003-8>
- 484 Angelovičová, L., Lodenius, M., Tulisalo, E., Fazekašová, D., 2014. Effect of heavy metals
485 on soil enzyme activity at different field conditions in middle spis mining area
486 (Slovakia). *Bull. Environ. Contam. Toxicol.* 93, 670–675.
487 <https://doi.org/10.1007/s00128-014-1397-0>
- 488 Ayangbenro, A.S., Olanrewaju, O.S., Babalola, O.O., 2018. Sulfate-reducing bacteria as an
489 effective tool for sustainable acid mine bioremediation. *Front. Microbiol.* 9, 1–10.
490 <https://doi.org/10.3389/fmicb.2018.01986>
- 491 Azarbad, H., Niklińska, M., Laskowski, R., van Straalen, N.M., van Gestel, C.A.M., Zhou, J.,
492 He, Z., Wen, C., Röling, W.F.M., 2015. Microbial community composition and
493 functions are resilient to metal pollution along two forest soil gradients. *FEMS*
494 *Microbiol. Ecol.* 91, 1–11. <https://doi.org/10.1093/femsec/fiu003>
- 495 Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L., Lilley, A.K., 2005. The contribution
496 of species richness and composition to bacterial services. *Nature* 436, 1157–1160.
497 <https://doi.org/10.1038/nature03891>

498 Berg, J., Brandt, K.K., Al-Soud, W.A., Holm, P.E., Hansen, L.H., Sørensen, S.J., Nybroe, O.,
499 2012. Selection for Cu-tolerant bacterial communities with altered composition, but
500 unaltered richness, via long-term cu exposure. *Appl. Environ. Microbiol.* 78, 7438–7446.
501 <https://doi.org/10.1128/AEM.01071-12>

502 Bravo, S., García-Ordiales, E., García-Navarro, F.J., Amorós, J.Á., Pérez-de-los-Reyes, C.,
503 Jiménez-Ballesta, R., Esbrí, J.M., García-Noguero, E.M., Higuera, P., 2019.
504 Geochemical distribution of major and trace elements in agricultural soils of Castilla-La
505 Mancha (central Spain): finding criteria for baselines and delimiting regional anomalies.
506 *Environ. Sci. Pollut. Res.* 26, 3100–3114. <https://doi.org/10.1007/s11356-017-0010-6>

507 Campos, J.A., Esbrí, J.M., Madrid, M.M., Naharro, R., Peco, J., García-Noguero, E.M.,
508 Amorós, J.A., Moreno, M.M., Higuera, P., 2018a. Does mercury presence in soils
509 promote their microbial activity? The Almadenejos case (Almadén mercury mining
510 district, Spain). *Chemosphere* 201, 799–806.
511 <https://doi.org/10.1016/j.chemosphere.2018.02.163>

512 Campos, J.A., Peco, J.D., De Toro, J.A., Moreno, C., Amorós, J.A., Moreno, M.M., García-
513 Noguero, E.M., Higuera, P., 2018b. Approach to the potential usage of two wood ashes
514 waste as soil amendments on the basis of the dehydrogenase activity and soil oxygen
515 consumption. *J. Soils Sediments* 18, 2148–2156. [https://doi.org/10.1007/s11368-017-](https://doi.org/10.1007/s11368-017-1840-z)
516 [1840-z](https://doi.org/10.1007/s11368-017-1840-z)

517 Caporaso, J. Gregory, Bittinger, K., Bushman, F.D., Desantis, T.Z., Andersen, G.L., Knight,
518 R., 2010. PyNAST: A flexible tool for aligning sequences to a template alignment.
519 *Bioinformatics* 26, 266–267. <https://doi.org/10.1093/bioinformatics/btp636>

520 Caporaso, J Gregory, Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,
521 E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G. a, Kelley, S.T.,

522 Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C. a, Mcdonald, D., Muegge, B.D.,
523 Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W. a, Widmann, J.,
524 Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-
525 throughput community sequencing data Intensity normalization improves color calling in
526 SOLiD sequencing. *Nat. Publ. Gr.* 7, 335–336. <https://doi.org/10.1038/nmeth0510-335>

527 Casida, L.E., 1977. Microbial metabolic activity in soil as measured by dehydrogenase
528 determinations. *Appl. Environ. Microbiol.* 34, 630–636.
529 <https://doi.org/10.1128/aem.34.6.630-636.1977>

530 Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., Bakken, L.R., Baylis, M., Behrenfeld,
531 M.J., Boetius, A., Boyd, P.W., Classen, A.T., Crowther, T.W., Danovaro, R., Foreman,
532 C.M., Huisman, J., Hutchins, D.A., Jansson, J.K., Karl, D.M., Koskella, B., Mark Welch,
533 D.B., Martiny, J.B.H., Moran, M.A., Orphan, V.J., Reay, D.S., Remais, J. V., Rich, V.I.,
534 Singh, B.K., Stein, L.Y., Stewart, F.J., Sullivan, M.B., van Oppen, M.J.H., Weaver, S.C.,
535 Webb, E.A., Webster, N.S., 2019. Scientists’ warning to humanity: microorganisms and
536 climate change. *Nat. Rev. Microbiol.* 17, 569–586. [https://doi.org/10.1038/s41579-019-](https://doi.org/10.1038/s41579-019-0222-5)
537 [0222-5](https://doi.org/10.1038/s41579-019-0222-5)

538 Chen, Y., Jiang, Y., Huang, H., Mou, L., Ru, J., Zhao, J., Xiao, S., 2018. Long-term and high-
539 concentration heavy-metal contamination strongly influences the microbiome and
540 functional genes in Yellow River sediments. *Sci. Total Environ.* 637–638, 1400–1412.
541 <https://doi.org/10.1016/j.scitotenv.2018.05.109>

542 Chu, H., Lin, X., Fujii, T., Morimoto, S., Yagi, K., Hu, J., Zhang, J., 2007. Soil microbial
543 biomass, dehydrogenase activity, bacterial community structure in response to long-term
544 fertilizer management. *Soil Biol. Biochem.* 39, 2971–2976.
545 <https://doi.org/10.1016/j.soilbio.2007.05.031>

546 Ciarkowska, K., Sołek-Podwika, K., Wieczorek, J., 2014. Enzyme activity as an indicator of
547 soil-rehabilitation processes at a zinc and lead ore mining and processing area. *J.*
548 *Environ. Manage.* 132, 250–256. <https://doi.org/10.1016/j.jenvman.2013.10.022>

549 Costello, E.K., Halloy, S.R.P., Reed, S.C., Sowell, P., Schmidt, S.K., 2009. Fumarole-
550 supported islands of biodiversity within a hyperarid, high-elevation landscape on
551 socompa volcano, puna de atacama, andes. *Appl. Environ. Microbiol.* 75, 735–747.
552 <https://doi.org/10.1128/AEM.01469-08>

553 Costello, E.K., Schmidt, S.K., 2006. Microbial diversity in alpine tundra wet meadow soil:
554 Novel Chloroflexi from a cold, water-saturated environment. *Environ. Microbiol.* 8,
555 1471–1486. <https://doi.org/10.1111/j.1462-2920.2006.01041.x>

556 De Santiago-Martín, A., Cheviron, N., Quintana, J.R., González, C., Lafuente, A.L., Mougín,
557 C., 2013. Metal contamination disturbs biochemical and microbial properties of
558 calcareous agricultural soils of the Mediterranean area. *Arch. Environ. Contam. Toxicol.*
559 64, 388–398. <https://doi.org/10.1007/s00244-012-9842-8>

560 Delgado-Baquerizo, M., Eldridge, D.J., Ochoa, V., Gozalo, B., Singh, B.K., Maestre, F.T.,
561 2017. Soil microbial communities drive the resistance of ecosystem multifunctionality to
562 global change in drylands across the globe. *Ecol. Lett.* 20, 1295–1305.
563 <https://doi.org/10.1111/ele.12826>

564 Delgado-Baquerizo, M., Giaramida, L., Reich, P.B., Khachane, A.N., Hamonts, K., Edwards,
565 C., Lawton, L.A., Singh, B.K., 2016a. Lack of functional redundancy in the relationship
566 between microbial diversity and ecosystem functioning. *J. Ecol.* 104, 936–946.
567 <https://doi.org/10.1111/1365-2745.12585>

568 Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D.,
569 Berdugo, M., Campbell, C.D., Singh, B.K., 2016b. Microbial diversity drives

570 multifunctionality in terrestrial ecosystems. *Nat. Commun.* 7, 1–8.
571 <https://doi.org/10.1038/ncomms10541>

572 Delgado-Baquerizo, M., Reich, P.B., Trivedi, C., Eldridge, D.J., Abades, S., Alfaro, F.D.,
573 Bastida, F., Berhe, A.A., Cutler, N.A., Gallardo, A., García-velázquez, L., Hart, S.C.,
574 Hayes, P.E., He, J., Hseu, Z., Hu, H., Kirchmair, M., Neuhauser, S., Perez, C.A., Reed,
575 S.C., Santos, F., Sullivan, B.W., Trivedi, P., Wang, J.-T., Weber-Grullon, L., Williams,
576 M.A., Singh, B.K., 2020. Multiple elements of soil biodiversity drive ecosystem
577 functions across biomes. *Nat. Ecol. Evol.* 4, 210–220. [https://doi.org/10.1038/s41559-](https://doi.org/10.1038/s41559-019-1084-y)
578 [019-1084-y](https://doi.org/10.1038/s41559-019-1084-y)

579 Doelman, P., Haanstra, L., 1984. Short-term and long-term effects of cadmium, chromium,
580 copper, nickel, lead and zinc on soil microbial respiration in relation to abiotic soil
581 factors. *Plant Soil* 79, 317–327. <https://doi.org/10.1007/BF02184325>

582 Domeignoz-Horta, L.A., Philippot, L., Peyrard, C., Bru, D., Breuil, M.-C., Bizouard, F.,
583 Justes, E., Mary, B., Léonard, J., Spor, A., 2018. Peaks of in situ N₂O emissions are
584 influenced by N₂O-producing and reducing microbial communities across arable soils.
585 *Glob. Chang. Biol.* 24, 360–370. <https://doi.org/10.1111/gcb.13853>

586 Domeignoz-Horta, L.A., Pold, G., Liu, X.-J.A., Frey, S.D., Melillo, J.M., DeAngelis, K.M.,
587 2020. Microbial diversity drives carbon use efficiency in a model soil. *Nat. Commun.* 11,
588 3684. <https://doi.org/10.1038/s41467-020-17502-z>

589 Domeignoz-Horta, L.A., Spor, A., Bru, D., Breuil, M.C., Bizouard, F., Léonard, J., Philippot,
590 L., 2015. The diversity of the N₂O reducers matters for the N₂O:N₂ denitrification end-
591 product ratio across an annual and a perennial cropping system. *Front. Microbiol.* 6.
592 <https://doi.org/10.3389/fmicb.2015.00971>

593 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST.

594 Bioinformatics 26, 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>

595 Eivazi, F., Tabatabai, M.A., 1988. Glucosidases and galactosidases in soils. *Soil Biol.*

596 *Biochem.* 20, 601–606. [https://doi.org/10.1016/0038-0717\(88\)90141-1](https://doi.org/10.1016/0038-0717(88)90141-1)

597 Elmayel, I., Elouear, Z., Jalel, B., Farieri, A., Roqueñ, N., Cienfuegos, P., 2020. Evolution of

598 the Speciation and Mobility of Pb , Zn and Cd in Relation to Transport Processes in a

599 Mining Environment. *Int. J. Environ. Res. Public Health* 17, 1–16.

600 Esbrí, J., Higuera, P., Martínez-Coronado, A., Naharro, R., 2020. 4D dispersion of total

601 gaseous mercury derived from a mining source: identification of criteria to assess risks

602 related with high concentrations of atmospheric mercury. *Atmos. Chem. Phys.* 1–21.

603 <https://doi.org/10.5194/acp-2019-1107>

604 Favas, P.J. de C., Martino, L.E., Prasad, M.N.V., 2018. Abandoned Mine Land Reclamation-

605 Challenges and Opportunities (Holistic Approach), in: *Bio-Geotechnologies for Mine*

606 *Site Rehabilitation*. Elsevier, pp. 3–31. [https://doi.org/10.1016/B978-0-12-812986-](https://doi.org/10.1016/B978-0-12-812986-9.00001-4)

607 [9.00001-4](https://doi.org/10.1016/B978-0-12-812986-9.00001-4)

608 Fernandes, C.C., Kishi, L.T., Lopes, E.M., Omori, W.P., Souza, J.A.M. de, Alves, L.M.C.,

609 Lemos, E.G. de M., 2018. Bacterial communities in mining soils and surrounding areas

610 under regeneration process in a former ore mine. *Brazilian J. Microbiol.* 49, 489–502.

611 <https://doi.org/10.1016/j.bjm.2017.12.006>

612 Fernández-Martínez, R., Esbrí, J.M., Higuera, P., Rucandio, I., 2019. Comparison of mercury

613 distribution and mobility in soils affected by anthropogenic pollution around chloralkali

614 plants and ancient mining sites. *Sci. Total Environ.* 671, 1066–1076.

615 <https://doi.org/10.1016/j.scitotenv.2019.03.348>

616 Gallego, S., Devers-Lamrani, M., Rousidou, K., Karpouzas, D.G., Martin-Laurent, F., 2019.

617 Assessment of the effects of oxamyl on the bacterial community of an agricultural soil
618 exhibiting enhanced biodegradation. *Sci. Total Environ.* 651, 1189–1198.
619 <https://doi.org/10.1016/j.scitotenv.2018.09.255>

620 Gao, Y., Zhou, P., Mao, L., Zhi, Y. er, Shi, W. jun, 2010. Assessment of effects of heavy
621 metals combined pollution on soil enzyme activities and microbial community structure:
622 Modified ecological dose-response model and PCR-RAPD. *Environ. Earth Sci.* 60, 603–
623 612. <https://doi.org/10.1007/s12665-009-0200-8>

624 García-Lorenzo, M.L., Crespo-Feo, E., Esbrí, J.M., Higuera, P., Grau, P., Crespo, I.,
625 Sánchez-Donoso, R., 2019. Assessment of Potentially Toxic Elements in Technosols by
626 Tailings Derived from Pb–Zn–Ag Mining Activities at San Quintín (Ciudad Real,
627 Spain): Some Insights into the Importance of Integral Studies to Evaluate Metal
628 Contamination Pollution Hazards. *Minerals* 9, 1–20. <https://doi.org/10.3390/min9060346>

629 Gerakis, A., Baer, B., 1999. A Computer Program for Soil Textural Classification. *Soil Sci.*
630 *Soc. Am. J.* 63, 807–808. <https://doi.org/10.2136/sssaj1999.634807x>

631 Gillard, B., Chatzievangelou, D., Thomsen, L., Ullrich, M.S., 2019. Heavy-metal-resistant
632 microorganisms in deep-sea sediments disturbed by mining activity: An application
633 toward the development of experimental in vitro systems. *Front. Mar. Sci.* 6, 1–12.
634 <https://doi.org/10.3389/fmars.2019.00462>

635 Giller, K.E., Witter, E., Mcgrath, S.P., 1998. Toxicity of heavy metals to microorganisms and
636 microbial processes in agricultural soils: A review. *Soil Biol. Biochem.* 30, 1389–1414.
637 [https://doi.org/10.1016/S0038-0717\(97\)00270-8](https://doi.org/10.1016/S0038-0717(97)00270-8)

638 Griffiths, B.S., Philippot, L., 2013. Insights into the resistance and resilience of the soil
639 microbial community. *FEMS Microbiol. Rev.* 37, 112–129.
640 <https://doi.org/10.1111/j.1574-6976.2012.00343.x>

641 Gülser, F., Erdoğan, E., 2008. The effects of heavy metal pollution on enzyme activities and
642 basal soil respiration of roadside soils. *Environ. Monit. Assess.* 145, 127–133.
643 <https://doi.org/10.1007/s10661-007-0022-7>

644 Guo, H., Nasir, M., Lv, J., Dai, Y., Gao, J., 2017. Understanding the variation of microbial
645 community in heavy metals contaminated soil using high throughput sequencing.
646 *Ecotoxicol. Environ. Saf.* 144, 300–306. <https://doi.org/10.1016/j.ecoenv.2017.06.048>

647 Higuera, P., Campos, J.A., Esbrí, J.M., García-Noguero, E.M., Elmayel, I., 2019a.
648 *Biogeochemical Mapping: A New Tool to Assess the Soil Quality and Health*. pp. 3–5.
649 https://doi.org/10.1007/978-3-030-01575-6_1

650 Higuera, P., Esbrí, J.M., García-Ordiales, E., González-Corrochano, B., López-Berdonces,
651 M.A., García-Noguero, E.M., Alonso-Azcárate, J., Martínez-Coronado, A., 2017.
652 Potentially harmful elements in soils and holm-oak trees (*Quercus ilex* L.) growing in
653 mining sites at the Valle de Alcudia Pb-Zn district (Spain)—Some clues on plant metal
654 uptake. *J. Geochemical Explor.* 182, 166–179.
655 <https://doi.org/10.1016/j.gexplo.2016.07.017>

656 Higuera, P., Oyarzun, R., Iraizoz, J.M., Lorenzo, S., Esbrí, J.M., Martínez-Coronado, A.,
657 2012. Low-cost geochemical surveys for environmental studies in developing countries:
658 Testing a field portable XRF instrument under quasi-realistic conditions. *J. Geochemical*
659 *Explor.* 113, 3–12. <https://doi.org/10.1016/j.gexplo.2011.02.005>

660 Higuera, P., Oyarzun, R., Lillo, J., Sánchez-Hernández, J.C., Molina, J.A., Esbrí, J.M.,
661 Lorenzo, S., 2006. The Almadén district (Spain): Anatomy of one of the world's largest
662 Hg-contaminated sites. *Sci. Total Environ.* 356, 112–124.
663 <https://doi.org/10.1016/j.scitotenv.2005.04.042>

664 Higuera, P., Sáez-Martínez, F.J., Lefebvre, G., Moilleron, R., 2019b. Contaminated sites,

665 waste management, and green chemistry: new challenges from monitoring to
666 remediation. *Environ. Sci. Pollut. Res.* 26, 3095–3099. [https://doi.org/10.1007/s11356-](https://doi.org/10.1007/s11356-018-3564-z)
667 018-3564-z

668 Hinojosa, M.B., Carreira, J.A., García-Ruíz, R., Dick, R.P., 2004. Soil moisture pre-treatment
669 effects on enzyme activities as indicators of heavy metal-contaminated and reclaimed
670 soils, in: *Soil Biology and Biochemistry*. Pergamon, pp. 1559–1568.
671 <https://doi.org/10.1016/j.soilbio.2004.07.003>

672 Hmid, A., Al Chami, Z., Sillen, W., De Vocht, A., Vangronsveld, J., 2014. Olive mill waste
673 biochar: a promising soil amendment for metal immobilization in contaminated soils.
674 *Environ. Sci. Pollut. Res.* 22, 1444–1456. <https://doi.org/10.1007/s11356-014-3467-6>

675 Hobman, J.L., Yamamoto, K., Oshima, T., 2007. Transcriptomic Responses of Bacterial Cells
676 to Sublethal Metal Ion Stress, in: *Molecular Microbiology of Heavy Metals*. Springer
677 Berlin Heidelberg, pp. 73–115. https://doi.org/10.1007/7171_2006_074

678 ISO-10390, 2005. Soil quality — Determination of pH, International organization for
679 Standardization (ISO).

680 ISO-11265, 1994. Soil quality — Determination of the specific electrical conductivity,
681 International organization for Standardization (ISO).

682 Jacquiod, S., Nunes, I., Brejnrod, A., Hansen, M.A., Holm, P.E., Johansen, A., Brandt, K.K.,
683 Priemé, A., Sørensen, S.J., 2018. Long-term soil metal exposure impaired temporal
684 variation in microbial metatranscriptomes and enriched active phages. *Microbiome* 6, 1–
685 14. <https://doi.org/10.1186/s40168-018-0606-1>

686 Jorfi, S., Maleki, R., Jaafarzadeh, N., Ahmadi, M., 2017. Pollution load index for heavy
687 metals in Mian-Ab plain soil, Khuzestan, Iran. *Data Br.* 15, 584–590.

688 <https://doi.org/10.1016/j.dib.2017.10.017>

689 Jung, M.C., Thornton, I., 1996. Heavy metal contamination of soils and plants in the vicinity
690 of a lead-zinc mine, Korea, in: *Applied Geochemistry*. Elsevier Ltd, pp. 53–59.
691 [https://doi.org/10.1016/0883-2927\(95\)00075-5](https://doi.org/10.1016/0883-2927(95)00075-5)

692 Karpouzas, D.G., Ntougias, S., Iskidou, E., Rousidou, C., Papadopoulou, K.K., Zervakis, G.I.,
693 Ehaliotis, C., 2010. Olive mill wastewater affects the structure of soil bacterial
694 communities. *Appl. Soil Ecol.* 45, 101–111. <https://doi.org/10.1016/j.apsoil.2010.03.002>

695 Kisková, J., Perháčová, Z., Vlčko, L., Sedláková, J., Kvasnová, S., Pristaš, P., 2018. The
696 Bacterial Population of Neutral Mine Drainage Water of Elizabeth's Shaft (Slovinky,
697 Slovakia). *Curr. Microbiol.* 75, 988–996. <https://doi.org/10.1007/s00284-018-1472-6>

698 Krzaklewski, W., Pietrzykowski, M., 2002. Selected physico-chemical properties of zinc and
699 lead ore tailings and their biological stabilisation. *Water. Air. Soil Pollut.* 141, 125–141.
700 <https://doi.org/10.1023/A:1021302725532>

701 Lair, G.J., Gerzabek, M.H., Haberhauer, G., 2007. Sorption of heavy metals on organic and
702 inorganic soil constituents. *Environ. Chem. Lett.* 5, 23–27.
703 <https://doi.org/10.1007/s10311-006-0059-9>

704 Lessard, I., Sauv e, S., Desch enes, L., 2014. Toxicity response of a new enzyme-based
705 functional diversity methodology for Zn-contaminated field-collected soils. *Soil Biol.*
706 *Biochem.* 71, 87–94. <https://doi.org/10.1016/J.SOILBIO.2014.01.002>

707 Li, X., Meng, D., Li, J., Yin, H., Liu, H., Liu, X., Cheng, C., Xiao, Y., Liu, Z., Yan, M., 2017.
708 Response of soil microbial communities and microbial interactions to long-term heavy
709 metal contamination. *Environ. Pollut.* 231, 908–917.
710 <https://doi.org/10.1016/j.envpol.2017.08.057>

711 Lindsay, W.L., 1991. Iron oxide solubilization by organic matter and its effect on iron
712 availability. *Plant Soil* 130, 27–34. <https://doi.org/10.1007/BF00011852>

713 Liu, K., Li, C., Tang, S., Shang, G., Yu, F., Li, Y., 2020. Heavy metal concentration, potential
714 ecological risk assessment and enzyme activity in soils affected by a lead-zinc tailing
715 spill in Guangxi, China. *Chemosphere* 251, 1–12.
716 <https://doi.org/10.1016/j.chemosphere.2020.126415>

717 Liu, X., Ma, A., Zhuang, G., Zhuang, X., 2018. Diversity of microbial communities
718 potentially involved in mercury methylation in rice paddies surrounding typical mercury
719 mining areas in China. *Microbiologyopen* 7, e00577. <https://doi.org/10.1002/mbo3.577>

720 Luo, L.Y., Xie, L.L., Jin, D.C., Mi, B.B., Wang, D.H., Li, X.F., Dai, X.Z., Zou, X.X., Zhang,
721 Z., Ma, Y.Q., Liu, F., 2019. Bacterial community response to cadmium contamination of
722 agricultural paddy soil. *Appl. Soil Ecol.* 139, 100–106.
723 <https://doi.org/10.1016/j.apsoil.2019.03.022>

724 Luo, Z., Ma, J., Chen, F., Li, X., Zhang, S., 2018. Effects of Pb smelting on the soil bacterial
725 community near a secondary lead plant. *Int. J. Environ. Res. Public Health* 15, 1–16.
726 <https://doi.org/10.3390/ijerph15051030>

727 Martín-Crespo, T., Gómez-Ortiz, D., Martín-Velázquez, S., Esbrí, J.M., de Ignacio-San José,
728 C., Sánchez-García, M.J., Montoya-Montes, I., Martín-González, F., 2015. Abandoned
729 mine tailings in cultural itineraries: Don Quixote Route (Spain). *Eng. Geol.* 197, 82–93.
730 <https://doi.org/10.1016/j.enggeo.2015.08.008>

731 Martín Duque, J.F., Zapico, I., Oyarzun, R., López García, J.A., Cubas, P., 2015. A
732 descriptive and quantitative approach regarding erosion and development of landforms
733 on abandoned mine tailings: New insights and environmental implications from SE
734 Spain. *Geomorphology* 239, 1–16. <https://doi.org/10.1016/j.geomorph.2015.02.035>

735 Martínez-Iñigo, M.J., Pérez-Sanz, A., Ortiz, I., Alonso, J., Alarcón, R., García, P., Lobo,
736 M.C., 2009. Bulk soil and rhizosphere bacterial community PCR-DGGE profiles and β -
737 galactosidase activity as indicators of biological quality in soils contaminated by heavy
738 metals and cultivated with *Silene vulgaris* (Moench) Garcke. *Chemosphere* 75, 1376–
739 1381. <https://doi.org/10.1016/j.chemosphere.2009.03.014>

740 Martinez, R.J., Wang, Y., Raimondo, M.A., Coombs, J.M., Barkay, T., Sobecky, P.A., 2006.
741 Horizontal gene transfer of PIB-type ATPases among bacteria isolated from
742 radionuclide- and metal-contaminated subsurface soils. *Appl. Environ. Microbiol.* 72,
743 3111–3118. <https://doi.org/10.1128/AEM.72.5.3111-3118.2006>

744 Mekki, A., Dhouib, A., Sayadi, S., 2006. Changes in microbial and soil properties following
745 amendment with treated and untreated olive mill wastewater. *Microbiol. Res.* 161, 93–
746 101. <https://doi.org/10.1016/j.micres.2005.06.001>

747 Mendez, M.O., Neilson, J.W., Maier, R.M., 2008. Characterization of a bacterial community
748 in an abandoned semiarid lead-zinc mine tailing site. *Appl. Environ. Microbiol.* 74,
749 3899–3907. <https://doi.org/10.1128/AEM.02883-07>

750 Meunier, L., Koch, I., Reimer, K.J., 2011. Effects of organic matter and ageing on the
751 bioaccessibility of arsenic. *Environ. Pollut.* 159, 2530–2536.
752 <https://doi.org/10.1016/j.envpol.2011.06.018>

753 Nagajyoti, P.C., Lee, K.D., Sreekanth, T.V.M., 2010. Heavy metals, occurrence and toxicity
754 for plants: A review. *Environ. Chem. Lett.* 8, 199–216. [https://doi.org/10.1007/s10311-](https://doi.org/10.1007/s10311-010-0297-8)
755 [010-0297-8](https://doi.org/10.1007/s10311-010-0297-8)

756 Naharro, R., Esbrí, J.M., Amorós, J.Á., García-Navarro, F.J., Higuera, P., 2018. Assessment
757 of mercury uptake routes at the soil-plant-atmosphere interface. *Geochemistry Explor.*
758 *Environ. Anal.* 19, 146–154. <https://doi.org/10.1144/geochem2018-019>

759 Naharro, R., Esbrí, J.M., Amorós, J.A., Higuera, P.L., 2020. Experimental assessment of the
760 daily exchange of atmospheric mercury in *Epipremnum aureum*. *Environ. Geochem.*
761 *Health* 1–14. <https://doi.org/10.1007/s10653-020-00557-8>

762 Nies, D.H., 2003. Efflux-mediated heavy metal resistance in prokaryotes. *FEMS Microbiol.*
763 *Rev.* 27, 313–339. [https://doi.org/10.1016/S0168-6445\(03\)00048-2](https://doi.org/10.1016/S0168-6445(03)00048-2)

764 Ntougias, S., Bourtzis, K., Tsiamis, G., 2013. The microbiology of olive mill wastes. *Biomed*
765 *Res. Int.* <https://doi.org/10.1155/2013/784591>

766 Nunes, I., Jacquiod, S., Brejnrod, A., Holm, P.E., Johansen, A., Brandt, K.K., Priemé, A.,
767 Sørensen, S.J., 2016. Coping with copper: legacy effect of copper on potential activity of
768 soil bacteria following a century of exposure. *FEMS Microbiol. Ecol.* 92, fiw175.
769 <https://doi.org/10.1093/femsec/fiw175>

770 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P., O'hara, R., Simpson, G.,
771 Solymos, P., Henry, M., Stevens, H., Wagner, H., 2018. Package 'vegan' community
772 ecology package [WWW Document].

773 Pacwa-Płociniczak, M., Płociniczak, T., Yu, D., Kurola, J.M., Sinkkonen, A., Piotrowska-
774 Seget, Z., Romantschuk, M., 2018. Effect of *Silene vulgaris* and Heavy Metal Pollution
775 on Soil Microbial Diversity in Long-Term Contaminated Soil. *Water. Air. Soil Pollut.*
776 229, 1–13. <https://doi.org/10.1007/s11270-017-3655-3>

777 Pająk, M., Błońska, E., Szostak, M., Gąsiorek, M., Pietrzykowski, M., Urban, O., Derbis, P.,
778 2018. Restoration of Vegetation in Relation to Soil Properties of Spoil Heap Heavily
779 Contaminated with Heavy Metals. *Water. Air. Soil Pollut.* 229, 1–15.
780 <https://doi.org/10.1007/s11270-018-4040-6>

781 Paredes, C., Cegarra, J., Roig, A., Sánchez-Monedero, M.A., Bernal, M.P., 1999.

782 Characterization of olive mill wastewater (alpechin) and its sludge for agricultural
783 purposes. *Bioresour. Technol.* 67, 111–115. <https://doi.org/10.1016/S0960->
784 [8524\(98\)00106-0](https://doi.org/10.1016/S0960-8524(98)00106-0)

785 Parelho, C., Rodrigues, A.S., Barreto, M.C., Ferreira, N.G.C., Garcia, P., 2016. Assessing
786 microbial activities in metal contaminated agricultural volcanic soils - An integrative
787 approach. *Ecotoxicol. Environ. Saf.* 129, 242–249.
788 <https://doi.org/10.1016/j.ecoenv.2016.03.019>

789 Peoples, M.S., Koide, R.T., 2012. Considerations in the storage of soil samples for enzyme
790 activity analysis. *Appl. Soil Ecol.* 62, 98–102.
791 <https://doi.org/10.1016/j.apsoil.2012.08.002>

792 Philippot, L., Spor, A., Hénault, C., Bru, D., Bizouard, F., Jones, C.M., Sarr, A., Maron, P.A.,
793 2013. Loss in microbial diversity affects nitrogen cycling in soil. *ISME J.* 7, 1609–1619.
794 <https://doi.org/10.1038/ismej.2013.34>

795 Price, M.N., Dehal, P.S., Arkin, A.P., 2009. Fasttree: Computing large minimum evolution
796 trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* 26, 1641–1650.
797 <https://doi.org/10.1093/molbev/msp077>

798 Rieuwerts, J.S., Thornton, I., Farago, M.E., Ashmore, M.R., 1998. Factors influencing metal
799 bioavailability in soils: Preliminary investigations for the development of a critical loads
800 approach for metals. *Chem. Speciat. Bioavailab.* 10, 61–75.
801 <https://doi.org/10.3184/095422998782775835>

802 Rodríguez, L., Ruiz, E., Alonso-Azcárate, J., Rincón, J., 2009. Heavy metal distribution and
803 chemical speciation in tailings and soils around a Pb-Zn mine in Spain. *J. Environ.*
804 *Manage.* 90, 1106–1116. <https://doi.org/10.1016/j.jenvman.2008.04.007>

805 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open
806 source tool for metagenomics. *PeerJ* 4, e2584. <https://doi.org/10.7717/peerj.2584>

807 Rohart, F., Gautier, B., Singh, A., Lê Cao, K.A., 2017. mixOmics: An R package for ‘omics
808 feature selection and multiple data integration. *PLoS Comput. Biol.* 13, e1005752.
809 <https://doi.org/10.1371/journal.pcbi.1005752>

810 Rouch, D.A., Lee, B.T.O., Morby, A.P., 1995. Understanding cellular responses to toxic
811 agents: a model for mechanism-choice in bacterial metal resistance. *J. Ind. Microbiol.* 14,
812 132–141. <https://doi.org/10.1007/BF01569895>

813 Ruiz, E., Alonso-Azcárate, J., Rodríguez, L., Rincón, J., 2009. Assessment of metal
814 availability in soils from a Pb-Zn mine site of South-central Spain. *Soil Sediment
815 Contam.* 18, 619–641. <https://doi.org/10.1080/15320380903113600>

816 Sánchez-Donoso, R., Martín-Duque, J.F., Crespo, E., Higuera, P.L., 2019. Tailing’s
817 geomorphology of the San Quintín mining site (Spain): landform catalogue, aeolian
818 erosion and environmental implications. *Environ. Earth Sci.* 78, 1–17.
819 <https://doi.org/10.1007/s12665-019-8148-9>

820 Schmidt, S.K., Gendron, E.M.S., Vincent, K., Solon, A.J., Sommers, P., Schubert, Z.R.,
821 Vimercati, L., Porazinska, D.L., Darcy, J.L., Sowell, P., 2018. Life at extreme elevations
822 on Atacama volcanoes: the closest thing to Mars on Earth? *Antonie van Leeuwenhoek*,
823 *Int. J. Gen. Mol. Microbiol.* 111, 1389–1401. <https://doi.org/10.1007/s10482-018-1066-0>

824 Schneider, A.R., Gommeaux, M., Duclercq, J., Fanin, N., Conreux, A., Alahmad, A., Lacoux,
825 J., Roger, D., Spicher, F., Ponthieu, M., Cancès, B., Morvan, X., Marin, B., 2017.
826 Response of bacterial communities to Pb smelter pollution in contrasting soils. *Sci. Total
827 Environ.* 605–606, 436–444. <https://doi.org/10.1016/j.scitotenv.2017.06.159>

828 Siles, J.A., Pérez-Mendoza, D., Ibáñez, J.A., Scervino, J.M., Ocampo, J.A., García-Romera,
829 I., Sampedro, I., 2014. Assessing the impact of biotransformed dry olive residue
830 application to soil: Effects on enzyme activities and fungal community. *Int. Biodeterior.*
831 *Biodegrad.* 89, 15–22. <https://doi.org/10.1016/j.ibiod.2014.01.001>

832 Singh, A., Shannon, C.P., Gautier, B., Rohart, F., Vacher, M., Tebbutt, S.J., Cao, K.-A.L.,
833 2019. DIABLO: An Integrative Approach for Identifying Key Molecular Drivers From
834 Multi-Omics Assays - PubMed. *Bioinformatics* 35, 3055–3062.
835 <https://doi.org/10.1093/bioinformatics/bty1054>

836 Sobolev, D., Begonia, M.F.T., 2008. Effects of heavy metal contamination upon soil
837 microbes: Lead-induced changes in general and denitrifying microbial communities as
838 evidenced by molecular markers, in: *International Journal of Environmental Research*
839 *and Public Health*. Multidisciplinary Digital Publishing Institute (MDPI), pp. 450–456.
840 <https://doi.org/10.3390/ijerph5050450>

841 Stefanowicz, A.M., Kapusta, P., Zubek, S., Stanek, M., Woch, M.W., 2020. Soil organic
842 matter prevails over heavy metal pollution and vegetation as a factor shaping soil
843 microbial communities at historical Zn–Pb mining sites. *Chemosphere* 240, 124922.
844 <https://doi.org/10.1016/j.chemosphere.2019.124922>

845 Stefanowicz, A.M., Niklińska, M., Laskowski, R., 2008. Metals affect soil bacterial and
846 fungal functional diversity differently. *Environ. Toxicol. Chem.* 27, 591.
847 <https://doi.org/10.1897/07-288.1>

848 Tabatabai, M.A., Bremner, J.M., 1970. Arylsulfatase Activity of Soils. *Soil Sci. Soc. Am. J.*
849 34, 225–229. <https://doi.org/10.2136/sssaj1970.03615995003400020016x>

850 Tabatabai, M.A., Bremner, J.M., 1969. Use of p-nitrophenyl phosphate for assay of soil
851 phosphatase activity. *Soil Biol. Biochem.* 1, 301–307. <https://doi.org/10.1016/0038->

852 0717(69)90012-1

853 Tang, J., Zhang, J., Ren, L., Zhou, Y., Gao, J., Luo, L., Yang, Y., Peng, Q., Huang, H., Chen,
854 A., 2019. Diagnosis of soil contamination using microbiological indices: A review on
855 heavy metal pollution. *J. Environ. Manage.* 242, 121–130.
856 <https://doi.org/10.1016/j.jenvman.2019.04.061>

857 Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K., Sutton, D.J., 2012. Heavy metal toxicity and
858 the environment. *EXS* 101, 133–164. https://doi.org/10.1007/978-3-7643-8340-4_6

859 Tebo, B.M., Davis, R.E., Anitori, R.P., Connell, L.B., Schiffman, P., Staudigel, H., 2015.
860 Microbial communities in dark oligotrophic volcanic ice cave ecosystems of Mt. Erebus,
861 Antarctica. *Front. Microbiol.* 6, 179. <https://doi.org/10.3389/fmicb.2015.00179>

862 Tejero, J., Higuera, P.L., Garrido, I., Esbrí, J.M., Oyarzun, R., Español, S., 2015. An
863 estimation of mercury concentrations in the local atmosphere of Almadén (Ciudad Real
864 Province, South Central Spain) during the twentieth century. *Environ. Sci. Pollut. Res.*
865 22, 4833–4841. <https://doi.org/10.1007/s11356-014-2860-5>

866 Thiele-Bruhn, S., Schloter, M., Wilke, B.-M., Beaudette, L.A., Martin-Laurent, F., Cheviron,
867 N., Mougin, C., Römbke, J., 2020. Identification of new microbial functional standards
868 for soil quality assessment. *SOIL* 6, 17–34. <https://doi.org/10.5194/soil-6-17-2020>

869 Tipayno, S.C., Truu, J., Samaddar, S., Truu, M., Preem, J.-K., Oopkaup, K., Espenberg, M.,
870 Chatterjee, P., Kang, Y., Kim, K., Sa, T., 2018. The bacterial community structure and
871 functional profile in the heavy metal contaminated paddy soils, surrounding a nonferrous
872 smelter in South Korea. *Ecol. Evol.* 8, 6157–6168. <https://doi.org/10.1002/ece3.4170>

873 Van Der Heijden, M.G.A., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority:
874 Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems.

875 Ecol. Lett. 11, 296–310. <https://doi.org/10.1111/j.1461-0248.2007.01139.x>

876 Wagg, C., Bender, S.F., Widmer, F., Van Der Heijden, M.G.A., 2014. Soil biodiversity and
877 soil community composition determine ecosystem multifunctionality. *Proc. Natl. Acad. Sci. U. S. A.* 111, 5266–5270. <https://doi.org/10.1073/pnas.1320054111>

878

879 Walkley, A.J., Black, I., 1934. Estimation of soil organic carbon by the chromic acid titration
880 method. *Soil Sci.* 37, 29–38.

881 Wang, S., Mulligan, C.N., 2006. Effect of natural organic matter on arsenic release from soils
882 and sediments into groundwater. *Environ. Geochem. Health.*
883 <https://doi.org/10.1007/s10653-005-9032-y>

884 Xian, Y., Wang, M., Chen, W., 2015. Quantitative assessment on soil enzyme activities of
885 heavy metal contaminated soils with various soil properties. *Chemosphere* 139, 604–608.
886 <https://doi.org/10.1016/j.chemosphere.2014.12.060>

887 Xie, Y., Fan, J., Zhu, W., Amombo, E., Lou, Y., Chen, L., Fu, J., 2016. Effect of heavy metals
888 pollution on soil microbial diversity and bermudagrass genetic variation. *Front. Plant*
889 *Sci.* 7, 1–12. <https://doi.org/10.3389/fpls.2016.00755>

890 Yang, J.S., Yang, F.L., Yang, Y., Xing, G.L., Deng, C.P., Shen, Y.T., Luo, L.Q., Li, B.Z.,
891 Yuan, H.L., 2016. A proposal of “core enzyme” bioindicator in long-term Pb-Zn ore
892 pollution areas based on topsoil property analysis. *Environ. Pollut.* 213, 760–769.
893 <https://doi.org/10.1016/j.envpol.2016.03.030>

894 Zhang, J., Kobert, K., Flouri, T., Stamatakis, A., 2014. PEAR: a fast and accurate Illumina
895 Paired-End reAd mergeR. *Bioinformatics* 30, 614–620.
896 <https://doi.org/10.1093/bioinformatics/btt593>

897 Zhao, X., Sun, Y., Huang, J., Wang, H., Tang, D., 2020. Effects of soil heavy metal pollution

898 on microbial activities and community diversity in different land use types in mining
899 areas. *Environ. Sci. Pollut. Res.* 27, 20215–20226. [https://doi.org/10.1007/s11356-020-](https://doi.org/10.1007/s11356-020-08538-1)
900 08538-1

901 Zhen, Z., Wang, S., Luo, S., Ren, L., Liang, Y., Yang, R., Li, Y., Zhang, Y., Deng, S., Zou,
902 L., Lin, Z., Zhang, D., 2019. Significant Impacts of Both Total Amount and Availability
903 of Heavy Metals on the Functions and Assembly of Soil Microbial Communities in
904 Different Land Use Patterns. *Front. Microbiol.* 10, 1–14.
905 <https://doi.org/10.3389/fmicb.2019.02293>

906

Figures




Figure 1. Geographical area and sampling sites (reference, green; contaminated, red; olive mill waste, yellow; tailings, blue and dumps, grey) in the study area.




Figure 2. Enzymatic activities (mean \pm S.D): dehydrogenase (DHA) ($\mu\text{g TPF g}^{-1} \text{DW day}^{-1}$), arylsulfatase (ARS) ($\mu\text{g PNF g}^{-1} \text{DW day}^{-1}$), acid phosphomonoesterase (PhA 6.5) ($\mu\text{g PNF g}^{-1} \text{DW day}^{-1}$), alkaline phosphomonoesterase (PhA 11) ($\mu\text{g PNF g}^{-1} \text{DW day}^{-1}$) and β -Galactosidase (β -Gal) ($\mu\text{g PNF g}^{-1} \text{DW day}^{-1}$) for reference-, contaminated-samples, olive mill waste (OMW), tailings and dumps. ANOVA followed by Tukey's test and Kruskal Wallis tests (*) were performed. For DHA, ANOVA was performed on log-transformed data. For each parameter, different letters within the same line indicate significant differences ($p < 0.05$). $n=4$






Figure 3. Bacterial α -diversity indices (mean values \pm standard deviation) derived from amplicon sequencing of the 16S rDNA gene copies from reference-, contaminated-samples, olive mill waste (OMW), tailings and dumps. ANOVA test followed by Tukey's test was performed. For each diversity index value, different letters indicate that samples are significantly different for each treatment and time. For dominance, ANOVA was performed on log-transformed data. n=4



a



b

Figure 4. Bacterial β -diversity analysis for reference-, contaminated-samples, olive mill waste (OMW), tailings and dumps using Bray Curtis distance matrix of 16S rRNA sequences and (a) PTEs concentrations or (b) enzymatic analysis as explanatory factors. The variance explained by each axis is given as percentage. For each sample type, the four replicates are represented with the same color.




Figure 5. Relative abundance of the major bacterial phyla (expressed as % of the total number of OTUs) for reference-, contaminated-samples, olive mill waste (OMW), tailings and dumps. Phyla whose relative abundance was below 5% were grouped as “others”. ANOVA followed by Tukey’s test and Kruskal Wallis tests (Proteobacteria, Chloroflexi, Acidobacteria and Cyanobacteria) were performed. For Firmicutes and TM7, ANOVA was performed on log-transformed data. n=4




Figure 6. Heatmap plot representing relative abundances of OTUs correlated to physicochemical and enzymatic analysis (correlation cut-off $r > 0.4$) in reference-, contaminated-samples, olive mill waste, tailings and dumps.

Tables

Table 1. Distribution (geographic coordinates) and main characteristics of samples

| Sample type | Sample | X (UTM) | Y (UTM) | Altitude | Other Characteristics |
|--------------------|---------------|----------------|----------------|-----------------|------------------------------|
| Reference | SQD-9 | 388986 | 4298003 | 666 | Recently fertilized |
| Reference | SQD-13 | 390156 | 4298068 | 664 | |
| Reference | SQD-18 | 390256 | 4297642 | 671 | Dry land, plowed |
| Reference | SQD-19 | 390207 | 4297137 | 669 | Fallow, unplowed |
| Contaminated | SQD-8 | 388935 | 4297593 | 687 | Agricultural |
| Contaminated | SQD-11 | 389261 | 4297675 | 675 | 2 meters from a plant |
| Contaminated | SQD-12 | 389392 | 4297805 | 678 | Developed soil |
| Contaminated | SQD-14 | 389798 | 4297560 | 663 | |
| Olive Mill Waste | SQD-2 | 389170 | 4297396 | 655 | |
| Olive Mill Waste | SQD-4 | 389103 | 4297409 | 659 | |
| Olive Mill Waste | SQD-6 | 389059 | 4297448 | 657 | |
| Olive Mill Waste | SQD-20 | 389238 | 4297521 | 671 | With visible dump debris |
| Tailings | SQD-1 | 389208 | 4297287 | 646 | Aeolic erosion |
| Tailing | SQD-15 | 389704 | 4297503 | 667 | |
| Tailing | SQD-16 | 389521 | 4297623 | 667 | |
| Tailing | SQD-17 | 389553 | 4297525 | 673 | |
| Dumps | SQD-3 | 389129 | 4297398 | 658 | Heterogeneous granulometry |
| Dumps | SQD-5 | 389081 | 4297558 | 658 | Heterogeneous granulometry |
| Dumps | SQD-7 | 389033 | 4297470 | 662 | Flooded area |
| Dumps | SQD-10 | 389032 | 4297607 | 674 | Heterogeneous granulometry |

Table 2. Physicochemical characteristics of reference-, contaminated-samples, olive mill waste, tailings and dumps. Values were mean \pm standard deviation. ANOVA followed by Tukey's test were performed. For EC and OM, ANOVA was performed on log-transformed data. For each parameter, different letters within the same line indicate significant differences ($p < 0.05$). For EC and OM, ANOVA performed on log-transformed data. $n=4$

| | Reference | Contaminated | Olive Mill Waste | Tailings | Dumps |
|--------------------|------------------|---------------------|-------------------------|---------------------|----------------------|
| pH | 6.1 \pm 0.3a | 6.8 \pm 1.0a | 5.2 \pm 0.6ab | 6.7 \pm 1.7a | 3.9 \pm 0.5b |
| EC (mV) | 62.4 \pm 27.8a | 90.6 \pm 49.9a | 651.3 \pm 489.7ab | 415.6 \pm 378.8ab | 2282.3 \pm 1786.4b |
| OM (%) | 2.9 \pm 1.9a | 8.1 \pm 8.9a | 60.6 \pm 19.7b | 2.0 \pm 0.4a | 1.6 \pm 0.5a |
| Sample type | sandy loam | sandy loam | loamy sand | silt loam | sandy loam |

EC: electric conductivity, OM: organic matter


Table 3. PTEs concentrations (mg/kg) (mean values \pm standard deviation) and Pollution Load Index (PLI) in reference-, contaminated-samples, olive mill waste, tailings and dumps. ANOVA test followed by Tukey's test and Kruskal Wallis followed by pairwise comparisons with Bonferroni correction (*) were performed. Values indicated by different letters are significantly different within the same line. For Ni and Cu ANOVA was performed on root square transformed data. For Sb, ANOVA was performed on inverse transformed data, respectively. For Zn, Hg and Th, ANOVA performed on log-transformed data, respectively.

n=4

| | Reference | Contaminated | Olive Mill Waste | Tailings | Dumps |
|------------|---------------------|-----------------------|-------------------------|-----------------------|------------------------|
| Pb | 315.8 \pm 332.6a | 3696.5 \pm 4406.2a | 38890 \pm 20687.0b | 4030.2 \pm 7170.8a | 48597.5 \pm 16835.1b |
| Zn | 204.8 \pm 153.8a | 1522.4 \pm 2165.2ab | 86050 \pm 99094.2c | 2024.5 \pm 2492.1ab | 4640.0 \pm 2259.5b |
| Hg | 0 \pm 0a | 55.4 \pm 92.1a | 3605.8 \pm 6752.0a | 557.1 \pm 430.8a | 130.8 \pm 142.5a |
| Sb | 16.8 \pm 1.8a | 37.4 \pm 31.0abc | 330.9 \pm 435.2bc | 65.5 \pm 87.7ab | 433.6 \pm 198.9c |
| Co | 221.7 \pm 76.4a | 268.3 \pm 80.0ab | 451.7 \pm 78.4bc | 192.5 \pm 27.9a | 503.9 \pm 180.6c |
| Cu | 17.3 \pm 10.0a | 165.6 \pm 297.6a | 963.7 \pm 466.2b | 113.4 \pm 196.7a | 1000.6 \pm 805.0b |
| Mn* | 688.2 \pm 238.9ab | 1572.5 \pm 908.6a | 1637.4 \pm 1866.4ab | 423.5 \pm 136.5ab | 210.4 \pm 90.8b |
| Ni | 31.4 \pm 18.7a | 38.9 \pm 31.4a | 132.1 \pm 95.6a | 37.0 \pm 14.0a | 27.6 \pm 8.1a |
| Th | 9.7 \pm 3.9ab | 4.6 \pm 4.1ab | 0 \pm 0b | 3.8 \pm 6.9ab | 35.1 \pm 23.7a |
| PLI | 1.0 \pm 0.1 | 3.2 \pm 2.8 | 21.1 \pm 23.9 | 10.5 \pm 4.8 | 7.4 \pm 2.4 |

PLI: Pollution Load Index

SAN QUINTÍN MINE



Analytical chemistry

Microbial enzyme activities

Composition and diversity of bacterial community

Ecotoxicological risk assessment of San Quintín mine area