



**HAL**  
open science

## Environmental impacts and their association with performance and excretion traits in growing pigs

Alessandra N. T. R. Monteiro, Ludovic Brossard, H el ene Gilbert, Jean-Yves  
Dourmad

### ► To cite this version:

Alessandra N. T. R. Monteiro, Ludovic Brossard, H el ene Gilbert, Jean-Yves Dourmad. Environmental impacts and their association with performance and excretion traits in growing pigs. *Frontiers in Veterinary Science*, 2021, 8, 10.3389/fvets.2021.677857 . hal-03311758

**HAL Id: hal-03311758**

**<https://hal.inrae.fr/hal-03311758>**

Submitted on 2 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche franais ou  trangers, des laboratoires publics ou priv es.



Distributed under a Creative Commons Attribution 4.0 International License



# Environmental Impacts and Their Association With Performance and Excretion Traits in Growing Pigs

Alessandra N. T. R. Monteiro<sup>1</sup>, Ludovic Brossard<sup>1</sup>, H el ene Gilbert<sup>2</sup> and Jean-Yves Dourmad<sup>1\*</sup>

<sup>1</sup> PEGASE, INRAE, Institut Agro, Saint-Gilles, France, <sup>2</sup> GenPhySE, Universit e de Toulouse, INRAE, INPT, ENSAT, Castanet-Tolosan, France

## OPEN ACCESS

### Edited by:

Aline Remus,  
Agriculture and Agri-Food Canada  
(AAFC), Canada

### Reviewed by:

Cormac John O'Shea,  
University of Nottingham,  
United Kingdom  
Nikola Puva ca,  
University Business Academy in Novi  
Sad, Serbia

### \*Correspondence:

Jean-Yves Dourmad  
jean-yves.dourmad@inrae.fr

### Specialty section:

This article was submitted to  
Animal Nutrition and Metabolism,  
a section of the journal  
Frontiers in Veterinary Science

**Received:** 08 March 2021

**Accepted:** 06 May 2021

**Published:** 21 June 2021

### Citation:

Monteiro ANTR, Brossard L, Gilbert H  
and Dourmad J-Y (2021)  
Environmental Impacts and Their  
Association With Performance and  
Excretion Traits in Growing Pigs.  
Front. Vet. Sci. 8:677857.  
doi: 10.3389/fvets.2021.677857

The selection of pigs for improved production traits has been, for a long time, the major driver of pig breeding. More recently, because of the increasing concern with the environment, new selection criteria have been explored, such as nitrogen (N) excretion. However, many studies indicate that life cycle assessment (LCA) provides much better indicators of environmental impacts than excretion. Therefore, the objective of this study was to investigate, using a modeling approach, the relationships between production traits and LCA impacts of individual growing pigs calculated at the farm gate for 1 kg of body weight gain. Performances of pigs were simulated for 2-phase (2P) and precision feeding (PR), using the InraPorc population model (on 1,000 pigs). Nitrogen excretion was positively correlated with feed conversion ratio (FCR;  $r = +0.96$ ), climate change (CC;  $r = +0.96$ ), acidification potential (AC;  $r = +0.97$ ), eutrophication potential (EU;  $r = +0.97$ ), and land occupation (LO;  $r = +0.96$ ), whatever the feeding program. However, FCR appeared to be a better indicator of LCA impacts, with very high and positive correlations ( $r > +0.99$ ) with CC, AC, EU, and LO for both feeding programs. The CC, AC, and EU impacts of pig production for PR feeding were 1.3, 10, and 7.5% lower than for 2P, respectively, but the correlations within each outcome were very similar among feeding programs. It was concluded that the use of FCR as a selection criterion in pig breeding seems to be a promising approach to associate improved performance and low environmental impact of pig fattening.

**Keywords:** feed efficiency, environmental impacts, life cycle assessment, growing pig, modeling

## INTRODUCTION

The selection of animals for improved production traits has been, for a long time, the major driver of pig breeding (1, 2). More recently, because of the increasing concern with environment, new selection criteria have been explored, such as nitrogen (N) or phosphorus (P) excretion, which are related to both feed efficiency and environmental impact (3). Residual feed intake (RFI) was also proposed as a possible selection criterion to simultaneously improve feed efficiency and reduce N and P excretion (1).

However, the pig supply chain involves a complex system, which requires production of fertilizers and pesticides; production of feed ingredients; feed processing; animal raising; transportation of animals and feed; water use for drinking and cleaning; energy use for light, heat, and ventilation; and waste management (4). Therefore, the environmental degradation is not the

consequence of only one process (e.g., the raising of pigs) or one element (e.g., N excretion) and, as reviewed by McAuliffe et al. (4), impacts are better evaluated through integrated methodologies such as life cycle assessment (LCA).

Recently, a comparative LCA showed that pigs selected for low RFI have, on average, 6% lower environmental impacts on climate change (CC), acidification (AC), eutrophication (EU), land occupation (LO), and water depletion than these selected for high RFI (5). However, in this study, RFI did not appear to be the optimum measure for efficient environmentally friendly selection, since it was rather poorly correlated to environmental impacts ( $r = 0.73$  for CC in the low RFI line).

The objective of the present study was thus to investigate, using a modeling approach, the relationships between different performance selection traits and LCA environmental impacts evaluated in individual growing pigs.

## MATERIALS AND METHODS

### Feeding Strategies and Animal Performance

This study considered a conventional growing–finishing pig unit located in West France, as described in detail by Monteiro et al. (6). Two feeds were formulated on the basis of net energy (NE, 9.6 MJ/kg), standardized ileal digestible (SID) amino acids, and digestible phosphorus (P): feed A to achieve 110% the mean population nutrient requirements at the beginning of the growing period (9.84 g/g SID lysine, 3.01 g/kg digestible P), and feed B to achieve 90% the mean population nutrient requirements at the end of the finishing period (4.55 g/kg SID lysine, 1.68 g/kg digestible P). The two feeds were blended according to two feeding programs: 2-phase feeding (2P) corresponding to the strategy used in French central test stations or precision feeding (PR). The 2P pigs were fed with feed A from 30 to 70 kg BW, and then with a blend of 50% of each feed until the end of fattening, to achieve 110% the mean population SID-lysine requirement at the start of the finishing period. For PR pigs, the blend of the two feeds was calculated according to a factorial approach in order that each pig received each day the exact amount of SID lysine required to achieve its potential of protein deposition, which was defined according to a Gompertz function, as described by van Milgen et al. (7).

Simulations for a virtual population of 1,000 female pigs were performed individually, from 30 to 115 kg of BW, for each feeding program to determine individual animal performance, nutrient balance, and excretion according to InraPorc population model (8). This virtual population was generated according to the method described by Brossard et al. (8), from a variance–covariance matrix with two parameters describing individual pig feed intake (the net energy intake at 50 and 100 kg BW:  $20.2 \pm 2.0$  and  $25.0 \pm 2.9$  MJ NE/day, respectively) and three parameters describing the Gompertz function of potential protein deposition (the BW at 70 days:  $30.0 \pm 2.9$  kg, the mean protein deposition rate between 70 days of age and 110 kg BW:  $142.8 \pm 15.2$  g, and the precocity  $b$ -value of the Gompertz function:  $0.0169 \pm 0.0103$ ).

The simulated performance and excretion data were then used to calculate gaseous emissions from animals and manure, according to Rigolot et al. (9). The pig production system considered was a conventional growing–finishing pig farm located in Brittany (West France) with indoor raising of animals on complete slatted floor, in a building with mechanical ventilation and collection and storage of manure as liquid slurry (6).

### Life Cycle Assessment

The LCA was performed for each pig, considering all the impacts associated with feed production, animal housing, and manure management (as described by 6). We based our analysis on the CML 2001 (baseline) method version 3.02 as implemented in SimaPro software version 8.05 (PRé Consultants) and added the category land occupation from CML 2001 (all categories) version 2.04. Thus, we considered the potential impacts of pig production on CC (kg CO<sub>2</sub>-eq), EU (g PO<sub>4</sub>-eq), AC (g SO<sub>2</sub>-eq), and LO (m<sup>2</sup>·year). The CC was calculated according to the 100-year global warming potential factors in kilograms CO<sub>2</sub>-eq. Impacts were calculated at the farm gate, and the functional unit considered was 1 kg of BW gain over the fattening period.

### Statistical Analysis

The LCA calculation model was implemented using SAS software (SAS Inst. Inc., Cary, NC). Performance and environmental impacts were subjected to variance analysis using GLM procedure with feeding strategy as main effect. Pearson correlations for each feeding strategy were calculated between performance and environmental impacts data using CORR procedure, and pigs were ranked according to their CC impact, considering the feeding strategy and using the RANK procedure. All analyses were conducted using SAS software version 9.1 (SAS Inst. Inc., Cary, NC).

All the data used in the statistical analysis are available in the INRAE data repository (10).

## RESULTS AND DISCUSSION

### Feeding Strategies, Animal Performance, and Environmental Impacts

Feeding strategies affected most of the parameters evaluated (Table 1); effects were more accentuated for N excretion and N retention efficiency, and for CC, EU, and AC environmental impacts, which are highly dependent on dietary crude protein (CP) content, which was on average lower for PR (144 g/kg) than for 2P (167 g/kg). Compared to 2P, with PR, ADG was slightly improved (by 1.3%), efficiency of N retention was increased (40.5 vs. 36.2%), N excretion was reduced (by 16%), and environmental impacts were decreased (CC, AC, EU, and LO impacts 1.3, 10.0, 7.5, and 0.8% lower than for 2P, respectively). These results are in agreement with previous studies indicating that PR feeding strategy allows the improvement of the performance of pigs, compared to phase feeding, by providing sufficient amount of amino acids even to the animals with the highest potential of protein retention, which may not be the case with phase feeding, especially at the beginning of each phase. In

**TABLE 1** | Effect of feeding strategy on pig performance, nitrogen excretion, and environmental impacts measured by life cycle assessment ( $n = 1,000$  pigs).

Item	Two-Phase feeding	Precision feeding	P-value <sup>a</sup>
<b>Performance</b>			
ADFI, g/day	2,310 ± 259	2,316 ± 261	ns
ADG, g/day	864 ± 112	876 ± 116	*
FCR, kg/kg	2.69 ± 0.29	2.67 ± 0.32	<i>t</i>
SID lysine intake, g/day	1.89 ± 0.20	1.47 ± 0.11	***
Protein intake, g/day	375 ± 42.2	340 ± 32.5	***
Protein retained, g/day	135 ± 18.4	138 ± 20.1	*
N retention efficiency, %	36.2 ± 4.8	40.5 ± 4.8	***
<b>Environmental impacts</b>			
N excreted, kg/pig	3.83 ± 0.69	3.20 ± 0.56	***
CC, kg CO <sub>2</sub> -eq/kg BW gain	2.34 ± 0.25	2.31 ± 0.28	*
EU, g PO <sub>4</sub> -eq/kg BW gain	17.4 ± 2.34	16.1 ± 2.22	***
AC, g SO <sub>2</sub> -eq/kg BW gain	48.1 ± 7.29	43.3 ± 6.60	***
LO, m <sup>2</sup> year/kg BW gain	3.77 ± 0.40	3.74 ± 0.45	<i>t</i>

ADG, average daily gain; ADFI, average daily feed intake; FCR, feed conversion ratio; CC, Climate Change; EU, Eutrophication potential; AC, Acidification Potential; LO, Land Occupation. Two-phases feeding = pigs received a "growing" diet from 30 to 70 kg of BW and a "finishing" diet from 70 to 115 kg of BW; Precision feeding = individual pigs were fed daily with a diet providing the exact amount of digestible amino acids they required.

<sup>a</sup>*t*:  $P < 0.10$ ; \* $P < 0.05$ ; \*\*\* $P < 0.001$ .

**TABLE 2** | Correlations<sup>a</sup> between performance traits, nitrogen excretion, and environmental impacts, for the precision (PR) and the two-phase (2P, in italic) feeding strategies.

		ADFI	ADG	FCR	NR	NEff	NE	CC	EU	AC
ADG	PR	0.683								
	2P	<i>0.576</i>								
FCR	PR	0.181	-0.583							
	2P	<i>0.246</i>	<i>-0.636</i>							
NR	PR	-0.692	-0.002	-0.643						
	2P	<i>-0.675</i>	<i>-0.112</i>	<i>-0.593</i>						
Neff	PR	-0.361	0.543	-0.986	0.683					
	2P	<i>-0.385</i>	<i>0.400</i>	<i>-0.959</i>	<i>0.669</i>					
NE	PR	0.187	-0.552	0.963	-0.428	-0.940				
	2P	<i>0.125</i>	<i>-0.682</i>	<i>0.956</i>	<i>-0.450</i>	<i>-0.953</i>				
CC	PR	0.188	-0.577	0.999	-0.647	-0.987	0.963			
	2P	<i>0.249</i>	<i>-0.633</i>	<i>0.999</i>	<i>-0.599</i>	<i>-0.960</i>	<i>0.955</i>			
EU	PR	0.225	-0.546	0.998	-0.625	-0.983	0.971	0.998		
	2P	<i>0.225</i>	<i>-0.645</i>	<i>0.997</i>	<i>-0.610</i>	<i>-0.973</i>	<i>0.966</i>	<i>0.997</i>		
AC	PR	0.241	-0.532	0.996	-0.621	-0.981	0.972	0.996	0.999	
	2P	<i>0.224</i>	<i>-0.643</i>	<i>0.996</i>	<i>-0.617</i>	<i>-0.977</i>	<i>0.968</i>	<i>0.996</i>	<i>0.997</i>	
LO	PR	0.181	-0.583	0.999	-0.643	-0.986	0.963	0.999	0.998	0.996
	2P	<i>0.245</i>	<i>-0.637</i>	<i>0.999</i>	<i>-0.593</i>	<i>-0.959</i>	<i>0.956</i>	<i>0.999</i>	<i>0.997</i>	<i>0.996</i>


ADFI, average daily feed intake; ADG, average daily gain; FCR, feed conversion ratio; NE, nitrogen excretion; CC, Climate Change; EU, Eutrophication potential; AC, Acidification Potential; LO, Land Occupation.

<sup>a</sup>All correlations were significantly different from 0 ( $P < 0.001$ ).

PR compared to 2P feeding strategy, protein and SID lysine intakes were reduced by 9.3 and 22.2%, respectively. Combined with the slightly improved protein retention in PR pigs, this resulted in a significant increase of N retention efficiency (from 36.2 to 40.5%) and a reduction of nutrient load in excreta, contributing to the lower CC, EU, and AC impacts with precision feeding, as already shown by Monteiro et al. (6) and Andretta et al. (11).

## Correlation Between Performance, Excretion, and Environmental Impacts

Correlations between performance, excretion and environmental impacts are shown in Table 2. The correlation values obtained for 2P and PR strategies were very close. Nitrogen excretion was highly and positively correlated with CC ( $r = +0.96$ , Figure 1A), AC ( $r = +0.97$ ), EU ( $r = +0.97$ ), and LO ( $r = +0.96$ ). Correlations between environmental impacts and NR were much



lower than with NE, with  $r$  values ranging between 0.42 and 0.64, depending on the category. Correlations between environmental impacts and N retention efficiency were similar to these obtained with N excretion.

Average daily feed intake (ADFI) presented much lower correlation with all the impact categories ( $r = +0.21$  on average). The weak correlation between ADFI and environmental impacts corroborated the 0.25–0.30 values obtained by Soleimani and Gilbert (5).

Feed conversion ratio appeared the best indicator of LCA impacts, with very high and positive correlations (Table 2,  $r > +0.99$ ) with CC (Figure 1B), AC, EU, and LO for both feeding programs. This is consistent with the major contribution of feed intake to most environmental impacts (more than 70% for CC, EU, and LO, and about 30% for AC; 6), as well as to FCR. Moreover, efficient pigs, with lower FCR, ingest less energy and protein per kilogram of gain, which results in reduced enteric and manure methane production, and reduced organic matter, N, and P excretion. Gaseous emissions of N compounds from excreta have an important contribution to CC (due to N<sub>2</sub>O emission) and to AC and EU (due to NH<sub>3</sub> emission). Moreover, NO<sub>3</sub><sup>-</sup> and PO<sub>4</sub><sup>-</sup> leaching after manure spreading also contributes to EU. This contributes to explain the close correlation between N retention efficiency and environmental impacts ( $r$  ranging from 0.96 to 0.98 depending on the impact category). These reductions in enteric emissions and emissions from excreta and manure from more efficient pigs (with low FCR) also contribute to explain the close relationship obtained between FCR and environmental impacts, both expressed per kilogram of body weight gain.

Despite the lower CC, AC, EU, and LO of pig production in the PR program, the correlations within each outcome were very similar among feeding programs.

## Between-Animal Variability

It has already been shown that precision feeding strategy removes a constraint on reaching maximum growth potential and allows all animals to express their maximum growth potential, whereas with phase-feeding strategy, the performance of the highest potential animals may be limited due to insufficient amino acid

supplies (8, 12). This explains that the variability of performance and environmental impacts may differ according to the feeding strategy. For instance, the coefficient of variation of CC impact was higher with PR than with 2P feeding strategy (12.1 and 10.7%, respectively). This affects the pigs' ranking, as illustrated in Figure 1C, which shows the correlation between the ranking of pigs according to CC impact with the two feeding strategies. Similar results were obtained for FCR.

## CONCLUSIONS

The results of this study indicate that FCR is better correlated with environmental impacts evaluated using LCA than nitrogen excretion or other performance criteria. This offers interesting perspectives for the improvement of both feed efficiency and environmental impacts. However, further studies are still required before implementing LCA environmental impacts (or FCR as a proxy of these impacts) in selection programs. The same approach as the one used in this study with simulated data could be carried out on real data collected from selection programs. This would allow the assessment of the genetic parameters of the different LCA impacts and would allow taking better account of all the biological phenomena influencing growth performance, nutrient excretion, and enteric emission, which are probably not completely represented in the growth simulation model. Moreover, the correlated effects on other important criteria, such as carcass lean percentage, meat quality, or animal health and behavior, should also be evaluated.

## DATA AVAILABILITY STATEMENT

All the data used in the statistical analysis are available in the INRAE data repository (10).

## AUTHOR CONTRIBUTIONS

AM and J-YD contributed to conception and design of the study. AM, LB, and J-YD organized the database. AM, LB,

HG, and J-YD wrote sections of the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

## REFERENCES

1. Saintilan R, Mérour I, Brossard L, Tribout T, Dourmad JY, Sellier P, et al. Genetics of residual feed intake in growing pigs: relationships with production traits, and nitrogen and phosphorus excretion traits. *J Anim Sci.* (2013) 91:2542–54. doi: 10.2527/jas.2012-5687
2. Niemi JK, Sevón-Aimonen ML, Stygar AH, Partanen K. The economic and environmental value of genetic improvements in fattening pigs: an integrated dynamic model approach. *J Anim Sci.* (2015) 93:4161–71. doi: 10.2527/jas.2015-9011
3. Shirali M, Duthie CA, Doeschl-Wilson A, Knap PW, Kanis E, van Arendonk JAM, et al. Novel insight into the genomic architecture of feed and nitrogen efficiency measured by residual energy intake and nitrogen excretion in growing pigs. *BMC Genomics.* (2013) 14:1–12. doi: 10.1186/1471-2156-14-121
4. McAuliffe GA, Chapman DV, Sage CL. A thematic review of life cycle assessment (LCA) applied to pig production. *Environ Impact Assess Rev.* (2016) 56:12–22. doi: 10.1016/j.eiar.2015.08.008
5. Soleimani Jevinani F, Gilbert H. Evaluating environmental impacts of selection for residual feed intake in pigs. *Animal.* (2020) 14:2598–608. doi: 10.1017/S175173112000138X
6. Monteiro ANTR, Garcia-Launay F, Brossard L, Wilfart A, Dourmad JY. Effect of feeding strategy on environmental impacts of pig fattening in different contexts of production: evaluation through life cycle assessment. *J Anim Sci.* (2016) 94:4832–47. doi: 10.2527/jas.2016-0529
7. van Milgen J, Valancogne A, Dubois S, Dourmad JY, Sève B, Noblet J. InraPorc: a model and decision support tool for the nutrition of growing pigs. *Anim Feed Sci Technol.* (2008) 143:387–405. doi: 10.1016/j.anifeedsci.2007.05.020
8. Brossard L, Vautier B, van Milgen J, Salaün Y, Quiniou N. Comparison of *in vivo* and *in silico* growth performance and variability in pigs when applying a feeding strategy designed by simulation to control the variability of slaughter weight. *Anim Prod Sci.* (2014) 54:1939–45. doi: 10.1071/AN14521
9. Rigolot C, Espagnol S, Pomar C, Dourmad JY. Modelling of manure production by pigs and NH<sub>3</sub>, N<sub>2</sub>O and CH<sub>4</sub> emissions. Part I: animal excretion and enteric CH<sub>4</sub>, effect of feeding and performance. *Animal.* (2010) 4:1401–12. doi: 10.1017/S1751731110000492
10. Dourmad JY. Article Data - Environmental impacts and their association with performance and excretion traits in growing pigs. *Front Vet Sci.* (2021). doi: 10.15454/IRDMYU
11. Andretta I, Hauschild L, Kipper M, Pires PGS, Pomar C. Environmental impacts of precision feeding programs applied in pig production. *Animal.* (2018) 12:1990–8. doi: 10.1017/S1751731117003159
12. Andretta I, Pomar C, Rivest J, Pomar J, Lovatto PA, Radünz Neto J. The impact of feeding growing pigs with daily tailored diets using precision feeding techniques on animal performance, nutrient utilization, and body and carcass composition. *J Anim Sci.* (2014) 92:3925–36. doi: 10.2527/jas.2014-7643

## FUNDING

This research was funded by INRAE.

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Monteiro, Brossard, Gilbert and Dourmad. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.