



First Complete Genome Sequences of *Xanthomonas citri* pv. *vignicola* Strains CFBP7111, CFBP7112, and CFBP7113 Obtained Using Long-Read Technology

Mylène Ruh, Martial Briand, Sophie Bonneau, Marie-Agnès Jacques, Nicolas Chen

► To cite this version:

Mylène Ruh, Martial Briand, Sophie Bonneau, Marie-Agnès Jacques, Nicolas Chen. First Complete Genome Sequences of *Xanthomonas citri* pv. *vignicola* Strains CFBP7111, CFBP7112, and CFBP7113 Obtained Using Long-Read Technology. *Genome Announcements*, 2017, 5 (36), pp.e00813-17. 10.1128/genomeA.00813-17 . hal-03314799

HAL Id: hal-03314799

<https://hal.inrae.fr/hal-03314799>

Submitted on 9 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



First Complete Genome Sequences of *Xanthomonas citri* pv. *vignicola* Strains CFBP7111, CFBP7112, and CFBP7113 Obtained Using Long-Read Technology

Mylène Ruh, Martial Briand, Sophie Bonneau, Marie-Agnès Jacques,

 Nicolas W. G. Chen

IRHS, INRA, Agrocampus Ouest, Université d'Angers, SFR4207 QUASAV, Beauzoué, France

ABSTRACT *Xanthomonas citri* pv. *vignicola* strains cause bacterial blight of the legume crop cowpea. We report whole-genome sequences of three *X. citri* pv. *vignicola* strains obtained using PacBio single-molecule real-time sequencing. Such genomic data provide new information on pathogenicity factors, such as transcription activator-like effectors.

Cowpea (*Vigna unguiculata*) is a legume crop grown in semiarid tropical and subtropical regions of Africa, America, and Asia (1). It is used as staple food in different forms (grain, green pods, roots, or leaves) and as animal feed (2). Cowpea bacterial blight (CoBB) is a major disease of cow pea that severely reduces crop yields (3) and is present wherever cowpea is grown (4). The causal agent of CoBB is *Xanthomonas citri* pv. *vignicola* (formerly *X. Axonopodis* pv. *vignicola*) (5). *X. Citri* pv. *vignicola* mainly produces symptoms on leaves, starting as tiny water-soaked spots that evolve into large necrotic areas surrounded by a yellow halo. This pathogen can also cause stem canker and water-soaking of pods, providing the opportunity for the bacteria to gain access to seeds (6). CoBB is seed-borne and can also be transmitted by aerosols and insects (7). Moreover, *X. citri* pv. *vignicola* is able to survive in infected plant debris, seeds, and soil, which constitute different inoculum sources (8, 9). Some genotypes of cowpea have been selected for resistance to CoBB (10); however, no information on the molecular basis for this resistance is available to date.

We sequenced three strains of *X. citri* pv. *vignicola* using PacBio single-molecule real-time (SMRT) sequencing (11). Genomic DNA was extracted using the Wizard genomic DNA purification kit (Promega) according to the manufacturer's recommendations. Whole genomes were sequenced using the PacBio RSII SMRT technology at the Icahn School of Medicine at Mount Sinai (New York, NY). One SMRT cell was used per strain to achieve ~100× coverage. *De novo* assembly was performed using Canu version 1.5 (12), circularization of the contigs was done using Circlator version 1.5.1 (13), and polishing was done using the Quiver algorithm (14). Annotation was performed with the NCBI Prokaryotic Genome Annotation Pipeline (15).

Each *X. citri* pv. *vignicola* genome comprised one chromosome plus zero to three circularized plasmids per strain. The genome of strain CFBP7111 comprised three plasmids, for a total assembly size of 5,710,499 bp, with a G+C content of 64.5% and a total of 5,095 predicted coding sequences (CDS). The genome of strain CFBP7112 comprised two plasmids, for a total assembly size of 5,155,996 bp, with a G+C content of 64.7% and a total of 4,460 predicted CDS. The genome of strain CFBP7113 comprised no plasmids, for a total assembly size of 5,105,205 bp, with a G+C content of 64.7% and a total of 4,446 predicted CDS. Interestingly, the content and localization of genes encoding transcription activator-like type III effectors (*tal*) differed among the three

Received 11 July 2017 Accepted 18 July

2017 Published 7 September 2017

Citation Ruh M, Briand M, Bonneau S, Jacques M-A, Chen NWG. 2017. First complete genome sequences of *Xanthomonas citri* pv. *vignicola* strains CFBP7111, CFBP7112, and CFBP7113 obtained using long-read technology. *Genome Announc* 5:e00813-17. <https://doi.org/10.1128/genomeA.00813-17>.

Copyright © 2017 Ruh et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Nicolas W. G. Chen, nicolas.chen@agrocampus-ouest.fr.

genomes. Strain CFBP7111 had no *tal* gene, while strain CFBP7112 contained seven *tal* genes located on plasmids, and strain CFBP7113 bore one *tal* gene located on the chromosome. This diversity may reflect different modes of adaptation of these strains to cowpea.

Accession number(s). The three whole-genome sequences have been deposited at GenBank under accession numbers [CP022263](#) to [CP022266](#) for strain CFBP7111, [CP022267](#) to [CP022269](#) for strain CFBP7112, and [CP022270](#) for strain CFBP7113.

ACKNOWLEDGMENTS

This work was supported by grants from the French National Research Agency (CROpTAL project, grant number ANR-14-CE19-0002-04) and Angers-Loire Metropole, France (X-HOT project).

We thank Laurent D. Noël, Boris Szurek, Lionel Gagnevin, and Ralf Koebnik for recurrent scientific exchanges on the project CROpTAL led by Laurent D. Noël.

PacBio sequencing was performed at the Icahn School of Medicine at Mount Sinai (NY, USA) under the supervision of Robert Sebra and preliminary quality control of DNA was performed at the ANAN platform from the SFR 4207 QUASAV (Angers, France).

We also thank the CIRM-CFBP (Angers, France) for strain preservation and supply.

REFERENCES

- Ehlers JD, Hall AE. 1997. Cowpea (*Vigna unguiculata* L. Walp.). Field Crops Res 53:187–204. [https://doi.org/10.1016/S0378-4290\(97\)00031-2](https://doi.org/10.1016/S0378-4290(97)00031-2).
- Singh BB, Ajeigbe HA, Tarawali SA, Fernandez-Rivera S, Abubakar M. 2003. Improving the production and utilization of cowpea as food and fodder. Field Crops Res 84:169–177. [https://doi.org/10.1016/S0378-4290\(03\)00148-5](https://doi.org/10.1016/S0378-4290(03)00148-5).
- de Lima-Primo HE, de L. Nechet K, de A. Halfeld-Vieira B, de Oliveira JR, Mizubuti ESG, de Souza GR. 2015. Epidemiological aspects of cowpea bacterial blight. Trop Plant Pathol 40:46–55. <https://doi.org/10.1007/s40858-015-0007-y>.
- Khatri-Chhetri GB, Wydra K, Rudolph K. 2003. Metabolic diversity of *Xanthomonas axonopodis* pv. *vignicola*, causal agent of cowpea bacterial blight and pustule. Eur J Plant Pathol 109:851–860. <https://doi.org/10.1023/A:1026104716650>.
- Constantin EC, Cleenwerck I, Maes M, Baeyens S, Van Malderghem C, De Vos P, Cottyn B. 2016. Genetic characterization of strains named as *Xanthomonas axonopodis* pv. *dieffenbachiae* leads to a taxonomic revision of the *X. axonopodis* species complex. Plant Pathol 65:792–806. <https://doi.org/10.1111/ppa.12461>.
- Williams RJ. 1975. Diseases of cowpea (*Vigna unguiculata* (L.) Walp.) in Nigeria. Trop Pest Manag 21:253–267. <https://doi.org/10.1080/09670877509411407>.
- Zandjanakou-Tachin M, Fanou A, Le Gall P, Wydra K. 2007. Detection, survival and transmission of *Xanthomonas axonopodis* pv. *manihotidis* and *X. axonopodis* pv. *vignicola*, causal agents of cassava and cowpea bacterial blight, respectively, in/by insect vectors. J Phytopathol 155: 159–169. <https://doi.org/10.1111/j.1439-0434.2007.01210.x>.
- Okechukwu R, Ekpo E. 2008. Survival of *Xanthomonas campestris* pv. *vignicola* in infested soil, cowpea seed and cowpea debris. Trop Agric Res Ext 11:45–48.
- Sikirou R, Wydra K. 2004. Persistence of *Xanthomonas axonopodis* pv. *vignicola* in weeds and crop debris and identification of *Sphenostylis stenocarpa* as a potential new host. Eur J Plant Pathol 110:939–947. <https://doi.org/10.1007/s10658-004-8949-9>.
- Duche TR, Iheukwumere CC, Omoigui L. 2015. Evaluation of selected cowpea genotypes for resistance to bacterial blight. Int J Curr Microbiol Appl Sci 4:257–270.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, Zhao P, Zhong F, Korlach J, Turner S. 2009. Real-time DNA sequencing from single polymerase molecules. Science 323:133–138. <https://doi.org/10.1126/science.1162986>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. Genome Biol 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Non-hybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat Methods 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity GM, Kodira CD, Kyprides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta) genomic annotation. OMICS 12:137–141. <https://doi.org/10.1089/omi.2008.0017>.