



**HAL**  
open science

## Has breed any effect on beef sensory quality?

Alexandre Conanec, Maria del Mar Campo, Ian Richardson, Per Ertbjerg, Sebastiana Failla, Begoña Panea, Marie Chavent, Jérôme Saracco, John L. Williams, Marie-Pierre Ellies-Oury, et al.

► **To cite this version:**

Alexandre Conanec, Maria del Mar Campo, Ian Richardson, Per Ertbjerg, Sebastiana Failla, et al.. Has breed any effect on beef sensory quality?. *Livestock Science*, 2021, 250, pp.104548. 10.1016/j.livsci.2021.104548 . hal-03316097

**HAL Id: hal-03316097**

**<https://hal.inrae.fr/hal-03316097v1>**

Submitted on 13 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# 1 Has breed any effect on beef sensory quality?

2 Conanec A. (1,2,3), Campo M. (4), Richardson I. (5), Ertbjerg P. (6), Failla S. (7), Panea B. (4,8),

3 Chavent M. (3), Saracco J. (3), Williams J.L. (9,10), Ellies-Oury M-P. (1,2). Hocquette J-F. (2)

4

5 (1) Bordeaux Science Agro, 1 cours du Général de Gaulle, CS 40201, 33175 Gradignan, France

6 (2) INRAE, UMR1213 Herbivores, 63122 Saint Genès Champanelle, France

7 (3) Université de Bordeaux, UMR5251, INRIA, 33400 Talence, France

8 (4) Department of Animal Production and Food Science, Agrifood Institute of Aragón (University of

9 Zaragoza - CITA). 50013 Zaragoza, Spain

10 (5) Bristol Veterinary School, University of Bristol, Langford, Bristol, UK

11 (6) Department of Food and Nutrition, University of Helsinki, 00014 Helsinki, Finland

12 (7) Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria (CREA), Centro di Zootecnia  
13 e Acquacoltura, Monterotondo, Roma, 00015, Italy

14 (8) Agrifood Institute of Aragon-IA2 (CITA-University of Zaragoza), 50059 Zaragoza, Spain.

15 (9) Davies Research Centre, School of Animal and Veterinary Sciences, The University of Adelaide,  
16 Adelaide, Australia

17 (10) Dipartimento di Scienze Animali, della Nutrizione e degli Alimenti, Università Cattolica del Sacro  
18 Cuore, Piacenza, Italy

## 19 Highlights

- 20
- The 15 studied beef breeds can be categorized in five groups of sensory quality.
- 21
- After ten days of ageing, beef from rustic breeds is slightly less tender than those from the
- 22
- other breeds.
- 23
- Fat breeds like Aberdeen Angus, Highland and Jersey produce meat with the highest beef
- 24
- flavor intensity.


- 25       • Mixed effect models including “panelist” as a random effect provide similar results as ANOVA  
26       based on average scores per animal.

## 27 Abstract

28 A total of 436 young cattle from 15 cattle breeds **were** reared in as similar conditions as possible to  
29 evaluate the impact of breed on sensory quality of beef from *longissimus* muscle determined by  
30 sensory analysis. Two statistical methods for processing the sensory data were compared. The  
31 analysis of variance with or without the panelist effect gave similar conclusions indicating that the  
32 robustness of the results was not dependent on the method chosen. The 4 meat descriptors  
33 (tenderness, juiciness, beef flavor and off-flavor) placed breeds into 5 groups using an unsupervised  
34 classification (hierarchical ascending classification). Aberdeen Angus, Highland and Jersey, that have  
35 a high lipid content in the muscle studied, differed from the other breeds **in that they had a higher**  
36 **beef flavour**. The dual-purpose and rustic breeds, Simmental, Casina and Marchigiana, produced  
37 significantly less juicy and less tender **meat** than that from breeds selected for meat production.  
38 Overall, **despite significant differences previously identified for animal, carcass, muscle and beef**  
39 **traits for the same animals**, differences in sensory scores between most of the breeds **were** small,  
40 with only significant differences between the few breeds that had extreme sensory profiles (such as  
41 Simmental and Pirenaica).

42

## 43 Graphical abstract



## 45 Keywords

46 beef; breed; sensory analysis; mixed effect models; fixed effect models

## 47 Abbreviations

- 48 • ME: Mixed Effect
- 49 • FE: Fixed Effect

## 50 1. Introduction

51 Given the significant heterogeneity between regions and territories and the selection of cattle breeds  
52 either for milk or meat production, there is a significant phenotypic and genetic diversity among  
53 cattle breeds in Europe. The meat production characteristics of these breeds is quite heterogeneous:  
54 conformation and fat cover of carcasses and meat yield vary greatly from one breed to another. The  
55 variation among breeds in size and carcass characteristics is well known (Albertí et al. 2008),  
56 however, the impact of the breed on the sensory quality of meat is still subject of debate, and  
57 therefore of interest.

58 Breed is a factor commonly taken into account when characterizing and studying sensory quality of  
59 meat. Significant differences in tenderness, juiciness and beef flavor intensities have been reported  
60 between Aberdeen Angus and Gasconne, Holstein and Simmental breeds (Bures et al. 2018). The  
61 tenderness of the Gasconne breed has been shown to be superior to that of the Holstein and  
62 Simmental breeds while the Angus breed has superior tenderness and juiciness compared to the Red  
63 Nordic (Huuskonen et al. 2016), although these breeds were not found to differ in beef flavor. The  
64 meat from the Charolais breed has been reported be more tender and have a higher and beef flavor  
65 than that from the Simmental and Eastern Anatolian Red, but not to have higher juiciness (Ozluturk  
66 et al. 2004). Avileña -Negra Ibérica breed is reported to have higher tenderness and beef flavor  
67 intensity than Bruna dels Pirineus and Morucha (Serra et al. 2008). Other studies, however, have  
68 reported no significant difference in beef characteristics between breeds e.g. Limousine vs Aberdeen  
69 Angus, (Pesonen et al. 2012) or Holstein vs Jersey (Nian et al. 2017). This raises the question as to  
70 whether the differences observed are variations among the breeds or a result of differences in  
71 management and feeding practices. Increasing ageing time of beef also seems to decrease  
72 differences in sensory quality between breeds. Better tenderness scores were obtained after ageing  
73 for one week for fast-growing breeds (Pirenaica and Rubia Gallega) compared with a double muscle  
74 Spanish breed (Asturiana de los Valles), a dual-purpose breed (Brown Swiss) and rustic breeds  
75 (Avileña-Negra Ibérica Morucha and Retinta), but the differences were no longer significant after 21  
76 days ageing (Campo et al. 1999).

77 Studies that have compared f beef eating quality between breeds have generally been limited in size,  
78 and in the number of breeds compared. While comparisons among studies is impaired by the  
79 different experimental protocols used. Work carried out so far has rarely studied more than 7 breeds  
80 under similar experimental conditions (Judge et al. 2021) and in larger studies comparisons could not  
81 be made, as rearing conditions were country specific (Panea et al. 2018). The differences observed  
82 between breeds are generally small and not always consistent between studies, which is most likely  
83 due to the different animal management and meat sensory protocols used. The European

84 consortium **GemQual (GEnetics of Meat Quality)** addressed these issues by studying the influence of  
85 breed on meat quality with all animals managed under similar husbandry conditions and as close as  
86 possible the same rearing conditions. The sensory quality of meat from young bulls from 15 European  
87 breeds was compared by testing in the same laboratories or by the same taste panels. **Data related**  
88 **to growth, carcass properties, muscle biochemical contents and physical measurements from these**  
89 **animals have been published (Alberti et al. 2008; Christensen et al. 2011) and these data are used**  
90 **here to test the grouping of breed by sensory parameters.**

91 The statistical treatment of the sensory data differs among studies, majority of which average scores  
92 provided by the panelists per animal before using them in statistical models (Pesonen et al. 2012 and  
93 Ozluturk et al. 2004 for example). Models used may include one or more fixed factors such as breed,  
94 diet, sex, etc., to assess their potential impact on meat quality. However, in other studies, sensory  
95 analysis is conducted using a statistical model that also include random effects, particularly to take  
96 into account variations in ratings between panelists. This approach is commonly used for food  
97 products, e.g. yoghurts (Saint-eve et al. 2006), cheese (Gierczynski et al. 2007) and wine (Vidal et al.  
98 2003), and has been used to study meat quality (Bures et al. 2018, Blanco et al. 2020). However, this  
99 is not the general case (Pesonen et al. 2012 and Ozluturk et al. 2004, Nian et al. 2017, Huuskonen et  
100 al. 2016).

101 The objective of this study was to evaluate the effect of breed on sensory quality of meat using the  
102 most appropriate statistical treatment. Two statistical treatments were tested, one of which included  
103 the effect of the panelist to assess if this would reduce the residual variability.

## 104 2. Materials & Methods

### 105 2.1. Experimental design

106 The study involved 436 **young bulls**, from 15 European breeds, reared under as similar conditions as  
107 possible (Albertí et al. 2008) in five European experimental stations: in France **(for the Limousin and**

108 Charolais breeds), the United Kingdom (for the Jersey, South Devon, Hereford, Aberdeen Angus and  
109 Galloway breeds), Spain (for the Casina, Avileña -Negra Ibérica, Pirenaica and Asturiana de los Valles  
110 breeds), Italy (for the Piemontese and Marchigiana breeds) and Denmark (for the Holstein, Danish  
111 Red and Simmental breeds). Animals with no direct relationship for two generations were selected to  
112 represent the genetic diversity of each breed. All management procedures were approved by the  
113 respective ethics committees of each research centre in accordance with the European Directive  
114 (U.S., 2010). The animals were slaughtered when they reached 75% of the average mature weight for  
115 their breed ( $15 \pm 1.3$  months of age, with a range from 398 and 511 days, Alberti et al., 2008). They  
116 were slaughtered by captive bolt pistol and exsanguination in commercial or experimental  
117 slaughterhouses depending on the infrastructure available in the different countries. Carcass  
118 dressing followed a standardized project protocol, without use of electrical stimulation, and with the  
119 removal of the remaining subcutaneous fat cover and testicles. Carcasses were split into two sides  
120 with tail on the right side of the carcass and chilled at  $4 \pm 1$  °C for 24 h. Temperature in the center of  
121 *M. longissimus thoracis*, at the 10<sup>th</sup> thoracic rib, was not allowed to fall below 10 °C within the first 10  
122 h. The *Longissimus thoracis* muscle was cut at 24h *post-mortem* between the 6<sup>th</sup> and 13<sup>th</sup> left ribs and  
123 vacuum-packed for maturation at  $2^{\circ}\text{C} \pm 1^{\circ}\text{C}$  for 10 days *post-mortem* and then frozen at  $-18^{\circ}\text{C}$  for  
124 preservation until analysis (Christensen et al. 2011).

## 125 2.2. Sensory analysis and texture measurements

126 Sensory assessment was as described by MacKintosh et al. (2017). In brief, after overnight thawing at  
127  $1^{\circ}\text{C}$ , 2 cm steaks were cooked under a conventional grill temperature turning every 2min until the  
128 internal temperature of the muscle reached  $74^{\circ}\text{C}$  as measured by a thermocouple probe (Testo  
129 Limited, Alton, UK). After cooking, cubes (2 x 2 x 1.9 cm) were then cut from the centre of the steak  
130 (avoiding incursions of connective tissue where present), one per panelist, wrapped in aluminum foil,  
131 coded with three-digit numbers, and kept warm for less than 10 min at  $55^{\circ}\text{C}$  before tasting took  
132 place. Samples were served hot to a 10-person trained professional taste panel, using the same  
133 people for the duration of each experiment.

134 Two panels composed of 10 trained panelists, one in the United Kingdom and the other in Spain,  
135 assessed the meat quality using 4 sensory descriptors: tenderness, juiciness, beef flavour and off-  
136 flavour, on an 8-point scale with 1 meaning less intense descriptor to 8 meaning most intense  
137 descriptor, as described by Wood et al. (1995). Each beef sample from between 29 and 31 animals  
138 per breed was assessed by 10 trained panelists, in sessions with 12 samples each in plates of 4  
139 samples, with random order between panelists to avoid first-order and carry-over effects.  
140 Assessments took place in a purpose-built panel room illuminated by red-light. Each booth contained  
141 a computer screen and optical mouse as part of the computerised sensory system, (Fizz, Version 2.20  
142 h, Biosystemes, Couternon, France), for direct entry of sensory responses. Assessors tasted the  
143 samples in an order based on the designs outlined by MacFie et al. (1989) for balancing carryover  
144 effects between samples.

145 A common set of samples were used for training and calibrating the panels. One extra sample was  
146 taken from 40 animals from 7 breeds that was tasted by both panels. This extra sample was used to  
147 compare the Spanish panel vs the UK panel to set the calibration values for each sensory attribute. A  
148 strong relationship was found between the scores although absolute values differed. Results for  
149 animals tested in common by both panels were used to derive a correction factor between both  
150 panels. Training consisted of 5 sessions of 8 samples each with discussion, reaching common  
151 agreement between panelists.

152 Values for Warner–Bratzler shear force and compressive force measured on raw and cooked meat  
153 obtained for samples from the same animals (Christensen et al. 2011) were used in the statistical  
154 analyses. For these measurements, frozen samples were thawed overnight and equilibrated to room  
155 temperature (25 °C) prior to texture analysis. For Warner–Bratzler shear force, slices were cooked in  
156 a water bath at 80 °C until the internal temperature reached 75 °C. The sample was then cooled for  
157 45 min in running tap water and stored at 4 °C until analysed. Shear force measurements for raw and  
158 cooked samples were performed on 10 blocks (2 cm in length and 1 cm by 1 cm of cross section) cut



159 perpendicular to the fiber direction. For the compression test, samples (1 cm<sup>2</sup> in cross section), were  
160 cut with muscle fibers, parallel to the longitudinal axis, and were analyzed using a modified  
161 compression device that avoids transversal elongation of the sample. Shear force and compression  
162 data are described in Christensen et al. (2011).

### 163 2.3. Statistical treatment

164 The statistical processing of sensory data compared a model that included a random effect to take  
165 into account the differences in scoring between panelists with a classical analysis of variance using  
166 the average scores for each sample.

167 The score  $y_{ijkl}^{(d)}$  of the descriptor  $d$  (tenderness, juiciness, beef flavor and off- flavor) for breed  $i$  given  
168 by the panelist  $j$  for the sample from animal  $k$  analyzed by the panel  $l$  can be decomposed by the  
169 following model:

$$170 \quad (1) \quad y_{ijkl}^{(d)} = \mu^{(d)} + \alpha_i^{(d)} + \beta_j^{(d)} + \gamma_{k(i)}^{(d)} + \lambda age_k + \delta_l^{(d)} + \varepsilon_{ijkl}^{(d)}$$

171 Where, for a given  $d$  descriptor,  $\mu^{(d)}$  is the average score of all animals,  $\alpha_i^{(d)}$  is the effect of breed  $i$ ,  
172  $\beta_j^{(d)}$  is the effect of panelist  $j$ ,  $\gamma_{k(i)}^{(d)}$  is the effect of animal  $k$  of breed  $i$ ,  $\lambda$  is the effect of age  
173 introduced as a continuous variable,  $\delta_l^{(d)}$  is the effect of panel  $l$  and  $\varepsilon_{ijkl}^{(d)}$  is a random error term  
174 whose distribution is assumed to be normal.

175 For simplicity, only the Spanish sensory panel data were used to compare the two modelling  
176 approaches with the presence or the absence of the  $\delta_l^{(d)}$  panel effect.

177 The first analysis of variance (ANOVA) model with mixed effects (ME) included all the effects of (1),  
178 with the exception of the panel effect. In this model, breed and age effects were considered as fixed  
179 effects because they were factors of interest in the study. Whereas the animal effect and the panel  
180 effect, resulting from the sampling process, were considered as random effects, i.e. characterized by  
181 a normal distribution (2):

182 
$$(2) \beta_j^{(d)} N(0, \sigma_a^{2(d)}) \text{ and } \gamma_{k(i)}^{(d)} N(0, \sigma_b^{2(d)})$$

183 where  $\sigma_a^{2(d)}$  and  $\sigma_b^{2(d)}$  are unknown variances and  $N(0, \sigma^2)$  is a normal distribution of mean 0 and  
184 standard variance  $\sigma^2$ .

185 The panelist random effect differs, however, from the animal random effect, which is nested (or  
186 hierarchical). Indeed, each panelist scored several beef samples of the  $I = 15$  breeds tested,  
187 whereas any  $k$  animal can only belong to a single  $i$  breed, hence the presence of the index  $i$  in the  
188 animal effect score namely  $k(i)$ .

189 To estimate the mixed effects (ME) parameters, the *lmer* function of the *lmerTest* package version  
190 3.1-0 was used with the REML optimization criterion (Kuznetsova et al., 2017) in R software version  
191 3.6.1.

192 The second analysis of variance (ANOVA) model had only fixed effects (FE), and is the most  
193 commonly used model in the literature to analyse meat quality sensory data:

194 
$$(3) y_{i.k}^{(d)} = \mu^{(d)} + \alpha_i^{(d)} + \lambda age_k + \varepsilon_{ik}^{(d)}$$

195 where  $y_{i.k}^{(d)}$  is the average of all the scores given by the  $J$  panelists for an animal  $k$ . For both models,  
196 the age of the animal at slaughter was introduced in the model as a continuous variable only when  
197 the effect was significant.

198 To evaluate the two models, we compared the p-values, testing the breed effect. The pairwise breed  
199 comparison was then tested using Tukey's *post-hoc* test when the previous null hypothesis was  
200 rejected. The *lsmean* function in the *emmeans* package (Lenth 2020) was used to perform this test,  
201 and a synthesis of significant differences was allowed by adding a superscript letter to the *cl*  
202 function in the *multcomp* package (Hothorn et al., 2008). A threshold of 5% was chosen to reject the  
203 H0 hypothesis.

204 Based on the previous comparison of the models, the breed effect was analyzed either by including a  
205 panelist random effect (ME) if the differences between the two models were significant, or based on  
206 the fixed effect model (FE) if no or little difference was observed between the models. This second  
207 choice is consistent with data in the literature and thus allowed the comparison of the results with  
208 published information. In the second analysis, the panel (country) effect was taken into account. As  
209 this effect has only two levels, it was not appropriate to model as a random effect with only two  
210 observations.

211 Finally, a multivariate analysis of all descriptors was conducted to identify similarities and differences  
212 between breeds. The averaged (lsmean) centered and reduced data were analyzed by hierarchical  
213 ascending classification (CAH), using the "Ward.D" agglomeration criterion.

214 A Principal Component Analysis (PCA) was then carried out according to Destefanis et al. (2000) to  
215 assess correlations between the **sensory scores** and the mean values of shear force and compressive  
216 force measured on raw and cooked meat previously obtained on the same samples (Christensen et  
217 al. 2011), which were included as supplementary variables.

## 218 3. Results

### 219 3.1. Statistical model comparison

220 Each of the 4 descriptors used for the sensory quality of meat discriminated at least one breed from  
221 the others, independent of the statistical model used (Table 1). The tenderness descriptor was had  
222 the lowest p-value discriminating breeds.

223 *Table 1: P-value testing the null hypothesis H0 of equality of the mean of each breed with the two models tested.*

Descriptors	<i>Mixed Effect Anova</i>	<i>Fixed Effect Anova</i>
Tenderness	8.5e-06	9.2e-06
Juiciness	1.9e-04	3.2e-04

Beef flavor	2.6e-03	2.5e-03
Off-flavor	2.8e-04	3.6e-04

224

225 Few differences were found between the two statistical approaches, although the model that  
 226 included panelist as a random effect (ME) had slightly lower p-values, on average, than those using  
 227 averaged data (FE), except for beef flavour (Table 1). Taking into account the panelist effect did not  
 228 change the discrimination between breeds, as p-values observed for both models were similar.

229 Pairwise comparisons of breeds showed that only the most significant differences changed between  
 230 the two models (Table 2). The Simmental breed differed significantly from the Pirenaica for  
 231 tenderness and juiciness with the model including a panelist random effect (ME) with p-values of  
 232 0.049 and 0.047 respectively, whereas for the model without the panelist effect, p-values of 0.065  
 233 and 0.058 were obtained. As the model based on mean scores (FE) is widely used, and does not bias  
 234 analysis of meat sensory quality data, it was used to assess the effect of breed on meat sensory  
 235 quality in subsequent analyses. Animal age (which varied from 398 and 511 days) had a significant  
 236 effect on beef flavour (p-value=0.019) but was not significant for the other descriptors. Age was  
 237 therefore omitted in the ANOVA analysis (Table 3).

238 The first test (Table 2) was done on the sensory scores from the Spanish panel only whereas later  
 239 analyses reported in Table 3 used all the animals. Apart from the English breeds, for which sensory  
 240 analysis was only done in the United Kingdom, the other breeds were tested in both the United  
 241 Kingdom and Spain. This may explain the differences obtained between the two tests.

242 *Table 2 : P-value of the Tukey test comparing the breed Pirenaica and Simmental with the two statistical models*

Descriptors	Mixed Effect Anova	Fixed Effect Anova
Tenderness	0.049	0.065
Juiciness	0.047	0.058

Beef flavor	0.003	0.007
Off-flavor	0.999	0.999

243

244 *Table 3 : Breed comparison based on LSMEAN ( $\pm$  standard deviation) for each descriptor of sensory quality of the meat*

Race	Tenderness	Juiciness	Beef flavor	Off-flavor
Aberdeen Angus	4.6 $\pm$ 1.0 <sup>abc</sup>	5.3 $\pm$ 0.9 <sup>e</sup>	5.5 $\pm$ 1.2 <sup>d</sup>	3.4 $\pm$ 1.8
Asturiana de los Valles	4.9 $\pm$ 1.2 <sup>bc</sup>	5.0 $\pm$ 0.8 <sup>bcde</sup>	4.5 $\pm$ 0.9 <sup>ab</sup>	3.2 $\pm$ 0.6
Avileña Negra Ibérica	5.0 $\pm$ 1.1 <sup>bc</sup>	4.8 $\pm$ 0.9 <sup>abcde</sup>	4.4 $\pm$ 0.9 <sup>ab</sup>	3.0 $\pm$ 0.5
Casina	4.0 $\pm$ 1.5 <sup>ab</sup>	4.5 $\pm$ 0.8 <sup>abcd</sup>	4.5 $\pm$ 0.7 <sup>ab</sup>	3.0 $\pm$ 0.5
Charolaise	4.8 $\pm$ 1.0 <sup>c</sup>	5.0 $\pm$ 0.7 <sup>bcde</sup>	4.3 $\pm$ 1.1 <sup>ab</sup>	2.8 $\pm$ 0.6
Danish Red cattle	4.7 $\pm$ 0.9 <sup>abc</sup>	4.6 $\pm$ 0.5 <sup>abcde</sup>	4.5 $\pm$ 0.7 <sup>ab</sup>	2.9 $\pm$ 0.5
Highland	4.1 $\pm$ 1.0 <sup>ab</sup>	5.0 $\pm$ 0.9 <sup>bcde</sup>	5.2 $\pm$ 1.2 <sup>cd</sup>	3.3 $\pm$ 2
Holstein	4.6 $\pm$ 0.9 <sup>abc</sup>	4.4 $\pm$ 0.6 <sup>ab</sup>	4.5 $\pm$ 0.7 <sup>ab</sup>	2.9 $\pm$ 0.5
Jersey	4.9 $\pm$ 1.79 <sup>bc</sup>	5.0 $\pm$ 0.7 <sup>bcde</sup>	4.8 $\pm$ 1.0 <sup>bc</sup>	3.6 $\pm$ 1.9
Limousin	4.8 $\pm$ 0.9 <sup>bc</sup>	5.0 $\pm$ 0.8 <sup>bcde</sup>	4.2 $\pm$ 1.0 <sup>a</sup>	2.9 $\pm$ 0.5
Marchigiana	3.8 $\pm$ 1.3 <sup>a</sup>	4.3 $\pm$ 0.8 <sup>abc</sup>	4.3 $\pm$ 0.9 <sup>ab</sup>	2.9 $\pm$ 0.6
Piemontese	4.9 $\pm$ 1.0 <sup>bc</sup>	4.7 $\pm$ 0.7 <sup>abcde</sup>	4.2 $\pm$ 0.9 <sup>ab</sup>	3.2 $\pm$ 0.6
Pireneica	5.3 $\pm$ 1.0 <sup>c</sup>	5.0 $\pm$ 0.9 <sup>de</sup>	4.5 $\pm$ 0.9 <sup>ab</sup>	3.1 $\pm$ 0.5
Simmental	3.7 $\pm$ 1.3 <sup>a</sup>	4.2 $\pm$ 0.7 <sup>l</sup>	4.2 $\pm$ 0.8 <sup>ab</sup>	3.0 $\pm$ 0.6
South Devon	4.9 $\pm$ 0.7 <sup>bc</sup>	5.1 $\pm$ 0.4 <sup>cde</sup>	4.8 $\pm$ 1.4 <sup>abcd</sup>	2.9 $\pm$ 1.4

245 *For a given descriptor, two breeds are significantly different from each other if none of the*  
 246 *superscripts letters between brackets are identical. Scores range from the least (1) to the highest (8)*  
 247 *intensity.*

248


### 249 3.2. Impact of cattle breed on sensory meat quality

250

251 The hierarchical classification of the sensory profile of the meat from the 15 breeds revealed that  
252 they form five groups (Figure 1).

253 **Aberdeen Angus, Highland and Jersey fall** into the first group (Figure 1). These breeds are  
254 characterized by high beef flavour intensity and juiciness (Table 3). Breeds in the second group  
255 include a dual-purpose and two rustic breeds (Simmental, Casina and Marchigiana) which are  
256 characterized by the toughest and the driest meat. The dairy breeds, Holstein and Danish Red cattle,  
257 form group 3, and are characterized by intermediate juiciness and tenderness scores (Figure 2). The  
258 double muscled breeds, Asturiana de los Valles and Piemontese and fast-growing beef breeds  
259 Pirenaica and Avileña Negra Ibérica, group 4, are characterized by a tender and juicy meat. The highly  
260 specialized French beef breeds (group 5), Limousine and Charolaise, have similar sensory scores and  
261 the lowest beef flavor intensity (Table 3).


262 Principal Component Analysis (PCA) that included previously published mean values of shear force  
263 and compressive force measured on raw and cooked meat (Christensen et al., 2011) showed three  
264 distinct groups (Figure 2) corresponding to the five discriminatory associations on the first two  
265 dimensions, which explain 84% of the variation. The groups which were closest together in the  
266 hierarchical agglomerative clustering (groups 3, 4 and 5) are also grouped together in the PCA.  
267 Breeds which produce the juiciest and the most tender beef (scores > 4.8), including Asturiana de los  
268 Valles, Avileña Negra Ibérica and South Devon are grouped in the upper right-hand corner of the  
269 PCA. Breeds producing beef with both intense beef flavour (> 4.8) and off flavor (> 3.3), Jersey,  
270 Aberdeen Angus, and Highland (Group 1), are grouped in the lower right-hand side (Figure 2)  
271 whereas breeds with the lowest beef tenderness scores, Casina, Marchigiana, Simmental (Group 2)  
272 are grouped in the lower left-hand side.



273

274 *Figure 1 Dendrogram of the hierarchical agglomerative clustering (HAC) of breed based on the scale mean data (lsmean)*

275 *given in the table 3, where five clusters were from each other.*




276

277 *Figure 2: Principal component analyses showing variability between breeds in sensory analysis descriptors (presented in*  
 278 *Table 3 and in Figure 1) and in mechanical descriptors (shear force and compressive force measured on raw and cooked*  
 279 *meat published by Christensen et al., 2011). The breed associations produced by the hierarchical ascending classification are*  
 280 *represented by symbols.*

281 Based on the Principal Component Analysis, a significant correlation was found between tenderness  
 282 and juiciness ( $r=0.63$ ,  $p\text{-value}=0.01$ ) (Figure 3). The negative correlation observed between  
 283 compression values and tenderness was not significant ( $r=-0.21$ ,  $p\text{-value}=0.44$ ), nor was the  
 284 correlation between raw shear force and tenderness ( $r=-0.33$ ,  $p\text{-value}=0.22$ ). However, the negative  
 285 correlation observed between tenderness score and shear force for cooked meat was significant ( $r=-$



286 0.60, p-value=0.02). This indicates that meat texture is affected by the cooking process. A positive  
287 correlation between beef flavor and off-flavor was also observed ( $r=0.60$ , p-value=0.02) meaning that  
288 the breeds with the highest beef flavor intensity also had the highest off-flavor, which may be  
289 because the sensory panels assess both of these descriptors in a similar way.



290  
291 *Figure 3: Correlation circle of sensory analysis descriptors (Table 3), illustrated by black arrows, combined with previously*  
292 *published rheological data (Christensen et al. 2011) as supplementary variables (in dotted arrows).*

293

## 294 4. Discussion

295

### 296 4.1. Breed clustering is trait-dependent

297 Hierarchical ascending classification placed breeds into groups according to their sensory traits, as  
298 did a Principal Component Analysis similar to those used in other studies (Destefanis et al. 2000).  
299 These analyses indicated the impact of the breed on the variations in meat sensory characteristics.  
300 This information can be used compare the relative standards of one breed with respect to other  
301 breeds, and by the whole supply chain to choose the breed that best suits their needs and  
302 consumers' expectations.

303 **Animal and carcass characteristics and their classification according to the European carcass (EUROP)**  
304 **grid placed the breeds assessed in the GemQual study, into 3 groups** (Albertí et al. 2008). Using  
305 muscle and mechanical characteristics of beef produced by these breeds and **on the chemical**  
306 **analysis of samples from the same animals**, the dairy breeds, Danish Red Cattle, Holstein, Jersey,  
307 were grouped together as they had the highest total and insoluble collagen contents (Christensen et  
308 al. 2011) whereas doubled muscle, Piemontese, Asturiana de los Valles, and the Limousine breeds  
309 have the lowest contents. Aberdeen Angus, Highland and the dairy breeds have been shown to have  
310 the highest intramuscular lipid content whereas Piemontese, Limousine and Asturiana de los Valles  
311 had the lowest (Christensen et al. 2011). From the classification of breeds based on animal and  
312 carcass characteristics, or muscle and beef characteristics (Albertí et al., 2008), it was expected that  
313 breeds would differ significantly for sensory traits. However, this study found only minor differences  
314 between breeds, which were much lower than expected.

315 The small differences **in sensory scores** among breeds maybe be due to the within breed **variability**  
316 **being high and that the subjective nature of evaluating** sensory traits leading to greater variation in  
317 values when compared with animal, carcass, muscle and beef characteristics. Results from sensory  
318 panels are known to be highly variable (Gagaoua et al., 2016a) which makes standardization and

319 accuracy of measuring the phenotype difficult, which is essential to investigate variation (Hocquette  
320 et al., 2012). In this study, we compared two statistical models to assess the technical variability  
321 associated with the experimental design. This comparison showed that the inclusion of panelist as  
322 factor does not improve significantly the robustness of the statistical model, and does not allow the  
323 detection of additional differences between breeds than the model without panelist as a factor.

324 The contribution of animal, carcass and muscle characteristics to explain the variability in sensory  
325 scores is known to be low or moderate as has been seen for carcass traits (Judge et al., 2021), the  
326 EUROP grid scores (Bonny et al., 2016a), marbling (Liu et al, 2020), muscle biochemical traits  
327 (Gagaoua et al., 2016b) or mechanical measurements (Destefanis et al., 2008).

328

#### 329 4.2. Muscle growth potential and muscle characteristics of breeds have some impact on 330 sensory quality

331 This work found that fast growing breeds such as Pirenaica (Campo et al. 1999) as well as the double-  
332 muscled breeds produce the most tender beef. This may be related to proportion of total and  
333 insoluble collagen, which is higher when muscle mass is low which contributes to beef toughness  
334 (Purslow, 2005), while low total and insoluble collagen contents are known to increase tenderness  
335 scores (Chriki et al., 2012). Higher collagen content per g of tissue has been reported in the  
336 longissimus muscle from Angus compared to Limousin steers (Chambaz et al., 2003) and in Limousin  
337 compared to double-muscled Belgian Blue cattle (Raes et al., 2003). Other differences in sensory  
338 traits between breeds may be explained by differences in intramuscular fat content. The high  
339 intramuscular lipid content of the Aberdeen Angus, Highland and Jersey breeds could explain in part  
340 the high flavour, whereas the low intramuscular fat content of the Limousin breed could be related  
341 to the less intense beef flavour(Gagaoua et al., 2016b). Differences in lipid, intramuscular collagen  
342 content and also in fibre type are thought to explain differences in sensory quality of beef (Chriki et

343 al. 2012, Chriki et al. 2013), although. there is inconsistency in these differences associated with  
344 muscle, animal type and the productive function related to the breed (dairy or meat type).

345 Tenderness of meat from Simmentals has been reported to be lower than that of other breeds  
346 (Shackelford et al., 1994; Chambaz et al., 2003; Zwambag et al., 2013, Xie et al. 2012) which is in  
347 agreement with our results. Interestingly, Chambaz et al. (2003) reported that the Simmental breed  
348 produced less tender meat than Angus and Limousin breeds when slaughtered at the same level of  
349 intramuscular fat. Beef from Angus has been reported to be more tender, juicy and flavorsome than  
350 that of Holsteins (Bures and Barton, 2018), which is also in agreement with our data.

351

#### 352 4.3. Knowing breed characteristics might be important at the consumer end

353 Comparing breeds based on their sensory characteristics is a factor, among others, on which  
354 consumers choose the beef. However, comparing breeds is difficult as there is little information  
355 available. Collecting reliable information has been hampered by animals being reared in different  
356 ways. Cattle management differs among breeds and is dependent both on breed, resources and  
357 practices in different regions. Breeds that originated in specific regions where particular feed,  
358 resources and condition were available, are now found internationally. Many studies have compared  
359 different types of cattle and systems, confounding breeds, sexes and management systems (Gagaoua  
360 et al., 2016b) rather than strictly comparing the breeds. Compare breed-management effects on beef  
361 sensory quality is important, but must be explained to the consumer (Panea et al. 2018). In order to  
362 make an unbiased comparison between a large number of breeds, the GemQual project established  
363 protocols that were as standardized as possible with entire bulls only. Nevertheless, some biases are  
364 likely to have persist. In addition to differences in management and breed, it is necessary to recruit  
365 members of sensory panels that give consistent results. Some descriptors have been shown to have  
366 moderate correlation, e.g. tenderness  $r=0.67$ , juiciness  $r=-0.14$ , flavour  $r=0.1$ , abnormal flavour  $r=0.2$ .  
367 Meta-analyses of several studies are also possible, although this requires that the diversity of

368 protocols is accounted for in the analysis. A recent work showed that standardization of scoring  
369 scales is only partially successful and a random effect associated with the experiment is necessary in  
370 the analysis model (Judge et al. 2021) Other sources of variability such as meat ageing time and sex  
371 of the animal make it difficult to determine of the contribution of breed to differences in sensory  
372 quality. Increasing the ageing period, reduces difference in meat between breeds. It has been shown  
373 that with 21 days aging differences between breeds cannot be detected (Campo et al. 1999). In the  
374 GemQual study, the meat was aged for 10-days, at which time the breed effects on sensory quality of  
375 beef should still have been observed.

376 At the consumer level, availability of information on breed and sex of the animal and on  
377 management, such as feeding is important as these have factors influence consumer assessments,  
378 irrespective of sensory quality. Generally, meat from traditional beef breeds is better appreciated by  
379 consumers over that from dairy breeds, although little differences in eating quality have been found  
380 for beef from dairy and beef breeds by untrained consumers (Bonny et al., 2016b).

## 381 5. Conclusion

382 We have shown that there is little variability in sensory quality of beef of young bulls from the  
383 diversity of European cattle breeds despite significant differences in animal, carcass, muscle and beef  
384 characteristics of the same animals. The choice of analysis methods, using averaged sensory values or  
385 including panelist as a random effect in a mixed model gave similar results.

386 Five groups were observed for the 15 breeds studied based on meat sensory attributes. The breeds  
387 having high lipid content are characterized by superior beef flavour intensity. The rustic breeds  
388 produce meat with lower tenderness and juiciness. Double muscled, fast growing and meat-type  
389 breeds tend to produce more tender meat. Breed groups formed based on sensory quality traits  
390 could help the consumers to choose among them depending on their quality expectations.

## 391 6. Ethics statement

392 All animal work was reviewed and approved by the ethics committees of the organizations involved.

## 393 Acknowledgements

394 Authors would like to thank Drs Carlos Sañudo, Geoff Nute, Pere Albertí for their valuable scientific  
395 contribution.

## 396 CRediT authorship contribution statement

397 Conanec Alexandre: Analysis, Writing-original draft. Ellies-Oury Marie-Pierre: Investigation, Writing -  
398 review & editing. Hocquette Jean-François: Investigation, Writing - review & editing. Chavent Marie:  
399 Methodology, Writing - review & editing. Saracco Jérôme: Methodology, Writing - review & editing.  
400 Campo Marimar and Ian Richardson: Investigation, Writing - review & editing. Others: Investigation,  
401 review & editing. John Williams project design and management.

## 402 Declaration of Competing Interest

403 The authors declare to have no conflicts of interest.

## 404 References

- 405 Albertí, P., Panea, B., Sañudo, C., Olleta, J. L., Ripoll, G., Ertbjerg, P., Christensen, M., Gigli, S., Failla, S.  
406 & Concetti, S. (2008). Live weight, body size and carcass characteristics of young bulls of fifteen  
407 European breeds, *Livestock Science* 114(1), 19–30. <https://doi.org/10.1016/j.livsci.2007.04.010>.
- 408 Blanco, M., Ripoll, G., Delavaud, C. & Casasús, I. (2020). Performance, carcass and meat quality of  
409 young bulls, steers and heifers slaughtered at a common body weight, *Livestock Science* p. 104156.  
410 <https://doi.org/10.1016/j.livsci.2020.104156>.
- 411 Bonny, S. P. F., Hocquette, J.-F., Pethick, D. W., Farmer, L. J., Legrand, I., Wierzbicki, J., Allen, P.,  
412 Polkinghorne, R. J. & Gardner, G. E. (2016). The variation in the eating quality of beef from different  
413 sexes and breed classes cannot be completely explained by carcass measurements, *Animal* 10(6),  
414 987–995. <https://doi.org/10.1017/S175173111500292X>.
- 415 Bureš, D. & Bartoň, L. (2018). Performance, carcass traits and meat quality of Aberdeen Angus,  
416 Gascon, Holstein and Fleckvieh finishing bulls, *Livestock Science* 214: 231–237.  
417 <https://doi.org/10.1016/j.livsci.2018.06.017>.
- 418 Campo, M. M., Sanudo, C., Panea, B., Albertí, P. & Santolaria, P. (1999). Assessment of breed type  
419 and ageing time effects on beef meat quality using two different texture devices. *Meat Science*, 55,  
420 371–378. [https://doi.org/10.1016/S0309-1740\(99\)00162-X](https://doi.org/10.1016/S0309-1740(99)00162-X).
- 421 Campo, M. M., Santolaria, P., Sanudo, C., Lepetit, J., Olleta, J. L., Panea, B. & Albertí, P. (1999). Breed  
422 type and ageing time effects on sensory characteristics of beef strip loin steaks. *Meat Science*, 51,  
423 383–390. [https://doi.org/10.1016/S0309-1740\(98\)00159-4](https://doi.org/10.1016/S0309-1740(98)00159-4).
- 424 Chambaz, A., Scheeder, M. R. L., Kreuzer, M., & Dufey, P.-A. (2003). Meat quality of Angus,  
425 Simmental, Charolais, and Limousine steers compared at the same intramuscular fat content. *Meat*  
426 *Science*, 63, 491–500.

427 Chriki, S., Gardner, G. E., Jurie, C., Picard, B., Micol, D., Brun, J.-P., Journaux, L. & Hocquette, J.-F.  
428 (2012). Cluster analysis application identifies muscle characteristics of importance for beef  
429 tenderness, *BMC Biochemistry* 13(1): 29. <https://doi.org/10.1186/1471-2091-13-29>.

430 Chriki, S., Renand, G., Picard, B., Micol, D., Journaux, L. & Hocquette, J.-F. (2013). Meta-analysis of the  
431 relationships between beef tenderness and muscle characteristics, *Livestock Science* 155(2-3): 424–  
432 434. <https://doi.org/10.1016/j.livsci.2013.04.009>.

433 Christensen, M., Ertbjerg, P., Failla, S., Sañudo, C., Richardson, R. I., Nute, G. R., Olleta, J. L., Panea, B.,  
434 Albertí, P. & Juárez, M. (2011). Relationship between collagen characteristics, lipid content and raw  
435 and cooked texture of meat from young bulls of fifteen European breeds, *Meat Science* 87(1): 61–65.  
436 <https://doi.org/10.1016/j.meatsci.2010.09.003>.

437 Destefanis G, Barge MT, Brugiapaglia A, Tassone S. (2000). The use of principal component analysis  
438 (PCA) to characterize beef. *Meat Science* 56, 255–259.

439 Destefanis G., Brugiapaglia A., Barge M.T., Dal Molin E. (2008). Relationship between beef consumer  
440 tenderness perception and Warner–Bratzler shear force. *Meat Science* 78, 153–156.

441 Dransfield, E., Martin, J.-F., Bauchart, D., Abouelkaram, S., Lepetit, J., Culioli, J., Jurie, C. & Picard, B.  
442 (2003). Meat quality and composition of three muscles from French cull cows and young bulls,  
443 *Animal Science* 76(3): 387–399. <https://doi.org/10.1017/S1357729800058616>.

444 Gagaoua, M., Micol, D., Picard, B., Terlouw, C. E., Moloney, A. P., Juin, H., Météau, K., Scollan, N.,  
445 Richardson, I. & Hocquette, J.-F. (2016a). Interlaboratory assessment by trained panelists from  
446 France and the United Kingdom of beef cooked at two different end-point temperatures, *Meat*  
447 *Science* 122: 90–96. <https://doi.org/10.1016/j.meatsci.2016.07.026>.

448 Gagaoua M., Terlouw E.M.C., Micol D., Hocquette J-F., Moloney A.P., Nuernberg K., Bauchart D.,  
449 Boudjellal A., Scollan N.D., Richardson R.I., Picard B., (2016b). Sensory quality of meat from eight  
450 different types of cattle in relation with their biochemical characteristics *Journal of Integrative*  
451 *Agriculture*, 15, 1550–1563. <http://prodinra.inra.fr/record/360882>

452 Gierczynski, I., Labouré, H., Sémon, E. & Guichard, E. (2007). Impact of hardness of model fresh  
453 cheese on aroma release: in vivo and in vitro study, *Journal of Agricultural and Food Chemistry* 55(8):  
454 3066–3073. <https://doi.org/10.1021/jf0633793>.

455 Hocquette J.F., Capel C., David V., Guéméné D., Bidanel J., Ponsart C., Gastinel P.L., Le Bail P.Y.,  
456 Monget P., Mormède P., Barbezant M., Guillou F, Peyraud J.L., 2012. Objectives and applications of  
457 phenotyping network set-up for livestock. *Animal Science Journal*, 83, 517–528.

458 Hothorn, T., Bretz, F. & Westfall, P. (2008). Simultaneous inference in general parametric models,  
459 *Biometrical Journal* 50(3): 346–363. <https://doi.org/10.1002/bimj.200810425>.

460 Huuskonen, A. K., Pesonen, M. & Honkavaara, M. (2016). Performance and meat quality of Nordic  
461 Red and Aberdeen Angus bulls offered faba bean or field pea based whole crop legume-cereal  
462 silages, *Agricultural and Food Science* 25(1): 1–12. <https://doi.org/10.23986/afsci.52311>.

463 Judge, M., Conroy, S., Hegarty, P., Cromie, A., Fanning, R., Kelly, D., Croften, E. & Berry, D. (2021).  
464 Eating quality of the longissimus thoracis muscle in beef cattle—Contributing factors to the underlying  
465 variability and associations with performance traits, *Meat Science* 172, 108371.  
466 <https://doi.org/10.1016/j.meatsci.2020.108371>

467 Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. (2017). lmerTest package: Tests in linear  
468 mixed effects models, Journal of Statistical Software 82(13): 1–26.  
469 <http://dx.doi.org/10.18637/jss.v082.i13>.

470 Lenth, R. (2020). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version  
471 1.4.5. URL: <https://CRAN.R-project.org/package=emmeans>

472 Liu J., Chriki S., Ellies-Oury M.P., Legrand I., Pogorzelski G., Wierzbicki J., Farmer L., Troy D.,  
473 Polkinghorne R., Hocquette J.F. (2020). European conformation and fat scores of bovine carcasses  
474 are not good indicators of marbling. Meat Science, 170, 108233.

475 MacFie H.J., Bratchell N., Greenhoff K. and Vallis L.V. (1989). Designs to balance the effect of order of  
476 presentation and first-order carry-over effects in hall tests. Journal of Sensory Studies 4, 129-148.

477 MacKintosh S.B., Richardson I., Kim E.J., Dannenberger D., Coilmier D. and Scollan N.D. (2017).  
478 Addition of an extract of Lucerne (*Medicago sativa* L.) to cattle diets – Effects on fatty acid profile,  
479 meat quality and eating quality of the *M. longissimus* muscle. Meat Science 130: 69-80 (2017).

480 Monsón, F., Sañudo, C. & Sierra, I. (2005). Influence of breed and ageing time on the sensory meat  
481 quality and consumer acceptability in intensively reared beef. Meat Science 71(3): 471–479.  
482 <https://doi.org/10.1016/j.meatsci.2005.04.026>.

483 Nian, Y., Kerry, J. P., Prendiville, R. & Allen, P. (2017). The eating quality of beef from young dairy  
484 bulls derived from two breed types at three ages from two different production systems, Irish Journal  
485 of Agricultural and Food Research 56(1): 31–44. <http://dx.doi.org/10.1515/ijafr-2017-0003>.

486 Özlütürk, A., Tüzemen, N., Yanar, M., Esenbuga, N. & Dursun, E. (2004). Fattening performance,  
487 carcass traits and meat quality characteristics of calves sired by Charolais, Simmental and Eastern  
488 Anatolian Red sires mated to Eastern Anatolian Red dams, Meat Science 67(3): 463–470.  
489 <https://doi.org/10.1016/j.meatsci.2003.11.022>.

490 Panea, B., Olleta, J., Sañudo, C., Campo, M., Olivier, M., Gispert, M., Serra, X., Renand, G., Oliván,  
491 M.C. & Jabet, S. (2018). Effects of breed-production system on collagen, textural, and sensory traits  
492 of 10 European beef cattle breeds, Journal of texture studies 49(5), 528–535.  
493 <https://doi.org/10.1111/jtxs.12350>

494 Pesonen, M., Honkavaara, M. & Huuskonen, A. K. (2012). Effect of breed on production, carcass traits  
495 and meat quality of Aberdeen Angus, Limousin and Aberdeen Angus x Limousin bulls offered a grass  
496 silage-grain-based diet, Agricultural and Food Science 21(4), 361–369.  
497 <https://doi.org/10.23986/afsci.6520>.

498 Purslow, P. P. (2005). Intramuscular connective tissue and its role in meat quality. Meat Science 70,  
499 435–447.

500 Raes, K., Balcaen, A., Dirinck, P., De Winne, A., Claeys, E., Demeyer, D., et al. (2003). Meat quality,  
501 fatty acid composition and flavor analysis in Belgian retail beef. Meat Science, 65, 1237–1246.

502 Renand, G., Picard, B., Touraille, C., Berge, P. & Lepetit, J. (2001). Relationships between muscle  
503 characteristics and meat quality traits of young Charolais bulls, Meat Science 59(1): 49–60.  
504 [https://doi.org/10.1016/S0309-1740\(01\)00051-1](https://doi.org/10.1016/S0309-1740(01)00051-1).

505 Saint-Eve, A., Martin, N., Guillemin, H., Sémon, E., Guichard, E. & Souchon, I. (2006). Flavored yogurt  
506 complex viscosity influences real-time aroma release in the mouth and sensory properties, Journal of  
507 Agricultural and Food Chemistry 54(20), 7794–7803. <https://doi.org/10.1021/jf060849k>.



508 Shackelford, S. D., Koohmaraie, M., Cundiff, L. V., Gregory, K. E., Rohrer, G. A., & Savell, J. W. (1994).  
509 Heritabilities and phenotypic and genetic correlations for bovine postrigor calpastatin activity,  
510 intramuscular fat content, Warner-Bratzler shear force, retail product yield, and growth rate. *Journal*  
511 *of Animal Science*, 72, 857–863.

512 Serra, X., Guerrero, L., Guàrdia, M., Gil, M., Sañudo, C., Panea, B., Campo, M., Olleta, J, García-  
513 Cachán, M. & Piedrafita, J. (2008). Eating quality of young bulls from three Spanish beef breed-  
514 production systems and its relationships with chemical and instrumental meat quality, *Meat Science*  
515 79(1), 98-104. <https://doi.org/10.1111/jtxs.12350>

516 Vidal, S., Francis, L., Guyot, S., Marnet, N., Kwiatkowski, M., Gawel, R., Cheynier, V. & Waters, E. J.  
517 (2003). The mouth-feel properties of grape and apple proanthocyanidins in a wine-like medium,  
518 *Journal of the Science of Food and Agriculture* 83(6): 564–573. <https://doi.org/10.1002/jsfa.1394>.

519 Wood, J., Nute, G., Fursey, G. & Cuthbertson, A. (1995). The effect of cooking conditions on the  
520 eating quality of pork, *Meat Science* 40(2): 127-135. [https://doi.org/10.1016/0309-1740\(94\)00051-8](https://doi.org/10.1016/0309-1740(94)00051-8)

521 Zwambag, A., Kelly, M., Schenkel, F., Mandell, I., Wilton, J., & Miller, S. (2013). Heritability of beef  
522 tenderness at different aging times and across breed comparisons. *Canadian Journal of Animal*  
523 *Science.*, 93, 307–312.

524

525