



HAL
open science

Seroprevalance of *Trichinella* spp. in wild boars (*Sus scrofa*) from Bihor county, western Romania

Zsolt Boros, Isabelle Vallée, Luciana-Catalina Panait, Calin M. Gherman, Aurélie Chevillot-Grasset, Pascal Boireau, Vasile Cozma

► **To cite this version:**

Zsolt Boros, Isabelle Vallée, Luciana-Catalina Panait, Calin M. Gherman, Aurélie Chevillot-Grasset, et al.. Seroprevalance of *Trichinella* spp. in wild boars (*Sus scrofa*) from Bihor county, western Romania. *Helminthologia*, 2020, 57 (3), pp.235-240. 10.2478/helm-2020-0032 . hal-03316855

HAL Id: hal-03316855

<https://hal.inrae.fr/hal-03316855>

Submitted on 6 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Seroprevalance of *Trichinella* spp. in wild boars (*Sus scrofa*) from Bihor county, western Romania

Z. BOROS^{1,*}, I. VALLÉE², L. C. PANAIT¹, C. M. GHERMAN¹, A. CHEVILLOT², P. BOIREAU², V. COZMA^{1,3†}

¹University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Faculty of Veterinary Medicine, Department of Parasitology and Parasitic Diseases, 3-5 Mănăstur street, 400372, Cluj-Napoca, Romania, E-mail: *zsolt.boros@usamvcluj.ro, luciana.rus@usamvcluj.ro, calin.gherman@usamvcluj.ro, vasile.cozma@usamvcluj.ro; ²UMR, BIPAR, ANSES, ENVA, INRA, Laboratory for Animal Health, Street 14 Pierre and Marie Curie, F-94 701 Maisons-Alfort, France, E-mail: isabelle.vallee@anses.fr, aurelie.chevillot@anses.fr, pascal.boireau@anses.fr; ³Academy of Agricultural and Forestry Sciences Gheorghe Ionescu-Sisești (A.S.A.S), 61 Mărăști Boulevard, 011464, Bucharest, Romania

Article info

Received February 6, 2020
 Accepted April 28, 2020

Summary

The wild boar (*Sus scrofa*) has a wide geographical distribution and can be an important source of *Trichinella* spp. infection in humans in Romania.

The objective of this study was to identify the presence of *Trichinella* spp. in the wild boar population in Bihor County, Romania.

Eighty four plasma and diaphragm samples, collected from wild boars, were included in this study. Artificial digestion, ELISA and Western blot were performed on these specimens. All diaphragm samples were negative for *Trichinella* larvae in artificial digestion, while in ELISA, 54 (64.2 %) plasma samples were positive and 6 (7.1 %) plasma samples were doubtful. Western blot was performed on 26 plasma samples from which only 6 (23.0 %) gave a positive result.

Serological evidences indicate the presence of *Trichinella* spp. in wild boars from western Romania. Therefore, human consumers might be at risk to ingest *Trichinella* larvae, even in low numbers.

Keywords: Wild boars; Romania; artificial digestion; ELISA; Western blot

Introduction

The wild boar (*Sus scrofa*) has one of the widest geographical distributions of all terrestrial wild mammals with increasing populations in all European countries (Massei & Genov, 2004). Moreover, they are considered to be an important source of *Trichinella* spp. infection in humans (Boadella *et al.*, 2011). In Romania, trichinellosis, is considered “a never-ending story” because of the local traditions and culinary customs that favor the parasites transmission to humans. Especially the consumption of undercooked meat and meat products has led to various outbreaks over the years (Neghina, 2010).

The magnetic stirrer digestion is the gold standard method for the direct detection of *Trichinella* larvae in meat samples (Gamble *et*

al., 2000). This method is mandatory for meat inspection according to EU regulation (EU 2015/1375). Enzyme linked immunosorbent assay (ELISA) using excretory-secretory (ES) antigen is a highly sensitive test for the detection of circulating antibodies against *Trichinella* spp. in blood samples from pigs and wild boars (Kapel & Gamble, 2000; Kapel, 2001). On the other hand, Western blot allows the identification of *Trichinella* infection by highlighting bands characteristic of the genus (Cuttell *et al.*, 2014). However, this method is time consuming and is mostly used as a confirmatory assay. Successful *Trichinella* spp. diagnosis and controls are important for ensuring food safety and for preventing future outbreaks (Gajadhar *et al.*, 2009).

Trichinelloscopy and artificial digestion were previously used for the identification of *Trichinella* spp. in wild animals in Romania

* – corresponding author




Fig. 1. The map of Romania with counties

(Blaga *et al.*, 2009a; Blaga *et al.*, 2009b; Ciobotă *et al.*, 2015). Therefore, the objectives of this study was to assess the presence and prevalence of *Trichinella* spp. in wild boars from Bihor County, Romania, by using artificial digestion and serological methods.

Materials and Methods

Study area and samples

Romania is located in Eastern Europe, in the north of the Balkan Peninsula. The country has a temperate-continental climate of transitional type, with four clearly defined seasons (Trusca & Alecu, 2005). The animals originated from Bihor County (Fig. 1), a high-altitude area covered by forest. During the hunting season, in 2018, 84 plasma and diaphragm samples were collected from wild boars for research purposes. The blood samples, from each animal, were collected by jugular venipuncture into EDTA tubes. Plasma was obtained after the centrifugation of blood samples at 2500 RPM for 15 min and frozen at -15 °C until further use.

Artificial digestion

For artificial digestion, 5 g of diaphragm tissues were used from each animal. Four digestion pools of 100 g and one digestion pool of 20 g were made. For each pool of 100g, 2 l of heated water, 10

g of pepsin powder (1: 10,000 NF) and 16 ml of hydrochloric acid 25 % were added. The artificial digestion method was performed on all the samples according to Gamble and others (2000), in order to detect and collect the larvae for species identification. The artificial digestion method was carried out as stated in the European community regulation EU 2015/1375.

ELISA

Eighty four samples of plasma were tested for the presence of anti-*Trichinella* (IgG) antibodies by an indirect ELISA commercial kit (IDEXX, Westbrook, Maine, USA). Plasma and controls were diluted to 1/20 in dilution buffer provided in the kit. Afterwards, 200 µl were transferred into the wells and incubated 30 minutes at room temperature. After 3 washes, a volume of 100 µl of diluted conjugate was introduced in each well and incubated 30 minutes at room temperature, followed by 3 repetitions of the washing step. Subsequently, 100 µl of TMB (Tetramethylbenzidine) were added and left in contact for 10 minutes at room temperature in darkness. Finally, 100 µl of stop solution were transferred into each well. The optical densities were read at 450 nm.

Samples with a cut off value (E/P- positivity index calculation formula) equal or lower than 30 % were considered negative, while values that ranged from 30 % to 40 % were doubtful. Moreover,

positive samples were associated with E/P values that were equal or higher than 40 % (IDEXX ELISA technical guide, 2007).

Western blot

Twenty six plasma samples were tested with the Western blot technique. The samples were chosen randomly from the total of 84 samples previously tested in ELISA. Antigens were separated on a 10 % SDS-PAGE and transferred onto a nitrocellulose membrane. Plasma, diluted 1/100 in TBS, with an addition of 5 % milk was incubated for 1 hour at room temperature under gentle shaking. After 3 washes in TBS-tween, secondary antibodies diluted 1/15 000 in TBS, with 5 % milk were incubated in the same conditions, followed by 3 repetitions of the washing step. The obtained bands were identified by chemiluminescence with Geldoc™ reader (Bio-Rad, Bulletin 6376).

Statistical analysis

The prevalence of *Trichinella* spp. and the associated confidence intervals at a confidence level of 95 % were established for each diagnosis tests. Furthermore, the difference in seroprevalence among females and males was analyzed by chi-squared independence test. A p value lower than 0.05 was considered as statistically significant. The agreement between the results obtained in artificial digestion, ELISA and Western blot, was evaluated by Kappa statistics method (Landis, & Koch, 1977). Analyses were performed with SPSS® 20 software for Windows.

Ethical Approval and/or Informed Consent

All applicable national and institutional guidelines for the care and use of animals were followed.

Results and Discussion

Among the 84 animals included in the study, 45 (53.6 %) were females (38 adults, 7 juveniles) and 39 (46.4 %) were males (30 adults, 9 juveniles). All the samples were negative when artificial digestion was conducted. In ELISA 54 samples (64.2 %; 95 % CI: 53.7 – 74.2) were positive and 6 samples (7.1 %; 95 % CI: 1.4 – 12.5) were doubtful. Western blot was performed on 26 random samples, from which only 6 samples (23.0 %; 95 % CI: 5.6 – 40.3) gave a positive result. All the positive samples had three bands which were situated in the margin of 46 – 56.8.kD. Furthermore, ELISA and Western blot showed different results in the 26 samples tested. Thus, from the six samples that were positive in Western blot, only five were positive in ELISA. In addition, 11 (42.3 %; 95 % CI: 31.1 – 78.8) of the samples that were negative in Western blot, were positive in ELISA, as shown in Table 1. A Kappa value of 0.18 was found between ELISA and Western blot results, which demonstrated a slight agreement between the two methods used. Also, a Kappa value of 0.0 was observed in case of Western blot and artificial digestion, but also in case of ELISA and artificial di-

gestion, respectively showing no inter-rated agreement between the methods chosen for diagnosis. Moreover, the seroprevalence of *Trichinella* spp. was statistically higher in females than males, when ELISA test was used, at a p value < 0.01.

In case of artificial digestion, the lack of other regional muscles samples that are recommended in EU regulation can be responsible for the negative results. Kapel (2001) suggested that diaphragm and tongue muscles were the predilection sites in wild boars, independent of *Trichinella* genotype and infection level. Blaga *et al.*, (2009a) observed that in Romania from 1997 to 2004, the prevalence rate of *Trichinella* spp. infection in wild boars tested by trichinelloscopy was 8.6 %. Another study conducted from 2010 to 2014 in Hunedoara county, Romania, indicated a prevalence rate of 0.82 % for *Trichinella* spp. infection in wild boars after the examination of intercostal muscle samples by trichinelloscopy and artificial digestion (Ciobotă *et al.*, 2015). Moreover, another study in which the preferential muscles samples were tested with trichinoscopy and artificial digestion, found the presence of *Trichinella* larvae in five wild boars. In three of the animals, *T. britovi* was identified, while the remaining two animals were diagnosed with *T. spiralis* infection (Blaga *et al.*, 2009b). Some recently collected larvae in Romania, during the artificial digestion of diaphragm samples from pigs and wild boars evidenced the circulation of *T. spiralis* and *T. britovi* in these animal species (personal communication, unpublished data).

Although artificial digestion was negative, ELISA and Western blot showed high rates of positivity. Indirect ELISA is commonly used for the surveillance of *Trichinella* spp. infection (Elbers *et al.*, 2000; Gamble *et al.*, 2004; Kim *et al.*, 2015) and it can detect infection levels as low as 1 larvae/ 100g of tissue (Gamble *et al.*, 1983). Møller *et al.* (2005), by using indirect ELISA, detected *Trichinella* specific antibodies in sera and muscle fluid samples from wild boars. Moreover, ELISA based on carbohydrate tyvelose or excretory/secretory antigens, have been considered the most suitable antigens available (Forbes *et al.*, 2004). However, the assay yields false positive results due to cross-reactivity of the secondary antibodies (Nöckler *et al.*, 2004; Davidson *et al.*, 2009; Széll *et al.*, 2012). Therefore, ELISA positive results are recommended to be confirmed by Western blot due to its specificity of 100 % and the capacity to eliminate ELISA false-positive sera results (Gómez-Morales *et al.*, 2012; Cuttell *et al.*, 2014).

Taken together, these results indicate the circulation of *Trichinella* larvae at a low level in wild boar population of Bihor County. The larvae burden might be too low to be detected by direct artificial digestion whereas serological approaches allow to detect circulating antibodies or the presence of *Trichinella* spp. specific antigens. However, other studies observed a higher seroprevalence percentages, than the positivity rates observed in Western Blot or muscle digestion (Davidson *et al.*, 2009).

In Romania, the human population can acquire trichinellosis mainly by consuming raw pork and wild boar meat. The meat from these animals are usually prepared accordingly to local habits

Table 1. The comparative results for artificial digestion, ELISA, and Western blot.

Nr. Ctr.	Gender	Artificial digestion	ELISA (E/P)	Western blot (bands)
1	Male	N	D (34.55%)	N
3	Male	N	D (38.66%)	N
4	Male	N	P (66.42%)	N
6	Female	N	P (41.85%)	N
8	Female	N	D (32.18%)	N
11	Female	N	N (29.51%)	N
15	Male	N	P (47.33%)	N
18	Female	N	P (52.52%)	N
25	Female	N	P (40.79%)	N
30	Female	N	D (30.58%)	P (47.4; 51.8; 55.6 kD)
32	Male	N	P (157.72%)	P (46.4; 50.7; 58.5 kD)
39	Female	N	P (62.04%)	N
68	Male	N	P (53.37%)	N
79	Male	N	N (27.20%)	N
81	Male	N	N (26.78%)	N
83	Male	N	P (171.46%)	P (48.7; 52.3; 57.4 kD)
92	Female	N	P (63.20%)	P (47.4; 51.8; 55.6 kD)
104	Male	N	P (57.03%)	P (47.0; 51.8; 56.7 kD)
115	Female	N	D (39.11%)	N
154	Female	N	N (25.70%)	N
156	Male	N	P (65.00%)	N
157	Male	N	P (49.52%)	P (46.0; 50.2; 55.7 kD)
158	Female	N	P (42.48%)	N
166	Female	N	D (37.12%)	N
169	Female	N	P (56.14%)	N
176	Female	N	P (50.10%)	N

and traditions. Commonly, the wild boar meat products (dried raw salami, raw dried sausage, smoked pastrami e.g.) are not subjected to thermal processing, exposing the population to parasitic infestations. In 1993, controlled pig meat was mixed with infested wild boar meat, causing an extensive outbreak in Romania (Ciro-neanu & Ispas, 2002). Moreover, in January 2007, in the city of Timisoara, 5 humans acquired trichinellosis after consuming sausages with infected wild boar meat mixed with pork (Neghina, unpublished data).

Epidemiological actions that limit the spread of the infestation to humans and animals should be widely available and correctly applied by pig breeders, hunters, and consumers to prevent future outbreaks. These measures should include hygiene conditions in backyard pig breeding and raising, testing of wild boar meat from hunters and the consumption of only controlled wild boar meat by the local population (Neghina, 2010). However, in countries with high percentages of *Trichinella* spp. infection, in both animals and humans, an effective diagnosis method is essential for providing

food safety and preventing the onset of human outbreaks. Even if trichinelloscopy is a widely used method for *Trichinella* spp. detection, its sensitivity is lower than that of artificial digestion (Li *et al.*, 2010). The digestion assay is considered to be the most reliable procedure and it can be widely applied in practice (Gamble *et al.*, 2000).

The EU Regulation 2015/1375 highlights the importance, in case of wild boar meat, of using a sufficient amount of samples from the predilection sites, meaning foreleg, tongue or diaphragm muscles. In this regard and giving the recommendation of using at least 10 g of muscle tissue from each animal, is essential to ensure a correct training for the personnel that is responsible for the sample collection and inspection.

Conclusions

Serological evidences confirmed the exposure of wild boars from Bihor County, to *Trichinella* spp. Therefore, the consumers might

be at risk to ingest low larvae burden as animals are misdiagnosed after the parasitological diagnosis by artificial digestion. To prevent future misdiagnosis by using artificial digestion, only trained personal should perform the sampling and the digestion assay protocol. Human trichinellosis is continuously reported in Romania, consequently, risk factor assessment and surveillance should be improved in wild boars and other wild species to monitor the parasite presence and prevalence.

Conflicts of Interest

Authors have no potential conflict of interest pertaining to this submission to Helminthologia.

Acknowledgements

The authors would like to thank to, Sivia Diana Borşan, Maria Emilia Nedişan, Cristian Magdaş, and Andra Toma Naic, for their help in sample collection.

This research was supported by COST FA 1408 action, French Agency for Food, Environmental and Occupational Health & Safety (ANSES) and the University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca (USAMV Cluj-Napoca).

References

Bio-Rad, Bulletin 6376, General Protocol for Western Blotting (2016): Bio-Rad Laboratories, Inc, 1-2. Retrieved January 10, 2019 from bio-rad.com database https://www.bio-rad.com/web-root/web/pdf/lsr/literature/Bulletin_6376.pdf

BLAGA, R., DURAND, B., STOICHICI, A., GHERMAN, C., STEFAN, N., COZMA, V., BOIREAU, P. (2009a): Animal *Trichinella* infection in Romania: Geographical heterogeneity for the last 8 years. *Vet. Parasitol.*, 159(3 – 4): 290 – 294. DOI: 10.1016/j.vetpar.2008.10.059

BLAGA, R., GHERMAN, C., COZMA, V., ZOCEVIC, A., POZIO, E., BOIREAU, P. (2009b): *Trichinella* species circulating among wild and domestic animals in Romania. *Vet. Parasitol.*, 159(3 – 4): 218 – 221. DOI: 10.1016/j.vetpar.2008.10.034

BOADELLA, M., GORTAZAR, C., ACEVEDO, P., CARTA, T., MARTIN-HERNANDO, M.P., DE LA FUENTE, J., VICENTE. (2011): Six recommendations for improving monitoring of diseases shared with wildlife: examples regarding mycobacterial infections in Spain. *Eur. J. Wildl. Res.*, 57: 697 – 706. DOI: 10.1007/s10344-011-0550-x

Commission implementing regulation (EU) 2015/1375 of 10 August (2015) laying down specific rules on official controls for *Trichinella* in meat, Official Journal of the European Union EN. DOI: 10.1007/s12161-016-0619-y

CIOBOTĂ, F. O., CRISTEA, G., IONIȚĂ, M., MITREA, I. L. (2015): Epidemiological study on *Trichinella* infection in pigs and wild boars in hunedoara county (romania), during of 2010 – 2014 period. *Proc. Rom. Acad., Series B, Supplement 1*, 45 – 47. Supplement 1/2015, 4th ISAA. Retrieved January 9, 2020 from academiario-

mana.ro database sectii2002/proceedingsChemistry/doc2015-3s/art11_45.pdf

CIRONEANU, I., ISPAS, A. T. (2002): Totul despre trichineloză. [All about trichinellosis]. Bucharest: MAST Publishing House, Bucharest, Romania. 9 – 64 pp. (In Romanian)

CUTTELL, L., GÓMEZ-MORALES, M. A., COOKSON, B., ADAMS, P. J., REID, S. A., VANDERLINDE, P. B., JACKSON, A. L., GRAY, C., TRAUB, R. J. (2014): Evaluation of ELISA coupled with western blot as a surveillance tool for *Trichinella* infection in wild boar (*Sus scrofa*). *Vet. Parasitol.*, 199(3 – 4): 179 – 190. DOI: 10.1016/j.vetpar.2013.10.012

DAVIDSON, R. K., ØRPETVEIT, I., MØLLER, L., KAPEL, C. M. (2009): Serological detection of anti-*Trichinella* antibodies in wild foxes and experimentally infected farmed foxes in Norway. *Vet. Parasitol.*, 163(1 – 2): 93 – 100. DOI: 10.1016/j.vetpar.2009.03.020

ELBERS, A.R.W., DEKKERS, L.J.M., VAN DER GIESSEN, J.W.B. (2000): Sero-surveillance of wild boar in the Netherlands, 1996 – 1999. *Rev. sci. tech. Off. int. Epiz.* 19(3), 848 – 854. Retrieved January 10, 2020 from Agris database http://agris.fao.org/agris_search/search.do?recordID=FR2000004841

FORBES, L. B., APPELYARD, G. D., GAJADHAR, A. A. (2004): Comparison of synthetic tyvelose antigen with excretory–secretory antigen for the detection of trichinellosis in swine using enzyme-linked immunosorbent assay. *J. Parasitol.*, 90(4): 835 – 840. DOI: 10.1645/GE-187R

GAMBLE, H. R., ANDERSON, W. R., GRAHAM, C. E., MURRELL, K. D. (1983): Diagnosis of swine trichinosis by enzyme-linked immunosorbent assay (ELISA) using an excretory-secretory antigen. *Vet. Parasitol.*, 13(4): 349 – 361. DOI: 10.1016/0304-4017(83)90051-1

GAMBLE, H. R., BESSONOV, A.S., CUPERLOVIC, K., GAJADHAR, A.A., VAN KNAPEN, F., NOECKLER, K., SCHENONE, H., ZHU, X. (2000): International commission on trichinellosis: recommendations on methods for the control of *Trichinella* in domestic and wild animals intended for human consumption. *Vet. Parasitol.*, 93: 393 – 408. DOI: 10.1016/S0304-4017(00)00354-X

GAMBLE, H. R., POZIO, E., BRUSCHI, F., NÖCKLER, K., KAPEL, C. M. O., GAJADHAR, A. A. (2004): International Commission on Trichinellosis: recommendations on the use of serological tests for the detection of *Trichinella* infection in animals and man. *Parasite*, 11(1): 3 – 13. DOI: 10.1051/parasite/20041113

GAJADHAR, A. A., POZIO, E., GAMBLE, H. R., NÖCKLER, K., MADDOX-HYTTEL, C., FORBES, L. B., VALLEE I., ROSSI, P., MARINCULIC, A., BOIREAU, P. (2009): *Trichinella* diagnostics and control: mandatory and best practices for ensuring food safety. *Vet. Parasitol.*, 159(3 – 4): 197 – 205. DOI: 10.1016/j.vetpar.2008.10.063

IDEXX ELISA technical guide. (2007): IDEXX Laboratories Inc, 10-18. Retrieved February 10, 2019 from idexx.com database. <https://www.idexx.com/files/elisa-technical-guide.pdf>

KAPEL, C.M.O., GAMBLE, H.R. (2000): Infectivity, persistence, and antibody response to domestic and sylvatic *Trichinella* spp. in experimentally infected pigs. *Int. J. Parasitol.*, 30: 215 – 221. DOI: 10.1016/S0020-7519(99)00202-7

KAPEL, C.M.O. (2001): Sylvatic and domestic *Trichinella* spp. in

- wild boars; infectivity, muscle larvae, distribution, and antibody response. *J. Parasitol.*, 87: 309 – 314. DOI: 10.2307/3285046
- KIM, H. J., JEONG, W. S., KIM, E. M., YEO, S. G., AN, D. J., YOON, H., KIM, E.L., PARK, C. K. (2015): Prevalence of *Trichinella* spp. antibodies in wild boars (*Sus scrofa*) and domestic pigs in Korea. *Vet. Med.*, 60(4): 181 – 185. DOI: 10.17221/8105-VETMED
- LANDIS, J.R., KOCH, G.G. (1977): An application of hierarchical kappa-type statistics in the assessment of majority agreement among multiple observers. *Biometrics*, 33 (2): 363 – 374. DOI: 10.2307/2529786
- LI, F., CUI, J., WANG, Z.Q., JIANG, P. (2010): Factors affecting the sensitivity of artificial digestion and its optimization for inspection of *Trichinella spiralis* in meat. *Foodborne Pathog. Dis.*, 7(8): 879 – 885. DOI:10.1089/fpd.2009.0445
- GÓMEZ-MORALES, M. A., LUDOVISI, A., AMATI, M., BLAGA, R., ZIVOJINOVIC, M., RIBICICH, M., POZIO, E. (2012): A distinctive Western blot pattern to recognize *Trichinella* infections in humans and pigs. *Int. J. Parasitol.*, 42(11): 1017 – 1023. DOI: 10.1016/j.ijpara.2012.08.003
- MASSEI, G., GENOV, P. V. (2004): The environmental impact of wild boar. *Galemys*. 16(1), 135 – 145. ISSN: 1137-8700. Retrieved July 20, 2019 from semanticscholar.org database pdfs.semanticscholar.org/1717/da1dd916df958802589654700bf80045029c.pdf
- MØLLER, L. N., PETERSEN, E., GAMBLE, H. R., KAPEL, C. M. (2005): Comparison of two antigens for demonstration of *Trichinella* spp. antibodies in blood and muscle fluid of foxes, pigs and wild boars. *Vet. Parasitol.*, 132(1 – 2): 81 – 84. DOI: 10.1016/j.vetpar.2005.05.032
- NEGHINA, R. (2010): Trichinellosis, a Romanian never-ending story. An overview of traditions, culinary customs, and public health conditions. *Foodborne Pathog. Dis.*, 7(9): 999 – 1003. DOI: 10.1089/fpd.2010.0546
- NÖCKLER, K., HAMIDI, A., FRIES, R., HEIDRICH, J., BECK, R., MARINCULIC, A. (2004): Influence of Methods for *Trichinella* Detection in Pigs from Endemic and Non-endemic European Region. *J. Vet. Med., Series B*, 51(6): 297 – 301. DOI: 10.1111/j.1439-0450.2004.00770.x
- SZÉLL, Z., MARUCCI, G., LUDOVISI, A., GÓMEZ-MORALES, M. A., SRÉTER, T., POZIO, E. (2012): Spatial distribution of *Trichinella britovi*, *T. spiralis* and *T. pseudospiralis* of domestic pigs and wild boars (*Sus scrofa*) in Hungary. *Vet. Parasitol.*, 183(3 – 4): 393 – 396. DOI: 10.1016/j.vetpar.2011.07.035
- TRUSCA, V., ALECU, M. (2005): Romania's Third National Communication on Climate Change under the United Nations Framework Convention on Climate Change. Grue and Hornstrup, Holstebro. 28. Retrieved January 5, 2020 from unfccc.int database https://unfccc.int/resource/docs/natc/romnc3.pdf