



**HAL**  
open science

# Identification of discriminating behavioural and movement variables in lameness scores of dairy cows at pasture from accelerometer and GPS sensors using a Partial Least Squares Discriminant Analysis

Lucie Riaboff, Anne Relun, Charles-Elie Petiot, Mathieu Feuilloy, Sébastien Couvreur, Aurélien Madouasse

## ► To cite this version:

Lucie Riaboff, Anne Relun, Charles-Elie Petiot, Mathieu Feuilloy, Sébastien Couvreur, et al.. Identification of discriminating behavioural and movement variables in lameness scores of dairy cows at pasture from accelerometer and GPS sensors using a Partial Least Squares Discriminant Analysis. Preventive Veterinary Medicine, 2021, 193, pp.105383. 10.1016/j.prevetmed.2021.105383 . hal-03321095

**HAL Id: hal-03321095**

**<https://hal.inrae.fr/hal-03321095v1>**

Submitted on 2 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Identification of discriminating behavioural and movement variables in lameness scores of dairy**  
2 **cows at pasture from accelerometer and GPS sensors using a Partial Least Squares**  
3 **Discriminant Analysis**

4 Riaboff L.<sup>1</sup>, Relun A.<sup>1</sup>, Petiot C-E.<sup>1</sup>, Feuilloy M.<sup>2</sup>, Couvreur S.<sup>3</sup>, Madouasse A<sup>1</sup>.

5 <sup>1</sup>INRAE, BIOEPAR, Oniris, 44307 Nantes, France

6 <sup>2</sup>ESEOTech-LAUM, Ecole Supérieure d'Electronique de l'Ouest, 49000 Angers, France

7 <sup>3</sup>USC ESA-INRAE 1481 URSE, Ecole Supérieure d'Agricultures, University Bretagne Loire, 49007  
8 Angers, France

9 Corresponding author:

10 Lucile Riaboff – lucile.riaboff@ucd.ie – 41 Belmont Avenue, Donnybrook, Dublin 4, D04RH74 –  
11 + 353 613 7495

12 **Abstract**

13 The behaviour and movement of lame dairy cows at pasture have been studied little, yet they could be  
14 relevant to improve the automatic detection of lameness in cows in pasture-based systems. Our aim in  
15 this study is to identify behavioural and movement variables of dairy cows at pasture that could  
16 discriminate lameness scores. Individual cow behaviours were predicted from accelerometer data and  
17 movements measured using GPS data. Sixty-eight dairy cows from three pasture-based commercial  
18 farms were equipped with a 3-D accelerometer and a GPS sensor fixed on a neck collar for 1 to 5  
19 weeks, depending on the farm, in spring and summer 2018. A lameness score was assigned to each  
20 cow by a trained observer twice a week. Behaviours were predicted every 10 seconds based on  
21 accelerometer data, and then combined with the GPS position. Segmentation on behavioural time  
22 series was used to delineate each behavioural bout within each outdoor period. Thirty-seven  
23 behavioural and movement variables were then calculated from the behavioural bouts for each cow . A  
24 partial least square discriminant analysis was performed to identify the variables that best discriminate  
25 lameness scores. Time spent grazing, grazing bout duration, duration before lying down in the pasture,  
26 time spent resting, number of resting bouts, distance travelled during grazing, and dispersion were the  
27 most discriminant variables in the PLS-DA (VIP > 1). Severely lame cows spent 4.5 times less time  
28 grazing and almost twice as much time resting as their sound congeners, especially in the lying

29 position. Exploratory behaviour was also reduced for both moderately and severely lame cows,  
30 resulting in 1.2 and 1.7 times less distance travelled respectively, especially during grazing, These  
31 variables could be used as additional variables to improve the performance of existing lameness  
32 detection devices in pasture-based systems.

### 33 **Keywords**

34 Behaviour and movement, dairy cow lameness, pasture, accelerometer, GPS, PLS-DA

### 35 **Highlights**

- 36 - Behavioural and movement variables can discriminate lameness scores using PLS-DA
- 37 - Lameness cows spend more time resting and less time grazing and exploring
- 38 - Accelerometer combined with GPS could help lameness detection in grazing systems

### 39 **Abbreviations**

- 40 - AHDB: Agriculture and Horticulture Development Board
- 41 - AMS: Automatic Milking System
- 42 - ANOVA: Analysis of Variance
- 43 - GPS: Global Positioning System
- 44 - HMM: Hidden Markov Model
- 45 - LS: Lameness Score
- 46 - PLS-DA: Partial Least Squares Discriminant Analysis
- 47 - VIP: Variable Importance in Projection

48

## 49 **1. Introduction**

50 Lameness is one of the main health disorders in dairy cattle (Huxley, 2013), affecting animal  
51 welfare (Whay and Shearer, 2017) and inducing economic losses for farmers (Willshire and Bell,  
52 2009). Lameness detection is usually done by visual inspection by the farmer, and lameness  
53 prevalence is often underestimated by farmers (Cutler et al., 2017), leading to delayed treatment of

54 lame cows. For these reasons, automatic lameness detection tools could be a relevant way to improve  
55 the early identification of lame cows in dairy farms in order to reduce associated costs (Green et al.,  
56 2002) and to increase chances of recovery (Leach et al., 2012).

57 Automatic detection tools are being developed for indoor systems. The main systems are  
58 based on gait change determined using 2D and 3D cameras, pressure mats or pedometers, or on  
59 changes in behaviour determined using accelerometers on the neck, feeding data or automatic milking  
60 system (AMS) data (Van Nuffel et al., 2015). The performance of these systems is promising,  
61 although their sensitivity and specificity should be improved in order to increase their use in  
62 commercial dairy farms (O’Leary et al., 2020).

63 Agroecology promotes the return of pasture-based systems for dairy cows (Dumont et al.,  
64 2013). Although lameness is less common in grazing systems, its prevalence in dairy farming remains  
65 non-negligible (Haskell et al., 2006). Furthermore, lameness may be favoured by certain grazing  
66 conditions, such as grazing on moist soils during periods of heavy rainfall (Politiek et al., 1986;  
67 Vermunt and Greenough, 1995), alternate grazing with sheep (Barker et al., 2010), or when cows have  
68 to walk on roads or concrete tracks between the parlour and grazing (Barker et al., 2009). In addition,  
69 dairy cows usually graze for only part of the year and are housed during the winter season (van den  
70 Pol-van Dasselaar et al., 2020), possibly leading to lameness prevalence similar to indoor-based  
71 systems in early spring. For these reasons, the development of automatic tools for lameness detection  
72 in dairy cows would also be relevant for pasture-based systems.

73 Previous studies have shown that the amount of time spent in standing and lying positions by  
74 dairy cows at pasture changed with lameness (Navarro et al., 2013), suggesting that behaviour could  
75 be a relevant indicator to help lameness detection at pasture, in the same way as in indoor systems  
76 (Almeida et al., 2008; Blackie et al., 2011; Weigele et al., 2018). A wide range of behaviours on  
77 pasture can now be collected using accelerometer sensors (Riaboff et al., 2020a), a technology already  
78 used in dairy farming for heat detection (Kamphuis et al., 2012), and one that is quite affordable for  
79 farmers (Delagarde and Lamberton, 2015). Furthermore, indicators evaluating the movement of dairy  
80 cows at grazing obtained with embedded Global Positioning System (GPS) sensors (Feldt and  
81 Schlecht, 2016; Riaboff et al., 2020b) could also be interesting for lameness detection, although this

82 has not yet been studied. The identification of behavioural variables (time-budget, duration and  
83 number of bouts) and movement variables (distance, dispersion, distance travelled during grazing,  
84 etc.) that discriminate lameness scores on pasture would thus be a first step in the development of a  
85 lameness prediction model for pasture-based systems.

86 In two previous studies, a methodology was developed to predict a wide range of behaviours  
87 in dairy cows at pasture from accelerometer data every 10 seconds, including grazing, walking, resting  
88 and ruminating both in lying and standing position (Riaboff et al., 2019; Riaboff et al., 2020a). These  
89 predicted behaviours can be combined with the cow position using GPS data collected on animals  
90 (Riaboff et al., 2020b). Behavioural and movement variables can then be calculated from the predicted  
91 behaviours and GPS position. In this study, we propose to identify behavioural variables predicted  
92 from accelerometer data as well as movement variables computed from GPS data that could be used to  
93 discriminate lameness scores in pasture-based systems, using a Partial Least Squares Discriminant  
94 Analysis (PLS-DA).

## 95 **2. Materials and methods**

96 An overview of the main steps in the Materials and methods section is provided in Figure 1. First,  
97 accelerometer and GPS data were obtained from 68 dairy cows from three pasture-based farms. For  
98 each cow, a lameness score was collected twice a week over the course of the experiment (*section*  
99 *2.1*). Prediction of behaviours was carried out using accelerometer data and combined with the  
100 position of dairy cows using GPS data (*section 2.2*). For each cow, a lameness score was assigned for  
101 each day the cow was fitted with the accelerometer and GPS device. For each day the cow was fitted,  
102 the lameness score was the score from the closest scoring session. Behavioural and movement  
103 variables were then computed for each cow on each day she was equipped after identifying  
104 behavioural bouts within each outdoor period (*section 2.3*). Finally, the identification of discriminant  
105 behavioural and movement variables was done using a PLS-DA (*section 2.4*).

106 < Figure 1.>

### 107 **2.1 Data collection**

108                    *2.1.1 Description of farms and animals*

109                    The experiment was carried out on three commercial dairy farms successively, with herd size  
110 ranging from 54 to 71 Holstein cows, located in the Pays-de-la-Loire region (France) from April to  
111 July of 2018. Dairy cows were milked using an AMS (milking count per day (mean  $\pm$  SD):  $2.7 \pm 0.8$ ;  
112  $2.4 \pm 0.7$  and  $2.0 \pm 0.8$ ). Cows were housed from November to March and had access to pasture  
113 continuously from April to October. During the grazing season, cows were free to stay in the barn.  
114 Barns were equipped with straw cubicles in all three farms. Cows received supplementation with  
115 maize silage and concentrates in the first and the third farm (theoretical supplementation: 40 %), and  
116 supplementation with hay and concentrates in the second farm (theoretical supplementation: 20 %).  
117 Most of the concentrates were delivered during milking with the AMS. Details on each farm are  
118 provided in Appendix 1.

119                    Cows with less than 50 or more than 250 days in milk at the start of the experiment were not  
120 included in order to avoid animals leaving the herd or being dried off during the experiment. Animals  
121 were then selected so that both lame cows (lameness score  $\geq 2$ ; Agriculture and Horticulture  
122 Development Board, Stoneleigh Park, Kenilworth, Warwickshire, CV8 2TL, 2020) and sound cows  
123 (lameness score  $\leq 1$ ; Agriculture and Horticulture Development Board, Stoneleigh Park, Kenilworth,  
124 Warwickshire, CV8 2TL, 2020) were equipped in each farm. For this purpose, each cow in the herd  
125 was scored three days before the start of the experiment in each farm according to the methodology  
126 described in *section 2.1.3.1*. The distribution of lameness scores in each farm is provided in  
127 Appendix 1. As parity may impact behaviour, we tried to balance the dataset in terms of parity. For  
128 this purpose, we chose the animals so that each lame cow had an equivalent sound cow with the same  
129 parity (1, 2 or  $\geq 3$ ) and with a difference in days in milk of less than 20 days. Based on these criteria,  
130 13/54 cows were equipped in the first farm, 27/71 cows in the second farm, and 26/61 in the last one,  
131 giving a total of 68 different cows used for the study. Among the equipped cows, one cow in Farm 1  
132 and eight cows in Farm 3 showed signs of heat on one day each during the experiment (AMS data).  
133 Details on the animals selected in each farm are provided in Table 1.

134                    < Table 1 >

135           2.1.2 *Description of sensors*

136   An RF-Track datalogger (RF-Track, Rennes, France) comprising an LSM9DS1 three-axis  
137   accelerometer (STMicroelectronics, Geneva, Switzerland)  $\pm 2$  g and a GPS sensor (part number EVA-  
138   7M-0,  $\mu$ -Blox, Thalwil, Switzerland) with a static position error estimated at  $\pm 1.72$  m was used. The  
139   sampling rate was 59.5 Hz and 1 Hz for the accelerometer and GPS data, respectively. Sensors were  
140   powered with two 3.7 V lithium batteries (2.6 Ah). A secure digital card was used for data storage.  
141   The dataloggers were 98.2 mm x 51.60 mm x 36.0 mm in size and weighed 250 g. The dataloggers  
142   were attached to a collar and positioned on the right side of the cow's neck. A counter-weight of 500 g  
143   was added to prevent the collars from turning around and they were tightly adjusted. The x-axis  
144   detected the up-down direction, the y-axis detected the backward-forward direction and the z-axis  
145   detected the left-right direction. The device used is shown in Figure 2.

146           2.1.3 *Experiment design*

147   The experiment was carried out from April 2018 to July 2018 using 68 cows from three pasture-based  
148   commercial farms. The course of the experiment is shown in Figure 2.

149   < Figure 2 >

150           2.1.3.1 *Lameness score collection*

151   Equipped cows were scored twice a week for the duration of the experiment. Each scoring session was  
152   carried out in the barn after supplementation delivery. Cows were made to walk along a 3 m x 30 m  
153   path marked with a tape. A camera was positioned near the path so that each animal was filmed as it  
154   walked along it. The videos were used to check intra-observer consistency. If necessary, a handler  
155   would stand behind the cows to encourage them to walk. Lameness scores (LS) were assigned to each  
156   cow by a single experimenter (C.E. Petiot, 5<sup>th</sup> year veterinary student at the time of the experiment)  
157   trained in the methodology developed by the Agriculture and Horticulture Development Board  
158   (Agriculture and Horticulture Development Board, Stoneleigh Park, Kenilworth, Warwickshire, CV8  
159   2TL, 2020). The AHDB method classifies lameness using a four-point scale (0 = good mobility (not  
160   lame), 1 = imperfect mobility, 2 = impaired mobility (lame), and 3 = severely impaired mobility

161 (severely lame); Table 2). The Cohen's Kappa was computed to evaluate the intra-reliability of  
162 observer scoring. An intra-observer Cohen's Kappa of 0.66 was obtained by scoring 50 cows from the  
163 three farms using videos watched twice several months apart, suggesting a strong agreement between  
164 the scoring sessions (Cohen, 1960). We did not interfere in the management of lameness by the  
165 farmers during the experiment. At the end of the data collection process, each farmer was provided  
166 with a detailed report on the prevalence of lameness on their farm, along with a list of  
167 recommendations to reduce this prevalence.

168 < Table 2 >

### 169 2.1.3.2 Accelerometer and GPS data collection

170 In each farm, the selected cows were continuously equipped with the device for a minimum of two  
171 days and a maximum of six days, the latter corresponding to the battery life of the sensors. Data were  
172 downloaded after each continuous period and batteries were recharged for eight hours before  
173 equipping the cows for the next period. Four, two and five continuous periods were carried out,  
174 leading to 21, 12 and 25 days of experiment in the first, the second and the third farm, respectively. In  
175 this way, accelerometer and GPS data were collected on a total of 58 different days (Table 1).

## 176 2.2 Prediction of dairy cow behaviour and combination with GPS data

177 This step aimed to (i) predict the behaviours successively expressed every 10 seconds by the cows  
178 from accelerometer data over the course of the experiment, and to then (ii) combine the predicted  
179 behaviours with GPS data. For this purpose, we used a methodology described in two previous studies  
180 (Riaboff et al., 2019; Riaboff et al., 2020a), which ensures the prediction of six behaviours in dairy  
181 cows on pasture from accelerometer data with high performance (accuracy: 98 %; Cohen's Kappa:  
182 0.96). We refer to Riaboff et al. (2020a, 2019) for a detailed explanation of the methodology. The six  
183 predicted behaviours are the following:

- 184 - Grazing: biting, taking frequent bites or chewing and searching without raising the head.
- 185 - Walking: moving from one location to another without lowering the head to ground level.

- 186 - Ruminating while lying: lying down with regurgitating ruminal bolus before chewing and then  
187 re-swallowing.
- 188 - Ruminating while standing: standing with regurgitating ruminal bolus before chewing and  
189 then re-swallowing.
- 190 - Resting while lying: lying down without rumination.
- 191 - Resting while standing: standing without movement or rumination.

192 The methodology developed was thus applied in this study to predict the behaviours as illustrated in  
193 Figure 3. Raw signal accelerometer sequences collected on the 68 cows over the 58 days of the  
194 experiment were divided into segments (*windows*) of 10 s, without data in common between two  
195 consecutive windows (without *overlap*). Sixty-one features were then calculated in each window. This  
196 pre-processing step was performed in Matlab R2018a. The eXtreme Gradient Boosting model fitted by  
197 Riaboff et al. (2020a) was then directly used to predict the behaviours from the calculated features  
198 using the xgboost package (Chen et al., 2018) in R 4.0.0 (*R Core Team*, 2020). The Hidden Markov  
199 Model (HMM)-based Viterbi algorithm reported by Riaboff et al. (2020a) was then applied with the R  
200 package HMM (Himmelmann, 2010) to smooth the predicted behaviours in successive windows from  
201 the same cow over the entire experiment. Each 10 s window of behaviour was finally associated with  
202 the position (longitude and latitude) of the cow, based on the time of the GPS and the accelerometer,  
203 which was previously synchronised. In this way, each 10 s window of behaviour was associated with  
204 the GPS coordinates of the first sample of the window and those of the last sample of the window. At  
205 the end of this step, we obtained the sequences of the 10 s windows of behaviour for each cow over  
206 time, with the associated GPS coordinates.

207 < Figure 3 >

208 2.3 Creation of the dataset with behavioural and movement variables and the associated  
209 lameness score

210 The aim of this step was to gather in the same dataset the variables calculated from the predicted  
211 behaviours and the movement of the cows and the corresponding lameness scores for each cow on  
212 each of the days she was equipped with the device.

### 213 *2.3.1 Lameness score assignment for each cow on each of the days she was equipped*

214 As explained in section 2.1.3.1, scoring sessions were carried out twice a week. The score assigned to  
215 each observation (cow X day) corresponded to the score obtained for a given cow in the scoring  
216 session closest to the day of the observation. If the scoring session was more than three days away  
217 from the day considered (i.e. if the cow missed a scoring session), a missing value was assigned for  
218 that observation. For 10 of the 58 days of the experiment, the number of days between the two  
219 successive scoring sessions was the same. If the cow was scored in both sessions, we chose to assign  
220 the lameness score from the second session as the default value.

### 221 *2.3.2 Behavioural and movement variables for each cow on each of the days she was* 222 *equipped*

#### 223 *2.3.2.1 Identification of behavioural bouts within outdoor periods*

224 The aim of this step was to identify behavioural bouts within each outdoor period, i.e. sessions  
225 during which an animal expressed the same behaviour continuously, with only brief interruptions. The  
226 principle used to identify behavioural bouts within each outdoor period is illustrated in Figure 4. For  
227 this purpose, we first identified periods spent outdoors for each cow from GPS data and the  
228 coordinates of the barn (French National Geographic Institute; “Géoportail,” 2006), using the R  
229 package *sp* (Pebesma and Bivan, 2005). We then applied Lavielle segmentation (Lavielle, 1999) on  
230 the “indoor-outdoor” periods time series in order to smooth potential GPS errors (Figure 4 a) using the  
231 R package *adehabitatLT* (Calenge, 2006). This method finds the best segmentation of a time series  
232 given a fixed maximum number of segments by minimising a contrast function (contrast between the  
233 actual time series and the segmented time series). Finally, the sequence of 10 s windows of predicted  
234 behaviour for each cow within each outdoor period was smoothed, also using Lavielle segmentation

235 (Figure 4 b). A more in-depth explanation of the use of this method and the parameters chosen is  
236 provided in Appendix 2.

237 Each segment resulting from the segmentation was considered as a bout. The behaviour  
238 associated with each bout was the behaviour most represented. It should be noted that in some  
239 segments, no behaviour could be identified as the most represented. In this case, the bout was  
240 annotated “heterogeneous bout” (16 % of the bouts). As the “heterogeneous bouts” were difficult to  
241 interpret from a behavioural point of view (Appendix 3), they were not considered in the rest of the  
242 analysis.

243

244 < Figure 4 >

#### 245 2.3.2.2 Calculation of movement and behavioural variables

246 The behavioural bouts previously obtained and the corresponding positions of dairy cows on pasture  
247 were used to compute different variables. These variables are presented in Table 3. The following  
248 behavioural variables were computed for each cow on each day she was equipped: the overall time  
249 spent on pasture (1), the time-budgets (2), the number of bouts (3), and the mean duration of bouts (4).  
250 The variables (2) to (4) were calculated for the six predicted behaviours described in section 2.2, as  
251 well as for the following grouped behaviours:

- 252 • Ruminating: grouping of “ruminating while lying” and “ruminating while standing”  
253 behaviours
- 254 • Resting: grouping of “resting while lying” and “resting while standing” behaviours
- 255 • Lying: grouping of behavioural variables from “ruminating while lying” and “resting while  
256 lying”
- 257 • Standing: grouping of behavioural variables from “ruminating while standing” and “resting  
258 while standing”

259 The following movement variables were also computed from the GPS data and behavioural bouts: the  
260 number of round-trips between the pasture and the barn (5), the overall distance travelled (6), the

261 overall dispersion (7), the distance travelled during grazing (8), the distance travelled during walking  
262 (9), the ratio of the distance travelled during grazing over the distance travelled during grazing and  
263 walking (10), the speed of walking (11), and the mean duration between the time the cow enters the  
264 pasture and the time she lies down (12). It should be noted that behavioural and movement variables  
265 were recorded as a missing value when no outdoor period was detected over a whole day for a given  
266 cow.

267 < Table 3 >

### 268 2.3.2.3 Weighting and standardisation

269 As explained in section 2.1, 68 cows from three commercial farms were used in this study, and data  
270 were collected over 58 days, which resulted in a considerable variation between cows (time spent  
271 outdoors, etc.) and between days (pasture access, area of pasture, etc.), thus preventing comparison of  
272 the behavioural variables from one cow to another and from one day to the next. Consequently, both  
273 behavioural and movement variables were corrected to make the observations comparable thereafter.  
274 A weighting was thus applied depending on the variable, using:

- 275 - The time spent outside by the cow on a given day ( $\text{Time}_{\text{out}}$ ).
- 276 - The maximum  $\text{time}_{\text{out}}$  recorded on a given day among all the cows equipped on that day  
277 ( $\text{Time}_{\text{out,max}}$ ).
- 278 - The distance travelled by the cow on a given day ( $\text{Distance}$ ).
- 279 - The maximum distance travelled on a given day among all the cows equipped on that day  
280 ( $\text{Distance}_{\text{max}}$ ).
- 281 - The maximum dispersion on a given day among all the cows equipped on that day ( $\text{R2n}_{\text{max}}$ ).

282 The weighting applied to each variable is provided in Table 3. A more detailed explanation of the  
283 weighting is provided in Appendix 4. Finally, the behavioural and movement variables were  
284 standardised.

### 285 2.3.3 Dataset cleaning

286 The resulting dataset was then cleaned to avoid including unusable observations and highly correlated  
287 variables in the analysis to follow. First, observations for which the nearest scoring session was more  
288 than three days away (the cow having missed a scoring session; section 2.3.1) were removed from the  
289 dataset, corresponding to 14 observations (1.5 % of the dataset). Missing values corresponding to  
290 observations in which the cow did not go to the pasture on a given day (section 2.3.2.2) were also  
291 removed from the dataset. This corresponded to the deletion of 88 observations (9.3 % of the dataset).  
292 Details on the observations removed are provided in Appendix 5 a. This resulted in a dataset of 863  
293 observations (Appendix 5 b). The coefficient of correlations between each behavioural and movement  
294 variable were calculated using the R package corrplot (Wei, 2017). No variable was highly correlated  
295 with another (correlation > 0.95; see Appendix 6), and the 37 quantitative variables were therefore  
296 retained.

#### 297 2.4 Identification of discriminating variables in lameness score with a PLS-DA

298 A one-way Analysis of Variance followed by a Tukey test was applied in a preliminary step,  
299 and showed that the group LS=0 was never significantly different from the group LS=1 for any of the  
300 variables studied (Appendix 7). For this reason, we combined LS=0 and LS=1 within the same group  
301 “LS0-1” before applying the PLS-DA.

302 We applied a PLS-DA model because (i) it is a Machine Learning method adapted to perform  
303 a first analysis on a large dataset about which we have limited *a priori* knowledge, and (ii) it is  
304 particularly relevant to identify the variables that have contributed to discrimination between groups  
305 using the Variable Importance in Projection (VIP) metric (Tenenhaus, 1998).

306 PLS-DA is a multivariate projection method for modelling a relationship between the  
307 quantitative variables X and a dummy matrix Y (Barker and Rayens, 2003). The target is to find PLS-  
308 components which both restore the variance of the matrix X and maximise the separation between the  
309 classes of Y. In this study, the independent variables X were the behavioural and movement variables,  
310 and the dummy matrix Y was a three-column matrix, representing the groups “LS0\_1”, “LS2” and  
311 “LS3”. For each column in Y, each sample was assigned to 0 or 1 depending on the group to which it

312 belonged. VIPs were then used to identify variables of X which are important in determining class  
313 membership of Y, also called discriminant variables (Chong and Jun, 2005). A more detailed  
314 description of the PLS-DA and the definition of VIP are provided in Appendix 8. The PLS-DA and  
315 VIP analysis were performed with the R package plsdeplot (Sanchez, 2016).

### 316 **3. Results**

317 Means and standard deviations obtained for the three lameness groups (LS0\_1, LS2 and LS3) for  
318 each variable are provided in Table 4. Plots of the included cow X day observations on the first two  
319 PLS-components with the associated performance metrics of the PLS-DA model, as well as the circle  
320 of correlations and the VIP of variables in descending order, are displayed in Figure 5. It should be  
321 noted that the VIPs used were those computed only on the first component, because it was the only  
322 significant component in the discrimination model.

323 < Table 4>

324 < Figure 5>

325 Three PLS-components were kept in the model as the Q2cum (predictive coefficient)  
326 continued to increase until the third component (Appendix 9 a), but only the first one was significant  
327 based on the Tenenhaus criteria (Tenenhaus, 1998; Appendix 9 b). The PLS-DA model using the first  
328 two components led to a Q2cum of 0.11, a R2Xcum of 0.42 and a R2Ycum of 0.11. Plots of  
329 observations (Figure 5 a) showed a discrimination between each LS on the first PLS component (R2X:  
330 20%), although there is an overlap between the 95 % confidence ellipses. The circle of correlations on  
331 the first two components between the most important variables (VIP > 0.8) and lameness scores is  
332 provided in Figure 5 b. The discrimination was explained by both the behavioural and movement  
333 variables. The LS0\_1 group was associated with higher “distance”, “R2n”, “distance grazing”,  
334 “distance ratio” and “duration before lying” and with longer “time<sub>grazing</sub>” and “duration bout<sub>grazing</sub>”  
335 than the two other groups (Figure 5 b; Table 4). The LS2 group was associated with longer  
336 “time<sub>ruminating\_standing</sub>”, “time<sub>standing</sub>” and “duration bout<sub>standing</sub>” (Figure 5 b; Table 4). The LS3 group  
337 was associated with an increase in resting behaviour (“time<sub>resting</sub>”, “number bouts<sub>resting</sub>”, “duration

338 bouts<sub>resting</sub>”), especially in the lying position (“time<sub>resting\_lying</sub>”, “duration bout<sub>resting\_lying</sub>”, “number  
339 bouts<sub>resting\_lying</sub>”) (Figure 5 b; Table 4).

340         Among the 37 variables tested, 17 contributed to discriminate the lameness scores (VIP > 0.8),  
341 among which 14 were very important for discrimination (VIP>1) (Figure 5 c). The most important  
342 behavioural variables (VIP > 1) included grazing behaviour (“time<sub>grazing</sub>”), resting behaviour  
343 (“time<sub>resting</sub>”, “duration bout<sub>resting</sub>”, “number bouts<sub>resting</sub>”) especially while lying (“time<sub>resting\_lying</sub>”,  
344 “duration bout<sub>resting\_lying</sub>”, “duration before lying”), and ruminating both while standing  
345 (“time<sub>ruminating\_standing</sub>”) and lying (“time<sub>ruminating\_lying</sub>”). The most important movement variables  
346 (VIP > 1) were the distance travelled (“distance grazing”, “distance”, “distance ratio”) and the  
347 dispersion (“dispersion R2n”). Other variables that contributed to discrimination (VIP > 0.8) were  
348 “number bouts<sub>resting</sub>”, “number bouts<sub>resting\_lying</sub>”, “duration bouts<sub>grazing</sub>” and “duration bouts<sub>resting</sub>”.

## 349         **4. Discussion**

### 350         4.1 Behavioural and movement variables to discriminate lameness scores

351 Instances where cows did not go out to pasture a given day (missing outdoor periods; *section 2.3.2.2*)  
352 occurred for both lame and sound cows (Appendix 5 a). Furthermore, the PLS-DA showed that the  
353 time spent outdoors (time<sub>out</sub>) was not a discriminating variable. Consequently, lame and sound cows  
354 spent a similar amount of time on pasture but the expressed behaviours and movement patterns are  
355 altered with the lameness score.

#### 356         4.1.1 *Discriminating variables related to grazing behaviour*

357 The time spent grazing was the variable that best discriminated the lameness scores (Figure 5 c).  
358 LS0\_1 (sound cows) spent 4.5 times longer grazing than LS3 (severely lame) and 1.6 times longer  
359 than LS2 (moderately lame), which was explained by longer grazing bouts (Figure 5 b; Table 4).  
360 These results contrast with those obtained in the study by Walker et al. (2008), where no difference  
361 was observed in grazing behaviour between lame and sound cows at pasture. It should be noted that  
362 the latter study was carried out on animals after oestrus synchronisation, whereas only nine cows  
363 showed signs of heat on one day each during the experiment in our study (AMS data; *section 2.1.1*),

364 which could explain this difference in results, as feeding behaviour is also modified during heat events  
365 (Dolecheck et al., 2015). Furthermore, our results are in accordance with the studies conducted in  
366 indoor-based systems, as reductions in feeding time (Weigele et al., 2018) and the duration of feeding  
367 bouts (Norrington et al., 2014) have already been shown.

#### 368 *4.1.2 Discriminating variables related to resting behaviour*

369 Time spent resting was the second most discriminating variable between lameness scores  
370 (Figure 5 c). Severely lame cows spent almost twice as much time resting as their sound congeners,  
371 especially in the lying position, which was explained by both more and longer resting while lying  
372 bouts (Figure 5 b; Table 4). These results are consistent with those obtained in indoor-based systems,  
373 as Weigele et al. (2018) found a decrease in activity in lame cows, while Blackie et al. (2011)  
374 observed an increase in lying bouts. The increase in the number of lying bouts was also noted by  
375 Navarro et al. (2013), while the increase in resting bout duration has already been observed in indoor-  
376 based systems (Weigele et al., 2018).

377 In our study, the duration before lying down was also one of the most important variables  
378 (Figure 5 c) and contributed to discriminating the sound group from the other two (Figure 5 b; Table  
379 4). Indeed, sound cows or those with a slight asymmetry in their gait lie down twice as late once they  
380 enter the pasture as lame cows (Table 4). Yunta et al., (2012) also showed that lame cows lay down  
381 earlier than sound cows once the ration had been distributed in indoor based-systems. Similarly, lame  
382 cows probably shortened their first grazing bout once on the pasture in our study. It is also possible  
383 that severely lame cows avoid lying down in the barn and therefore lie down more quickly once on  
384 pasture.

#### 385 *4.1.3 Discriminant variables related to exploratory behaviour*

386 The total distance travelled contributes to the discrimination between lameness scores (Figure 5 c).  
387 Indeed, sound cows travelled about 1.5 times further than their lame congeners (Figure 5 b; Table 4).  
388 This finding is in agreement with the study by Blackie et al. (2011) in indoor-based systems. In our  
389 study, the distance travelled during grazing as well as the distance ratio were also 1.6 times and 2.3

390 times greater for the sound and slightly asymmetric gait cows than for the moderately and severely  
391 lame cows, respectively (Figure 5 b; Table 4), explaining why these variables were also important in  
392 discriminating between lameness scores (Figure 5 c). Similarly, the dispersion of the sound group was  
393 1.6 times greater than for the severely lame cows, and also slightly greater than for the moderately  
394 lame cows (Figure 5 b; Table 4). All of these results suggest a more pronounced exploratory dynamic  
395 in sound and slightly asymmetric gait cows than in lame animals, especially during grazing behaviour.  
396 Nevertheless, walking time did not discriminate between lameness scores (Figure 5 c), which is in  
397 accordance with the study by Beer et al. (2016) in indoor-based systems. However, speed has not  
398 emerged as a discriminant variable in our study, contrary to what was observed in indoor-based  
399 systems (Beer et al., 2016; Blackie et al., 2011). It should be noted that in our study, speed was  
400 measured by averaging the distances travelled calculated from the GPS data across all the 10 s  
401 windows in which the walking behaviour was predicted. It is therefore possible that the chosen  
402 window size was too small to calculate a representative walking speed.

#### 403 *4.1.4 Discriminant variables related to ruminating and standing position*

404 The moderately lame group was associated with a slightly longer time ruminating while in standing  
405 position than sound cows (Figure 5 b; Table 4). These results differ from those obtained by Walker et  
406 al. (2008) on pasture, as lame cows spent more time ruminating while lying down and less time  
407 ruminating while standing compared to sound cows. However, the results are difficult to compare, as  
408 the latter study was carried out with cows after oestrus synchronization, contrary to our study (section  
409 2.1.1), and rumination may be modified during this period (Dolecheck et al., 2015). It would be  
410 interesting to conduct our experiment again with other cows in order to confirm the results observed  
411 on the standing and lying positions adopted by lame cows while ruminating. The moderately lame  
412 group also spent 1.7 times longer standing than the sound group (Figure 5 b; Table 4). This result is  
413 quite unexpected, as time spent standing on pasture is somewhat reduced in lame cows in other studies  
414 (Walker et al., 2008; Navarro et al., 2013). This could be explained by the difference in the grazing  
415 systems used (size of the herd, duration of pasture access, milking system, etc.). In particular, it should  
416 be noted that in the second and the third farm, a single AMS was used for 71 and 61 cows,

417 respectively. This led to AMS saturation at certain times, forcing animals to wait on the access road  
418 from the pasture to the AMS before being milked. As it has already been shown that lame cows avoid  
419 aggressive behaviours with others (Galindo and Broom, 2002) and move to the back of the herd  
420 (Walker et al., 2008), it is possible that moderately lame animals waited longer than sound animals,  
421 explaining the increase in the total time spent standing. In contrast, severely lame cows may postpone  
422 milking and wait lying down in the pasture as long as possible before being milked, which could also  
423 explain why they spend more time lying down. Experiments in farms using a milking parlour rather  
424 than an AMS should be done to confirm the discriminatory status of the standing position between the  
425 moderately lame group and the other two groups.

## 426 4.2 Behavioural and movement variables as relevant indicators to improve the performance of 427 existing lameness detection devices in pasture-based systems

### 428 *4.2.1 Variables providing additional information in pasture-based systems*

429 Most existing lameness detection systems use data collected from housed cows, such as computer  
430 vision based on digital cameras (Van Hertem et al., 2014) or the indoor behaviour of dairy cows from  
431 accelerometer data (Beer et al., 2016). In this regard, combining different types of data is often a  
432 means to obtain better performance of prediction, whatever the classification problem (Wang et al.,  
433 2018). Especially for lameness prediction, Beer et al. (2016) combined data from leg-attached  
434 accelerometers and from a noseband sensor to detect lameness in indoor-based systems, and reached  
435 good performance of prediction (sensitivity: 90.2 %; specificity: 91.7 %). In the same way, de Mol et  
436 al. (2013) combined activity data from accelerometers, milking data and data from computerised  
437 concentrate feeders, and obtained satisfactory performance of lameness prediction (sensitivity: 85.5 %;  
438 specificity: 88.8 %). Use of discriminant behavioural and movement variables is therefore certainly a  
439 potential way to enhance the performance of lameness prediction in pasture-based systems. However,  
440 it should be noted that lameness scores 0 and 1 cannot be discriminated (Appendix 7), probably  
441 because it is difficult for the observer to distinguish between a cow with a lameness score 0 (not lame)  
442 and a cow with a lameness score 1 (slight asymmetry, unidentifiable affected leg) during the scoring  
443 sessions. Furthermore, as explained in the previous section, discriminating variables between lameness

444 scores 2 and 3 should be confirmed by carrying out experiments in other farms that do not use an  
445 AMS. On the basis of this study, it thus seems possible to discriminate sound cows or cows with a  
446 slight asymmetry from severely lame cows using behavioural and movement variables at pasture, but  
447 the relevance of these variables for discriminating moderately lame cows from the other two groups  
448 has yet to be confirmed.

#### 449 *4.2.2 A potentially transferable approach in the field*

450 Behavioural and movement variables were obtained from (1) the collection of accelerometer  
451 and GPS data from dairy cows on commercial farms, (2) the application of a robust methodology to  
452 predict a wide range of behaviours every 10 seconds (Riaboff et al., 2019; Riaboff et al., 2020a) and  
453 then combination with the GPS position (Riaboff et al., 2020b), (3) an unsupervised segmentation on  
454 time series (Lavielle, 1999) to first isolate the outdoor periods and then to identify the behavioural  
455 bouts within each of these, and (4) a weighting of variables per cow or per day. Steps (1) and (2)  
456 provide automatic monitoring of cattle behaviour at pasture every 10 s. Step (3) ensures smoothing of  
457 the predicted behaviours to obtain continuous behavioural bouts without the need to set arbitrary  
458 criteria, as are often applied in commercial systems (Hendriks et al., 2020; Werner et al., 2018). Step  
459 (4) is a way of adapting these variables to each animal, each day of grazing and each farm, which is  
460 necessary for use in commercial farms. The combination of these four steps thus ensures a robust  
461 collection of behavioural and movement variables, which is easily transferable to the field.

462 In addition, neck-mounted accelerometer sensors are already used for other applications  
463 (“Medria Solutions,” 2020), which could facilitate transfer to the field if a centralised collection of raw  
464 accelerometer data is produced to respond to several issues (heat detection, welfare monitoring,  
465 lameness detection, etc.). In the same way, embedded GPS sensors can be used for several  
466 applications in pasture-based systems, including the development of targeted preventive treatments  
467 based on the identification of risk areas visited by cows (Agoulon et al., 2012), or the reduction of  
468 environmental impacts at farm level based on the identification of overused areas (Lush et al., 2018). It  
469 should therefore be possible to integrate the overall methodology used in our study into automatic  
470 lameness detection devices for pasture-based systems.

471                   4.3 Limits and perspectives

472                   The discriminant behavioural and movement variables identified can be used to enhance the  
473 performance of lameness detection systems in pasture-based farming, but the creation of a high-  
474 performance prediction model based exclusively on these variables seems difficult to achieve at the  
475 moment. The two main reasons for this are (i) the poor performance of the PLS-DA model ( $Q^2_{cum}$ :  
476 0.11; criterion of validity of the predictive model:  $Q^2_{cum} > 0.5$  (Tenenhaus, 1998)), and (ii) the need  
477 for the animals to go on pasture to collect data. Concerning the first point (i), other statistical models  
478 should be used to confirm our results. In our study, we chose a machine learning method (PLS-DA) to  
479 identify discriminant variables because (i) we had a large dataset of 863 observations (Appendix 5 b)  
480 and 37 innovative variables for which we had limited *a priori* knowledge, and (ii) cows were not  
481 followed long enough to ensure sufficient transitions between scores, especially for lameness score 3,  
482 nor were there enough animals (Appendix 5 b) to be able to distinguish the random cow effect from  
483 the fixed lameness effect in a probabilistic model. As this study has highlighted the relevance of 17  
484 behavioural and movement variables at pasture to discriminate between lameness scores, it would now  
485 be interesting to carry out this experiment again with an adapted protocol to apply a statistical model,  
486 such as an ordinal logistic regression, in order to conclude on the significance of these 17 variables,  
487 and to assess their ability to predict lameness scores at pasture. The second point (ii) is a key limitation  
488 because we can only predict behaviours on pasture, since the model developed in Riaboff et al. (2020a,  
489 2019) was fitted on behaviours observed for grazing exclusively. Consequently, additional data on  
490 cows in the barn is needed to monitor lameness throughout the year if (1) cows only have access to  
491 pasture for part of the year, and (2) cows could have access to pasture but prefer to remain in the barn,  
492 as observed in our study with farms using AMS (9.3 % of the data were discarded for this reason;  
493 *section 2.3.3*). The behavioural and movement variables calculated in this way are therefore relevant in  
494 addition to other variables measured on the animals in the barn, but are not sufficient to develop a  
495 prediction model based solely on these variables.

496                   Furthermore, from a technological point of view, the device used to automatically report GPS  
497 and accelerometer data is not usable in its current state for a field application. The major drawback is

498 the battery life, limited to six days, which requires the sensors to be removed every six days for  
499 recharging. As the sampling rate of both the GPS (1 Hz) and the accelerometer (59.5 Hz) sensors are  
500 high, a possible solution is to (i) reduce the sampling rate of the GPS sensor to compute the movement  
501 variables with the desired minimum precision, and (ii) to reduce the sampling rate of the accelerometer  
502 sensor to find an appropriate trade-off between battery life and the performance of the prediction of  
503 behaviours. Another solution would be using solar energy, as recently proposed for virtual fences  
504 (Acosta et al., 2020). Data extraction is also a key concern, as it is currently done manually. Automatic  
505 data transfer is also required for field use. In view of the relevance of such an approach for the  
506 diagnosis of lameness in pasture, these technological limitations are certainly worth removing to make  
507 the developed system functional.

## 508 **5. Conclusion**

509 This study aimed to identify behavioural and movement variables for dairy cows at pasture, calculated  
510 automatically from a methodology based on accelerometer and GPS sensors, to discriminate between  
511 lameness scores. To the authors' knowledge, this is one of the most comprehensive studies focusing on  
512 the relationship between the behaviour and the movement of dairy cows on pasture and lameness. We  
513 found that grazing and resting behaviours and the position (standing/lying) adopted by the cows while  
514 ruminating were modified with lameness, as were the exploratory dynamics of the cows. Therefore, 17  
515 variables derived from these behaviours (time spent grazing and resting, duration of grazing and  
516 resting bouts, duration before lying down, etc.) and related to the exploratory dynamics (distance,  
517 dispersion, distance travelled during grazing, etc.) may be relevant additional indicators to improve the  
518 performance of automatic lameness detection devices in pasture-based systems, although  
519 methodological and technological challenges still need to be addressed.

520 **Funding:** This study was funded by Terrena Innovation (Ancenis, France), Région Pays-de-la-Loire  
521 and BIOEPAR (INRAE, Oniris, Nantes, France).

522 **Acknowledgments:** We would like to thank the commercial farms in which the experiment was  
523 conducted. We also thank Marie Roig-Pons for her assistance in data collection and Romain Feron for  
524 the provision of his computer.

525 **Ethical considerations:** This study did not require approval from an ethics committee under French  
526 legislation. The only procedures performed on the cows consisted in visual locomotion scoring and  
527 fitting and removing the collars on the cows twice a week. Farmers were informed about lameness  
528 prevalence just after the experiment carried out in their farm, and a list of recommendations was  
529 provided to reduce the lameness prevalence in their farm.

530

## 531 **References**

532 Acosta, N., Barreto, N., Caitano, P., Marichal, R., Pedemonte, M., Oreggioni, J., 2020. Research  
533 platform for cattle virtual fences, in: 2020 IEEE International Conference on Industrial  
534 Technology (ICIT). Presented at the 2020 IEEE International Conference on Industrial  
535 Technology (ICIT), IEEE, Buenos Aires, Argentina, pp. 797–802.  
536 <https://doi.org/10.1109/ICIT45562.2020.9067313>

537 Agoulon, A., Malandrin, L., Lepigeon, F., Vénisse, M., Bonnet, S., Becker, C.A.M., Hoch, T., Bastian,  
538 S., Plantard, O., Beaudeau, F., 2012. A Vegetation Index qualifying pasture edges is related to  
539 *Ixodes ricinus* density and to *Babesia divergens* seroprevalence in dairy cattle herds. *Vet.*  
540 *Parasitol.* 185, 101–109. <https://doi.org/10.1016/j.vetpar.2011.10.022>

541 Agriculture and Horticulture Development Board, Stoneleigh Park, Kenilworth, Warwickshire, CV8  
542 2TL, 2020. AHDB: mobility score.

543 Almeida, P.E., Weber, P.S.D., Burton, J.L., Zanella, A.J., 2008. Depressed DHEA and increased  
544 sickness response behaviors in lame dairy cows with inflammatory foot lesions. *Domest.*  
545 *Anim. Endocrinol.* 34, 89–99. <https://doi.org/10.1016/j.domaniend.2006.11.006>

546 Barker, M., Rayens, W., 2003. Partial least squares for discrimination. *J. Chemom.* 17, 166–173.  
547 <https://doi.org/10.1002/cem.785>

548 Barker, Z.E., Amory, J.R., Wright, J.L., Mason, S.A., Blowey, R.W., Green, L.E., 2009. Risk factors  
549 for increased rates of sole ulcers, white line disease, and digital dermatitis in dairy cattle from  
550 twenty-seven farms in England and Wales. *J. Dairy Sci.* 92, 1971–1978.  
551 <https://doi.org/10.3168/jds.2008-1590>

552 Barker, Z.E., Leach, K.A., Whay, H.R., Bell, N.J., Main, D.C.J., 2010. Assessment of lameness  
553 prevalence and associated risk factors in dairy herds in England and Wales. *J. Dairy Sci.* 93,  
554 932–941. <https://doi.org/10.3168/jds.2009-2309>

555 Beer, G., Alsaad, M., Starke, A., Schuepbach-Regula, G., Müller, H., Kohler, P., Steiner, A., 2016.  
556 Use of Extended Characteristics of Locomotion and Feeding Behavior for Automated  
557 Identification of Lamé Dairy Cows. *PLOS ONE* 11, e0155796.  
558 <https://doi.org/10.1371/journal.pone.0155796>

559 Blackie, N., Bleach, E., Amory, J., Scaife, J., 2011. Impact of lameness on gait characteristics and  
560 lying behaviour of zero grazed dairy cattle in early lactation. *Appl. Anim. Behav. Sci.* 129,  
561 67–73. <https://doi.org/10.1016/j.applanim.2010.10.006>

562 Calenge, C., Dray, S., Royer-Carenzi, M., 2009. The concept of animals' trajectories from a data  
563 analysis perspective. *Ecol. Inform.* 4, 34–41. <https://doi.org/10.1016/j.ecoinf.2008.10.002>

564 Calenge, C., 2006. The package adehabitat for the R software: tool for the analysis of space and  
565 habitat use by animals. *Ecol. Model.* 197, 516–519.

566 Chen, T., He, T., Benesty, M., Khotilovich, V., Tang, Y., Cho, H., Chen, K., Mitchell, R., Cano, I.,  
567 Zhou, T., Li, M., Xie, J., Lin, M., Geng, Y., Li, Y., 2018. xgboost: Extreme Gradient  
568 Boosting.

569 Chong, I.-G., Jun, C.-H., 2005. Performance of some variable selection methods when  
570 multicollinearity is present. *Chemom. Intell. Lab. Syst.* 78, 103–112.  
571 <https://doi.org/10.1016/j.chemolab.2004.12.011>

572 Cohen, J., 1960. A Coefficient of Agreement for Nominal Scales. *Educ. Psychol. Meas.* 20, 37–46.  
573 <https://doi.org/10.1177/001316446002000104>

574 Cutler, J.H.H., Rushen, J., de Passillé, A.M., Gibbons, J., Orsel, K., Pajor, E., Barkema, H.W., Solano,  
575 L., Pellerin, D., Haley, D., Vasseur, E., 2017. Producer estimates of prevalence and perceived

576 importance of lameness in dairy herds with tiestalls, freestalls, and automated milking  
577 systems. *J. Dairy Sci.* 100, 9871–9880. <https://doi.org/10.3168/jds.2017-13008>

578 de Mol, R.M., André, G., Bleumer, E.J.B., van der Werf, J.T.N., de Haas, Y., van Reenen, C.G., 2013.  
579 Applicability of day-to-day variation in behavior for the automated detection of lameness in  
580 dairy cows. *J. Dairy Sci.* 96, 3703–3712. <https://doi.org/10.3168/jds.2012-6305>

581 Delagarde, R., Lamberton, P., 2015. Daily grazing time of dairy cows is recorded accurately using the  
582 Lifecorder Plus device. *Appl. Anim. Behav. Sci.* 165, 25–32.  
583 <https://doi.org/10.1016/j.applanim.2015.01.014>

584 Dolecheck, K.A., Silvia, W.J., Heersche, G., Chang, Y.M., Ray, D.L., Stone, A.E., Wadsworth, B.A.,  
585 Bewley, J.M., 2015. Behavioral and physiological changes around estrus events identified  
586 using multiple automated monitoring technologies. *J. Dairy Sci.* 98, 8723–8731.  
587 <https://doi.org/10.3168/jds.2015-9645>

588 Dumont, B., Fortun-Lamothe, L., Jouven, M., Thomas, M., Tichit, M., 2013. Prospects from  
589 agroecology and industrial ecology for animal production in the 21st century. *animal* 7, 1028–  
590 1043. <https://doi.org/10.1017/S1751731112002418>

591 Feldt, T., Schlecht, E., 2016. Analysis of GPS trajectories to assess spatio-temporal differences in  
592 grazing patterns and land use preferences of domestic livestock in southwestern Madagascar.  
593 *Pastoralism* 6. <https://doi.org/10.1186/s13570-016-0052-2>

594 Galindo, F., Broom, D.M., 2002. The Effects of Lameness on Social and Individual Behavior of Dairy  
595 Cows. *J. Appl. Anim. Welf. Sci.* 5, 193–201.  
596 [https://doi.org/10.1207/S15327604JAWS0503\\_03](https://doi.org/10.1207/S15327604JAWS0503_03)

597 Géoportail [WWW Document], 2006. . Portail Natl. Connaiss. Territ. Mis En Oeuvre Par IGN. URL  
598 <https://www.geoportail.gouv.fr/>

599 Green, L.E., Hedges, V.J., Schukken, Y.H., Blowey, R.W., Packington, A.J., 2002. The Impact of  
600 Clinical Lameness on the Milk Yield of Dairy Cows. *J. Dairy Sci.* 85, 2250–2256.  
601 [https://doi.org/10.3168/jds.S0022-0302\(02\)74304-X](https://doi.org/10.3168/jds.S0022-0302(02)74304-X)

602 Haskell, M.J., Rennie, L.J., Bowell, V.A., Bell, M.J., Lawrence, A.B., 2006. Housing System, Milk  
603 Production, and Zero-Grazing Effects on Lameness and Leg Injury in Dairy Cows. *J. Dairy*  
604 *Sci.* 89, 4259–4266. [https://doi.org/10.3168/jds.S0022-0302\(06\)72472-9](https://doi.org/10.3168/jds.S0022-0302(06)72472-9)

605 Hendriks, S.J., Phyn, C.V.C., Huzzey, J.M., Mueller, K.R., Turner, S.-A., Donaghy, D.J., Roche, J.R.,  
606 2020. Graduate Student Literature Review: Evaluating the appropriate use of wearable  
607 accelerometers in research to monitor lying behaviors of dairy cows. *J. Dairy Sci.* 103, 12140–  
608 12157. <https://doi.org/10.3168/jds.2019-17887>

609 Himmelmann, L., 2010. HMM: HMM - Hidden Markov Models.

610 Huxley, J.N., 2013. Impact of lameness and claw lesions in cows on health and production. *Livest.*  
611 *Sci.* 156, 64–70. <https://doi.org/10.1016/j.livsci.2013.06.012>

612 Kamphuis, C., Delarue, B., Burke, C.R., Jago, J., 2012. Field evaluation of 2 collar-mounted activity  
613 meters for detecting cows in estrus on a large pasture-grazed dairy farm. *J. Dairy Sci.* 95,  
614 3045–3056. <https://doi.org/10.3168/jds.2011-4934>

615 Lavielle, M., 1999. Detection of multiple changes in a sequence of dependent variables. *Stoch.*  
616 *Process. Their Appl.* 83, 79–102. [https://doi.org/10.1016/S0304-4149\(99\)00023-X](https://doi.org/10.1016/S0304-4149(99)00023-X)

617 Leach, K.A., Tisdall, D.A., Bell, N.J., Main, D.C.J., Green, L.E., 2012. The effects of early treatment  
618 for hindlimb lameness in dairy cows on four commercial UK farms. *Vet. J.* 193, 626–632.  
619 <https://doi.org/10.1016/j.tvjl.2012.06.043>

620 Lush, L., Wilson, R.P., Holton, M.D., Hopkins, P., Marsden, K.A., Chadwick, D.R., King, A.J., 2018.  
621 Classification of sheep urination events using accelerometers to aid improved measurements  
622 of livestock contributions to nitrous oxide emissions. *Comput. Electron. Agric.* 150, 170–177.  
623 <https://doi.org/10.1016/j.compag.2018.04.018>

624 McSweeney, D., Foley, C., Halton, P., O'Brien, B., 2015. Calibration of an automated grass  
625 measurement tool to enhance the precision of grass measurement in pasture based farming  
626 systems. Presented at the Teagasc Ag conference, Tullamore.

627 Medria Solutions [WWW Document], 2020. . Farmlife Bouquet Enter New Era Monit. URL  
628 <https://www.medria.fr/en/solutions/herd-monitoring.html> (accessed 11.9.20).

629 Navarro, G., Green, L.E., Tadich, N., 2013. Effect of lameness and lesion specific causes of lameness  
630 on time budgets of dairy cows at pasture and when housed. *Vet. J.* 197, 788–793.  
631 <https://doi.org/10.1016/j.tvjl.2013.05.012>

632 Norring, M., Häggman, J., Simojoki, H., Tamminen, P., Winckler, C., Pastell, M., 2014. Short  
633 communication: Lameness impairs feeding behavior of dairy cows. *J. Dairy Sci.* 97, 4317–  
634 4321. <https://doi.org/10.3168/jds.2013-7512>

635 O’Leary, N.W., Byrne, D.T., O’Connor, A.H., Shalloo, L., 2020. Invited review: Cattle lameness  
636 detection with accelerometers. *J. Dairy Sci.* 103, 3895–3911. <https://doi.org/10.3168/jds.2019-17123>

637

638 Pebesma, E.J., Bivan, R.S., 2005. Classes and methods for spatial data in R.

639 Politiek, R.D., Distl, O., Fjeldaas, T., Heeres, J., McDaniel, B.T., Nielsen, E., Peterse, D.J., Reurink,  
640 A., Strandberg, P., 1986. Importance of claw quality in cattle: review and recommendations to  
641 achieve genetic improvement. Report of the e.a.a.p. working group on “claw quality in cattle.”  
642 *Livest. Prod. Sci.* 15, 133–152.

643 R Core Team, 2020. . R Foundation for Statistical Computing, Vienna, Austria.

644 Riaboff, L., Aubin, S., Bédère, N., Couvreur, S., Madouasse, A., Goumand, E., Chauvin, A., Plantier,  
645 G., 2019. Evaluation of pre-processing methods for the prediction of cattle behaviour from  
646 accelerometer data. *Comput. Electron. Agric.* 165, 104961.  
647 <https://doi.org/10.1016/j.compag.2019.104961>

648 Riaboff, L., Poggi, S., Madouasse, A., Couvreur, S., Aubin, S., Bédère, N., Goumand, E., Chauvin, A.,  
649 Plantier, G., 2020a. Development of a methodological framework for a robust prediction of  
650 the main behaviours of dairy cows using a combination of machine learning algorithms on  
651 accelerometer data. *Comput. Electron. Agric.* 169, 105179.  
652 <https://doi.org/10.1016/j.compag.2019.105179>

653 Riaboff, L., Couvreur, S., Madouasse, A., Roig-Pons, M., Aubin, S., Massabie, P., Chauvin, A.,  
654 Bédère, N., Plantier, G., 2020b. Use of Predicted Behavior from Accelerometer Data  
655 Combined with GPS Data to Explore the Relationship between Dairy Cow Behavior and  
656 Pasture Characteristics. *Sensors* 20, 4741. <https://doi.org/10.3390/s20174741>

657 Sanchez, G., 2016. Partial Least Squares (PLS) Data Analysis Methods.

658 Tenenhaus, M., 1998. Régression PLS (LA) Théorie et pratique, Editions Technip. ed. Paris.

659 van den Pol-van Dasselaar, A., Hennessy, D., Isselstein, J., 2020. Grazing of Dairy Cows in Europe—  
660 An In-Depth Analysis Based on the Perception of Grassland Experts. *Sustainability* 12, 1098.  
661 <https://doi.org/10.3390/su12031098>

662 Van Hertem, T., Viazzi, S., Steensels, M., Maltz, E., Antler, A., Alchanatis, V., Schlageter-Tello,  
663 A.A., Lokhorst, K., Romanini, E.C.B., Bahr, C., Berckmans, D., Halachmi, I., 2014.  
664 Automatic lameness detection based on consecutive 3D-video recordings. *Biosyst. Eng.* 119,  
665 108–116. <https://doi.org/10.1016/j.biosystemseng.2014.01.009>

666 Van Nuffel, A., Zwertvaegher, I., Van Weyenberg, S., Pastell, M., Thorup, V., Bahr, C., Sonck, B.,  
667 Saeys, W., 2015. Lameness Detection in Dairy Cows: Part 2. Use of Sensors to Automatically  
668 Register Changes in Locomotion or Behavior. *Animals* 5, 861–885.  
669 <https://doi.org/10.3390/ani5030388>

670 Vermunt, J.J., Greenough, P.R., 1995. Structural characteristics of the bovine claw: Horn growth and  
671 wear, horn hardness and claw conformation. *Br. Vet. J.* 151, 157–180.  
672 [https://doi.org/10.1016/S0007-1935\(95\)80007-7](https://doi.org/10.1016/S0007-1935(95)80007-7)

673 Walker, S.L., Smith, R.F., Routly, J.E., Jones, D.N., Morris, M.J., Dobson, H., 2008. Lameness,  
674 Activity Time-Budgets, and Estrus Expression in Dairy Cattle. *J. Dairy Sci.* 91, 4552–4559.  
675 <https://doi.org/10.3168/jds.2008-1048>

676 Wang, J., He, Z., Zheng, G., Gao, S., Zhao, K., 2018. Development and validation of an ensemble  
677 classifier for real-time recognition of cow behavior patterns from accelerometer data and  
678 location data. *PLOS ONE* 13, e0203546. <https://doi.org/10.1371/journal.pone.0203546>

679 Wei, T., 2017. R package “corrplot”: Visualization of a Correlation Matrix.

680 Weigele, H.C., Gygax, L., Steiner, A., Wechsler, B., Burla, J.-B., 2018. Moderate lameness leads to  
681 marked behavioral changes in dairy cows. *J. Dairy Sci.* 101, 2370–2382.  
682 <https://doi.org/10.3168/jds.2017-13120>

683 Werner, J., Leso, L., Umstatter, C., Niederhauser, J., Kennedy, E., Geoghegan, A., Shalloo, L., Schick,  
684 M., O'Brien, B., 2018. Evaluation of the RumiWatchSystem for measuring grazing behaviour  
685 of cows. *J. Neurosci. Methods* 300, 138–146. <https://doi.org/10.1016/j.jneumeth.2017.08.022>


686 Whay, H.R., Shearer, J.K., 2017. The Impact of Lameness on Welfare of the Dairy Cow. *Lameness*  
687 *Cattle* 33, 153–164. <https://doi.org/10.1016/j.cvfa.2017.02.008>

688 Willshire, J.A., Bell, N.J., 2009. An Economic Review of Cattle Lameness 17, 136–141.

689 Yunta, C., Guasch, I., Bach, A., 2012. Short communication: Lying behavior of lactating dairy cows is  
690 influenced by lameness especially around feeding time. *J. Dairy Sci.* 95, 6546–6549.  
691 <https://doi.org/10.3168/jds.2012-5670>

692

693




Legend:

- Main sections of Materials and Methods
- Method applied for the step
- Steps carried out in each section

695

696 Figure 1. Overview of the main steps applied in the Material and methods section.


697




698

699 Figure 2. Time sequence of the experiment carried out to collect sensor data from dairy cows and the

700 associated lameness score. The device used to obtain accelerometer and GPS data is displayed.



701 Figure 3. Description of the successive steps carried out to obtain the predicted behaviours for every  
 702 10 s time-window and the corresponding GPS coordinates.



703

704 Figure 4. Principle of the identification of (a) outdoor periods for each cow on each day she was

705 equipped and (b) bouts of behaviours within each outdoor period using Lavielle segmentation.



	Farm 1	Farm 2	Farm 3	Total
Number of equipped cows / Herd size	15/54	27/71	26/61	68
Parity (mean $\pm$ sd)	1.7 $\pm$ 0.7	2.3 $\pm$ 1.2	1.5 $\pm$ 0.6	
Days in milk (mean $\pm$ sd)	145 $\pm$ 56	151 $\pm$ 63	176 $\pm$ 92	
Average milk yield	14,066 kg	9,912 kg	8,542 kg	
Number of days of experiment	21	12	25	58
Lameness score at the start of the experiment <sup>1</sup>				
Lameness score 0	1 (6%)	6 (22%)	0 (0%)	7 (10%)
Lameness score 1	7 (46%)	17 (63%)	12 (44%)	36 (53%)
Lameness score 2	6 (40%)	4 (15%)	12 (44%)	22 (32%)
Lameness score 3	1 (6%)	0 (0%)	2 (7.4%)	3 (4%)

710 Table 1. Dairy cows under study and duration of animal monitoring

711 <sup>1</sup> Number of equipped cows (percentage of the herd)

712

713 Table 2. Criteria to assign lameness score using the Agriculture and Horticulture Development Board

Lameness score (LS)	Clinical description	Decision criteria
LS 0	Satisfactory mobility	Walking has an even distribution of weight and rhythm is regular between the four legs. The back is straight.
LS 1	Imperfect mobility	Asymmetric strides (rhythm or weight transfer) or shorter. The affected member is not identifiable.
LS 2	Altered mobility	The lame leg is identifiable and/or the strides are clearly shorter. The back is usually arched.
LS 3	Severely altered mobility	The speed is lower than those of the herd. The lame leg is clearly identifiable as the cow hardly stands on this member. The back is arched both when standing and walking.

714 method.

715

716 Table 3. Variables calculated from the behavioural bouts and movement of cows on pasture.

Variable	Definition	Calculation	Weighting	Additional explanation
Time <sub>out</sub> (1)	Time spent outside	$\sum_{k=1}^n \text{Time}_k(c,d)$ for each outdoor period $k$ for the cow $c$ on day $d$ .	Time <sub>out,max</sub>	
Time <sub>b</sub> (2)	Time spent expressing behaviour $b$	$\sum_{k=1}^n \sum_{j=1}^m \text{Time}_{j,k}(c,d)$ for any bout $j$ of the behaviour $b$ within the outdoor period $k$ , for the cow $c$ on day $d$ .	Time <sub>out</sub>	Time <sub>b</sub> was computed for each predicted and grouped behaviour.
Number bouts <sub>b</sub> (3)	Number of bouts with the behaviour $b$	$\sum_{k=1}^n \sum_{j=1}^m \mathbb{1}_B\{\text{Bout}_{j,k}(c,d)\}$ where $\mathbb{1}_B = 1$ if $\text{bout}_{j,k}(c,d)$ is associated with the behaviour $b$ , 0 otherwise, for any bout $j$ of the behaviour $b$ within the outdoor period $k$ , for the cow $c$ on day $d$ .	Time <sub>out</sub>	Number_bouts <sub>b</sub> was computed for each predicted and grouped behaviour except for <i>walking</i> , as the number of occurrences was very low and not representative.
Duration bouts <sub>b</sub> (4)	Average bout duration for behaviour $b$	$\frac{\text{Time}_b}{\text{Number\_Bouts}_b}$ with Time <sub>b</sub> and Number_Bouts <sub>b</sub> defined in N°2 and N°3.	Time <sub>out</sub>	Duration_bouts <sub>b</sub> was computed for each predicted and grouped behaviour defined in section 2.2 except for <i>walking</i> , as the number of occurrences was too low to calculate a representative average duration.
Round trips (5)	Number of round trips between pasture and farm	$s-1$ for the $s$ outdoor/indoor segments obtained with the Lavielle function for the cow $c$ on day $d$ .	Time <sub>out</sub>	
Distance (6)	Total Euclidean distance travelled	$\sum_{k=1}^n \sum_{t=1}^T \text{Distance}_{t,k}(c,d)$ for any 10 s time-window $t$ within the outdoor period $k$ , for the cow $c$ on day $d$ .	Distance <sub>max</sub>	The distance within each 10 s time-window was calculated using the first and last GPS coordinates of the window. The GPS error was considered by (i) replacing by 0 the distance of time windows with stationary behaviour, and (ii) removing time windows with a speed greater than 25 m.s <sup>-1</sup> to avoid considering aberrant GPS coordinates in the distance calculation.
Dispersion (R2n) (7)	Sum of net squared displacement between each location and the	$\sum_{k=1}^n \sum_{t=1}^T \text{R2n}_{t,k}(c;d)$	R2n <sub>max</sub>	R2n was strongly related to the dispersion. The more a cow explored the pasture from the pasture entrance, the higher the R2n was. We refer to Calenge et al. (2009) for a more detailed

	first location in the pasture	<sup>1</sup> for any 10s time-window $t$ within the outdoor period $k$ , for the cow $c$ on day $d$ .		explanation. The GPS error was considered by (i) replacing by 0 the R2n in time windows with stationary behaviour, and (ii) removing time windows with a speed greater than 25 m.s <sup>-1</sup> to avoid considering aberrant GPS coordinates in the distance calculation.
Distance Grazing (8)	Euclidean distance travelled during grazing bouts	$\sum_{k=1}^n \sum_{j=1, b=\text{Grazing}}^m \sum_{t=1}^T \text{Distance}_{t,k}(b;c;d)$ for any 10s time-window $t$ within the bout $j$ with the behaviour $b = \text{Grazing}$ within the outdoor period $k$ , for the cow $c$ on day $d$ .	Distance	Distance Grazing was calculated in the same way as the distance (6) considering only the grazing bouts.
Distance Walking (9)	Euclidean distance travelled during walking	$\sum_{k=1}^n \sum_{t=1, b=\text{Walking}}^T \text{Distance}_{t,k}(b;c;d)$ for any 10s time-window $t$ where the behaviour $b = \text{Walking}$ has been predicted, within the outdoor period $k$ , for the cow $c$ on day $d$ .	Distance	Distance Walking was calculated in the same way as the distance (6) considering only windows where the walking behaviour was predicted. The distance travelled during each walking time window was only considered if the animal's movement corresponded to a minimum speed of 2 m.s <sup>-1</sup> .
Distance ratio (10)	Distance travelled during grazing over the distance travelled during grazing and walking	$\frac{\text{Distance\_Grazing}}{\text{Distance\_Grazing} + \text{Distance\_Walking}}$ with Distance_Grazing and Distance_Walking defined in (8) and (9), respectively.		
Speed (11)	Average speed during walking	$\frac{\text{Distance\_Walking}}{\text{Time\_Walking}}$ with Distance_Walking and Time_Walking defined in (9) and (2), respectively.		When no walking was predicted during an outdoor period, the missing speed for the cow $c$ on day $d$ was replaced by the average speed of the group of cows with the same lameness score as the cow $c$ .
Duration before lying (12)	Average duration before lying down after entrance to pasture	$\frac{\sum_{k=1}^n \text{Time\_first\_lying}_{k(c,d)} - \text{Time\_entrance}_{k(c,d)}}{\sum_{k=1}^n \mathbb{1}_B\{\text{Period}_{k(c,d)}\}}$ where Time_first_lying <sub>k(c,d)</sub> is the GPS time when the cow $c$ first lay down in the outdoor period $k$ on day $d$ . Time_entrance <sub>k(c,d)</sub> is the time recorded with the GPS when the cow $c$ entered the pasture in the outdoor period $k$ on day $d$ . $\mathbb{1}_B = 1$ if there is at least one lying bout over the outdoor period $k$ for the cow $c$ on day $d$ .		When no lying down was predicted during an outdoor period, the missing duration before lying down for the cow $c$ on day $d$ was replaced by the average duration before lying down of the group of cows with the same lameness score as the cow $c$ . It should be noted that the information “absence of lying” has already been provided by the behavioural variables.

718 Table 4. Mean and standard errors obtained for each weighted behavioural and movement variable

	LS = 0_1 (mean ± se)	LS = 2 (mean ± se)	LS = 3 (mean ± se)
Time <sub>out</sub>	0.76 ± 0.01	0.73 ± 0.01	0.69 ± 0.03
Round trips	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Time <sub>grazing</sub>	0.27 ± 0.01	0.17 ± 0.01	0.06 ± 0.01
Time <sub>ruminating lying</sub>	0.29 ± 0.01	0.22 ± 0.01	0.16 ± 0.03
Time <sub>resting lying</sub>	0.18 ± 0.01	0.22 ± 0.01	0.34 ± 0.03
Time <sub>walking</sub>	0.01 ± 0.00	0.01 ± 0.00	0.00 ± 0.00
Time <sub>ruminating standing</sub>	0.06 ± 0.00	0.11 ± 0.01	0.13 ± 0.02
Time <sub>resting standing</sub>	0.04 ± 0.00	0.06 ± 0.01	0.07 ± 0.02
Time <sub>lying</sub>	0.47 ± 0.01	0.44 ± 0.02	0.50 ± 0.03
Time <sub>standing</sub>	0.10 ± 0.01	0.17 ± 0.01	0.20 ± 0.02
Time <sub>ruminating</sub>	0.35 ± 0.01	0.33 ± 0.01	0.29 ± 0.03
Time <sub>resting</sub>	0.22 ± 0.01	0.28 ± 0.01	0.41 ± 0.03
Nb bouts <sub>grazing</sub>	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Nb bouts <sub>ruminating lying</sub>	0.01 ± 0.00	0.008 ± 0.00	0.007 ± 0.00
Nb bouts <sub>resting lying</sub>	0.01 ± 0.00	0.01 ± 0.00	0.03 ± 0.01
Nb bouts <sub>ruminating standing</sub>	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Nb bouts <sub>resting standing</sub>	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
Nb bouts <sub>lying</sub>	0.02 ± 0.00	0.02 ± 0.00	0.03 ± 0.01
Nb bouts <sub>standing</sub>	0.01 ± 0.00	0.01 ± 0.00	0.01 ± 0.00
Nb bouts <sub>ruminating</sub>	0.01 ± 0.00	0.01 ± 0.00	0.01 ± 0.00
Nb bouts <sub>resting</sub>	0.01 ± 0.00	0.02 ± 0.00	0.03 ± 0.01
Duration bouts <sub>grazing</sub>	0.11 ± 0.01	0.08 ± 0.01	0.04 ± 0.01
Duration bouts <sub>ruminating lying</sub>	0.10 ± 0.00	0.08 ± 0.01	0.08 ± 0.01
Duration bouts <sub>resting lying</sub>	0.06 ± 0.00	0.07 ± 0.01	0.15 ± 0.03
Duration bouts <sub>ruminating standing</sub>	0.02 ± 0.00	0.04 ± 0.01	0.04 ± 0.01
Duration bouts <sub>resting standing</sub>	0.01 ± 0.00	0.02 ± 0.00	0.03 ± 0.01
Duration bouts <sub>lying</sub>	0.08 ± 0.00	0.02 ± 0.0	0.12 ± 0.01
Duration bouts <sub>standing</sub>	0.02 ± 0.00	0.03 ± 0.00	0.04 ± 0.01
Duration bouts <sub>ruminating</sub>	0.06 ± 0.00	0.06 ± 0.00	0.06 ± 0.01
Duration bouts <sub>resting</sub>	0.04 ± 0.00	0.04 ± 0.00	0.09 ± 0.03
Distance	0.50 ± 0.01	0.43 ± 0.02	0.30 ± 0.03
R2n	0.48 ± 0.01	0.43 ± 0.02	0.29 ± 0.03
Distance Walking	0.13 ± 0.01	0.13 ± 0.01	0.13 ± 0.03
Distance Grazing	0.49 ± 0.02	0.34 ± 0.02	0.21 ± 0.03
Distance Ratio	0.55 ± 0.02	0.43 ± 0.03	0.30 ± 0.05
Speed	3.42 ± 0.10	3.43 ± 0.08	3.10 ± 0.05
Duration Before Lying	63.5 ± 2.08	55.4 ± 3.56	34.6 ± 4.13

719 before

standardisation

according to

lameness score (LS).

Notations are those

used in Table 3.