



Current Understanding of Leaf Senescence in Rice

Sichul Lee, Celine Masclaux-Daubresse

► To cite this version:

Sichul Lee, Celine Masclaux-Daubresse. Current Understanding of Leaf Senescence in Rice. International Journal of Molecular Sciences, 2021, 22 (9), pp.1-19. 10.3390/ijms22094515 . hal-03329779

HAL Id: hal-03329779

<https://hal.inrae.fr/hal-03329779>

Submitted on 31 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Review

Current Understanding of Leaf Senescence in Rice

Sichul Lee ^{1,*}  and Celine Masclaux-Daubresse ²

¹ Center for Plant Aging Research, Institute for Basic Science (IBS), Daegu 42988, Korea

² Institut Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, 78000 Versailles, France; celine.masclaux-daubresse@inrae.fr

* Correspondence: scironlee@gmail.com; Tel.: +82-53-785-1848

Abstract: Leaf senescence, which is the last developmental phase of plant growth, is controlled by multiple genetic and environmental factors. Leaf yellowing is a visual indicator of senescence due to the loss of the green pigment chlorophyll. During senescence, the methodical disassembly of macromolecules occurs, facilitating nutrient recycling and translocation from the sink to the source organs, which is critical for plant fitness and productivity. Leaf senescence is a complex and tightly regulated process, with coordinated actions of multiple pathways, responding to a sophisticated integration of leaf age and various environmental signals. Many studies have been carried out to understand the leaf senescence-associated molecular mechanisms including the chlorophyll breakdown, phytohormonal and transcriptional regulation, interaction with environmental signals, and associated metabolic changes. The metabolic reprogramming and nutrient recycling occurring during leaf senescence highlight the fundamental role of this developmental stage for the nutrient economy at the whole plant level. The strong impact of the senescence-associated nutrient remobilization on cereal productivity and grain quality is of interest in many breeding programs. This review summarizes our current knowledge in rice on (i) the actors of chlorophyll degradation, (ii) the identification of stay-green genotypes, (iii) the identification of transcription factors involved in the regulation of leaf senescence, (iv) the roles of leaf-senescence-associated nitrogen enzymes on plant performance, and (v) stress-induced senescence. Compiling the different advances obtained on rice leaf senescence will provide a framework for future rice breeding strategies to improve grain yield.

Keywords: rice; leaf senescence; productivity; chlorophyll breakdown; stay-green; transcription factors; phytohormones; nitrogen remobilization



Citation: Lee, S.;

Masclaux-Daubresse, C. Current Understanding of Leaf Senescence in Rice. *Int. J. Mol. Sci.* **2021**, *22*, 4515. <https://doi.org/10.3390/ijms22094515>

Academic Editor: Yasuhito Sakuraba

Received: 31 March 2021

Accepted: 24 April 2021

Published: 26 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The leaf is the primary photosynthetic organ for energy production and nutrient assimilation at the growth and mature development stages [1]. Senescence occurs in a coordinated manner, starting from the tip and margins of the leaf toward the base and petiole in many plant species [2]. In these tissues, cells undergo a sequential disorganization of cellular organelles, and changes in gene expression and metabolism are orderly managed [1–3]. Leaf senescence is the last step of organ life and the last development stage in their life history for many plant species. Senescence events are basically controlled by age information and occur irrespective of environmental conditions. However, leaf senescence onset and senescence intensity can be modulated by many internal cues that perceive and integrate phytohormonal signals, nutrient status, water usage, light quality and length, and climate change [1]. Although leaf senescence is mainly described by the visual yellowing phenotype related to chlorophyll breakdown, as it can be seen by autumnal color changes of leaves, chlorophyll degradation is not the first event in the long leaf senescence program [4].

Basically, plants follow monocarpic senescence or polycarpic senescence [5,6]. Monocarpic plants such as annual cereal crops have only a single reproductive event in their life cycle. They produce a high number of mature offspring, such as seeds, and will initiate

the next life cycle upon germination [5,6]. On the other hand, polycarpic plants (also known as perennial plants) such as trees and shrubs undergo many reproduction cycles and continue growing across many seasons [6]. Leaf senescence usually precedes plant death in monocarpic species and facilitates nutrient mobilization to the seeds, i.e., to the next generation. In perennial plants, leaf senescence facilitates whole plant lifespan through nutrient recycling, control of water status, and adaptation to the environment and seasonal changes. In both cases, the main role of leaf senescence is for nutrient management and export to growing sinks.

Oryza sativa (rice) is one of the major staple crops and a key nutritional source for half of the world's population [7]. Then, increased grain yield in rice is crucial to cope with the rising food demands due to global population growth. Rice is a monocarpic plant in which leaf senescence overlaps the reproductive stage; thus, premature senescence can reduce grain yield under adverse environmental conditions [6]. In cereal crops such as rice, all the nutrients are transported predominantly from leaves to grains [8]. Leaves capture solar energy for photosynthetic carbon (C) assimilation and provide grains with C in the form of sugars; decreased photosynthetic activity during senescence reduces C fixation and sugar export to grains. By contrast, nitrogen (N) can be transported to grains through N remobilization, and increased protein breakdown during senescence promotes N remobilization from leaves to grains [9,10]. Subsequently, leaf senescence influences the C/N balance of resource mobilization and allocation to the grains [11]. Delayed leaf senescence can increase grain yield due to continued photosynthetic activity and C fixation, whereas it decreases grain protein content due to delayed N remobilization from the source leaves [11]. Accelerated leaf senescence can increase grain protein content [12]. Thus, the timing of leaf senescence is crucial for regulating grain yield and quality during the grain-filling period in rice.

As genome-wide alterations in gene expression occur during leaf senescence, genes that are controlling leaf senescence in plants are designated as senescence-associated genes (SAGs), and many SAGs also have been isolated and characterized in rice [13–16]. They are involved in the degradation of macromolecules, nutrient relocation, defense mechanisms, transcriptional regulation, and signal transduction [1,16]. This review summarizes the recent research developments in rice leaf senescence. It includes knowledge about the chlorophyll degradation pathway, description of stay-green traits, transcriptional regulations, impacts of leaf senescence on N metabolism, the relationship between senescence and stress, and finally, it proposes perspectives for the future.

2. Chlorophyll Degradation Pathways in Rice

Leaf senescence is monitored by the gradual loss of green pigments, which is mainly due to chlorophyll (Chl) degradation [4]. Most Chl are present in protein complexes in leaves because it is critical to prevent cells from the photooxidative damages that may be caused by the free Chl molecules if disassociated from the light-harvesting Chl-binding complex proteins (LHCs) of the thylakoids [17]. As Chl degradation is preceded by the degradation of LHCs in senescent leaves [18], the biochemical pathway of Chl breakdown is tightly controlled, and the released Chl is converted to non-phototoxic colorless catabolites (phyllobilins) via the PAO (Pheophorbide *a* Oxygenase)/phyllobilin pathway, and stored in the vacuole (Figure 1a) [19]. This catabolic pathway starts in senescing chloroplasts by the conversion of Chl *b* to Chl *a*, which is a degradable form and is then followed by the removal of the Mg²⁺ ion and the phytol moiety. Subsequently, the cyclic tetrapyrrole is linearized and catabolized to a primary fluorescent Chl catabolite (*p*FCC), which is colorless (Figure 1a). All of these reactions are common in higher plant species and rice, and all Chl catabolic enzymes (CCEs) have been well characterized [17].




Figure 1. The PAO/phyllobilin pathway of chlorophyll (Chl) breakdown in leaves and in silico analysis of the expression of Chl catabolic enzymes (CCEs) in rice flag leaves during the grain-filling stage. **(a)** Enzymatic reactions of the first part of the Chl degradation pathway, from Chl *b* to primary fluorescent Chl catabolite (pFCC), that take place inside the chloroplast. See the text for abbreviations of enzyme names. Expression data were obtained from RNA-seq analysis **(b)** [20] and public microarray data **(c)** [21]. DAH; days after heading. Each expression data is normalized to starting time points **(b)**, 4 DAH; **(c)**, 1 DAH, respectively). Based on the transcript levels of starting time points, the relative gene expression levels of other time points were quantified. The blue or red colors indicate the lower or higher transcript abundances than those of relevant control (white), respectively.

The conversion of Chl *b* to Chl *a* requires the action of two enzymes, Chl *b* Reductase (CBR) and 7-hydroxymethyl Chl *a* Reductase (HCAR, Figure 1a). Rice has two CBR, named NONYELLOW COLORING1 (OsNYC1) and NYC1-LIKE (OsNOL) [22,23]. The co-localization of OsNOL and OsNYC1 in the thylakoid membrane, as a heterodimer is required for Chl *b* reductase activity for the conversion of Chl *b* to 7-hydroxymethyl Chl *a* (HMChl *a*) and degradation of light-harvesting complex II during senescence [23]. The second step of the reduction of Chl *b* is the conversion of HMChl *a* to Chl *a*, which is catalyzed by HMChl *a* reductase (HCAR) [17]. The rice genome contains a single HCAR homolog (OsHCAR), and its knockout mutant displays persistent green leaves during senescence, accumulating HMChl *a* and pheophytin *a* (Pheo *a*), which is a downstream intermediate of the Chl breakdown pathway [24]. These results indicate that HCAR may play a pivotal role in protecting plants by inhibiting the accumulation of HMChl *a* and Pheo *a* in leaves [24].

Mendel's green cotyledon gene, the *Stay-Green* (SGR) gene, encodes Mg^{2+} -dechelatase, which is responsible for the removal of Mg^{2+} from Chl *a* to yield Pheo *a* [25]. SGR functions

as a key positive regulator of Chl degradation by physically interacting with the CCEs and LHCII proteins [26,27]. The rice *ossgr* mutants typically show a strong stay-green phenotype during both natural and dark-induced senescence because of the stable Chl-protein complex and thylakoid membrane structures [28,29]. Despite their greener phenotypes, the *ossgr* mutants show the same decrease in photosynthetic capability during senescence as wild type (WT), leading to no yield advantage [28–30]. An SGR rice homolog, SGR-like (SGRL), also displays Mg^{2+} -dechelatase activity, but against the chlorophyllide *a* precursor and not against Chl *a* [25]. The *OsSGRL* gene is downregulated during senescence, and its overexpression promotes Chl degradation and reduces the Chl-binding protein levels [31].

The removal of the phytol chain of Pheo *a* increases the solubility of the further breakdown products and is catalyzed by the Pheophytinase (PPH), which yields the highly phototoxic intermediate pheophorbide *a* (Pheide *a*) [32]. The Non-Yellow Coloring 3 (OsNYC3) pheophytinase is responsible for this step in rice [33]. The PAO converts the toxic Pheide *a* to a red Chl catabolite (RCC), and further to the primary fluorescent Chl catabolite (*p*FCC) that is generated by the Red Chl Catabolite Reductase (RCCR) [34]. Since the colored intermediates Pheide *a* and RCC are phototoxic, the knockdown and knockout mutations in *OsPAO* and *OsRCCR1* generate cell death symptoms [35,36]. The *p*FCC catabolites are exported from the chloroplast, modified in the cytosol/ER, and then transported into the vacuole where they accumulate as non-fluorescent Chl catabolites [37].

Analyses of CCE expression patterns can infer their functions during senescence. Their transcript levels in rice were analyzed using publically available RNA-seq data [20] and microarray data [21] derived from flag leaves grown under natural field conditions (Figure 1b, c). The results show that the expressions of *OsNYC1*, *OsNOL*, *OsSGR*, *OsNYC3*, *OsPAO*, and *OsRCCR1* were higher during the grain-filling stage, but *OsHCAR* transcripts were more abundant in the early grain-filling stage. Most of the CCEs rice mutants display stay-green phenotypes during natural- and/or artificially induced-senescence that are discussed in the next section.

3. Stay-Green Traits in Rice

Leaf senescence is easily observed because Chl degradation leads to leaf yellowing [38]. Stay-green is the heritable delayed senescence phenotype observed in comparison with a reference genotype, and *stay-green* mutants refer to genotypes that retain green leaves during senescence [39,40]. Maintaining leaf Chl levels and active photosynthetic capacity extends the time for assimilation processes and grain development, thereby increasing crop yield [41,42]. The genetic and physiological ways to obtain seemingly similar stay-green phenotypes are diverse, and five types of stay-green plants have been reported [39]. Their differences are based on the Chl behavior during senescence and their photosynthetic activities [39,43].

The Type A stay-green plants initiate senescence late but lose Chl at a normal rate. In Type B stay-green plants, senescence is started on schedule, but thereafter, the decline of the photosynthetic activity is comparatively slower than in WT. In Type C stay-green plants, Chl may be retained more or less indefinitely because of defects in pigment breakdown, but functional senescence occurs on a normal time-scale beneath the cosmetic surface of retained pigmentation. Therefore, Type C stay-greens are named cosmetic stay-green. The Type D stay-greens are described as pseudo-stay-greens because their leaves die before or in the middle of the senescence process as if freezing, and subsequently, the plant still appears green after death. The Type E stay-green plants accumulate higher levels of Chl in photosynthetic tissues, resulting in a delay of senescence and the conservation of green tissue. However, the photosynthetic capacity of Type E stay-greens follows the normal ontogenetic pattern. Both Type A and B stay-green plants are termed functional stay-greens as the duration of the photosynthetic capacity is extended. These stay-green types are sought by breeders as they can result in higher yield. Several studies have reported a synergistic effect by combining functional stay-green and other useful traits such as tolerance to drought and heat stress [44]. On the contrary, the Type C, D, and E plants are

categorized as non-functional stay-green that retain greenness due to the impairment of Chl catabolism but lack the maintenance of leaf functionality during senescence [39,45]. Most CCE mutants in rice display a stay-green phenotype during senescence but lack an extended photosynthetic capacity and grain yield advantage; therefore, these mutants are regarded as non-functional stay-green plants.

Due to the high importance of maintaining photosynthesis longer and/or in promoting nutrient recycling and mobilization throughout the plant, the stay-green traits were investigated in many plant species, including rice, maize, Arabidopsis, soybean, barley, tomato, pepper, and wheat, and they were used as markers in plant breeding programs [46–52].

Several studies report that the Chl content in rice is regulated by a quantitative trait locus (QTL) [47,53–55]. Analyses of Chl content and the degree of greenness in the flag and second leaves identified 46 QTLs that were associated with delayed senescence [53]. Cha et al. mapped the *stay-green* (*sgr*) locus, which is responsible for maintaining greenness during leaf senescence, but not photosynthetic activity, to the long arm of chromosome 9 [47]. Further studies identified *SGR* as a senescence-associated gene encoding a novel chloroplast protein and Mg^{2+} -dechetalase mediating Chl breakdown, which confirmed that *sgr* was a cosmetic stay-green locus [28,30].

A total of six QTLs (*csfl2/tcs2*, *tcs4*, *tcs5*, *csfl6*, *csfl9/tcs9*, and *csfl12*) were identified in recombinant inbred line populations derived from intra- and inter-subspecific crosses of rice varieties [56]. The functional stay-green *japonica* rice ‘SNU-SG1’ is mapped on chromosome 9. The presence of SNU-SG1 improved grain yield by maintaining photosynthetic activity in the flag and second leaves during grain filling, and by increasing sink strength due to high seed-setting rate [54]. Fine mapping performed by Lim et al. identified eleven main-effect loci in the *japonica* rice SNU-SG1 that were responsible for functional SG phenotypes [57]. Fifteen strong candidate genes were identified that explained these main-effect QTLs using the whole-genome sequence [57]. All these genes represent interesting candidates for further plant breeding. The *qCC2* major QTL for Chl content was identified using a population derived from an interspecific cross between *O. sativa* and *O. grandiglumi* [55]. The *GW2* gene that encodes an E3 ubiquitin ligase is located in the *qCC2* region and is responsible for leaf senescence through the transcriptional regulation of phytohormone signaling pathways [55]. Whether the *qCC2* QTL is explained by *GW2* allelic variations in *O. sativa* and *O. grandiglumi* remains to be determined. Ten stable QTL regions for six delayed senescence traits have been identified [58]. Expression analysis of five candidate genes displayed the differential transcript levels, suggesting their strong association with the senescence pattern in the flag and second leaves and possible contribution in enhancing grain yield through genomics-assisted breeding.

A genome-wide association study (GWAS) of a diverse worldwide collection of rice accessions identified forty-six significant association loci in natural variation of Chl content (CC) [59]. Among them, the *Grain number, plant height, and heading date7* (*Ghd7*) was a major locus for natural variations of Chl content that was involved in the repression of Chl and chloroplast biosynthetic genes [59]. As indicated by its name, this locus is a functional stay-green locus that controls also grain number, plant height, and heading date. Another candidate revealed by GWAS is *Narrow leaf1* (*NAL1*) [59]. A high degree of polymorphism in the 5′ UTR and four non-synonymous single nucleotide polymorphisms (SNPs) in the coding region of *NAL1* was shown to confer natural variations of Chl content [59]. Whether pleiotropic effects of *NAL1* in regulating flag leaf width and spikelet number per panicle might be helpful for rice breeding remains to be determined. In another GWAS study, 25 known genes for CC and stay-green (SG) traits were identified in natural rice variations [60]. Non-synonymous SNPs were detected in six of the genes and three SNPs in the promoter region of *OsSG1* [60]. Another GWAS revealed fourteen regions associated with CC and strong SG phenotypes [60]. From them, twenty-five candidate genes identified around the GWAS signals are genes with known important roles in CC and SG phenotypes. This suggests that these genes could be involved in the genetic variation of CC or SG.

Non-synonymous SNPs within six of these genes, and three SNPs in the promoter of *OsSG1*, were unveiled. The impact of each of these genes on plant performance remains to be explored.

Two representative rice subspecies, *japonica* and *indica*, display distinct morphological and physiological features as well as clear variation in genomic sequences [61]. Furthermore, *indica* displays early leaf senescence, whereas *japonica* displays late leaf senescence [62]. Then, QTL mapping was conducted to determine the genetic factors responsible for the differential senescence patterns between *indica* and *japonica* [30]. Results showed that allelic polymorphisms in the *OsSGR* promoter in *indica* result in higher and earlier induction of *OsSGR*, thereby triggering earlier senescence. The introgression of *japonica* *OsSGR* alleles into elite *indica* cultivars produced near isogenic lines (NILs) with delayed leaf senescence and extended photosynthetic competence, leading to improved grain filling rate and yield. Then, the *japonica* *OsSGR* alleles represent technical solutions for a beneficial breeding strategy in rice.

In addition to QTL and GWAS approaches, experiments leading to the characterization of mutants were carried out. The screen of ethyl methansulphonate (EMS)-induced rice mutants led to the identification of three stay-green mutants that also displayed drought tolerance [63]. Amongst them, *SGM-3* was characterized as a novel functional stay-green mutant with extended photosynthetic capacity during senescence, increased harvest index, and higher grain yield under irrigated as well as drought conditions which can be further used for the development of high-yielding lines [63]. The exact function of *SGM-3* remains to be determined.

In summary, we can see that both the quantitative genetics and mutagenesis approaches successfully identified several functional stay-green genes that play a role in plant productivity. Although the function of many of these genes and loci still remains to be elucidated, their positive alleles could be used in marker-assisted plant breeding programs.

4. Transcriptional Regulation of Leaf Senescence

Leaf senescence is an age-dependent process and occurs irrespective of environmental conditions. Its initiation is tightly regulated by endogenous factors including phytohormones and metabolic status. Nevertheless, exogenous factors such as light, drought, nutrient deficiency, and pathogen can modulate the onset and the intensity of the leaf senescence symptoms [64]. Nutrient and water deficiencies, as well as shading, are known to accelerate leaf senescence as the absence of nutrients promotes proteolysis and the recycling of plant resources. As such, dark stress and nitrogen-starvation have been used as senescence-inducing factors to screen leaf senescence mutants in *Arabidopsis*.

One informative way to understand leaf senescence and the different factors that control it is to identify the senescence-related transcription factors (TFs), which can act as nodes in gene expression pathways of the senescence-associated genes (*SAGs*) and modulate *SAG* expressions directly or indirectly by binding to the *cis*-elements in their promoter regions [1,65]. Phytohormones are endogenous factors involved in complex interactions that depend on the successive stages of leaf development. Ethylene, abscisic acid (ABA), jasmonic acid (JA), auxin, and salicylic acid (SA) promote senescence, while cytokinin and gibberellin delay senescence [66]. Thus, it is crucial to identify TFs acting downstream of the phytohormonal signaling networks and the interactions between TFs and phytohormones to understand the molecular mechanisms regulating senescence. Several of the TFs cited in this review and listed in Table 1 indeed integrate phytohormonal signals to modulate leaf senescence [67].

The NAC (NAM, ATAF1/2 and CUC2) gene family is one of the largest group of plant-specific TFs, with more than 150 genes in rice, and a large set of NAC factors are involved in leaf senescence [68]. These NAC proteins contain a conserved NAC domain at the N-terminal region for DNA-binding at the N-terminal region and variable transcription regulatory regions (TRRs) at the C-terminal region acting as a transcription activator or repressor [69]. In *Arabidopsis*, 57.5% of the NAC genes are differentially expressed during

leaf senescence, and they positively or negatively regulate leaf senescence [70]. In other than Arabidopsis, including rice, the senescence-associated NACs remain largely unknown. However, the few examples cited below show that several of the rice *OsNAC* factors control leaf senescence and respond or participate in hormone signaling, and especially in response to ABA. As such, the *OsNAP* gene, which is the homologous gene of the Arabidopsis *AtNAP*, is highly expressed in the senescing tissues and directly regulates SAGs in response to ABA [71–73]. The *OsNAP*-overexpressing transgenic rice show an early leaf senescence phenotype, whereas the RNA-interference (RNAi) lines display delayed leaf senescence with an increase in grain yield, indicating that *OsNAP* functions as a senescence-promoting TF [71].

ONAC2, which is homologous of *AtORE1*, promotes leaf senescence by inducing ABA biosynthetic genes, downregulating ABA catabolic genes, and regulating the Chl degradation genes [74]. Therefore, the overexpression of *ONAC2* causes early leaf senescence, while *ONAC2* RNAi lines exhibit delayed senescence phenotype along with a 10% increase in seed productivity. *ONAC096* positively regulates Chl degradation and the expression of several SAGs during leaf senescence and mediates ABA-induced leaf senescence [65]. *ONAC54* is also tightly associated with the regulation of ABA-induced leaf senescence; *ONAC054*-overexpressing plants display early leaf yellowing, whereas *onac054* knockout mutants maintain green leaves longer [64]. Intriguingly, *ONAC054* nuclear import is regulated by the cleavage of a C-terminal putative transmembrane domain through alternative 3' splicing, showing that *ONAC054* is important for ABA-induced leaf senescence and is itself controlled by multilayered regulatory processes. *OsNAC109* regulates the transcription of senescence- and hormone-associated genes by binding to NAC recognition sequence element (CNTCSSNNSCAVG) within promoter regions [75]. The knockout mutants of *ONAC109* show premature senescence with altered expressions of senescence and photosynthesis-related genes. The *ONAC106* that negatively regulates natural senescence, by directly modulating the expression of several SAGs such as *OsSGR*, *OsNYC1*, and also *OsNAC5*, was shown to respond not to ABA but to salt stress [76]. The overexpression of *ONAC106* delays leaf senescence under natural and dark-induced conditions, and its role in response to salt stress remains to be investigated. The *ONAC011* is one of the positive regulators of leaf senescence. Although the *ONAC011* regulatory pathway, including signals and target genes, remains unknown, it was shown that *ONAC011* accelerates heading time during the reproductive phase and then possibly modulates plant productivity and acts as a positive regulator of leaf senescence in rice [77].

The transcriptome analysis of field-grown flag leaves during senescence has revealed six other NAC TFs that are upregulated from vegetative to senescence stages (*OsNAC1*, LOC_Os02g36880; *ONAC39*, LOC_Os03g21030; *OsNAP*, LOC_Os03g21060; *ONAC010*, LOC_Os07g37920; *OsNAC18*, LOC_Os07g48450; *ONAC121*, LOC_Os10g421300) [78]. However, except for *OsNAP*, their functional roles in leaf senescence are still unknown [78].

The MYB TFs form the largest TF family in rice (at least 197 members) and function as key regulators of plant development in response to biotic and abiotic stress [79]. Similar to the senescence-associated NAC TFs, the MYB TFs are not only influenced by age but also by hormones and for some by abiotic stresses. The MYB-related rice TF RADIALIS-LIKE3 (*OsRL3*) promotes dark-induced leaf senescence and delays the response of rice plants to salt stress via the ABA signaling pathways [80]. Under dark-induced senescence conditions, the *osrl3* mutants exhibit a stay-green phenotype due to the reduced expressions of Chl degradation and SAGs. *OsMYB102* functions as a negative regulator of leaf senescence [81]. The overexpression of *OsMYB102* delays senescence under natural, dark, and ABA conditions, while the *osmyb102* knockout mutant shows accelerated senescence [81]. The transcript levels of several SAGs and ABA-related genes are altered in the *OsMYB102*-activation tagging lines, indicating that *OsMYB102* plays a critical role in leaf senescence by downregulating ABA accumulation and ABA signaling responses.

The plant-specific WRKY TFs are involved in many biological processes, and they particularly respond to and mediate stress responses [82]. According to microarray data

from rice flag leaves, we can see that among the 89 WRKY members identified so far, several are induced during leaf senescence [82,83]. The relationship to senescence was investigated in depth for a few of them. For example, *OsWRKY80* is upregulated under dark-induced senescence and drought stress, as well as by ABA treatment [84]. The overexpression of *OsWRKY23* in Arabidopsis was shown to enhance dark-induced leaf senescence [85]. *OsWRKY42* represses *OsMT1d*-mediated scavenging of reactive oxygen species (ROS), thereby promoting leaf senescence [86]. *OsWRKY5* is upregulated at the onset of leaf senescence, and its overexpressing line displays early leaf yellowing under aging and dark treatment, while the *oswrky5*-knockdown mutants show the opposite phenotype [87]. *OsWRKY5* acts as a positive regulator of leaf senescence in rice and regulates the expression of Chl degradation genes, SAGs, senescence-associated NAC genes (*OsNAP* and *OsNAC2*), and ABA biosynthetic genes.

Shading or dark treatment are known to enhance leaf senescence. The plant-specific phytochrome-interacting factors (PIFs) are basic helix–loop–helix (bHLH)-type TFs that regulate various biologic processes in a red-light and phytochrome (phy)-dependent manner, including leaf senescence and Chl biosynthesis [88,89]. *OsPIL1* is promoting Chl biosynthesis via trifurcate feed-forward regulatory loops that involve two GOLDEN2-LIKE (*OsGLK*) TFs [90]. The *ospil1* mutants exhibit earlier senescence during dark treatment, indicating that *OsPIL1* negatively regulates leaf senescence in rice [90]. Microarray analyses unveiled that several SAGs were upregulated in *ospil1* mutants, while the *OsGLKs* negative regulators of leaf senescence were strongly repressed [91].

OsTZF1 (also known as *OsDOS*; Delay of the Onset of Senescence) is a CCCH-tandem zinc finger protein that acts as a negative regulator of leaf senescence under various stress conditions as salt stress, drought, and dark [92]. The overexpression of *OsTZF1/OsDOS* delays leaf senescence, whereas RNAi knockdown causes accelerated age-dependent leaf senescence [93]. *OsTZF1* confers abiotic stress tolerance through the downregulation of several stress-related genes [92] and controls leaf senescence in a JA-dependent manner [93].

In addition to the identification of the different TFs involved in leaf senescence, one of the main challenges is to understand their interconnections and networks. If network analyses are well advanced in Arabidopsis, we still are at the beginnings in rice. However, a few studies performed in rice reveal such interconnections. As such, the ETHYLENE RESPONSE FACTOR 101 (*OsERF101*) that positively regulates leaf senescence was shown to enhance the expression of *OsNAP* and the JA-responsive *OsMYC2* TF [94]. The overexpression of *OsMYC2* promoted leaf senescence and reduced Chl content under darkness, indicating that *OsMYC2* functions as a positive regulator of leaf senescence [95]. The *oserf101* knockout mutant accordingly exhibits delayed leaf senescence with higher Chl content during dark-induced and natural senescence [92]. After JA treatment, the leaves of *oserf101* still contain more Chl than WT, indicating that *OsERF101* is involved in promoting JA-induced leaf senescence. The expression of the JA signaling genes is indeed downregulated in *oserf101*.

Although not exhaustive, the several examples presented above show that the different TFs identified so far in rice can regulate positively or negatively the onset of leaf senescence, in response to one or several endogenous and/or exogenous signals. Many of these TFs are implicated in the response to ABA, JA, ethylene, SA, and dark. While their interconnections and networks remain poorly known and deserve further studies, it is already noticeable that several of the TFs listed in Table 1 have some senescence-related TFs as direct downstream target genes. Strikingly, we can see in Table 1 that *OsNAP* is a downstream target of many of them [71–73]. Dissecting the leaf senescence regulatory network will permit the identification of the master regulators and then facilitate the choice of candidate genes for further investigations. Nevertheless, a better characterization of the phenotypes of the senescence-TFs overexpressors and knock-down lines, based on the evaluation of plant performances, flowering dates, heading time, yield, and stress resistance, should point to the best candidates for plant breeding strategies.

Table 1. Summary of transcription factors associated with rice leaf senescence.

Gene	Locus Number	Gene Family	Role in Senescence	Pathways Involved	Direct Downstream Targets	References
<i>OsNAP</i>	LOC_Os03g21060	NAC TF	Positive	Age, ABA, dark, salt, drought	OsSGR, OsNYC1, OsNYC3, OsRCCR1, Osh36, OsI57, Osh69, OsI85	[71–73]
<i>OsNAC2</i>	LOC_Os04g38720	NAC TF	Positive	Age, ABA, dark	OsSGR, OsNYC3, OsNCED3, OsZEP1	[74]
<i>ONAC106</i>	LOC_Os08g33670	NAC TF	Negative	Age, dark, salt	OsSGR, OsNYC1, OsNAC5, OsNAP, OsEIN3, OsS3H, OsDREB2A, OsLEA3, OsbZIP23, LPA1	[76]
<i>ONAC011 (OsY37)</i>	LOC_Os06g46270	NAC TF	Positive	Age		[77]
<i>ONAC096</i>	LOC_Os07g04560	NAC TF	Positive	Age, ABA	OsSGR, OsPAO, OsNYC3, OsRCCR1, OsI85, OsI2, OsI57, OsNAP, ABI5, OsEEL	[65]
<i>ONAC054</i>	LOC_Os03g02800	NAC TF	positive	Age, ABA, JA	OsNYC1, OsABI5,	[64]
<i>OsNAC109 (OsYL3)</i>	LOC_Os09g38000	NAC TF	Negative	Age, ABA, dark	OsNAP, OsNYC3, OsEATB, OsAMTR1, OsZFP185, OsMPS, OsGA2ox3	[75]
<i>OsRL3</i>	LOC_Os02g47744	MYB TF	Positive	ABA, salt, dark	OsSGR, OsNYC1, OsRCCR1, OsI2, OsI43, OsSAG12-2, OsRK1, OsRAB16C, OsRAB16D	[80]
<i>OsMYB102</i>	LOC_Os06g43090	MYB TF	Negative	Age, ABA, dark	OsSGR, OsNYC1, OsABF4, OsNAP, OsCYP707A6	[81]
<i>OsWRKY80</i>	LOC_Os09g30400	WRKY TF	Positive	Dark, ABA, drought,		[84]
<i>OsWRKY23</i>	LOC_Os01g53260	WRKY TF	Positive	Age, dark, SA, pathogen		[85]
<i>OsWRKY42</i>	LOC_Os02g26430	WRKY TF	Positive	Age, ROS	OsMT1d	[86]
<i>OsWRKY5</i>	LOC_Os05g04640	WRKY TF	Positive	Age, dark, ABA	CCEs, SAGs, OsNAP, OsNAC2, OsNCED3, OsNCED4, OsNCED5	[87]
<i>OsPIL1</i>	LOC_Os03g56950	bHLH TF	Negative	Age, dark	OsPORB, OsCAO1, OsGLK1, OsGLK2	[90,91]
<i>OsMYC2</i>	LOC_Os10g42430	MYC TF	Positive	JA, dark	Several SAGs.	[95]
<i>OsERF101</i>	LOC_Os04g32620	ERF TF	Positive	Age, ABA, drought, JA	OsNAP, OsMYC2, OsJAI1, OsCOI1a, OsSGR, OsNYC1, OsNYC3	[94]
<i>OsTZF1</i>	LOC_Os01g09620	CCCH-type zinc finger TF	Negative	Age, ABA, JA, SA, salt, drought, dark	Stress-responsive genes	[92]
<i>OsTZF2 (OsDOS)</i>	LOC_Os05g10670	CCCH-type zinc finger TF	Negative	Age, JA		[93]

5. Leaf Senescence and Nitrogen Metabolism

Nitrogen (N) is essential for all living organisms and is predominantly remobilized from source to sink organs during leaf senescence [96]. During the vegetative growth stage, chloroplasts contain up to 75–80% of the leaf N, primarily as photosynthetic proteins including thylakoid proteins, light harvesting complexes, photosystems, Calvin cycle proteins, and Rubisco [2,97]. In rice, 70–90% of total panicle N is originating from the remobilization of N from vegetative organs [9]. Chl degradation during leaf senescence enables or signals further catabolic processes in chloroplasts [18]. The subsequent dismantling of chloroplasts is a major N source for recycling and remobilization [18]. Rubisco and other photosynthetic proteins are degraded during senescence, and the released N is remobilized step by step from source leaves to sink leaves, and then ultimately to grains [97]. The remobilization of

N from aging tissues to the seeds is an important determinant of productivity and yield, especially under N limitation [98].

Autophagy (meaning self-eating) facilitates the degradation of unwanted cytoplasmic components inside the lytic vacuoles [99]. Two types of autophagy have been described in plant leaves so far. The macro-autophagy is a vesicular process that involves eighteen ATG proteins in the formation of a double membrane vacuole named the autophagosome. Autophagosomes are formed in the cytosol where they engulf unwanted cytoplasmic materials (called cargoes) and drive them to the lytic vacuole. By fusing their external membranes to the tonoplast, autophagosomes release their inner membranes sequestering cargoes inside the vacuole for degradation [99]. Micro-autophagy consists of the direct engulfment of cytoplasmic material inside the vacuole through tonoplast invagination. Macro-autophagy is involved in the degradation of the stroma proteins released from the chloroplast stromules in budding structures named Rubisco-containing bodies (RBC). The trafficking of RBC to the lytic vacuole is dependent on the macro-autophagy machinery [100]. Entire damaged chloroplasts can also be degraded in the vacuole through a micro-autophagy process [101]. In Arabidopsis, it was shown that macro-autophagy is essential for the remobilization of several nutrients such as N, sulfur, and iron [102–104]. The ATG8 protein is essential for autophagosome membrane formation and for cargo sequestration. The ATG8 proteins interact with the cargoes directly, or indirectly through receptors, and facilitate their sequestration in the autophagosomes. The constitutive overexpressing of *ATG8* isoforms in Arabidopsis has been shown to stimulate the autophagic activity and to increase N remobilization from the rosette leaves to the seeds [105].

In rice, thirty-three Autophagy-related genes (*OsATGs*) have been identified [106]. *OsATG7* is essential for autophagosome formation and participates in efficient N utilization and remobilization [97]. The *osatg7-1* mutant display poor biomass and NUE during vegetative growth. Unable to export N from senescing leaves, this mutant is sterile. As the transcript levels of *OsATG8a*, *OsATG8b*, and *OsATG8c* are increased under N starvation, their overexpression was assayed in rice [107–110]. Irrespective of the *OsATG8* isoforms, rice transformants display higher NUE and yield [107–110]. The *OsATG8a*-overexpressing lines accumulate more N in their grains and less in their dry remains compared to WT, which indicates a better N remobilization to the seeds that increase grain yield [107]. The overexpression of *OsATG8b* could improve NUE and increased grain yield by stimulating autophagy flux and by enhancing the activities of enzymes related to N metabolism [110]. The higher N remobilization efficiency of *OsATG8b*-overexpressors was demonstrated using ¹⁵N labeling and tracing assays. On the contrary, the *osatg8b* knockout mutants showed the opposite phenotypes [110]. The specific roles of the different ATG8 isoforms in Arabidopsis and rice remain to be explored, as well as their respective involvement in nutrient recycling. The contribution and the nature of the plant proteases involved in the last steps of the autophagy process are still largely unknown [111]. Nevertheless, autophagy and senescence-related proteases are essential for the remobilization of N, as they facilitate the release of a pool of free amino acids from the protein cargoes that can be exported through the phloem to the sinks and up to the seeds.

Glutamine (Gln) and asparagine (Asn), which contain two N atoms per molecule, are considered the main amino acids involved in N translocation in the phloem sap [112]. From the bulk of amino acids originating from autophagic protein degradation, it is likely that amino-acid interconversions and transaminations occur to facilitate phloem loading [112]. Especially, the synthesis of Gln and Asn in senescing organs appears essential for N remobilization [113,114]. The increase in the Gln/glutamate (Glu) and Asn/aspartate (Asp) ratios with leaf senescence is in line with this assumption [2]. Therefore, attention has been paid for a long time to the role of the glutamine synthetases (GS) and asparagine synthetases (ASN) for the re-assimilation of ammonium in old leaves [115]. The lack of *OsGS1;1*, which is the major cytosolic isoform of GS in rice, caused a shortage of N resources in senescing source organs and limited N remobilization into sink organs, resulting in severe growth retardation and lower grain yield [115]. Glutamate synthetase (GOGAT,

also known as glutamine-2-oxoglutarate aminotransferase) catalyzes the transfer of the amide group of Gln (formed by GS) to 2-oxoglutarate (2-OG) to synthesize two molecules of Glu [116]. The GS/GOGAT cycle is the major route for ammonium assimilation and sustains N metabolism in higher plants [116]. It was recently shown that when the expression of NADH-dependent *OsGOGAT1* is enhanced, the remobilization of N from young leaves to old leaves is increased, which causes early leaf senescence but increases N remobilization, leading to higher N content in grains [96]. Interestingly, the mutation in the Ferredoxin-dependent *OsFd-GOGAT* also leads to premature leaf senescence and facilitates N remobilization [117,118]. This shows that the NADH- and Fd-dependent GOGATs play opposite roles in rice, the former being involved in N recycling and remobilization, the latter being involved in N primary assimilation and photorespiration. Asn synthesis is catalyzed by Asparagine synthetase (ASN) through the transfer of the Glu-amide group to the amide position of Asp. Asn has an essential role for N storage and transport in plants [119]. Among the two rice *ASN* genes, *OsASN1* is essential for plant growth under both N-sufficient and N-limiting conditions, and for biomass and grain yield [120]. During the grain-filling stage, the *osasn1* mutants accumulate more N in their flag leaves and less N in panicles than the wild type, indicating reduced N remobilization from source to sink organs.

One important step for N-remobilization during leaf senescence is phloem loading and amino acid transport [112]. Many senescence-associated amino acid transporters have been identified in Arabidopsis and rice, but their specific roles remain largely unknown, even in Arabidopsis [11,121]. It was shown in Arabidopsis that the AAP8 (Amino Acid Permease 8) plays a key role in amino acid phloem loading and its function strongly affects sink size and number [122]. Interestingly, a recent paper shows that increasing the expression of the rice *OsAAP3* that facilitates the transport of arginine and lysine into the mesophyll cells influences leaf senescence [123]. The overexpression of *OsAAP3* decreases grain yield and leads to lesion mimic and leaf senescence in rice flag leaves by an increase of nitric oxide. In a previous study, the authors have shown that blocking *OsAAP3* promotes rice tillering, grain yield, and tiller bud elongation [124]. Together, these studies show that enhancing or inhibiting amino acid influxes or effluxes in leaves impacts senescence program, leaf longevity, and plant productivity. The rice genome harbors nineteen *AAP* genes [125], and previous reports had shown the amino acid transporter activities of *OsAAP1*, *OsAAP4*, *OsAAP5*, and *OsAAP6*, which play important roles for grain quality and yield [126–129]. Using publically available expression data, we analyzed the transcript levels of the rice *AAP* genes (Figure 2). The expressions of *OsAAP4*, *OsAAP5*, *OsAAP6*, *OsAAP7*, and *OsAAP8* are upregulated during the grain-filling stage in rice, suggesting their possible role for N-remobilization during senescence. Then, their function during leaf senescence may deserve attention.




Figure 2. Heat map of the expression of the rice *AAPs* (amino acid permease, *OsAAPs*) during leaf senescence. Expression data were from RNA-seq analysis (a) [20] and microarray data (b) [21]. Gene nomenclature of each gene was followed in the previous report [120]. Relative expression levels were quantified as describes in Figure 1.

6. Stress-Induced Senescence

As plants are sessile, they are frequently exposed to adverse environmental conditions and are unavoidably challenged by various biotic and abiotic stresses during the growing season. Abiotic stresses such as drought, floods, salinity, radiation, darkness/shading, and temperature changes, nutrient deficiency, and mineral toxicity adversely affect plant growth and thereby cause the loss of productivity [5]. Attacks by diverse pathogens such as fungi, bacteria, nematodes, oomycetes, and insects are included in biotic stress [130]. Therefore, narrowing the yield gap between optimal growth and adverse environmental conditions throughout the globe is an urgent challenge to ensure food security and safety in the coming years [131].

Under harsh and unfavorable conditions, plants are acclimated using various ways, including conferring tolerance to environmental stresses or accelerating the senescence (stress-induced senescence) to ensure their survival [132]. Although levels of gene regulation and transcriptome changes by developmental and stress-induced senescence are not entirely overlapping, age-dependent and stress-induced senescence share many signaling pathways via TFs, the involvement of phytohormones such as ABA, JA, and SA, which also synthesize during stress conditions, and physiological, biochemical, and molecular mechanisms [5,94,132]. For example, the expression of *OsNAP* was significantly induced by ABA and abiotic stress, and hence, the overexpression of *OsNAP* results in improved tolerance to abiotic stresses such as high salinity, drought, and low temperature, indicating that *OsNAP* function as a transcriptional activator in mediating abiotic stress responses as well as leaf senescence [71,73]. *OsNAC2*, which functions as a positive regulator of senescence, directly activates the expression of Chl degradation genes and *SAGs*, and hence, its overexpression causes accelerated leaf senescence [74]. The knockdown of *OsNAC2* leads to increased tolerance to drought and salinity stresses by the upregulation of stress-related and abscisic acid (ABA)-signaling genes [133]. The expression of *OsNAC066* is significantly enhanced by multiple abiotic factors, and its overexpressing transgenic lines improve tolerance to drought and oxidative stress, indicating a positive regulator of them [134]. However, its role for senescence remains still unknown. It has been shown that *OsWRKY93* was involved in both flag leaf senescence and response to biotic stress [135]. *OsWRKY93*-overexpressing plants show increased resistance to blast disease by the enhancement of ROS production and PAMP-triggered immune response, indicating that *OsWRKY93* can be a favorable candidate for the breeding of high-yield and disease-resistant rice [135].

Global climate change that might affect crop production drastically is regarded as one of the future challenges, and it affects agriculture in different abiotic and biotic ways such as increasing the average temperature, heat stress, variations in annual rainfall, pests, or microbes, global change of atmospheric CO₂ or ozone level, and fluctuations in sea level [136]. Therefore, the development of climate-resilient crop species that will have enhanced and sustained productivity traits is becoming important for ensuring global food security [137]. For instance, high temperature during the grain-filling stage can cause a nearly 50% reduction in rice yield [138], and heat-tolerance studies in rice have mainly focused on the reproductive stage due to its high sensitivity and critical influence on grain yield [139]. Plants respond to high temperature through the activation of the heat shock TFs (HSFs) or other stress-related genes [140,141]. Heat stress-responsive two NAC TFs, *ONAC127*, and *ONAC129* are specifically expressed in the pericarp of rice seeds [141]. They are involved in the apoplasmic transport of photosynthates for starch accumulation during grain filling as a heterodimer.

Due to climate warming, plants are allowed to grow for a longer time during each growing season. Rice especially in Southeast Asia may increase the growth potential grown in the open paddy fields by absorbing more CO₂ for photosynthesis and hence increase the grain yield, but plant growth is also restricted by shorter day length in autumn, regardless of warming [142,143]. Thus, it is important to analyze how to adapt the plant according to the variable growing season length by adjusting the length of vegetative and reproductive growth, senescence duration, and timing of heading [5]. For example,

adoption of the functional stay-green trait may be one of the ways to deal with future warming conditions by extending the time for photosynthesis and further minimizing the sink limitation, leading to higher yield.

7. Breeding Strategies to Improve Rice Yield

The previous sections present different approaches to identify senescence genes that could be candidates for plant breeding strategies. Amongst these genes, we can distinguish the TFs that regulate positively or negatively the onset of the leaf senescence, and the executioners that work to the preservation of cell longevity or in the successive degradation events that allow the recycling of nutrients. Modulating both kinds of genes could influence leaf longevity and therefore impact plant productivity.

The first attempt to identify regulators or executioners in Arabidopsis has been done by analyzing the transcriptomic data of senescing leaves [144,145]. This facilitated the identification of senescence repressed or senescence-associated genes that were either regulators or executioners. In rice, senescence-associated genes were also identified [13,16], and studies of their roles were performed using reverse genetics. Studies of mutants or overexpressors mainly focused on executioners or TFs that had also been identified in other plant species; for example, the senescence-associated N enzymes and the autophagy proteins are known to participate in N remobilization, and the homologous of *OsNAP* and *OsNAC2* control leaf senescence.

The identification of new regulators/actors involved in leaf senescence was carried out using quantitative genetics (either QTL or GWAS). Quantitative traits considered were chlorophyll content, leaf senescence onset, and agronomic performances. Although such a forward genetic approach provided many cosmetic stay green candidates, it also revealed functional stay green candidates that might serve in traditional marker-assisted breeding programs, such as for example in the case of the introgression of the *japonica* *OsSGR* alleles into the elite *indica* cultivars that resulted in a higher grain-filling rate and better yield [30]. Nevertheless, the characterization of many of the QTLs and underlying genes still necessitates efforts. Indeed, the use of GWAS and QTL results for plant engineering requires that the genes underlying the QTLs are identified and confirmed. In addition, as leaf senescence and plant yield are multigenic traits, gene stacking might be considered.

8. Conclusions

Senescence and stay-green phenotypes are important traits in rice breeding programs. During the last few years, there have been significant advances in understanding the mechanisms and processes underlying rice leaf senescence. These include the dissection of the Chl degradation pathways, the characterization of stay-green traits, the investigation of the transcriptional regulations and interaction between TFs and phytohormones, and the dissection of impacts on nutrient remobilization and rice yield production. In rice, as in many other monocarpic cereals, grain production is concomitant with plant senescence, and the seed maturation ends with plant death. Therefore, the timing of the senescence onset, the duration of the senescence period, and the intensity of the senescence-related cell degradation are important traits that control the number of offspring in rice, and then grain quality and quantity. Up to now, studies have mainly considered leaf senescence onset. New studies on the duration of leaf senescence may identify additional important genes involved in leaf senescence and plant productivity. Dissecting the genetic bases of the different steps of leaf senescence in rice is of great importance, as it can provide genetic technical solutions for the improvement of rice performances.

The developments of high-throughput transcriptome analyses on mutants and over-expressors of senescence-related TFs should generate insights into the functional and regulatory aspects of leaf senescence in rice. The proteomic and metabolic approaches, which are still less developed, should, in the combination of multi-omics analyses, help to unveil better characterization of the different steps of the leaf senescence program. Functional stay-green traits should be dissected in depth to unveil the mechanisms that

cause the extension of photosynthetic capacity. Co-operation of the recent development of non-invasive phenotyping approaches and CRISPR/Cas9-based genome editing can be used to manipulate candidate genes and identify the new regulatory genes and design networks more easily. Phenomic approaches will also facilitate the development of forward genetics approaches to dissect leaf senescence regulation and identify novel genes.

Author Contributions: S.L. and C.M.-D. designed and wrote the article. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Institute for Basic Science (IBS-R013-D1) to S.L.; The IJPB benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007) to M.C.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors dedicate this review to the memory of Stefan Hörtensteiner whose pioneer work paved the way for all the studies dealing with chlorophyll degradation process during leaf senescence.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Woo, H.R.; Kim, H.J.; Lim, P.O.; Nam, H.G. Leaf Senescence: Systems and dynamics aspects. *Annu. Rev. Plant Biol.* **2019**, *70*, 347–376. [\[CrossRef\]](#)
2. Avila-Ospina, L.; Moison, M.; Yoshimoto, K.; Masclaux-Daubresse, C. Autophagy, plant senescence, and nutrient recycling. *J. Exp. Bot.* **2014**, *65*, 3799–3811. [\[CrossRef\]](#)
3. Gan, S. Concepts and types of senescence in plants. *Methods Mol. Biol.* **2018**, *1744*, 3–8. [\[PubMed\]](#)
4. Gan, S.; Hörtensteiner, S. Frontiers in plant senescence research: From bench to bank. *Plant Mol. Biol.* **2013**, *82*, 503–504. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Gregersen, P.L.; Culetic, A.; Boschian, L.; Krupinska, K. Plant senescence and crop productivity. *Plant Mol. Biol.* **2013**, *82*, 603–622. [\[CrossRef\]](#)
6. Yadava, P.; Singh, A.; Kumar, K.; Singh, I. Plant senescence and agriculture. In *Senescence Signalling and Control in Plants*; Academic Press: Cambridge, MA, USA, 2019; pp. 283–302.
7. Wing, R.A.; Purugganan, M.D.; Zhang, Q. The rice genome revolution: From an ancient grain to Green Super Rice. *Nat. Rev. Genet.* **2018**, *19*, 505–517. [\[CrossRef\]](#)
8. Evans, J.R. Improving photosynthesis. *Plant Physiol.* **2013**, *162*, 1780–1793. [\[CrossRef\]](#)
9. Mae, T. Physiological nitrogen efficiency in rice: Nitrogen utilization, photosynthesis, and yield potential. *Plant Soil* **1997**, *196*, 201–210. [\[CrossRef\]](#)
10. Masclaux-Daubresse, C.; Reisdorf-Cren, M.; Orsel, M. Leaf nitrogen remobilisation for plant development and grain filling. *Plant Biol.* **2008**, *10*, 23–36. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Havé, M.; Marmagne, A.; Chardon, F.; Masclaux-Daubresse, C. Nitrogen remobilization during leaf senescence: Lessons from Arabidopsis to crops. *J. Exp. Bot.* **2017**, *68*, 2513–2529.
12. Uauy, C.; Distelfeld, A.; Fahima, T.; Blechl, A.; Dubcovsky, J.A. A NAC Gene Regulating Senescence Improves Grain Protein, Zinc, and Iron Content in Wheat. *Science* **2006**, *314*, 1298–1301. [\[CrossRef\]](#)
13. Lee, R.; Wang, C.; Huang, L.; Chen, S.G. Leaf senescence in rice plants: Cloning and characterization of senescence up-regulated genes. *J. Exp. Bot.* **2001**, *52*, 1117–1121. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Liu, L.; Zhou, Y.; Zhou, G.; Ye, R.; Zhao, L.; Li, X.; Lin, Y. Identification of early senescence-associated genes in rice flag leaves. *Plant Mol. Biol.* **2008**, *67*, 37–55. [\[CrossRef\]](#)
15. Woo, H.R.; Kim, H.J.; Nam, H.G.; Lim, P.O. Plant leaf senescence and death—regulation by multiple layers of control and implications for aging in general. *J. Cell Sci.* **2013**, *126*, 4823–4833. [\[CrossRef\]](#)
16. Leng, Y.; Ye, G.; Zeng, D. Genetic Dissection of Leaf Senescence in Rice. *Int. J. Mol. Sci.* **2017**, *18*, 2686. [\[CrossRef\]](#)
17. Kuai, B.; Chen, J.; Hörtensteiner, S. The biochemistry and molecular biology of chlorophyll breakdown. *J. Exp. Bot.* **2018**, *69*, 751–767. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Tamary, E.; Nevo, R.; Naveh, L.; Levin-Zaidman, S.; Kiss, V.; Savidor, A.; Levin, Y.; Eyal, Y.; Reich, Z.; Adam, Z. Chlorophyll catabolism precedes changes in chloroplast structure and proteome during leaf senescence. *Plant Direct.* **2019**, *3*, e00127. [\[CrossRef\]](#)
19. Hörtensteiner, S. Update on the biochemistry of chlorophyll breakdown. *Plant Mol. Biol.* **2013**, *82*, 505–517. [\[CrossRef\]](#) [\[PubMed\]](#)

20. Lee, S.; Jeong, H.; Lee, S.; Lee, J.; Kim, S.J.; Park, J.W.; Woo, H.R.; Lim, P.O.; An, G.; Nam, H.G.; et al. Molecular bases for differential aging programs between flag and second leaves during grain-filling in rice. *Sci. Rep.* **2017**, *7*, 8792. [[CrossRef](#)] [[PubMed](#)]
21. Sato, Y.; Takehisa, H.; Kamatsuki, K.; Minami, H.; Namiki, N.; Ikawa, H.; Ohyanagi, H.; Sugimoto, K.; Antonio, B.A.; Nagamura, Y. RiceXPro version 3.0: Expanding the informatics resource for rice transcriptome. *Nucleic Acids Res.* **2013**, *41*, D1206–D1213. [[CrossRef](#)]
22. Kusaba, M.; Ito, H.; Morita, R.; Lida, S.; Sato, Y.; Fujimoto, M.; Kawasaki, S.; Tanaka, R.; Hirochik, H.; Nishimura, M.; et al. Rice Non-Yellow Coloring1 Is Involved in Light-Harvesting Complex II and Grana Degradation during Leaf Senescence. *Plant Cell* **2007**, *19*, 1362–1375. [[CrossRef](#)] [[PubMed](#)]
23. Sato, Y.; Morita, R.; Katsuma, S.; Nishimura, M.; Tanaka, A.; Kusaba, M. Two short-chain dehydrogenase/reductases, Non-Yellowcoloring1 and NYC1-LIKE, are required for chlorophyll *b* and light-harvesting complex II degradation during senescence in rice. *Plant J.* **2009**, *57*, 120–131. [[CrossRef](#)]
24. Piao, W.; Han, S.H.; Sakuraba, Y.; Paek, N.-C. Rice 7-Hydroxymethyl Chlorophyll *a* Reductase Is Involved in the Promotion of Chlorophyll Degradation and Modulates Cell Death Signaling. *Mol. Cells* **2017**, *40*, 773–786. [[PubMed](#)]
25. Shimoda, Y.; Ito, H.; Tanaka, A. Arabidopsis *STAY-GREEN*, Mendel's Green Cotyledon Gene, Encodes Magnesium-Dechelataase. *Plant Cell* **2016**, *28*, 2147–2160. [[CrossRef](#)] [[PubMed](#)]
26. Sakuraba, Y.; Schelbert, S.; Park, S.Y.; Han, S.H.; Lee, B.D.; Andr  s, C.B.; Kessler, F.; H  rtensteiner, S.; Paek, N.-C. *STAY-GREEN* and Chlorophyll Catabolic Enzymes Interact at Light-Harvesting Complex II for Chlorophyll Detoxification during Leaf Senescence in *Arabidopsis*. *Plant Cell* **2012**, *24*, 507–518. [[CrossRef](#)]
27. Sakuraba, Y.; Park, S.Y.; Paek, N.-C. The Divergent Roles of *STAYGREEN* (*SGR*) Homologs in Chlorophyll Degradation. *Mol. Cells* **2015**, *38*, 390–395. [[CrossRef](#)]
28. Park, S.Y.; Yu, J.W.; Park, J.S.; Li, J.; Yoo, S.C.; Lee, N.Y.; Lee, S.K.; Jeong, S.W.; Seo, H.S.; Koh, H.J.; et al. The Senescence-Induced Staygreen Protein Regulates Chlorophyll Degradation. *Plant Cell* **2007**, *19*, 1649–1664. [[CrossRef](#)]
29. Jiang, H.; Li, M.; Liang, N.; Yan, H.; Wei, Y.; Xu, X.; Liu, J.; Xu, Z.; Chen, F.; Wu, G. Molecular cloning and function analysis of the *stay green* gene in rice. *Plant J.* **2007**, *52*, 197–209. [[CrossRef](#)]
30. Shin, D.; Lee, S.; Kim, T.H.; Lee, J.H.; Park, J.; Lee, J.; Lee, J.Y.; Cho, L.H.; Choi, J.Y.; Lee, W.; et al. Natural variations at the *Stay-Green* gene promoter control lifespan and yield in rice cultivars. *Nat. Commun.* **2020**, *11*, 2819. [[CrossRef](#)]
31. Rong, H.; Tang, Y.; Zhang, H.; Wu, P.; Chen, Y.; Li, M.; Wu, G.; Jiang, H. The *Stay-Green Rice like* (*SGRL*) gene regulates chlorophyll degradation in rice. *J. Plant Physiol.* **2013**, *170*, 1367–1373. [[CrossRef](#)]
32. Schelbert, S.; Aubry, S.; Burla, B.; Agne, B.; Kessler, F.; Krupinska, K.; H  rtensteiner, S. Pheophytin Pheophorbide Hydrolase (Pheophytinase) Is Involved in Chlorophyll Breakdown during Leaf Senescence in *Arabidopsis*. *Plant Cell* **2009**, *21*, 767–785. [[CrossRef](#)]
33. Morita, R.; Sato, Y.; Masuda, Y.; Nishimura, M.; Kusaba, M. Defect in non-yellow coloring 3, an α/β hydrolase-fold family protein, causes a stay-green phenotype during leaf senescence in rice. *Plant J.* **2009**, *59*, 940–952. [[CrossRef](#)]
34. Das, A.; Christ, B.; H  rtensteiner, S. Characterization of the pheophorbide *a* oxygenase/phyllobilin pathway of chlorophyll breakdown in grasses. *Planta* **2018**, *248*, 875–892. [[CrossRef](#)]
35. Tang, Y.; Li, M.; Chen, Y.; Wu, P.; Wu, G.; Jiang, H. Knockdown of *OsPAO* and *OsRCCR1* cause different plant death phenotypes in rice. *J. Plant Physiol.* **2011**, *168*, 1952–1959. [[CrossRef](#)]
36. Zhang, Z.; He, Y.; Li, L.; Zhang, X.; Xu, X.; Shi, Y.; Wu, J.L. Characterization of a novel allele encoding pheophorbide *a* oxygenase in rice. *Plant Signal Behav.* **2021**, *16*, 1864606. [[CrossRef](#)] [[PubMed](#)]
37. Balazadeh, S. *Stay-Green* Not Always Stays Green. *Mol. Plant.* **2014**, *7*, 1264–1266. [[CrossRef](#)] [[PubMed](#)]
38. H  rtensteiner, S. Chlorophyll degradation during senescence. *Annu. Rev. Plant Biol.* **2006**, *57*, 55–77. [[CrossRef](#)] [[PubMed](#)]
39. Thomas, H.; Howarth, C.J. Five ways to stay green. *J. Exp. Bot.* **2000**, *51*, 329–337. [[CrossRef](#)]
40. Thomas, H.; Ougham, H. The stay-green trait. *J. Exp. Bot.* **2014**, *65*, 3889–3900. [[CrossRef](#)] [[PubMed](#)]
41. Borrell, A.K.; Hammer, G.L.; Oosterom, E.V. Stay-green: A consequence of the balance between supply and demand for nitrogen during grain filling. *Ann. Appl. Biol.* **2001**, *138*, 91–95. [[CrossRef](#)]
42. Morales, F.; Anc  n, M.; Fakhret, D.; Gonz  lez-Torralba, J.; G  mez, A.L.; Seminario, A.; Soba, D.; Ben, M.S.; Garriga, M.; Aranjuelo, I. photosynthetic metabolism under stressful growth conditions as a bases for crop breeding and yield improvement. *Plants* **2020**, *9*, 88. [[CrossRef](#)]
43. Kusaba, M.; Tanaka, A.; Tanaka, R. Stay-green plants: What do they tell us about the molecular mechanism of leaf senescence. *Photosynth. Res.* **2013**, *117*, 221–234. [[CrossRef](#)]
44. Kamal, N.M.; Alnor Gorafi, Y.S.; Abdelrahman, M.; Abdellatef, E.; Tsujimoto, H. Stay-Green trait: A prospective approach for Yield Potential, and Drought and Heat Stress Adaptation in Globally Important Cereals. *Int. J. Mol. Sci.* **2019**, *20*, 5837. [[CrossRef](#)]
45. Tian, F.; Gong, J.; Zhang, J.; Zhang, M.; Wang, G.; Li, A.; Wang, W. Enhanced stability of thylakoid membrane proteins and antioxidant competence contribute to drought stress resistance in the *tasg1* wheat stay-green mutant. *J. Exp. Bot.* **2013**, *64*, 1509–1520. [[CrossRef](#)] [[PubMed](#)]
46. Barry, C.S.; McQuinn, R.P.; Chung, M.Y.; Besuden, A.; Giovannoni, J.J. Amino Acid Substitutions in Homologs of the *STAY-GREEN* Protein Are Responsible for the *green-flesh* and chlorophyll *retainer* Mutations of Tomato and Pepper. *Plant Physiol.* **2008**, *147*, 179–187. [[CrossRef](#)] [[PubMed](#)]

47. Cha, K.W.; Lee, Y.J.; Koh, H.J.; Lee, B.M.; Nam, Y.W.; Paek, N.-C. Isolation, characterization, and mapping of the stay green mutant in rice. *Theor. Appl. Genet.* **2002**, *104*, 526–532. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Duvick, D.; Smith, J.; Cooper, M. Long-term selection in a commercial hybrid maize breeding program. *Plant Breed. Rev.* **2004**, *24*, 109–152.
49. Fang, C.; Li, C.; Li, W.; Wang, Z.; Zhou, Z.; Shen, Y.; Wu, M.; Wu, Y.; Li, G.; Kong, L.A.; et al. Concerted evolution of *D1* and *D2* to regulate chlorophyll degradation in soybean. *Plant J.* **2014**, *77*, 700–712. [\[CrossRef\]](#)
50. Grbić, V.; Bleecker, A.B. Ethylene regulates the timing of leaf senescence in *Arabidopsis*. *Plant J.* **1995**, *8*, 595–602. [\[CrossRef\]](#)
51. Rampino, P.; Spano, G.; Pataleo, S.; Mita, G.; Napier, J.A.; di Fonzo, N.; Shewry, P.R.; Perrotta, C. Molecular analysis of a durum wheat ‘stay green’ mutant: Expression pattern of photosynthesis-related genes. *J. Cereal Sci.* **2006**, *43*, 160–168. [\[CrossRef\]](#)
52. Gous, P.W.; Warren, F.; Gilbert, R.; Fox, G.P. Drought-proofing barley (*Hordeum vulgare*): The effects of stay green on starch and amylose structure. *Cereal Chem.* **2017**, *94*, 873–880. [\[CrossRef\]](#)
53. Jiang, G.H.; He, Y.Q.; Xu, C.G.; Li, X.H.; Zhang, Q. The genetic basis of stay-green in rice analyzed in a population of doubled haploid lines derived from an *indica* by *japonica* cross. *Theor. Appl. Genet.* **2004**, *108*, 688–698. [\[CrossRef\]](#)
54. Yoo, S.-C.; Cho, S.-H.; Zhang, H.; Paik, H.-C.; Lee, C.-H.; Li, J.; Yoo, J.-H.; Koh, H.-J.; Seo, H.S.; Paek, N.-C. Quantitative trait loci associated with functional stay-green SNU-SG1 in rice. *Mol. Cells* **2007**, *24*, 83–94. [\[PubMed\]](#)
55. Shim, K.-C.; Kim, S.H.; Jeon, Y.-A.; Lee, H.-S.; Adeva, C.; Kang, J.-W.; Kim, H.-J.; Tai, T.H.; Ahn, S.-N. A RING-Type E3 Ubiquitin Ligase, OsGW2, Controls Chlorophyll Content and Dark-Induced Senescence in Rice. *Int. J. Mol. Sci.* **2020**, *21*, 1704. [\[CrossRef\]](#)
56. Fu, J.D.; Yan, Y.F.; Kim, M.Y.; Lee, S.H.; Lee, B.W. Population-specific quantitative trait loci mapping for functional stay-green trait in rice (*Oryza sativa* L.). *Genome* **2011**, *5*, 235.
57. Lim, J.H.; Yang, H.J.; Jung, K.H.; Yoo, S.C.; Paek, N.-C. Quantitative Trait Locus Mapping and Candidate Gene Analysis for Plant Architecture Traits Using Whole Genome Re-Sequencing in Rice. *Mol. Cells* **2014**, *37*, 149–160. [\[PubMed\]](#)
58. Singh, U.M.; Sinha, P.; Dixit, S.; Abbai, R.; Venkateshwarlu, C.; Chitkineni, A.; Singh, V.K.; Varshney, R.K.; Kumar, A. Unraveling candidate genomic regions responsible for delayed leaf senescence in rice. *PLoS ONE* **2020**, *15*, e0240591. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Wang, Q.; Xie, W.; Xing, H.; Yan, J.; Meng, X.; Li, X.; Fu, X.; Xu, J.; Lian, X.; Yu, S.; et al. Genetic Architecture of Natural Variation in Rice Chlorophyll Content Revealed by a Genome-Wide Association Study. *Mol. Plant* **2015**, *8*, 946–957. [\[PubMed\]](#)
60. Zhao, Y.; Qiang, C.; Wang, X.; Chen, Y.; Deng, J.; Jiang, C.; Sun, X.; Chen, H.; Li, J.; Piao, W.; et al. New alleles for chlorophyll content and stay-green traits revealed by a genome wide association study in rice (*Oryza sativa*). *Sci. Rep.* **2019**, *9*, 2541. [\[CrossRef\]](#)
61. Hu, C.; Shi, J.; Quan, S.; Cui, B.; Kleessen, S.; Nikoloski, Z.; Tohge, T.; Alexander, D.; Guo, L.; Lin, H.; et al. Metabolic variation between *japonica* and *indica* rice cultivars as revealed by non-targeted metabolomics. *Sci. Rep.* **2014**, *4*, 5067. [\[CrossRef\]](#)
62. Abdelkhalik, A.F.; Shishido, R.; Nomura, K.; Ikehashi, H. QTL-based analysis of leaf senescence in an *indica/japonica* hybrid in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* **2005**, *110*, 1226–1235. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Ramkumar, M.K.; Senthil Kumar, S.; Gaikwad, K.; Pandey, R.; Chinnusamy, V.; Singh, N.K.; Singh, A.K.; Mohapatra, T.; Sevanthi, A.M. A Novel Stay-Green Mutant of Rice with Delayed Leaf Senescence and Better Harvest Index Confers Drought Tolerance. *Plants* **2019**, *8*, 375. [\[CrossRef\]](#)
64. Sakuraba, Y.; Kim, D.; Han, S.H.; Kim, S.H.; Piao, W.; Yanagisawa, S.; An, G.; Paek, N.-C. Multilayered Regulation of Membrane-Bound ONAC054 is Essential for Absciscic Acid-induced Leaf Senescence in Rice. *Plant Cell* **2020**, *32*, 630–649. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Kang, K.; Shim, Y.; Gi, E.; An, G.; Paek, N.-C. Mutation of ONAC096 Enhances Grain Yield by Increasing Panicle Number and Delaying Leaf Senescence during Grain Filling in Rice. *Int. J. Mol. Sci.* **2019**, *20*, 5241. [\[CrossRef\]](#)
66. Sarwat, M.; Naqvi, A.R.; Ahmad, P.; Ashraf, M.; Akram, N.A. Phytohormones and microRNAs as sensors and regulators of leaf senescence: Assigning macro roles to small molecules. *Biotechnol. Adv.* **2013**, *31*, 1153–1171. [\[CrossRef\]](#)
67. Bengoa Luoni, S.; Astigueta, F.H.; Nicosia, S.; Moschen, S.; Fernandez, P.; Heinz, R. Transcription factors associated with leaf senescence in crops. *Plants* **2019**, *8*, 411. [\[CrossRef\]](#) [\[PubMed\]](#)
68. Singh, S.; Koyama, H.; Bhati, K.K.; Alok, A. The biotechnological importance of the plant-specific NAC transcription factor family in crop improvement. *J. Plant Res.* **2021**. [\[CrossRef\]](#)
69. Nuruzzaman, M.; Manimekalai, R.; Shari, A.M.; Satoh, K.; Kondoh, H.; Ooka, H.; Kikuchi, S. Genome-wide analysis of NAC transcription factor family in rice. *Gene* **2010**, *465*, 30–44. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Kim, H.J.; Park, J.H.; Kim, J.; Kim, J.J.; Hong, S.; Kim, J.; Kim, J.H.; Woo, H.R.; Hyeon, C.; Lim, P.O.; et al. Time-evolving genetic networks reveal a NAC troika that negatively regulates leaf senescence in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, E4930–E4939. [\[CrossRef\]](#)
71. Liang, C.; Wang, Y.; Zhu, Y.; Tang, J.; Hu, B.; Liu, L.; Ou, S.; Wu, H.; Sun, X.; Chu, J.; et al. OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 10013–10018. [\[CrossRef\]](#)
72. Zhou, Y.; Huang, W.; Liu, L.; Chen, T.; Zhou, F.; Lin, Y. Identification and functional characterization of a rice NAC gene involved in the regulation of leaf senescence. *BMC Plant Biol.* **2013**, *13*, 132. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Chen, X.; Wang, Y.; Lv, B.; Li, J.; Luo, L.; Lu, S.; Zhang, X.; Ma, H.; Ming, F. The NAC family transcription factor OsNAP confers abiotic stress response through the ABA pathway. *Plant Cell Physiol.* **2014**, *55*, 604–619. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Mao, C.; Lu, S.; Lv, B.; Zhang, B.; Shen, J.; He, J.; Luo, L.; Xi, D.; Chen, X.; Ming, F. A Rice NAC transcription factor promotes leaf senescence via ABA biosynthesis. *Plant Physiol.* **2017**, *174*, 1747–1763. [\[CrossRef\]](#) [\[PubMed\]](#)

75. Li, L.; He, Y.; Zhang, Z.; Shi, Y.; Zhang, X.; Xu, X.; Wu, J.L.; Tang, S. OsNAC109 regulates senescence, growth and development by altering the expression of senescence- and phytohormone-associated genes in rice. *Plant Mol. Biol.* **2021**, *105*, 637–654. [\[CrossRef\]](#)
76. Sakuraba, Y.; Piao, W.; Lim, J.H.; Han, S.H.; Kim, Y.S.; An, G.; Paek, N.-C. Rice ONAC106 Inhibits leaf senescence and increases salt tolerance and tiller angle. *Plant Cell Physiol.* **2015**, *56*, 2325–2339. [\[CrossRef\]](#)
77. El Mannai, Y.; Akabane, K.; Hiratsu, K.; Satoh-Nagasawa, N.; Wabiko, H. The NAC Transcription factor gene OsY37 (ONAC011) promotes leaf senescence and accelerates heading time in rice. *Int. J. Mol. Sci.* **2017**, *18*, 2165. [\[CrossRef\]](#)
78. Sato, Y.; Antonio, B.; Namiki, N.; Motoyama, R.; Sugimoto, K.; Takehisa, H.; Minami, H.; Kamatsuki, K.; Kusaba, M.; Hirochika, H.; et al. Field transcriptome revealed critical developmental and physiological transitions involved in the expression of growth potential in japonica rice. *BMC Plant Biol.* **2011**, *11*, 10. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Katiyar, A.; Smita, S.; Lenka, S.K.; Rajwanshi, R.; Chinnusamy, V.; Bansal, K.C. Genome-wide classification and expression analysis of MYB transcription factor families in rice and Arabidopsis. *BMC Genom.* **2012**, *13*, 544. [\[CrossRef\]](#) [\[PubMed\]](#)
80. Park, D.-Y.; Shim, Y.; Gi, E.; Lee, B.-D.; An, G.; Kang, K.; Paek, N.-C. The MYB-related transcription factor RADIALIS-LIKE3 (OsRL3) functions in ABA-induced leaf senescence and salt sensitivity in rice. *Environ. Exp. Bot.* **2018**, *156*, 86–95. [\[CrossRef\]](#)
81. Piao, W.; Kim, S.H.; Lee, B.D.; An, G.; Sakuraba, Y.; Paek, N.-C. Rice transcription factor OsMYB102 delays leaf senescence by down-regulating abscisic acid accumulation and signaling. *J. Exp. Bot.* **2019**, *70*, 2699–2715. [\[CrossRef\]](#)
82. Sahebi, M.; Hanafi, M.M.; Rafii, M.Y.; Mahmud, T.M.M.; Azizi, P.; Osman, M.; Abiri, R.; Taheri, S.; Kalhori, N.; Shabanimofrad, M.; et al. Improvement of Drought Tolerance in Rice (*Oryza sativa* L.): Genetics, Genomic tools, and the WRKY Gene Family. *Biomed. Res. Int.* **2018**, *2018*, 3158474. [\[CrossRef\]](#)
83. Liu, L.; Xu, W.; Hu, X.; Liu, H.; Lin, Y. W-box and G-box elements play important roles in early senescence of rice flag leaf. *Sci. Rep.* **2016**, *6*, 20881. [\[CrossRef\]](#)
84. Ricachenevsky, F.K.; Sperotto, R.A.; Menguer, P.K.; Fett, J.P. Identification of Fe-excess-induced genes in rice shoots reveals a WRKY transcription factor responsive to Fe, drought and senescence. *Mol. Biol. Rep.* **2010**, *37*, 3735–3745. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Jing, S.; Zhou, X.; Song, Y.; Yu, D. Heterologous expression of OsWRKY23 gene enhances pathogen defense and dark-induced leaf senescence in *Arabidopsis*. *Plant Growth Regul.* **2009**, *58*, 181–190. [\[CrossRef\]](#)
86. Han, M.; Kim, C.Y.; Lee, J.; Lee, S.K.; Jeon, J.S. OsWRKY42 represses *OsMT1d* and induces reactive oxygen species and leaf senescence in rice. *Mol. Cells* **2014**, *37*, 532–539. [\[CrossRef\]](#)
87. Kim, T.; Kang, K.; Kim, S.-H.; An, G.; Paek, N.-C. OsWRKY5 Promotes Rice Leaf Senescence via Senescence-Associated NAC and Absciscic Acid Biosynthesis Pathway. *Int. J. Mol. Sci.* **2019**, *20*, 4437. [\[CrossRef\]](#)
88. Huq, E.; Al-Sady, B.; Hudson, M.; Kim, C.; Apel, K.; Quail, P.H. Phytochrome-interacting factor 1 is a critical bHLH regulator of chlorophyll biosynthesis. *Science* **2004**, *305*, 1937–1941. [\[CrossRef\]](#)
89. Sakuraba, Y.; Jeong, J.; Kang, M.Y.; Kim, J.; Paek, N.-C.; Choi, G. Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf senescence in *Arabidopsis*. *Nat. Commun.* **2014**, *5*, 4636. [\[CrossRef\]](#)
90. Sakuraba, Y.; Kim, E.Y.; Paek, N.-C. Roles of rice PHYTOCHROME-INTERACTING FACTOR-LIKE1 (OsPIL1) in leaf senescence. *Plant Signal Behav.* **2017**, *12*, e1362522. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Sakuraba, Y.; Kim, E.Y.; Han, S.H.; Piao, W.; An, G.; Todaka, D.; Yamaguchi-Shinozaki, K.; Paek, N.-C. Rice Phytochrome-Interacting Factor-Like1 (OsPIL1) is involved in the promotion of chlorophyll biosynthesis through feed-forward regulatory loops. *J. Exp. Bot.* **2017**, *68*, 4103–4114. [\[CrossRef\]](#) [\[PubMed\]](#)
92. Jan, A.; Maruyama, K.; Todaka, D.; Kidokoro, S.; Abo, M.; Yoshimura, E.; Shinozaki, K.; Nakashima, K.; Yamaguchi-Shinozaki, K. OsTZF1, a CCCH-tandem zinc finger protein, confers delayed senescence and stress tolerance in rice by regulating stress-related genes. *Plant Physiol.* **2013**, *161*, 1202–1216. [\[CrossRef\]](#)
93. Kong, Z.; Li, M.; Yang, W.; Xu, W.; Xue, Y. A novel nuclear-localized CCCH-type zinc finger protein, OsDOS, is involved in delaying leaf senescence in rice. *Plant Physiol.* **2006**, *141*, 1376–1388. [\[CrossRef\]](#) [\[PubMed\]](#)
94. Lim, C.; Kang, K.; Shim, Y.; Sakuraba, Y.; An, G.; Paek, N.-C. Rice ETHYLENE RESPONSE FACTOR 101 Promotes Leaf Senescence Through Jasmonic Acid-Mediated Regulation of *OsNAP* and *OsMYC2*. *Front. Plant Sci.* **2020**, *11*, 1096. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Uji, Y.; Akimitsu, K.; Gomi, K. Identification of OsMYC2-regulated senescence-associated genes in rice. *Planta* **2017**, *245*, 1241–1246. [\[CrossRef\]](#)
96. Lee, S.; Marmagne, A.; Park, J.; Fabien, C.; Yim, Y.; Kim, S.J.; Kim, T.H.; Lim, P.O.; Masclaux-Daubresse, C.; Nam, H.G. Concurrent activation of *OsAMT1;2* and *OsGOGAT1* in rice leads to enhanced nitrogen use efficiency under nitrogen limitation. *Plant J.* **2020**, *103*, 7–20. [\[CrossRef\]](#)
97. Wada, S.; Hayashida, Y.; Izumi, M.; Kurusu, T.; Hanamata, S.; Kanno, K.; Kojima, S.; Yamaya, T.; Kuchitsu, K.; Makino, A.; et al. Autophagy supports biomass production and nitrogen use efficiency at the vegetative stage in rice. *Plant Physiol.* **2015**, *168*, 60–73. [\[CrossRef\]](#)
98. Masclaux-Daubresse, C.; Danielvedele, F.; Dechorgnat, J.; Chardon, F.; Gaufichon, L.; Suzuki, A. Nitrogen uptake, assimilation and remobilization in plants: Challenges for sustainable and productive agriculture. *Ann. Bot.* **2010**, *105*, 1141–1157. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Chen, Q.; Shinozaki, D.; Luo, J.; Pottier, M.; Havé, M.; Marmagne, A.; Reisdorf-Cren, M.; Chardon, F.; Thomine, S.; Yoshimoto, K.; et al. Autophagy and Nutrients Management in Plants. *Cells* **2019**, *8*, 1426. [\[CrossRef\]](#)

100. Ishida, H.; Yoshimoto, K.; Izumi, M.; Reisen, D.; Yano, Y.; Makino, A.; Ohsumi, Y.; Hanson, M.R.; Mae, T. Mobilization of rubisco and stroma-localized fluorescent proteins of chloroplasts to the vacuole by an ATG gene-dependent autophagic process. *Plant Physiol.* **2008**, *148*, 142–155. [[CrossRef](#)]
101. Nakamura, S.; Hidema, J.; Sakamoto, W.; Ishida, H.; Izumi, M. Selective Elimination of Membrane-Damaged Chloroplasts via Microautophagy. *Plant Physiol.* **2018**, *177*, 1007–1026. [[CrossRef](#)]
102. Guiboileau, A.; Yoshimoto, K.; Soulay, F.; Bataillé, M.P.; Avice, J.C.; Masclaux-Daubresse, C. Autophagy machinery controls nitrogen remobilization at the whole-plant level under both limiting and ample nitrate conditions in Arabidopsis. *New Phytol.* **2012**, *194*, 732–740. [[CrossRef](#)]
103. Pottier, M.; Dumont, J.; Masclaux-Daubresse, C.; Thomine, S. Autophagy is essential for optimal translocation of iron to seeds in Arabidopsis. *J. Exp. Bot.* **2019**, *70*, 859–869. [[CrossRef](#)]
104. Lornac, A.; Havé, M.; Chardon, F.; Soulay, F.; Clément, G.; Avice, J.C.; Masclaux-Daubresse, C. Autophagy Controls Sulphur Metabolism in the Rosette Leaves of Arabidopsis and Facilitates S Remobilization to the Seeds. *Cells* **2020**, *9*, 332. [[CrossRef](#)] [[PubMed](#)]
105. Chen, Q.; Soulay, F.; Saudemont, B.; Elmayan, T.; Marmagne, A.; Masclaux-Daubresse, C. Overexpression of ATG8 in Arabidopsis Stimulates Autophagic Activity and Increases Nitrogen Remobilization Efficiency and Grain Filling. *Plant Cell Physiol.* **2019**, *60*, 343–352. [[CrossRef](#)]
106. Xia, K.F.; Liu, T.; Ouyang, J.; Wang, R.; Fan, T.; Zhang, M.Y. Genome-wide identification, classification, and expression analysis of autophagy-associated gene homologues in rice (*Oryza sativa* L.). *DNA Res.* **2011**, *18*, 363–377. [[CrossRef](#)]
107. Yu, J.; Zhen, X.; Li, X.; Li, N.; Xu, F. Increased Autophagy of Rice Can Increase Yield and Nitrogen Use Efficiency (NUE). *Front. Plant Sci.* **2019**, *10*, 584. [[CrossRef](#)]
108. Fan, T.; Yang, W.; Zeng, X.; Xu, X.; Xu, Y.; Fan, X.; Luo, M.; Tian, C.; Xia, K.; Zhang, M. A Rice Autophagy gene *OsATG8b* is involved in nitrogen remobilization and control of grain quality. *Front. Plant Sci.* **2020**, *11*, 588. [[CrossRef](#)] [[PubMed](#)]
109. Zhen, X.; Li, X.; Yu, J.; Xu, F. *OsATG8c*-mediated increased autophagy regulates the yield and nitrogen use efficiency in rice. *Int. J. Mol. Sci.* **2019**, *20*, 4956. [[CrossRef](#)] [[PubMed](#)]
110. Zhen, X.; Zheng, N.; Yu, J.; Bi, C.; Xu, F. Autophagy mediates grain yield and nitrogen stress resistance by modulating nitrogen remobilization in rice. *PLoS ONE* **2021**, *16*, e0244996. [[CrossRef](#)] [[PubMed](#)]
111. Havé, M.; Balliau, T.; Cottyn-Boitte, B.; Déron, E.; Cuff, G.; Soulay, F.; Lornac, A.; Reichman, P.; Dissmeyer, N.; Avice, J.C.; et al. Increases in activity of proteasome and papain-like cysteine protease in Arabidopsis autophagy mutants: Back-up compensatory effect or cell-death promoting effect? *J. Exp. Bot.* **2018**, *69*, 1369–1385. [[CrossRef](#)] [[PubMed](#)]
112. Yao, X.; Nie, J.; Bai, R.; Sui, X. Amino Acid Transporters in Plants: Identification and Function. *Plants* **2020**, *9*, 972. [[CrossRef](#)]
113. Hayashi, H.; Chino, M. Chemical composition of phloem sap from the upper most internode of the rice plant. *Plant Cell Physiol.* **1990**, *31*, 247–251.
114. Lea, P.J.; Sodek, L.; Parry, M.A.J.; Shewry, P.R.; Halford, N.G. Asparagine in plants. *Ann. Appl. Biol.* **2007**, *150*, 1–26. [[CrossRef](#)]
115. Tabuchi, M.; Sugiyama, K.; Ishiyama, K.; Inoue, E.; Sato, T.; Takahashi, H.; Yamaya, T. Severe reduction in growth rate and grain filling of rice mutants lacking *OsGS1;1*, a cytosolic glutamine synthetase1;1. *Plant J.* **2005**, *42*, 641–651. [[CrossRef](#)] [[PubMed](#)]
116. Xu, G.; Fan, X.; Miller, A.J. Plant nitrogen assimilation and use efficiency. *Annu. Rev. Plant Biol.* **2012**, *63*, 153–182. [[CrossRef](#)] [[PubMed](#)]
117. Yang, X.; Nian, J.; Xie, Q.; Feng, J.; Zhang, F.; Jing, H.; Zhang, J.; Dong, G.; Liang, Y.; Peng, J.; et al. Rice ferredoxin-dependent glutamate synthase regulates nitrogen-carbon metabolomes and is genetically differentiated between *japonica* and *indica* Subspecies. *Mol. Plant.* **2016**, *9*, 1520–1534. [[CrossRef](#)] [[PubMed](#)]
118. Zeng, D.D.; Qin, R.; Li, M.; Alamin, M.; Jin, X.L.; Liu, Y.; Shi, C.H. The ferredoxin-dependent glutamate synthase (*OsFd-GOGAT*) participates in leaf senescence and the nitrogen remobilization in rice. *Mol. Genet. Genom.* **2017**, *292*, 385–395. [[CrossRef](#)] [[PubMed](#)]
119. Gaufichon, L.; Reisdorf-Cren, M.; Rothstein, S.J.; Chardon, F.; Suzukia, A. Biological functions of asparagine synthetase in plants. *Plant Sci.* **2010**, *179*, 141–153. [[CrossRef](#)]
120. Lee, S.; Park, J.; Lee, J.; Shin, D.; Marmagne, A.; Lim, P.O.; Masclaux-Daubresse, C.; An, G.; Nam, H.G. *OsASN1* overexpression in rice increases grain protein content and yield under nitrogen-limiting conditions. *Plant Cell Physiol.* **2020**, *61*, 1309–1320. [[CrossRef](#)]
121. Zhao, H.; Ma, H.; Yu, L.; Wang, X.; Zhao, J. Genome-wide survey and expression analysis of amino acid transporter gene family in rice (*Oryza sativa* L.). *PLoS ONE* **2012**, *7*, e49210. [[CrossRef](#)]
122. Santiago, J.P.; Tegeder, M. Connecting Source with Sink: The Role of Arabidopsis AAP8 in Phloem Loading of Amino Acids. *Plant Physiol.* **2016**, *171*, 508–521. [[CrossRef](#)]
123. Wei, Q.; Yan, Z.; Xiong, Y.; Fang, Z. Altered Expression of *OsAAP3* Influences Rice Lesion Mimic and Leaf Senescence by Regulating Arginine Transport and Nitric Oxide Pathway. *Int. J. Mol. Sci.* **2021**, *22*, 2181. [[CrossRef](#)] [[PubMed](#)]
124. Lu, K.; Wu, B.; Wang, J.; Zhu, W.; Nie, H.; Qian, J.; Huang, W.; Fang, Z. Blocking amino acid transporter *OsAAP3* improves grain yield by promoting outgrowth buds and increasing tiller number in rice. *Plant Biotechnol. J.* **2018**, *16*, 1710–1722. [[CrossRef](#)] [[PubMed](#)]
125. Taylor, M.R.; Reinders, A.; Ward, J.M. Transport Function of Rice Amino Acid Permeases (AAPs). *Plant Cell Physiol.* **2015**, *56*, 1355–1363. [[CrossRef](#)]

126. Peng, B.; Kong, H.; Li, Y.; Wang, L.; Zhong, M.; Sun, L.; Gao, G.; Zhang, Q.; Luo, L.; Wang, G.; et al. OsAAP6 functions as an important regulator of grain protein content and nutritional quality in rice. *Nat. Commun.* **2014**, *5*, 4847. [[CrossRef](#)] [[PubMed](#)]
127. Wang, J.; Wu, B.; Lu, K.; Wei, Q.; Qian, J.; Chen, Y.; Fang, Z. The Amino Acid Permease 5 (OsAAP5) Regulates Tiller Number and Grain Yield in Rice. *Plant Physiol.* **2019**, *180*, 1031–1045. [[CrossRef](#)] [[PubMed](#)]
128. Ji, Y.; Huang, W.; Wu, B.; Fang, Z.; Wang, X. The amino acid transporter AAP1 mediates growth and grain yield by regulating neutral amino acid uptake and reallocation in *Oryza sativa*. *J. Exp. Bot.* **2020**, *71*, 4763–4777. [[CrossRef](#)] [[PubMed](#)]
129. Fang, Z.; Wu, B.; Ji, Y. The Amino Acid Transporter OsAAP4 Contributes to Rice Tillering and Grain Yield by Regulating Neutral Amino Acid Allocation through Two Splicing Variants. *Rice* **2021**, *14*, 2. [[CrossRef](#)]
130. Gull, A.; Lone, A.A.; Wani, N.U.I. Biotic and Abiotic Stresses in Plants. In *Abiotic and Biotic Stress in Plants*; IntechOpen: London, UK, 2019.
131. Pradhan, P.; Fischer, G.; van Velthuisen, H.; Reusser, M.E.; Kropp, J.P. Closing yield gaps: How sustainable can we be? *PLoS ONE* **2015**, *10*, e0129487. [[CrossRef](#)]
132. Gepstein, S.; Glick, B.R. Strategies to ameliorate abiotic stress-induced plant senescence. *Plant Mol. Biol.* **2013**, *82*, 623–633. [[CrossRef](#)]
133. Shen, J.; Lv, B.; Luo, L.; He, J.; Mao, C.; Xi, D.; Ming, F. The NAC-type transcription factor OsNAC2 regulates ABA-dependent genes and abiotic stress tolerance in rice. *Sci. Rep.* **2017**, *7*, 40641. [[CrossRef](#)] [[PubMed](#)]
134. Yuan, X.; Wang, H.; Cai, J.; Bi, Y.; Li, D.; Song, F. Rice NAC transcription factor ONAC066 functions as a positive regulator of drought and oxidative stress response. *BMC Plant Biol.* **2019**, *19*, 278. [[CrossRef](#)] [[PubMed](#)]
135. Li, Y.; Liao, S.; Mei, P.; Pan, Y.; Zhang, Y.; Zheng, X.; Xie, Y.; Miao, Y. OsWRKY93 Dually Functions Between Leaf Senescence and in Response to Biotic Stress in Rice. *Front. Plant Sci.* **2021**, *12*, 643011. [[CrossRef](#)] [[PubMed](#)]
136. Raza, A.; Razzaq, A.; Mehmood, S.S.; Zou, X.; Zhang, X.; Lv, Y.; Xu, J. Impact of Climate Change on Crops Adaptation and Strategies to Tackle Its Outcome: A Review. *Plants* **2019**, *8*, 34. [[CrossRef](#)]
137. Dhankher, O.P.; Foyer, C.H. Climate resilient crops for improving global food security and safety. *Plant Cell Environ.* **2018**, *5*, 877–884. [[CrossRef](#)]
138. Hu, H.; Xiong, L. Genetic engineering and breeding of drought-resistant crops. *Annu. Rev. Plant Biol.* **2014**, *65*, 715–741. [[CrossRef](#)]
139. Kilasi, N.L.; Singh, J.; Vallejos, C.E.; Ye, C.; Jagadish, S.V.K.; Kusolwa, P. Heat Stress Tolerance in Rice (*Oryza sativa* L.): Identification of Quantitative Trait Loci and Candidate Genes for Seedling Growth Under Heat Stress. *Front. Plant Sci.* **2018**, *9*, 1578. [[CrossRef](#)]
140. Schramm, F.; Larkindale, J.; Kiehlmann, E.; Ganguli, A.; Englich, E.; Vierling, E.; von Koskull-Döring, P. A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3 regulates the heat stress response of *Arabidopsis*. *Plant J.* **2008**, *53*, 264–274. [[CrossRef](#)] [[PubMed](#)]
141. Ren, Y.; Huang, Z.; Jiang, H.; Wang, Z.; Wu, F.; Xiong, Y.; Yao, J. A heat stress responsive NAC transcription factor heterodimer plays key roles in rice grain filling. *J. Exp. Bot.* **2021**, *72*, 2947–2964. [[CrossRef](#)]
142. Liu, L.; Zhou, Y.; Szczerba, M.W.; Li, X.; Lin, Y. Identification and application of a rice senescence-associated promoter. *Plant Physiol.* **2010**, *153*, 1239–1249. [[CrossRef](#)]
143. Zani, D.; Crowther, T.W.; Mo, L.; Renner, S.S.; Zohner, C.M. Increased growing-season productivity drives earlier autumn leaf senescence in temperate trees. *Science* **2020**, *370*, 1066–1071. [[CrossRef](#)] [[PubMed](#)]
144. Breeze, E.; Harrison, E.; McHattie, S.; Hughes, L.; Hickman, R.; Hill, C. High-Resolution Temporal Profiling of Transcripts during *Arabidopsis* Leaf Senescence Reveals a Distinct Chronology of Processes and Regulation. *Plant Cell* **2011**, *23*, 873–894. [[CrossRef](#)] [[PubMed](#)]
145. Woo, H.R.; Koo, H.J.; Kim, J.; Jeong, H.; Yang, J.O.; Lee, I.H.; Jun, J.H.; Choi, S.H.; Park, S.J.; Kang, B.; et al. Programming of plant leaf senescence with temporal and inter-organellar coordination of transcriptome in *Arabidopsis*. *Plant Physiol.* **2016**, *171*, 452–467. [[CrossRef](#)] [[PubMed](#)]