HAL
open science

Opinion: Using eDNA fingerprinting in high mountain environments to support soil restoration and hazard control

Amaury Frankl, O. Evrard, Elie Verleyen, Erik Cammeraat, Alexia Stokes

To cite this version:

Amaury Frankl, O. Evrard, Elie Verleyen, Erik Cammeraat, Alexia Stokes. Opinion: Using eDNA fingerprinting in high mountain environments to support soil restoration and hazard control. LASOSU2021: Online International Conference for Land Degradation, Soil Conservation and Sustainable Development, World Association of Soil and Water Conservation (WASWAC); Italian Association of Agricultural Engineering (AIIA); Soil Erosion Division of CSWCS (China), Aug 2021, Dalian, China. hal-03335309

HAL Id: hal-03335309

https://hal.inrae.fr/hal-03335309v1

Submitted on 6 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LASOSUD 2021
International Forum on
Land Degradation, Soil Conservation and Sustainable Development
Dalian, China | 21-23 August 2021

Opinion: Using eDNA fingerprinting in high mountain environments to support soil restoration and hazard control

A. Frankl*[1,2], O. Evrard[3], E. Verleyen[4], E. Cammeraat[5], A. Stokes[1]
[1] INRAE, AMAP, IRD, CIRAD, CNRS, University Montpellier, France
[2] Department of Geography, Ghent University, Belgium
[3] Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL), Unité Mixte de Recherche 8212 (CEA/CNRS/UVSQ), Université Paris-Saclay, France
[4] Department of Biology, Ghent University, Belgium
[5] Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, The Netherlands
*Corresponding A. Frankl, e-mail: Amaury.Frankl@cirad.fr

Keywords: Landslides, Sediment source fingerprinting, Soil Erosion, Vegetation.

Abstract

Mitigating erosion, mass movements, and geohazards in high mountains is increasingly conceived within frameworks of ecological restoration, that is, recovering the form and function of ecosystems that have been damaged by degradation (Hubble et al. 2017). From a geomorphological point of view, ecological restoration involves both the prevention and control of slope and riverbank instabilities as well as the confinement of runoff and sediment regimes to the capacity river channels. In this regard, the practice of soil and water bioengineering is rapidly emerging as a short-term hazard control that can enable long-term ecological recovery (Rey et al. 2019). Vegetation, as a chief ecological engineer, is key to soil and water bioengineering applications. However, the application of soil and water bioengineering in high mountains is limited by severe ecological conditions, making plant establishment and ecological recovery times are slow in high mountains (Dupin et al. 2019).

While applications of sediment source fingerprinting using for example geochemical or radionuclide soil signatures yield a rough distinction between sediment sources, they cannot reflect the multiple vegetation covers that are relevant source types and should be discriminated in high mountain environments to prioritize restoration works. Vegetation may be the most distinctive feature of high mountains, where the underlying lithology is heterogeneous and soils are mainly shallow and poorly developed. Because there are strong interrelations between land cover and geomorphological processes in high mountain environments (Geertsema and Pojar 2007; Giaccone et al. 2019; Lizaga et al. 2019), the use of land cover-based sediment tracers would be particularly meaningful. eDNA has the highest source discrimination potential in that regard, providing information up to the species level and reflecting changes in vegetation on over short timescales. Furthermore, eDNA signals in sediments will be strongest from areas experiencing higher erosion rates and which are highly connected with the hydrographic network. The use of eDNA sediment source fingerprinting would thus allow the investigation of complex and often poorly understood

relationships between vegetation cover, restoration activities, and geomorphological response at the catchment scale.

To improve the success rates of restoration activities, collaboration between scientists and stakeholders can accelerate technology transfer rates (Stokes et al. 2014; Giupponi et al. 2019; Rey et al. 2019). However, time and budget constraints often hamper in-situ monitoring of soil and water bioengineering applications, and very few monitoring programs exist (Giupponi et al. 2019). Knowledge of success rates is, however, essential for restoration (Frankl et al. 2021). To this end, sediment source fingerprinting has been shown to provide a valid framework for supporting soil restoration activities (Mukundan et al. 2012). Environmental DNA has already been used to successfully monitor restoration programs, but with a focus on fungal species (Yan et al., 2018). We opinionate that eDNA fingerprinting – as an emerging technique – could be particularly useful to support soil restoration and hazard control in high mountain environments.

References

Dupin B, Malaval S, Couëron G, et al (2019) Restauration écologique de prairies et de pelouses Pyrénéennes: un guide technique pour régénérer les sols et les végétations dégradés en montagne. écovars, OPCC, Bagnères-de-Bigorre

Frankl A, Nyssen J, Vanmaercke M, Poesen J (2021) Gully prevention and control: techniques, failures and effectiveness. Earth Surf Process Landforms 46:220-238. doi: 10.1002/esp.5033

Geertsema M, Pojar JJ (2007) Influence of landslides on biophysical diversity - A perspective from British Columbia. Geomorphology 89:55–69. doi: 10.1016/j.geomorph.2006.07.019

Giaccone E, Luoto M, Vittoz P, et al (2019) Influence of microclimate and geomorphological factors on alpine vegetation in the Western Swiss Alps. Earth Surf Process Landforms 44:3093–3107. doi: 10.1002/esp.4715

Giupponi L, Borgonovo G, Giorgi A, Bischetti GB (2019) How to renew soil bioengineering for slope stabilization: some proposals. Landsc Ecol Eng 15:37–50. doi: 10.1007/s11355-018-0359-9

Hubble T, Clarke S, Stokes A, Phillips C (2017) 4th International Conference on soil bio- and eco-engineering (SBEE2016) 'The Use of Vegetation to Improve Slope Stability.' Ecol Eng 109:141–144. doi: 10.1016/j.ecoleng.2017.11.003

Lizaga I, Gaspar L, Blake WH, et al (2019) Fingerprinting changes of source apportionments from mixed land uses in stream sediments before and after an exceptional rainstorm event. Geomorphology 341:216–229. doi: 10.1016/j.geomorph.2019.05.015

Mukundan R, Walling DE, Gellis AC, et al (2012) Sediment Source Fingerprinting: Transforming From a Research Tool to a Management Tool. J Am Water Resour Assoc 48:1241–1257. doi: 10.1111/j.1752-1688.2012.00685.x

Rey F, Bifulco C, Bischetti GB, et al (2019) Soil and water bioengineering: Practice and research needs for reconciling natural hazard control and ecological restoration. Sci Total Environ 648:1210–1218. doi: 10.1016/j.scitotenv.2018.08.217

Stokes A, Douglas GB, Fourcaud T, et al (2014) Ecological mitigation of hillslope instability: ten key issues facing researchers and practitioners. Plant Soil 377:1–23. doi: 10.1007/s11104-014-2044-6

Yan DF, Mills JG, Gellie NJC, et al (2018) High-throughput eDNA monitoring of fungi to track functional recovery in ecological restoration. Biol Conserv 217:113–120. doi: 10.1016/j.biocon.2017.10.035