



**HAL**  
open science

# Haploid Embryos: Being Like Mommy or Like Daddy?

Thomas Widiez

► **To cite this version:**

Thomas Widiez. Haploid Embryos: Being Like Mommy or Like Daddy?. Trends in Plant Science, 2021, 26 (5), pp.425-427. 10.1016/j.tplants.2021.02.010 . hal-03335618

**HAL Id: hal-03335618**

**<https://hal.inrae.fr/hal-03335618>**

Submitted on 6 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Haploid embryos: Being like Mommy or like Daddy?

Thomas Widiez<sup>1\*</sup>

<sup>1</sup> Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, UCB  
Lyon 1, CNRS, INRAE, F-69342, Lyon, France.

\* corresponding author: [thomas.widiez@ens-lyon.fr](mailto:thomas.widiez@ens-lyon.fr) (T. Widiez)

Twitter handle: @ThomasWidiez

Website: <http://www.ens-lyon.fr/RDP/spip.php?rubrique15&lang=en>

PMID: 33678580 DOI: 10.1016/j.tplants.2021.02.010

## Trends in Plant Science

Volume 26, Issue 5, May 2021, Pages 425-427



Spotlight

## Haploid Embryos: Being Like Mommy or Like Daddy?

Thomas Widiez<sup>1</sup>   

### ABSTRACT:

*In planta* haploid embryo induction is a powerful plant breeding tool, but is limited to few crops. Two studies reported seed-based haploid systems producing paternal haploid embryos by engineering CENTROMERIC-HISTONE3. Along with recent translation of maize maternal haploid induction ability in wheat and rice, it extends haploid inducer lines collection.

### Background

Haploid embryos contain only one set of parental chromosomes ( $n$ ), instead of the classical situation with two sets of chromosomes ( $2n$ ), one from the mother and one from the father. Nevertheless, haploid embryos and their subsequent haploid plantlets represent the basis of the Doubled Haploid (DH) technology which is a significant plant breeding tool [1,2]. DH technology could be simply summarized by: (i) the production of haploid embryos, and (ii) duplication (copy-paste) of the haploid genome to recover normal ploidy state. DH technology allows efficient plant breeding cycles, mainly by shortening the time for creating fixed genetic material (inbred line), since only two generations are needed to obtain homozygous plants, instead of six or more generations when using conventional crosses [1,2]. Thus, DH pipelines can offer rapid evaluation of phenotypic traits in

30 homogenous progeny, and the production of haploid embryos is central to this process. Although *in*  
31 *vitro* methods are most widely used to produce haploid embryos starting from the maternal or paternal  
32 germline (gametes or their precursors), they are costly and genotype-dependent [1]. The use of haploid  
33 inducer lines represents an attractive way to induce haploid embryos entirely *in planta*. These *in planta*  
34 methods produce haploid embryos within seeds, circumventing a major bottleneck of both *in vitro* and  
35 wide-hybridization methods, namely embryo rescue. A simple intraspecific cross between a plant of  
36 interest and a haploid inducer line allows the production of viable seeds, among which a certain  
37 proportion have haploid embryos (**Figure 1**). The uniparental haploid embryos lack the haploid inducer  
38 genome, making either paternal- or maternal haploid (**Figure 1**). Two main types of *in planta* haploid  
39 inducer lines were reported: (i) the CENH3-based induction system based on engineering of  
40 Centromeric Histone 3 (CENH3) (**Figure 1A**), and (ii) the maternal maize haploid induction ability which  
41 is based on mutation of the *MATRILINEAL (MATL) / NOT LIKE DAD (NLD) / PHOSPHOLIPASE-A (PLA)*  
42 gene (**Figure 1B**) [1–5]. Few additional haploid inducer lines were reported in sorghum and potato for  
43 example, but the molecular bases remain unknown [1].  
44

### 45 **A long-awaited discovery: efficient CENH3-based haploid induction in crops**

46 The CENH3-based haploid induction system was born in 2010 thanks to a discovery using the  
47 model plant *Arabidopsis thaliana* [6]. Although this pioneering work has held great potential for plant  
48 breeders, efforts to translate this technology to crops have been hindered by the low haploid induction  
49 rate [1]. However, by engineering CENH3 genes in both maize and wheat crops, two recent studies  
50 reported paternal haploid induction with a rather high haploid induction rate (~5-8%) (**Figure 1A**) [7,8].  
51 Wang *et al.* [7] initially failed to translate *Arabidopsis* CENH3-based haploid induction system to maize,  
52 but then surprisingly identified a new and simple way to induce haploid embryos. Crossing a female  
53 maize line heterozygous for the *cenh3* null mutation (*cenh3/WT*) with wild-type pollen led to ~5% of  
54 seeds with haploid embryos [7]. In wheat the situation is more complex since there are two *CENH3*  
55 genes ( $\alpha$  and  $\beta$ ), and because bread wheat is a hexaploid crop with three sub-genomes (named A, B  
56 and D), making a total of 12 *CENH3* alleles. Thanks to huge efforts in wheat transformation (more than  
57 1150 events screened!) and rigorous analyses of gene editing events, Lv *et al.* were able to isolate a  
58 specific mutagenic scenario that led to haploid embryo induction [8]. They found that a specific  
59 mutation type on *CENH3 $\alpha$ -A*, called restored frameshift (RFS), coupled with knockout alleles for both  
60 *CENH3 $\alpha$ -B* and *CENH3 $\alpha$ -D* triggered haploid induction (the 6 *CENH3 $\beta$*  alleles were left at wild-type  
61 state). The RFS consists of a wild-type protein sequence with the replacement of few amino acids from  
62 the N-terminal domain by unrelated ones [8]. They demonstrated the RFS on *CENH3 $\alpha$ -A* gene is critical  
63 for haploid induction but, unexpectedly, the haploid inducer lines heterozygous for *CENH3 $\alpha$ -A* RFS  
64 (RFS/WT) outperformed the homozygous ones (RFS/RFS) [8]. To sum-up, good paternal haploid  
65 induction systems based on CENH3 engineering were obtained in both maize and wheat (**Figure 1A**).  
66 Of note, maternal haploid embryos were reported at a low rate (0.5%) using the maize CENH3-based  
67 induction system [7], although the maternal origin was not double-checked.  
68

### 69 **What can we learn from maize and wheat CENH3-based haploid inducer lines?**

70 The novelty brought by these two studies is the importance of heterozygosity to induce  
71 haploid embryos, indicating that one copy of the CENH3 wild-type allele is required to obtain the  
72 highest haploid induction rates [7,8]. However, the wheat situation differs from what has been  
73 reported in maize, since the *CENH3 $\alpha$ -A* allele in a “knock-out/wild-type” configuration does not induce  
74 haploid embryos, but needs to be heterozygous in a “RFS/wild-type” state [8]. This contradictory  
75 situation could lie in the fact that the “RFS/wild-type” wheat haploid inducer line still has three copies  
76 of *CENH3 $\beta$*  gene in a wild-type state, knowing that wheat has a complex centromere organization with  
77 the two versions of CENH3 ( $\alpha$  and  $\beta$ ) localizing in complementary centromeric domains.

78 Interestingly, Wang *et al.* [8] took advantage of the relative large size of maize kernels to  
79 genotype endosperm from seeds with haploid embryos. They were able to deduce that haploid

80 induction occurred when the embryo sacs (and thus egg cells) carried the *cenh3* null mutation [8]. The  
81 proposed working model is that these *cenh3* null embryo sacs have a diluted quantity of the wild-type  
82 CENH3 protein leading to defective centromeres. Indeed, just before meiosis, the diploid megaspore  
83 mother cell is heterozygous in a *cenh3* null/WT state. After meiosis, half of the resulting megaspores  
84 have the *cenh3* null allele in their haploid genome, but should have nevertheless inherited wild-type  
85 CENH3 protein in their centrosomes. This initial CENH3 protein pool is then diluted by the three post-  
86 meiotic cell divisions, creating egg cells with defective centromeres. Thus, when these “CENH3 diluted”  
87 egg cells are fertilized by wild-type sperm cells, it could lead to postzygotic maternal genome  
88 incompatibility that generates paternal haploid embryos [7]. A similar scenario could be envisioned in  
89 the wheat CENH3 engineered system, but with RFS allele instead of *cenh3* null allele. Nevertheless, the  
90 mode of action of wheat RFS alleles remains to be investigated. Since the RFS alters putative histone  
91 post-translational modification sites, it can be speculated that these CENH3 modifications have an  
92 important role in centromere stability. Experimental testing of RFS mutation on diploid plants with a  
93 single-copy *CENH3* gene should also help to clarify the situation.

94

### 95 **Maize and wheat now have many ways to induce haploid embryos and multiple breeding** 96 **applications**

97 The discovery of efficient CENH3-based paternal haploid inducer lines places maize and wheat  
98 in a unique situation because different methods are now available to induce both maternal and  
99 paternal haploid embryos. The two historical wheat haploidization methods are: interspecific crosses,  
100 and *in vitro* culture of male haploid cells, the latter method being also reported for maize [1].  
101 Furthermore, wheat, together with rice, have recently benefited from successful translation of maize  
102 *in planta* haploid induction system [9,10] (**Figure 1B**). Whereas all these methods could be used to  
103 speed-up the production of fixed material (DH lines), the CENH3-based haploid induction system offers  
104 the possibility to obtain an unusual cellular arrangement in which the paternal genome is surrounded  
105 by maternal cytoplasm. This allows cytoplasm swapping in a single-step, simplifying hybrid crop  
106 management [1].

107 Finally, haploid inducer lines have recently been elegantly used to deliver genome edit  
108 machinery to crop genotypes that are recalcitrant to genetic transformation [11,12]. This *trans* editing  
109 system allows the production of genome edited elite cultivars that are both transgene-free plants  
110 (because CRISPR-cassette is unstable like the haploid inducer genome) and homozygous (because they  
111 are DH plants).

112 To sum-up, *in planta* haploid induction systems allow not only convenient production of  
113 haploid embryos within viable seeds, but represent the entry point for increasing numbers of breeding  
114 applications, not forgetting that haploid inducer lines also represent entry points to better characterize  
115 plant reproduction.

116

117

### 118 **Figure 1. Two different kind of haploid inducer lines to trigger either paternal (A) or maternal (B)** 119 **haploid embryos through seeds in wheat.**

120 (A) Engineering of CENH3 allows creation of haploid inducer lines which are able to generate viable  
121 seeds containing embryos with a paternal haploid genome. These CENH3-based haploid inducer lines  
122 trigger paternal haploid embryos when used as a female parent in outcrossing with wild-type pollen.

123 (B) Translation of maize haploid induction system to wheat allows induction of viable seeds containing  
124 embryos with a maternal haploid genome. In this situation, the haploid inducer line is used as a male  
125 and outcrosses with wild-type female. In all cases, the haploid embryos lack the genome from the  
126 haploid inducer line. The main breeding advantages of these *in planta* inducer lines are (i) shortening  
127 the creation of inbred lines, (ii) shortening the creation cytoplasm swapping (for CENH3-based haploid  
128 inducer lines only) and (iii) allowing *trans* editing of non-inducer genome. This figure was created using  
129 BioRender (<https://biorender.com/>).

130

## 131 Acknowledgments

132 I thank Douglas Pyott for helpful comments on the manuscript and I apologize to all authors whose  
133 important work we could not be discussed due to length limitations. T.W was supported by INRAE, by  
134 the ANR grant (ANR-19-CE20-0012, “Not-Like-Dad”), and by the “pack ambition recherche” from the  
135 Région Auvergne-Rhone-Alpes (“DH-INNOV”).

136


137

## 138 References

- 139 1 Jacquier, N.M.A. *et al.* (2020) Puzzling out plant reproduction by haploid induction for innovations in plant breeding.  
140 *Nature Plants* DOI: 10.1038/s41477-020-0664-9
- 141 2 Gilles, L.M. *et al.* (2017) Haploid induction in plants. *Current Biology* 27, R1095–R1097
- 142 3 Kelliher, T. *et al.* (2017) MATRILINEAL, a sperm-specific phospholipase, triggers maize haploid induction. *Nature* 542,  
143 105–109
- 144 4 Gilles, L.M. *et al.* (2017) Loss of pollen-specific phospholipase NOT LIKE DAD triggers gynogenesis in maize. *The EMBO*  
145 *Journal* 36, 707–717
- 146 5 Liu, C. *et al.* (2017) A 4-bp Insertion at ZmPLA1 Encoding a Putative Phospholipase A Generates Haploid Induction in  
147 Maize. *Molecular Plant* 10, 520–522
- 148 6 Ravi, M. and Chan, S.W.L. (2010) Haploid plants produced by centromere-mediated genome elimination. *Nature* 464,  
149 615–618
- 150 7 Wang, N. *et al.* (2021) Haploid induction by a maize cenH3 null mutant. *Science Advances* 7, eabe2299  
151 DOI: <https://doi.org/10.1126/sciadv.abe2299>
- 152 8 Lv, J. *et al.* (2020) Generation of paternal haploids in wheat by genome editing of the centromeric histone CENH3.  
153 *Nature Biotechnology* 38, 1397–1401
- 154 9 Liu, H. *et al.* (2020) Efficient induction of haploid plants in wheat by editing of TaMTL using an optimized Agrobacterium-  
155 mediated CRISPR system. *J Exp Bot* 71, 1337–1349
- 156 10 Yao, L. *et al.* (2018) OsMATL mutation induces haploid seed formation in indica rice. *Nature Plants* 4, 530–533
- 157 11 Kelliher, T. *et al.* (2019) One-step genome editing of elite crop germplasm during haploid induction. *Nature*  
158 *Biotechnology* 37, 287–292
- 159 12 Wang, B. *et al.* (2019) Development of a Haploid-Inducer Mediated Genome Editing System for Accelerating Maize  
160 Breeding. *Molecular Plant* 12, 597–602

161


### (A) Haploid embryo induction by CENH3 engineering



#### Breeding applications

- One-step cytoplasm swapping
- Shortening inbred line creation
- *trans* editing

### (B) Haploid embryo induction by translation of maize MATL/NLD/PLA1 system



#### Breeding applications

- Shortening inbred line creation
- *trans* editing