



HAL
open science

Precise detection of antibiotic resistance genes in chicken microbiota

Anne-Carmen Sanchez, Guillaume Kon Kam King, Sylvie Baucheron, Fanny Calenge, H el ene Chiapello, Pierre P. Nicolas, S ebastien Leclercq, Anne-Laure Abraham

► To cite this version:

Anne-Carmen Sanchez, Guillaume Kon Kam King, Sylvie Baucheron, Fanny Calenge, H el ene Chiapello, et al.. Precise detection of antibiotic resistance genes in chicken microbiota. Journ ees Ouvertes en Biologie, Informatique et Math ematiques - JOBIM - colloque virtuel, Jul 2021, Paris, France. . hal-03337234

HAL Id: hal-03337234

<https://hal.inrae.fr/hal-03337234>

Submitted on 7 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



Distributed under a Creative Commons Attribution 4.0 International License

Precise detection of antibiotic resistance genes in chicken microbiota

Anne-Carmen SANCHEZ¹, Guillaume KON KAM KING¹, Sylvie BAUCHERON²,
Fanny CALENGE³, H el ene CHIAPELLO¹, Pierre NICOLAS¹,
S ebastien LECLERCQ² and Anne-Laure ABRAHAM¹

¹ Universit e Paris-Saclay, INRAE, MaIAGE, 78350, Jouy-en-Josas, France

² Unit e Infectiologie et Sant e publique, INRAE Val de Loire, 37380 Nouzilly, France

³ Universit e Paris-Saclay, INRAE, AgroParisTech, GABI, Jouy-en-Josas, France

Corresponding Author: anne-carmen.sanchez@inrae.fr

Antibiotic resistance genes (ARGs) presence is largely documented in gut microbiomes of farm animals and represent a major risk for animal and human health but the routes and mechanisms of their dissemination are yet only partially understood. At the epidemiological level, it would be interesting to better understand the respective roles of contamination from the environment and transmission between generations of animals. These transmission events, at the genetic level, may be mediated by horizontal transfer of ARGs and clonal propagation of ARG harbouring bacterial lineages. To study these questions, an experiment was conducted aiming at following ARGs by shotgun metagenomics across 3 generations of broiler chicken in two separate buildings.

The study of ARG in metagenomic samples is challenging, because of sequencing errors, short read length, and variable abundance of genes, which suggest that specific methods may be needed to properly identify ARG flows. Moreover, preliminary results have shown that several haplotypes can co-exist for each gene, and that these haplotypes may be shared or not between samples. Analyzing this micro-evolutionary diversity is the key idea of our work. The identification of haplotypes in metagenomic data is a challenge that has been recently addressed with methods to identify strains relying on SNP frequencies in selected core genome marker genes (such as ConStrains[1], or DESMAN[2]), or with methods using reads as a phasing information to resolve haplotypes on individual genes (Hansel & Gretel [3]).

In order to accurately detect ARG flows between samples, we are working on a new workflow based on aligning metagenomic reads on a non redundant database of reference ARG sequences (Resfinder4.1 [4] clustered at 95% of identity) with BWA-MEM, and stringent quality filters are applied on reads and alignments using samtools. Nucleotide frequencies at each position of these reference sequences are then computed and statistics (number of polymorphic positions, number of alleles...) are recorded in order to quantify the diversity in ARGs within samples and the distance between samples. Our first results indicate a huge variability between genes in terms of number of polymorphic positions and number of haplotypes. Future steps will consist in identifying shared ARG haplotypes between samples by using approaches derived from those used for strain resolution to resolve the different haplotypes and determine their abundances in each sample, or by trying to infer ARG fluxes by comparing samples based on diversity and distance statistics only, without haplotype resolution.

References

- [1] Chengwei Luo, Rob Knight, Heli Siljander, Mikael Knip, Ramnik J Xavier, Dirk Gevers, ConStrains identifies microbial strains in metagenomic datasets. *Nat Biotechnology*, 33:1045–52, 2015.
- [2] Christopher Quince, Tom O. Delmont, S ebastien Raguideau, Johannes Alneberg, Aaron E. Darling, Gavin Collins & A. Murat Eren, DESMAN: a new tool for de novo extraction of strains from metagenomes. *Genome Biology*, 18, 181, 2017.
- [3] Samuel M. Nicholls, Wayne Aubrey, Arwyn Edwards, Kurt de Grave, Sharon Huws, Leander Schietgat, Andr e Soares, Christopher J. Creevey, Amanda Clare, Recovery of gene haplotypes from a metagenome. *BioRxiv*, 223404, 2019.
- [4] Valeria Bortolaia, Rolf S. Kaas, Etienne Ruppe, Marilyn C. Roberts, Stefan Schwarz, Vincent Cattoir, Alain Philippon, Rosa L. Allesoe, Ana Rita Rebelo, Alfred Ferrer Florensa, Linda Fagelhauer, Trinad Chakraborty, Bernd Neumann, Guido Werner, Jennifer K. Bender, Kerstin Stingl, Minh Nguyen, Jasmine Coppens, Basil Britto Xavier, Surbhi Malhotra-Kumar, Henrik Westh, Mette Pinholt, Muna F. Anjum, et al, ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy*, 75(12), 2020.