

Special Issue "Microorganisms in Recycling and Valorization of Organic Waste for Sustainable Soil Health and Management"

José A Siles, Mercedes García-Sánchez, María Gómez-Brandón

▶ To cite this version:

José A Siles, Mercedes García-Sánchez, María Gómez-Brandón. Special Issue "Microorganisms in Recycling and Valorization of Organic Waste for Sustainable Soil Health and Management". Microorganisms, 2021, 9 (8), pp.1682. 10.3390/microorganisms9081682 . hal-03338900

HAL Id: hal-03338900 https://hal.inrae.fr/hal-03338900

Submitted on 9 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License





Editorial Special Issue "Microorganisms in Recycling and Valorization of Organic Waste for Sustainable Soil Health and Management"

José A. Siles ^{1,2}, Mercedes García-Sánchez ³ and María Gómez-Brandón ^{4,*}

- ¹ Department of Plant and Microbial Biology, University of California at Berkeley, Berkeley, CA 94720, USA; josesimartos@gmail.com
- ² Department of Soil and Water Conservation and Waste Management, CEBAS-CSIC, 30100 Murcia, Spain

³ Eco & Sols, INRAE, Institut Agro, Université de Montpellier, CIRAD, IRD, CEDEX 02,

- 34060 Montpellier, France; garcia.sanchez.mercedes@gmail.com
- Grupo de Ecoloxía Animal (GEA), Universidad de Vigo, 36310 Vigo, Spain
- * Correspondence: mariagomez@uvigo.es or mariagomezbrandon@gmail.com

Anthropogenic activity generates huge amounts of solid organic wastes. Their incorrect handling may result in soil, water and air contamination. Therefore, finding strategies to reuse and treat organic wastes is needed. Their use as soil organic amendments has been proposed as an effective way of improving soil quality and fertility, while at the same time, protecting the environment. However, if not properly disposed of and treated, the use of raw wastes as soil inputs may lead to harmful environmental effects. In this context, the adoption of biological approaches, such as composting, vermicomposting or methanogenic processes offers the possibility of stabilizing the organic wastes before their use as soil amendments, achieving a dual purpose: environmental protection and production of amendments with improved characteristics. The role of microorganisms in these processes is crucial since they mediate the conversion of organic matter through a variety of chemical reactions. Thus, their study is fundamental, as is that of microbial communities in organically amended soils due to their usefulness as bio-indicators of soil quality. Altogether, it will help to reach integrative assessments on the suitability of waste transformation processes and promote sustainable agricultural practices.

The objective of this Special Issue on "Microorganisms in Recycling and Valorization of Organic Waste for Sustainable Soil Health and Management" was to provide a platform to researchers for sharing their new studies on the characterization of microorganisms mediating the transformation of organic wastes into appropriate soil amendments and on the impact of the application of stabilized wastes on soil microbial communities, nutrient cycling and plant yields. The present Special Issue comprises 7 research articles and 2 reviews, contributed by 47 authors.

Three articles were focused on the study of microbial communities during processes of waste transformation. Santos et al. [1] provided evidence about the efficient use of industrial alkaline wastes including CaO as chemical conditioners to sanitize sewage sludge by drastically reducing its pathogenic load. Thermal treatment assays were also tested as feasible alternatives to reach this goal. Jeong et al. [2] evaluated the effectiveness of the composting process on the inactivation of avian influenza virus (H9N2) and the encephalomyocarditis virus (EMCV) present in poultry and pig manures, respectively. The vitality of H9N2 and EMCV significantly declined as a result of composting. However, at the completion of the composting process after 168 h, inactivation effect appeared to be more sensitive in H9N2 than in EMCV, since it occurred within 1 h and 24 h of the start of the composting processes in H9N2 and EMCV, respectively, being more effective under mechanical aeration conditions. Nowadays, high-throughput sequencing technologies have become an indispensable tool to deeply characterize microbiomes from diverse environmental compartments. Using this technology, Gómez-Brandón et al. [3] revealed that the bacterial communities during the vermicomposting of grape marc over a period of



Citation: Siles, J.A.; García-Sánchez, M.; Gómez-Brandón, M. Special Issue "Microorganisms in Recycling and Valorization of Organic Waste for Sustainable Soil Health and Management". *Microorganisms* **2021**, *9*, 1682. https://doi.org/10.3390/ microorganisms9081682

Received: 2 July 2021 Accepted: 27 July 2021 Published: 8 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). 91 days were dominated by *Proteobacteria* and *Bacteriodetes*. They also found that the higher taxonomical diversity and richness were achieved after 14 days followed by significant increases in functional diversity of bacterial communities. Their findings evidenced that the valorization of distilled grape marc via vermicomposting led to a rich end product with higher bacterial diversity and functional attributes, which may be used as suitable plant growth promoter and/or soil organic amendment.

Six articles dealt with the characterization of the soil microbial responses to organic amendment or to contrasting agricultural practices. In this way, Neher et al. [4] assessed the impact of poultry litter compost (PLC) and dairy manure compost (DMC) on microbial communities of soils planted with spinach in the long term. Using metabarcoding, they evidenced that bacteria in the phylum Bacteriodetes, and classes Flavobacteriia and Spingobacteriales, were more abundant in soils amended with PLC than in those amended with DMC or in the unamended ones. However, fungi did not differ among treatments. Tang et al. [5] tested the performance of several plant growth-promoting rhizobacteria (PGPR), belonging to the genera Paraburkholderia, Burkholderia and Serratia, on maize growth, soil nutrient contents and nutrient use efficiency in soils subjected to different regimes of inorganic fertilization and organic amendment. They concluded that the amount of fertilizer applied into soil can be reduced through the combined use of fertilization and PGPR. On the other hand, Kim et al. [6], using 16S rRNA gene metabarcoding, aimed at identifying microbial groups involved in important soil processes as bio-indicators of improved soil health under different cover cropping and tillage practices. Bacteria were found to be more suitable than fungi and archaea for this purpose. Hence, the bacterial genera *Cellulomonas*, Solirubacter, Alererythrobacter and Massilia were found to be potential bio-indicators of enhanced aerobic decomposition, the family Nitrosomonadaceae of ammonia oxidation, and the bacterial genera Nitrospira and Reyranella of nitrite oxidation and nitrate reduction, respectively. Bai et al. [7] compared the effects of long-term and non-long-term fertilization on root-associated soils of walnut trees. This study pointed out that bacterial and fungal communities were negatively affected by long-term fertilization treatments as the result of an over-accumulation of ammonium-nitrogen (NH_4^+) and available phosphorus, which led to soil acidification. Meanwhile, soils non-subjected to long-term chemical fertilization resulted in the improvement of microbial communities involved in nutrient mobilization and plant growth.

This Special Issue includes two review articles. That of Lazcano et al. [8] provided a comprehensive overview on the use of organic fertilizers (OFs) for preserving soil health and sustainability. A special focus was given to the effects of OFs, including raw manure and processed materials, such as digestate, compost, vermicompost and biochar on the community structure and function of nitrifying and denitrifying soil microorganisms. Nitrification and denitrification are among the most important soil N transformations within the soil-plant interface, but there is still a lack of information about the underlying mechanisms related to the influence of OFs on these microbial processes and the consequences on nitrous oxide production. Putting the accent on this matter and with the aid of multi-omics approaches, it will be fundamental for the prediction of N2O emissions under different scenarios of land use. Finally, Siles et al. [9] provided a thorough review about the use of co-occurrence network approaches as a proxy to gain insights into the interactions among the members of a microbial community and the identification of its keystone members during the processes of composting and anaerobic digestion and following the application of the respective end products (compost and digestate) into soil. Depending on the degree of these interactions and the species involved, the dynamics of these processes may vary with further consequences on the quality of the end product and its potential usefulness as an organic amendment.

Author Contributions: The three guest editors contributed equally to this Special Issue and all the aspects of this editorial. The three authors have read and agreed to the published version of the editorial. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the "Xunta de Galicia" (grant number ED431F 2018/05).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: J.A.S. thanks the support of the program "Juan de la Cierva Incorporación" of the "Ministerio de Ciencia, Innovación y Universidades" (agreement no. IJC2018-034997-I). M.G.-B. acknowledges support by the Programa Ramón y Cajal (RYC-2016-21231; Ministerio de Economía y Competitividad). We would like to thank all the authors who contributed their papers to this Special Issue and the reviewers for their invaluable revisions. We are also grateful to all the members of the Editorial Office of Microorganisms, and specially to Josephine Xue, for giving us the opportunity of editing this Special Issue and their continuous support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Santos, A.F.; Santos, C.P.; Matos, A.M.; Cardoso, O.; Quina, M.J. Effect of Thermal Drying and Chemical Treatments with Wastes on Microbiological Contamination Indicators in Sewage Sludge. *Microorganisms* **2020**, *8*, 376. [CrossRef] [PubMed]
- Jeong, K.-H.; Lee, D.-J.; Lee, D.-H.; Ravindran, B.; Chang, S.W.; Mupambwa, H.A.; Lee, M.-G.; Ahn, H.-K. Composting Reduces the Vitality of H9N2 in Poultry Manure and EMCV in Pig Manure Allowing for an Environmentally Friendly Use of These Animal Wastes: A Preliminary Study. *Microorganisms* 2020, *8*, 829. [CrossRef] [PubMed]
- 3. Gómez-Brandón, M.; Aira, M.; Santana, N.; Pérez-Losada, M.; Domínguez, J. Temporal Dynamics of Bacterial Communities in a Pilot-Scale Vermireactor Fed with Distilled Grape Marc. *Microorganisms* **2020**, *8*, 642. [CrossRef] [PubMed]
- Neher, D.A.; Limoges, M.A.; Weicht, T.R.; Sharma, M.; Millner, P.D.; Donnelly, C. Bacterial Community Dynamics Distinguish Poultry Compost from Dairy Compost and Non-Amended Soils Planted with Spinach. *Microorganisms* 2020, *8*, 1601. [CrossRef] [PubMed]
- Tang, A.; Haruna, A.O.; Majid, N.M.A.; Jalloh, M.B. Effects of Selected Functional Bacteria on Maize Growth and Nutrient Use Efficiency. *Microorganisms* 2020, *8*, 854. [CrossRef] [PubMed]
- 6. Kim, N.; Zabaloy, M.C.; Riggins, C.W.; Rodríguez-Zas, S.; Villamil, M.B. Microbial Shifts Following Five Years of Cover Cropping and Tillage Practices in Fertile Agroecosystems. *Microorganisms* **2020**, *8*, 1773. [CrossRef] [PubMed]
- Bai, Y.-C.; Chang, Y.-Y.; Hussain, M.; Lu, B.; Zhang, J.-P.; Song, X.-B.; Lei, X.-S.; Pei, D. Soil Chemical and Microbiological Properties Are Changed by Long-Term Chemical Fertilizers That Limit Ecosystem Functioning. *Microorganisms* 2020, *8*, 694. [CrossRef] [PubMed]
- 8. Lazcano, C.; Zhu-Barker, X.; Decock, C. Effects of Organic Fertilizers on the Soil Microorganisms Responsible for N2O Emissions: A Review. *Microorganisms* **2021**, *9*, 983. [CrossRef] [PubMed]
- Siles, J.A.; García-Sánchez, M.; Gómez-Brandón, M. Studying Microbial Communities through Co-Occurrence Network Analyses during Processes of Waste Treatment and in Organically Amended Soils: A Review. *Microorganisms* 2021, 9, 1165. [CrossRef] [PubMed]