

Genetic parameters of maritime pine radial growth during a specific drought event

Laurent Bouffier, Raphael Segura, François Ehrenmann, Jean-Marc Gion

► **To cite this version:**

Laurent Bouffier, Raphael Segura, François Ehrenmann, Jean-Marc Gion. Genetic parameters of maritime pine radial growth during a specific drought event. Forest Genetics 2017 - Forest Health and Productivity in Changing Environments, Jun 2017, Edmonton, Canada. hal-03341365

HAL Id: hal-03341365

<https://hal.inrae.fr/hal-03341365>

Submitted on 10 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conference Proceedings



Forest Genetics 2017

Forest Health and Productivity in Changing Environments

June 26-29, 2017 in Edmonton, Alberta

A joint meeting of WFGA & CFGA

Conference Sponsors



Genetic parameters of maritime pine radial growth during a specific drought event

Laurent Bouffier^{1,2,*}, *Raphaël Segura*^{1,2}, *François Ehrenmann*^{1,2}, *Jean-Marc Gion*³

¹) INRA, BIOGECO, UMR 1202, 33610 Cestas, France

²) University Bordeaux, BIOGECO, UMR 1202, 33400 Talence, France

³) CIRAD, UMR AGAP, 69 Route Arcachon, Cestas, France

Abstract: Main selection criteria of French maritime pine (*Pinus pinaster*) breeding program are growth at 12 years old and stem straightness. Scenarios of climate changes forecast an increase in frequency and intensity of droughts in southwestern France where maritime pine plantations are currently established. In that context, a key issue for breeding is thus to be able to evaluate growth in the future climatic conditions characterized by low precipitations during the growth period.

A maritime pine genetic trial was established 19 years ago on two contrasting sites (humid and dry sites) with half-sib families. 150 high resolution dendrometers have been installed on 25 families selected to represent a large range of growth performance (based on growth data collected at 12 years old). Radial growth was measured every hour during two successive years. The two years considered (2015 and 2016) were characterized by contrasting climate (dry spring in 2015 and wet spring in 2016). Soil humidity and climatic data were also recorded on this trial during the same period.

Genetic analyses (estimation of genetic parameters and GxE interactions) were carried out on radial growth data related to short periods of time (few weeks) selected either for low or for high precipitation level. This methodology allows to estimate the genetic variability for growth in dry versus wet conditions and to identify the genotypes best adapted to the future climate.

*) Corresponding author's email: laurent.bouffier@inra.fr