



**HAL**  
open science

## Home-field advantage of litter decomposition differs between leaves and fine roots

Dunmei Lin, Pengpeng Dou, Guangrong Yang, Shenhua Qian, Hongjuan Wang, Liang Zhao, Yongchuan Yang, Xiangcheng Mi, Keping Ma, Nicolas Fanin

► **To cite this version:**

Dunmei Lin, Pengpeng Dou, Guangrong Yang, Shenhua Qian, Hongjuan Wang, et al.. Home-field advantage of litter decomposition differs between leaves and fine roots. *New Phytologist*, 2020, 227 (4), pp.995-1000. 10.1111/nph.16517 . hal-03345983

**HAL Id: hal-03345983**

**<https://hal.inrae.fr/hal-03345983>**

Submitted on 16 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Letters

# Home-field advantage of litter decomposition differs between leaves and fine roots

### Introduction

Litter decomposition is a fundamental process influencing carbon dynamics and nutrient cycling in terrestrial ecosystems. In the 20<sup>th</sup> century, it was commonly accepted that climate, litter quality and soil biota hierarchically influenced the decomposition process from the large to the local scale (Swift *et al.*, 1979; Couëteux *et al.*, 1995). However, recent studies have suggested that decomposer communities are much more important in driving litter decomposition than previously thought, mainly because aggregate data at large spatial scales are falsely assumed to represent causative relationships at small spatial scales (Bradford *et al.*, 2014, 2017). In particular, microbial functioning may depend on complex interactions between microbial decomposers and their substrates such as plant litter (Jackrel *et al.*, 2019; Lin *et al.*, 2019). The home-field advantage (HFA) hypothesis predicts that decomposers are more efficient with the plant litters they most frequently encounter (Gholz *et al.*, 2000; Ayres *et al.*, 2009b; Austin *et al.*, 2014; Palozzi & Lindo, 2018). This hypothesis has been empirically tested in various ecosystems via reciprocal litter transplant experiments (Ayres *et al.*, 2009a; Milcu & Manning, 2011; Fanin *et al.*, 2016; Lu *et al.*, 2017; Palozzi & Lindo, 2017; Veen *et al.*, 2018; Jackrel *et al.*, 2019; Lin *et al.*, 2019), and HFA was shown to increase decomposition rates by 7.5% on average (Veen *et al.*, 2015a). However, the vast majority of studies demonstrating HFA effects used leaf litter, and whether the results are consistent with root litter has seldom been tested.

Fine roots account for a substantial amount of litter inputs in terrestrial ecosystems, and play a major role in carbon and nutrient cycling through their afterlife ‘effects’ (Jackson *et al.*, 1997, 2017; Freschet *et al.*, 2013). Because HFA may arise due to the specialization of microbial communities towards their substrates, especially when the substrate is recalcitrant (Milcu & Manning, 2011), root litter should generate important HFA effects because of the necessity to synthesize specific oxidative enzymes to breakdown complex organic compounds. Yet, the very limited empirical experiments evaluating HFA effects using roots have reported positive, neutral or negative effects (Freschet *et al.*, 2012a; Jacobs *et al.*, 2018; Minerovic *et al.*, 2018), perhaps as a consequence of an important variability in litter chemistry between ‘home’ and ‘away’ sites (Veen *et al.*, 2015a). Hence, it is difficult to generalize the specialization of microbial communities towards root litter

decomposition. Furthermore, whether HFA effects are coherent between different plant organs originating from the same species (e.g. leaves and roots), has never been evaluated in the same experiment. Because leaves and roots of the same species commonly differ substantially in their chemical compositions (Freschet *et al.*, 2012b; Ma *et al.*, 2016; Sun *et al.*, 2018), different substrate qualities may stimulate contrasting microbial communities characterized by varying metabolic requirements and catabolic capacities (Sauvadet *et al.*, 2019). Therefore, we hypothesize that the concordance in HFA effects between aboveground and belowground plant parts depends on the (dis)similarity in litter chemistry between leaves and roots, that is, plant species having a greater similarity in their chemical traits between leaves and roots should present relatively similar HFA effects (hypothesis 1). Furthermore, because litter recalcitrance is thought to be one of the main drivers controlling HFA due to a higher degree of specialization of microbial communities (Milcu & Manning, 2011), we hypothesize that root litters should display greater HFA effects than leaf litters because they are often more recalcitrant (hypothesis 2).

### Materials and Methods

We performed an 1132-d reciprocal litter transplant experiment among three forest sites using litters collected from the dominant tree species in each site: broadleaf (*Castanopsis eyrei*), coniferous (*Cunninghamia lanceolata*) and bamboo (*Phyllostachys heterocycla* cv *Pubescens*) (see Supporting Information Methods S1 for more details of the study sites). Freshly fallen leaf litters were collected using litter traps to prevent infection by soil biota of the ‘home’ site. Roots were extracted from the soil and washed on a sieve under running water. Fresh fine roots ( $\Phi < 2$  mm) were picked out and sterilized using chloroform vapour. Litter quality was characterized by 14 chemical traits related to element concentration, stoichiometry and carbon quality (Table S1; details about trait measurements are in Methods S1). In each site, six randomized blocks, with a distance of at least 10 m between each block, were selected for litterbag incubation. Each litterbag (15 cm × 10 cm with a mesh size of 25- $\mu$ m, which allows microbial decomposers to enter the litterbags), was filled with *c.* 3.0 g litters, and labelled with a plastic tag. Although soil fauna may also significantly affect the magnitude of the HFA effects in some ecosystems (Milcu & Manning, 2011), previous results in our study sites showed that soil fauna did not substantially contribute to driving the HFA (Lin *et al.*, 2019). In total, we placed six litterbags per litter type in each block for six successive harvests during the first year (94 and 222 d), second year (392 and 583 d) and third year (827 and 1132 d) after the start of the incubation, resulting in a total of 648 litterbags used in this study (three sites × six blocks × six litter types × six harvest times). Leaf litterbags were anchored on the forest floor and root litterbags were buried in the soil. Following harvest, the litter samples were

cleaned under tap water, oven-dried at 60°C for 48 h and weighed. Litter mass loss (%) was calculated as  $(M_i - M_f)/M_i \times 100$ , where  $M_i$  and  $M_f$  are initial and final litter dry mass, respectively.


We ran nonmetric multidimensional scaling (NMDS) on the 14 chemical litter traits to visualize the variation in litter quality among the different litter types, and then performed a permutational multivariate analysis of variance (PERMANOVA) to test the significant effects of species, organ type and their interaction on litter chemical traits. We also calculated dissimilarity among the different litter types using the Euclidean distance. The effects of litter species, incubation site, harvest time and their interactions on mass loss of leaves or fine roots, respectively, were analysed using linear mixed effects models with a first-order auto-regressive variance structure to account for temporal pseudo-replication. To improve the reliability of our statistical inference, we selected the most parsimonious models based on the lowest AICc (Akaike's information criterion corrected for small sample sizes). In addition, the HFA index was estimated at each harvest time by running the regression model proposed by Keiser *et al.* (2014) (see Methods S1 for detailed description of the statistical analyses).

## Results and Discussion


The NMDS ordination separated relatively well the six litter types along the two first NMDS axes (Fig. 1), which were associated to chemical traits known to directly affect litter decomposition rates such as nutrients (i.e. nitrogen (N), phosphorus (P), potassium (K), manganese (Mn)), carbon forms (i.e. total phenols, lignin, tannins) and stoichiometric ratios (N : P, C : P, lignin : N) (Swift *et al.*, 1979; Melillo *et al.*, 1982; Keiluweit *et al.*, 2015; Chomel *et al.*,

2016; Sun *et al.*, 2018). Litter chemistry differed significantly between different species ( $P < 0.001$ ; Table S2). Litter chemical traits also differed significantly between plant organs ( $P < 0.001$ ; Table S2), especially for the bamboo and coniferous species (Fig. 1). Specifically, bamboo litters were the most labile among the six litter types (Table S1), and both leaves and roots were generally located at the high quality end of the NMDS axes (Fig. 1). By contrast, broadleaf litters were more recalcitrant in comparison to bamboo litters and were generally located at the low quality end of the NMDS axes (Fig. 1), especially for the fine root litter (Table S1). In line with our expectations, bamboo and broadleaf litters generally exhibited higher and lower litter mass loss, respectively (Fig. 2). Coniferous was intermediate between bamboo and broadleaf litters. Overall, coniferous fine root litter was more recalcitrant than leaf litter, mainly because of higher concentrations in total phenols and tannins (Table S1), which may explain the lower mass loss of fine root litter compared with leaf litter over the course of the experiment (Fig. 2).

We found a significant tree species  $\times$  incubation site interaction effect on leaf litter mass loss ( $F_{4,40} = 13.23$ ,  $P < 0.001$ ; Table 1), suggesting that leaf litter decomposition depends on the environment in which it decomposes (Ayres *et al.*, 2009a; Milcu & Manning, 2011; Lu *et al.*, 2017; Veen *et al.*, 2018; Lin *et al.*, 2019). This is mainly the result of lower leaf litter mass loss of broadleaf species in its 'away' site, especially in the bamboo forest (Fig. 2). By contrast, there is no significant tree species  $\times$  incubation site interaction effect on fine root litter mass loss ( $F_{4,40} = 0.78$ ,  $P = 0.546$ ; Table 1). Accordingly, we found that HFA effects were generally inconsistent between leaves and fine roots over 1132 d of decomposition (Fig. 2). In addition, although the similarity in litter



**Fig. 1** Nonmetric multidimensional scaling (NMDS) ordination based on Bray–Curtis similarity matrices depicting litter chemical traits. Different organs were grouped by polygons (leaf in green; fine root in blue). Litter quality is reflected by the direction of the different nutrient components.



**Fig. 2** Litter mass loss and home-field advantage (HFA) index of the six harvest times. At each harvest time (a–f), the left panels show the litter mass loss (%) of different organs (leaves and fine roots) for the three species that were reciprocally decomposed in the three forest sites and the right panel shows the HFA index estimated using the equation developed by Keiser *et al.* (2014). Ba, Bamboo species; Co, coniferous species; Br, broadleaf species. In the left panels, boxes represent the central 50% of the data and the whiskers on the boxes represent the 95% quantiles ( $n = 6$ ). Within each panel, the different boxes from the left to the right represent the litter that was decomposed in the bamboo, coniferous and broadleaf sites, respectively. The grey closed boxes denote the litter that was decomposed in its 'home' site. Different lowercase letters indicate significant differences in litter mass loss among different incubation sites within each of the six litter types, while uppercase letters indicate significant differences among the six litter types (Tukey's HSD test,  $P < 0.05$ ). In the right panels, a positive bar value means a 'home-field advantage', that is, litter decomposes faster beneath the plant species it derived from than beneath other species. Error bars represent  $\pm$  SE ( $n = 6$ ), asterisks indicate estimates significantly different from zero with: \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ .

**Table 1** Results from the most parsimonious linear mixed effects model to test for the effects of species, incubation site, harvest time and their interactions on mass loss of leaf and fine root litter, respectively (see Supporting Information Table S4 for model selection procedure).

Variable	Leaf litter		Fine root litter	
	F-value	P-value	F-value	P-value
Species	$F_{2,40} = 36.24$	<b>&lt;0.001</b>	$F_{2,40} = 71.16$	<b>&lt;0.001</b>
Site	$F_{2,40} = 20.77$	<b>&lt;0.001</b>	$F_{2,40} = 1.05$	0.358
Time	$F_{1,269} = 2327.41$	<b>&lt;0.001</b>	$F_{1,269} = 2275.74$	<b>&lt;0.001</b>
Species × site	$F_{4,40} = 13.23$	<b>&lt;0.001</b>	$F_{4,40} = 0.78$	0.546
<i>Model performance</i>				
logLik	−1097.78		−1071.02	
AICc	2224.90		2171.40	
$\omega_i$	1.00		0.96	

Values in bold typeface represent significant effect with  $P < 0.05$ ;  $\omega_i$ , model weight.

chemistry between leaves and roots varied in the order broadleaf > bamboo > coniferous (Table S3), the broadleaf species showed the highest degree of difference in HFA effects between leaves and roots, that is leaf litter exhibited significant HFA effects throughout the 1132-d of decomposition, but fine root litter did not (Fig. 2). The only case for which both leaf and root litters displayed significant HFA effects was for the bamboo litter after 222-d of decomposition (Fig. 2). Yet, this species showed a relatively low degree of similarity in litter chemistry between leaves and roots (Table S3). Therefore, these results contradict our first hypothesis predicting that plant species having a greater similarity in their chemical traits between leaves and roots should present a greater similarity in HFA effects, and suggest that the interactions between microbes and their substrates depend on the plant organ considered.

Interestingly, we found that root litter rarely showed significant HFA effects compared with leaf litter (Fig. 2). Contrary to our second hypothesis, these findings highlight that HFA effects are not higher for low litter quality compared with relatively high quality litter (Veen *et al.*, 2015b; Palozzi & Lindo, 2017; Lin *et al.*, 2019). In line with these results, previous microcosm incubation and field studies have also reported that root litter decomposition did not exhibit significant HFA effects (Jacobs *et al.*, 2018; Minerovic *et al.*, 2018). One potential explanation could be due to the differences in microhabitats along the soil profile. Leaf litter decomposes on the forest floor that is covered by freshly fallen litter, which is relatively rich in nutrients and labile carbon compounds, whereas root litter decomposes within the soil, in which the organic matter is heavily decomposed and more recalcitrant (Adl, 2003). These different resource conditions may generate significant differences in the composition and activity of microbial communities along the litter–soil continuum (Sterkenburg *et al.*, 2018). In particular, the microbial communities from recalcitrant environments may have a wider functional capacity than those originating from richer environments, meaning that they more efficiently decompose root litters that vary widely in their chemical characteristics (functional breadth

hypothesis; Keiser *et al.*, 2011, 2014; Fanin *et al.*, 2016), thereby decreasing their specialization toward the litters they most frequently encounter. Alternatively, although we buried fine root litter in the soil close to the adult trees of targeted species, the soil environment is likely more heterogeneous than the forest floor because it includes fine root litters from many other coexisting tree species. Therefore, the adaptation of specialist decomposers to individual species might be less important in an environment in which the resources are highly diverse and mixed. Furthermore, we cannot exclude that the difference in microclimatic conditions along the soil profile, with notably higher water content in the soil, may also affect decomposer communities and reduce the differences in decomposition rates for root litter (Fanin *et al.*, 2019a). Additional studies simultaneously manipulating leaves and roots on the forest floor and in the soil will be necessary to disentangle the effects of microclimatic conditions from those of resource environment on HFA effects.

## Conclusions

By using a long-term reciprocal litter transplant experiment, we sought to test whether HFA patterns were consistent between leaves and fine roots. We found that HFA effects were generally inconsistent between plant organs, either across or within species, thereby underlining that leaf and root litters need to be considered separately when evaluating the adaptation of decomposers to their substrates. These results have several implications. First, they emphasize that HFA is not more pronounced for recalcitrant (i.e. roots) than labile litter (i.e. leaves) (Veen *et al.*, 2015b; Palozzi & Lindo, 2017), and they demonstrate that HFA may also occur when litters are relatively rich in nutrients and labile carbon compounds (Lin *et al.*, 2019). Second, the results highlight that there is often some decoupling between the aboveground and belowground subsystems in terrestrial ecosystems (Cameron *et al.*, 2019; Delgado-Baquerizo *et al.*, 2019; Fanin *et al.*, 2019b), and this finding begs the question of whether the position at which soil communities perform decomposition is important to understand the local adaptation of microbes to their substrates. Finally, our results over a thousand days of incubation demonstrate an important temporal variability in HFA effects (Ayres *et al.*, 2009a; Fanin *et al.*, 2016; Veen *et al.*, 2018), indicating that the succession in microbial communities may be also an important factor to predict the variability in plant–soil interactions over time. We conclude that plant–microbe interactions contribute significantly to nutrient and carbon cycling, but these interactions may strongly depend on the plant organ considered and the environmental conditions at the local scale.







## Acknowledgements

This research was funded by the National Natural Science Foundation of China (31500356). The authors thank Mei Pan, Fang Wang, Yan Liu, Pei Wang for the laboratory and field assistance. The authors thank Tania Maxwell for her help with English and useful remarks. Three anonymous reviewers provided helpful suggestions to improve the manuscript.

## Author contributions

DL designed the research. DL, PD, GY and HW performed the experiments with input from SQ, LZ, YY, XM and KM. DL carried out all of the statistical analyses with input from NF. DL and NF wrote the first draft of the manuscript. All authors contributed to manuscript completion and revision.

## ORCID

Nicolas Fanin  <https://orcid.org/0000-0003-4195-855X>  
 Dunmei Lin  <https://orcid.org/0000-0001-6112-7783>  
 Keping Ma  <https://orcid.org/0000-0001-9112-5340>  
 Shenhua Qian  <https://orcid.org/0000-0003-1565-8748>  
 Guangrong Yang  <https://orcid.org/0000-0002-5335-0501>  
 Yongchuan Yang  <https://orcid.org/0000-0001-7627-7776>

**Dunmei Lin<sup>1\*</sup>** , **Pengpeng Dou<sup>1</sup>** ,  
**Shenhua Qian<sup>1</sup>** , **Hongjuan Wang<sup>2</sup>**, **Liang Zhao<sup>1</sup>**,  
**Yongchuan Yang<sup>1</sup>** , **Xiangcheng Mi<sup>3</sup>**, **Keping Ma<sup>3</sup>**  and  
**Nicolas Fanin<sup>4</sup>** 

<sup>1</sup>Key Laboratory of the Three Gorges Reservoir Region's Eco-Environment, Ministry of Education, Chongqing University, Chongqing, 400045 China;

<sup>2</sup>Biotechnology Research Centre, Chongqing Academy of Agricultural Sciences, Chongqing, 401329 China;

<sup>3</sup>State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, The Chinese Academy of Sciences, Beijing, 100093 China;

<sup>4</sup>Interaction Soil Plant Atmosphere (ISPA), UMR 1391, INRAE – Bordeaux Sciences Agro, 71 Avenue Edouard Bourlaux, 33882 Villenave-d'Ornon cedex, France

(\*Author for correspondence: tel +86 023 65120750; email lindunmei@cqu.edu.cn)

## References

- Adl SM. 2003. *The ecology of soil decomposition*. Wallingford, UK: CABI Publishing.
- Austin AT, Vivanco L, Gonzalez-Arzac A, Perez LI. 2014. There's no place like home? An exploration of the mechanisms behind plant litter-decomposer affinity in terrestrial ecosystems. *New Phytologist* **204**: 307–314.
- Ayres E, Steltzer H, Berg S, Wall DH. 2009a. Soil biota accelerate decomposition in high-elevation forests by specializing in the breakdown of litter produced by the plant species above them. *Journal of Ecology* **97**: 901–912.
- Ayres E, Steltzer H, Simmons BL, Simpson RT, Steinweg JM, Wallenstein MD, Mellor N, Parton WJ, Moore JC, Wall DH. 2009b. Home-field advantage accelerates leaf litter decomposition in forests. *Soil Biology & Biochemistry* **41**: 606–610.
- Bradford MA, Veen GFC, Bonis A, Bradford EM, Classen AT, Cornelissen JHC, Crowther TW, De Long JR, Freschet GT, Kardol P *et al.* 2017. A test of the hierarchical model of litter decomposition. *Nature Ecology and Evolution* **1**: 1836–1845.
- Bradford MA, Warren IJ, Baldrian P, Crowther TW, Maynard DS, Oldfield EE, Wieder WR, Wood SA, King JR. 2014. Climate fails to predict wood decomposition at regional scales. *Nature Climate Change* **4**: 625.
- Cameron EK, Martins IS, Lavelle P, Mathieu J, Tedersoo L, Bahram M, Gottschall F, Guerra CA, Hines J, Patoine G *et al.* 2019. Global mismatches in aboveground and belowground biodiversity. *Conservation Biology* **33**: 1187–1192.
- Chomel M, Guittonny-Larcheveque M, Fernandez C, Gallet C, DesRochers A, Pare D, Jackson BG, Baldy V. 2016. Plant secondary metabolites: a key driver of litter decomposition and soil nutrient cycling. *Journal of Ecology* **104**: 1527–1541.
- Coûteaux MM, Bottner P, Berg B. 1995. Litter decomposition, climate and litter quality. *Trends in Ecology & Evolution* **10**: 63–66.
- Delgado-Baquerizo M, Bardgett RD, Vitousek PM, Maestre FT, Williams MA, Eldridge DJ, Lambers H, Neuhauser S, Gallardo A, Garcia-Velazquez L *et al.* 2019. Changes in belowground biodiversity during ecosystem development. *Proceedings of the National Academy of Sciences, USA* **116**: 6891–6896.
- Fanin N, Bezaud S, Sarneel JM, Cecchini S, Nicolas M, Augusto L. 2019a. Relative importance of climate, soil and plant functional traits during the early decomposition stage of standardized litter. *Ecosystems*. doi: 10.1007/s10021-019-00452-z.
- Fanin N, Fromin N, Bertrand I. 2016. Functional breadth and home-field advantage generate functional differences among soil microbial decomposers. *Ecology* **97**: 1023–1037.
- Fanin N, Kardol P, Farrell M, Kempel A, Ciobanu M, Nilsson MC, Gundale MJ, Wardle DA. 2019b. Effects of plant functional group removal on structure and function of soil communities across contrasting ecosystems. *Ecology Letters* **22**: 1095–1103.
- Freschet GT, Aerts R, Cornelissen JHC. 2012a. Multiple mechanisms for trait effects on litter decomposition: moving beyond home-field advantage with a new hypothesis. *Journal of Ecology* **100**: 619–630.
- Freschet GT, Aerts R, Cornelissen JHC. 2012b. A plant economics spectrum of litter decomposability. *Functional Ecology* **26**: 56–65.
- Freschet GT, Cornwell WK, Wardle DA, Elumeeva TG, Liu W, Jackson BG, Onipchenko VG, Soudzilovskaia NA, Tao J, Cornelissen JHC *et al.* 2013. Linking litter decomposition of above- and below-ground organs to plant–soil feedbacks worldwide. *Journal of Ecology* **101**: 943–952.
- Gholz HL, Wedin DA, Smitherman SM, Harmon ME, Parton WJ. 2000. Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition. *Global Change Biology* **6**: 751–765.
- Jackrel SL, Gilbert JA, Wootton JT. 2019. The origin, succession, and predicted metabolism of bacterial communities associated with leaf decomposition. *mBio* **10**: e01703–e01719.
- Jackson RB, Lajtha K, Crow SE, Hugelius G, Kramer MG, Pineiro G. 2017. The ecology of soil carbon: pools, vulnerabilities, and biotic and abiotic controls. *Annual Review of Ecology, Evolution and Systematics* **48**: 419–445.
- Jackson RB, Mooney HA, Schulze ED. 1997. A global budget for fine root biomass, surface area, and nutrient contents. *Proceedings of the National Academy of Sciences, USA* **94**: 7362.
- Jacobs LM, Sulman BN, Brzostek ER, Feighery JJ, Phillips RP. 2018. Interactions among decaying leaf litter, root litter and soil organic matter vary with mycorrhizal type. *Journal of Ecology* **106**: 502–513.
- Keiluweit M, Nico P, Harmon ME, Mao JD, Pett-Ridge J, Kleber M. 2015. Long-term litter decomposition controlled by manganese redox cycling. *Proceedings of the National Academy of Sciences, USA* **112**: E5253–E5260.
- Keiser AD, Keiser DA, Strickland MS, Bradford MA, de Vries F. 2014. Disentangling the mechanisms underlying functional differences among decomposer communities. *Journal of Ecology* **102**: 603–609.
- Keiser AD, Strickland MS, Fierer N, Bradford MA. 2011. The effect of resource history on the functioning of soil microbial communities is maintained across time. *Biogeosciences* **8**: 1477–1486.
- Lin D, Pang M, Fanin N, Wang H, Qian S, Zhao L, Yang Y, Mi X, Ma K. 2019. Fungi participate in driving home-field advantage of litter decomposition in a subtropical forest. *Plant and Soil* **434**: 467–480.
- Lu WJ, Liu N, Zhang YJ, Zhou JQ, Guo YP, Yang X. 2017. Impact of vegetation community on litter decomposition: evidence from a reciprocal transplant study with C-13 labeled plant litter. *Soil Biology & Biochemistry* **112**: 248–257.
- Ma C, Xiong Y, Li L, Guo D. 2016. Root and leaf decomposition become decoupled over time: implications for below- and above-ground relationships. *Functional Ecology* **30**: 1239–1246.
- Melillo JM, Aber JD, Muratore JF. 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* **63**: 621–626.
- Milcu A, Manning P. 2011. All size classes of soil fauna and litter quality control the acceleration of litter decay in its home environment. *Oikos* **120**: 1366–1370.

- Minerovic AJ, Valverde-Barrantes OJ, Blackwood CB. 2018. Physical and microbial mechanisms of decomposition vary in importance among root orders and tree species with differing chemical and morphological traits. *Soil Biology & Biochemistry* 124: 142–149.
- Palozzi JE, Lindo Z. 2017. Pure and mixed litters of *Sphagnum* and *Carex* exhibit a home-field advantage in Boreal peatlands. *Soil Biology & Biochemistry* 115: 161–168.
- Palozzi JE, Lindo Z. 2018. Are leaf litter and microbes team players? Interpreting home-field advantage decomposition dynamics. *Soil Biology & Biochemistry* 124: 189–198.
- Sauvadet M, Fanin N, Chauvat M, Bertrand I. 2019. Can the comparison of above- and below-ground litter decomposition improve our understanding of bacterial and fungal successions? *Soil Biology & Biochemistry* 132: 24–27.
- Sterkenburg E, Clemmensen KE, Ekblad A, Finlay RD, Lindahl BD. 2018. Contrasting effects of ectomycorrhizal fungi on early and late stage decomposition in a boreal forest. *The ISME Journal* 12: 2187–2197.
- Sun T, Hobbie SE, Berg B, Zhang HG, Wang QK, Wang ZW, Hattenschwiler S. 2018. Contrasting dynamics and trait controls in first-order root compared with leaf litter decomposition. *Proceedings of the National Academy of Sciences, USA* 115: 10392–10397.
- Swift MJ, Heal OW, Anderson JM. 1979. *Decomposition in terrestrial ecosystems*. Berkeley, CA, USA: University of California Press.
- Veen GF, Freschet GT, Ordonez A, Wardle DA. 2015a. Litter quality and environmental controls of home-field advantage effects on litter decomposition. *Oikos* 124: 187–195.
- Veen GFC, Keiser AD, van der Putten WH, Wardle DA, Hart M. 2018. Variation in home-field advantage and ability in leaf litter decomposition across successional gradients. *Functional Ecology* 32: 1563–1574.
- Veen GF, Sundqvist MK, Wardle DA. 2015b. Environmental factors and traits that drive plant litter decomposition do not determine home-field advantage effects. *Functional Ecology* 29: 981–991.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

**Methods S1** Descriptions of the study site, litter traits measurements and statistical analyses.

**Table S1** Initial litter chemical traits of leaf and fine root litters for each of the three species used in this study.

**Table S2** PERMANOVA statistics for the effects of species, organs and their interaction on litter chemical traits.

**Table S3** Dissimilarity of chemical traits between leaf and root litters calculated with the Euclidean distance index among different litter types.

**Table S4** Results of the model selection procedure of the linear mixed effects models used to test for the effects of species, incubation site, harvest time and their interactions on mass loss of leaf and fine root litter, respectively.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.

**Key words:** aboveground–belowground, home-field advantage, litter decomposition, local adaptation, plant–soil interactions.

Received, 25 October 2019; accepted, 20 February 2020.



## About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews and Tansley insights.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit [www.newphytologist.com](http://www.newphytologist.com) to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office ([np-centraloffice@lancaster.ac.uk](mailto:np-centraloffice@lancaster.ac.uk)) or, if it is more convenient, our USA Office ([np-usaoffice@lancaster.ac.uk](mailto:np-usaoffice@lancaster.ac.uk))
- For submission instructions, subscription and all the latest information visit [www.newphytologist.com](http://www.newphytologist.com)