



**HAL**  
open science

## Serological evidence of equine infectious anaemia, West Nile fever, surra and equine piroplasmiasis in a herd of horses in northern Argentina

Laurent Hébert, Gonzalo Polledo, Fanny Lecouturier, Mariana Giorgi, Cécile Beck, Steeve Lowenski, Karine Laroucau, Philippe Büscher, Aymeric Hans, Teótimo Becù

### ► To cite this version:

Laurent Hébert, Gonzalo Polledo, Fanny Lecouturier, Mariana Giorgi, Cécile Beck, et al.. Serological evidence of equine infectious anaemia, West Nile fever, surra and equine piroplasmiasis in a herd of horses in northern Argentina. *Veterinary Parasitology: Regional Studies and Reports*, 2021, 24, pp.100566. 10.1016/j.vprsr.2021.100566 . hal-03350214

**HAL Id: hal-03350214**

**<https://hal.inrae.fr/hal-03350214>**

Submitted on 24 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Serological evidence of equine infectious anemia, West Nile fever, surra and equine**  
2 **piroplasmiasis in a herd of horses in northern Argentina**

3

4 **Running title:** Serological survey of a herd in Argentina

5

6 Laurent Hébert<sup>1,\*</sup>, Gonzalo Polledo<sup>2</sup>, Fanny Lecouturier<sup>1</sup>, Mariana Giorgi<sup>2</sup>, Cécile Beck<sup>3</sup>,  
7 Steeve Lowenski<sup>3</sup>, Karine Laroucau<sup>4</sup>, Philippe Büscher<sup>5</sup>, Aymeric Hans<sup>1</sup>, Teótimo Becù<sup>2</sup>

8

9 <sup>1</sup> Unité PhEED, Laboratoire de santé animale, site de Normandie, ANSES, RD675, 14430,  
10 Goustranville, France.

11 <sup>2</sup> Laboratorio de Diagnóstico, Clínica Equina SRL, Alem 1698 - Capitan Sarmiento, Buenos  
12 Aires, Argentina.

13 <sup>3</sup> UMR 1161 Virologie, Laboratoire de santé animale, site de Maisons-Alfort, ANSES,  
14 INRAE, ENVA, Maisons-Alfort, France.

15 <sup>4</sup> Unité Zoonoses bactériennes, Laboratoire de santé animale, site de Maisons-Alfort, ANSES,  
16 94700 Maisons-Alfort, France.

17 <sup>5</sup> Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155,  
18 2000 Antwerpen, Belgium.

19

20 #Corresponding author. Tel.: +33 231792276; Fax: +33 231392137. *E-mail address:*  
21 laurent.hebert@anses.fr (Laurent Hébert). *Complete address:* ANSES, Unité PhEED,  
22 Laboratoire de santé animale, site de Normandie, RD675, 14430, Goustranville, France.

23 **Abstract**

24 Northern Argentina hosts equine populations living under preserved natural areas and  
25 extensive breeding conditions, with limited access to veterinary care. Horses can be in contact  
26 with *i*) wildlife considered to be a potential reservoir of horse pathogens (e.g. capybara, coatis  
27 and pampas deer) and/or *ii*) potential disease vectors such as ticks, horse flies, *Culicidae* and  
28 vampire bats. In this context, the aim of this study was to assess the exposure of horses from a  
29 herd in northern Argentina to different vector-borne pathogens.

30 Serum samples were collected from 20 horses on a farm in Chaco province. Most of  
31 these horses were in good health, but a few showed clinical signs such as fever, neurological  
32 signs or emaciation. Potential vectors (ticks, horse flies and *Culicidae*) were present and a  
33 fresh bite of a vampire bat (*Desmodus rotundus*) was observed on one horse. This serological  
34 survey revealed that 100% (20/20) were positive for equine infectious anaemia (EIA), 100%  
35 (18/18) for West Nile fever (WNF), 53% (10/19) for surra and 45% (9/20) for equine  
36 piroplasmiasis (*Babesia equi*). Among these horses, four were found seropositive for all four  
37 infections. On the other hand, all the tested horses were seronegative for equine viral arteritis  
38 (EVA), Eastern equine encephalomyelitis (EEE), Venezuelan equine encephalitis (VEE),  
39 Western equine encephalomyelitis (WEE) and glanders.

40 The data from this survey conducted on a small number of animals illustrate the need for  
41 an effective application of surveillance programmes and control measures for equine diseases  
42 in northern Argentina and constitute, to our knowledge, the first report of horses  
43 simultaneously seropositive for EIA, WNF, surra and equine piroplasmiasis.

44

45 **Keywords**

46 Argentina, equine infectious anaemia, piroplasmiasis, surra, West Nile fever

47

48 **Short communication**

49 The province of Chaco in Argentina is characterised by subtropical weather with a dry season  
50 in a landscape composed of humid forest, savannas, gallery forests and flooded forests. In this  
51 rural environment, the economy is based on agriculture and extensive livestock activities  
52 (predominantly cattle and horses), but access to veterinary care such as vaccinations and  
53 treatment or confirmatory diagnosis remains limited for some horse populations. This farming  
54 method implies that domestic animals are frequently in contact with wild animals and exposed  
55 to numerous vectors. The reported wildlife includes numerous animals considered to be  
56 potential reservoirs for horse pathogens, including capybara (*Hydrochoerus hydrochaeris*)  
57 (Herrera and Castro, 2017), ring-tailed coati (*Nasua nasua*) (Santos et al., 2018), pampas deer  
58 (*Ozotoceros bezoarticus*) (Silveira et al., 2013), black howler monkeys (*Alouatta caraya*)  
59 (Martínez et al., 2016) and various bird species. The presence in this geographical area of  
60 disease vectors such as ticks (*Ixodidae*) (Nava et al., 2006), horse flies (*Tabanidae* and  
61 *Stomoxys* flies), *Culicidae* (Stein et al., 2013) and vampire bats (*Desmodus rotundus*) (Hoare,  
62 1965) is also documented. Although the number of field studies dedicated to the pathogens  
63 circulating in this province remains low, it may be noted that numerous vector-borne diseases  
64 have already been reported. These include equine infectious anaemia (EIA) (De la Sota et al.,  
65 2005), West Nile fever (WNF) (Oria et al., 2018), surra (Santa Cruz et al., 2013), equine  
66 piroplasmosis (*Babesia equi*) (Holman et al., 1998), Eastern equine encephalomyelitis (EEE),  
67 Venezuelan equine encephalitis (VEE) and Western equine encephalomyelitis (WEE)  
68 (Monath et al., 1985).

69 In a context where domestic and wild animals cohabit with pathogens and vectors, our  
70 objective was to evaluate the prevalence of various major equine diseases notifiable to the  
71 OIE (World Organisation for Animal Health) in a herd of horses from the northern

72 Argentinean province of Chaco. The diseases investigated were EIA, WNF, surra, equine  
73 piroplasmosis, glanders, EVA, EEE, VEE and WEE.

74 The study was conducted on a 3,000-hectare farm in the municipality of Laguna Limpia in the  
75 province of Chaco, Argentina (Fig. 1). The area comprises native forest, ponds and flooded  
76 areas, and a diverse fauna including capybaras, coatis, deers, howler monkeys and cattle,  
77 together with the presence of common vectors including ticks, tabanids, stomoxes, *Culicidae*  
78 and vampire bats. Following concerns expressed by farmers about the general health of their  
79 animals, serum samples were collected in June 2018 from 20 horses randomly selected in a  
80 herd in preserved natural areas and extensive breeding condition made up of 48 mixed-breed  
81 horses of all ages (Fig. 2). These horses have regular access to food but limited access to  
82 veterinary care. The only pharmaceutical products administered to our knowledge are  
83 mebendazole, and vitamin B12 and iron supplements. None of the horses were vaccinated.  
84 They were sampled regardless of their physical condition. Their general health, rectal  
85 temperature and clinical signs such as oedema and ataxia were recorded. Blood samples were  
86 collected aseptically by jugular venipuncture and either collected in glass tubes and  
87 centrifuged at 1600 xg for 15 min for serum collection or in EDTA tubes to monitor the  
88 packed cell volume (PCV), white blood cells (WBC), neutrophils and lymphocytes  
89 (Supporting Information Table S1).

90 The serological methods used in this study are listed in Table 1, and were performed in  
91 accordance with the corresponding chapters in the OIE Manual of Diagnostic Tests and  
92 Vaccines for Terrestrial Animals (OIE, 2015). Inconclusive results (cytotoxicity and  
93 anticomplementary) were excluded for calculation of seropositivity rates.

94 Of the 20 horses sampled in this study, 60% (12/20) were of average or poor general health,  
95 and 90% (7/8; the temperature of the other animals was not recorded) had a fever ( $\geq 38.5$  °C)  
96 (Supporting Information Table S1). The observed clinical signs (Fig. 2) included oedema,

97 nasal discharge, emaciation, depression, recumbency and neurological signs (Fig. 2 and  
98 Supporting Information Table S1). Moreover, cases of abortion had previously been reported  
99 on this farm. An abundant population of haematophagous flies (*Tabanidae* and *Stomoxys*),  
100 *Culicidae* and ticks (*Amblyomma cajennense*) were observed at the same time in the horses'  
101 environment, and a fresh vampire bat bite was observed on one horse's neck (Fig. 2F).  
102 In this environment conducive to the dissemination of vector-borne diseases, we evaluated the  
103 prevalence of antibodies against EIA, WNF, surra, equine piroplasmosis, glanders, EVA,  
104 EEE, VEE and WEE. Results are presented in Supporting Information Table S1 and  
105 summarised in Fig 3.

106 The seroprevalence values were 100% for EIA (20/20) and WNF (18/18), 53% (10/19) for  
107 surra and 45% (9/20) for equine piroplasmosis. Interestingly, four horses were seropositive for  
108 all four diseases. All the animals tested were seronegative for glanders, EVA, EEE, VEE and  
109 WEE (Fig. 3A and Supporting Information Table S1).

110 EIA is widespread in northern Argentina, with infection rates of 39% and 76% reported in the  
111 provinces of Chaco and Formosa, respectively (De la Sota et al., 2005). However, this is the  
112 first time to our knowledge that a seroprevalence value for EIA of 100% is reported for a  
113 given herd. This potential EIA reservoir should be of great concern for the horse-breeding  
114 industry in a context where EIA is covered by strict regulatory control measures in many  
115 countries.

116 WNF was first detected in Argentina from diseased and dead horses in 2006 (Chancey et al.,  
117 2015) and, since then, its circulation has been reported in different provinces in Argentina,  
118 including Chaco (Oria et al., 2018). Given the abundance of *Culicidae* (vectors of WNF) and  
119 birds (hosts, natural reservoir) in this region (Campbell et al., 2002), there appears to be a  
120 major risk of this virus being transmitted to incidental dead-end hosts represented by horses  
121 (and potentially humans). This concern is illustrated by our results, which revealed

122 seroprevalence for WNF of 100% within our panel. Even if such a level of seroprevalence has  
123 already been reported for WNF in equids in Morocco, Pakistan, Palestine and Iran (Eyboosh  
124 et al., 2019), this is, to our knowledge, the first time that a whole group of equids have been  
125 detected as seropositive for WNF in Argentina. Besides, even if the WNF microneutralization  
126 test is highly specific for WNF, WNF belonging to Japanese encephalitis serocomplex,  
127 serological cross reaction with Saint Louis encephalitis virus can't be totally ruled out.

128 Surra (formerly referred to as “mal de caderas” in South America), caused by *T. evansi*, has  
129 been circulating in Argentina for decades. It not only affects horses but also wildlife such as  
130 capybaras and domestic animals such as dogs (Aregawi et al., 2019). In this study, we  
131 determined the percentage of horses seropositive for surra by immune trypanolysis (IT) RoTat  
132 1.2, which detects only antibodies that recognise a single multi-copy epitope at the surface of  
133 a *T. evansi* type A clone expressing the RoTat 1.2 variant surface glycoprotein (Verloo et al.,  
134 2001). This test is the reference test for surra antibody detection due to its very high  
135 specificity and thus positive prediction value (Holland et al., 2002). Given the severity of this  
136 disease and the absence of treatment or a vaccination strategy, the high level of seropositivity  
137 detected in this herd (53%) is a finding of serious concern not only for this herd but for the  
138 whole equine industry in the province of Chaco.

139 Equine piroplasmiasis is known to be endemic in Chaco (Holman et al., 1998). As  
140 complement fixation test (CFT) signals generally disappear 2 to 3 months post infection in  
141 infected horses (Weiland, 1986) the (CFT) results obtained in this study, with 9/20 animal  
142 tested positive, indicate that the disease is currently circulating in this area. Furthermore, we  
143 can assume that we would have obtained much higher prevalence rates if we had tested the  
144 sera with the indirect fluorescent antibody test, a method that gives positive results during  
145 latent infection (Weiland, 1986), and we could even expect 100% seroprevalence as  
146 previously described (Holman et al., 1998).

147 No obvious correlation between a specific disease and the general health of these horses was  
148 detected (Supporting Information Table S1).

149

150 The main findings of our study were the confirmation of the presence of antibodies against  
151 EIA, WNF, equine trypanosomiasis and equine piroplasmiasis in horses indigenous to the  
152 province of Chaco, Argentina. In this context, the presence of vectors such as ticks, horse  
153 flies, *Culicidae* and vampire bats constitute an important potential source of pathogen  
154 dissemination that should raise significant concern among the Argentinean horse-breeding  
155 industry. Interestingly, four horses appeared simultaneously seropositive for EIA, WNF, surra  
156 and equine piroplasmiasis. All the animals tested were seronegative for glanders, EVA, EEE,  
157 VEE and WEE. Despite the small scale of our study, our results illustrate the need for  
158 effective implementation of surveillance programmes and control measures to tackle equine  
159 diseases in northern Argentina.

160

## 161 **Acknowledgements**

162 The authors would like to thank Julieta Vargas and Jose Margosa for serum sampling. We  
163 also wish to thank Delphine Libby-Claybrough, a professional translator and native English  
164 speaker, for carefully reading and reviewing the paper. This work was supported by ANSES  
165 and the European Commission through DG SANTE funding for the Reference Laboratory for  
166 Equine Diseases other than African Horse Sickness. The funders had no role in study design,  
167 data collection and interpretation, or the decision to submit the work for publication.  
168 Laboratoire de santé animale, site de Normandie is a member of the GIS Centaure equine  
169 research and authors thank Regional Council of Normandy for providing an excellent  
170 scientific environment.

171

172

173 **Conflict of interest statement**

174 The authors declare no potential conflicts of interest with respect to the research, authorship,  
175 publication of this article and/or financial and personal relationships that could  
176 inappropriately influence this work.

177

178 **Ethical statement**

179 The authors confirm their adherence to the ethical policies of the journal, as noted on the  
180 journal's author guidelines page, and that the appropriate ethical review committee approval  
181 has been received. Procedures involving horses were in accordance with the ethical standards  
182 of the institution or practice at which the studies were conducted. The protocol for the culture  
183 of trypanosomes in mice was approved by the Veterinary Ethical Committee of the Institute  
184 of Tropical Medicine Antwerp, Belgium (ITM) (BM2013-7).

185

186 **Data Availability Statement**

187 The data that support the findings of this study are available from the corresponding author  
188 upon reasonable request.

189

190 **References**

- 191 Aregawi, W.G., Agga, G.E., Abdi, R.D., Büscher, P., 2019. Systematic review and meta-  
192 analysis on the global distribution, host range, and prevalence of *Trypanosoma evansi*.  
193 Parasit Vectors 12, 67.
- 194 Campbell, G.L., Marfin, A.A., Lanciotti, R.S., Gubler, D.J., 2002. West Nile virus. Lancet  
195 Infect Dis 2, 519-529.
- 196 Chancey, C., Grinev, A., Volkova, E., Rios, M., 2015. The global ecology and epidemiology  
197 of West Nile virus. Biomed Res Int 2015, 376230.
- 198 De la Sota, M.D., Gonzáles, R., Chiricosta, A., 2005. Contribución a la prevalencia de la  
199 Anemia Infecciosa Equina en la República Argentina. Revista Colegio de Médicos  
200 Veterinarios de la Provincia de Buenos Aires 10, 52-60.
- 201 Eybpoosh, S., Fazlalipour, M., Baniyadi, V., Pouriayevali, M.H., Sadeghi, F., Ahmadi  
202 Vasmehjani, A., Karbalaie Niya, M.H., Hewson, R., Salehi-Vaziri, M., 2019.  
203 Epidemiology of West Nile Virus in the Eastern Mediterranean region: A systematic  
204 review. PLoS neglected tropical diseases 13, e0007081.
- 205 Herrera, E.A., Castro, Y., 2017. *Trypanosoma evansi* (Kinetoplastida: Trypanosomatidae) in  
206 capybaras (*Hydrochoerus hydrochaeris*, Rodentia: Hydrochoeridae): prevalence,  
207 effect and sexual selection. Rev Biol Trop 65, 229-237.
- 208 Hoare, C.A., 1965. Vampire bats as vectors and hosts of equine and bovine trypanosomes.  
209 Acta tropica 22, 204-216.
- 210 Holland, W.G., Thanh, N.G., My, L.N., Magnus, E., Verloo, D., Büscher, P., Goddeeris, B.,  
211 Vercruyse, J., 2002. Evaluation of whole fresh blood and dried blood on filter paper  
212 discs in serological tests for *Trypanosoma evansi* in experimentally infected water  
213 buffaloes. Acta tropica 81, 159-165.

214 Holman, P.J., Becú, T., Bakos, E., Polledo, G., Cruz, D., Wagner, G.G., 1998. *Babesia equi*  
215 field isolates cultured from horse blood using a microcentrifuge method. J Parasitol  
216 84, 696-699.

217 Lazić, S., Lupulović, D., Gaudaire, D., Petrovic, T., Lazić, G., Hans, A., 2017. Serological  
218 evidence of equine arteritis virus infection and phylogenetic analysis of viral isolates  
219 in semen of stallions from Serbia. BMC Vet Res 13, 316.

220 Martínez, M.F., Kowalewski, M.M., Salomón, O.D., Schijman, A.G., 2016. Molecular  
221 characterization of trypanosomatid infections in wild howler monkeys (*Alouatta*  
222 *caraya*) in northeastern Argentina. Int J Parasitol Parasites Wildl 5, 198-206.

223 Monath, T.P., Sabattini, M.S., Pauli, R., Daffner, J.F., Mitchell, C.J., Bowen, G.S., Cropp,  
224 C.B., 1985. Arbovirus investigations in Argentina, 1977–1980. The American Journal  
225 of Tropical Medicine and Hygiene 34, 966-975.

226 Nava, S., Caparros, J.A., Mangold, A.J., Guglielmone, A.A., 2006. Ticks (*Acari: Ixodida:*  
227 *Argasidae, Ixodidae*) infesting humans in Northwestern Cordoba Province, Argentina.  
228 Medicina (B Aires) 66, 225-228.

229 Oria, G.I., Spinsanti, L.I., Pirota, V.L., Martinez, F., Stechina, O.S., Etchepare, E., Contigiani,  
230 M.S., Stein, M., 2018. Seroprevalence of *Flavivirus* in horses in Chaco, Argentina.  
231 Circulation during 2013-2014. Revista Brasileira de Medicina Veterinaria 40.

232 Santa Cruz, A., Comolli, J., Ortiz, J., González, J., González, A., 2013. Morphometric data of  
233 *Trypanosoma evansi* in capybaras (*Hydrochaeris hydrochaeris*) from Chaco,  
234 Argentina, Vol 24, 60-62 pp.

235 Santos, F.M., de Macedo, G.C., Barreto, W.T.G., Oliveira-Santos, L.G.R., Garcia, C.M.,  
236 Mourão, G.M., Edith de Oliveira Porfírio, G., Domenis Marino, E., Rogério André,  
237 M., Perles, L., Elisei de Oliveira, C., Braziliano de Andrade, G., Jansen, A.M.,  
238 Miraglia Herrera, H., 2018. Outcomes of *Trypanosoma cruzi* and *Trypanosoma evansi*

239 infections on health of Southern coati (*Nasua nasua*), crab-eating fox (*Cerdocyon*  
240 *thous*), and ocelot (*Leopardus pardalis*) in the Brazilian Pantanal. PLoS One 13,  
241 e0201357.

242 Silveira, J.A., Rabelo, E.M., Lacerda, A.C., Borges, P.A., Tomás, W.M., Pellegrin, A.O.,  
243 Tomich, R.G., Ribeiro, M.F., 2013. Molecular detection and identification of  
244 hemoparasites in pampas deer (*Ozotoceros bezoarticus* Linnaeus, 1758) from the  
245 Pantanal Brazil. Ticks Tick Borne Dis 4, 341-345.

246 Stein, M., Zalazar, L., Willener, J.A., Almeida, F.L., Almirón, W.R., 2013. Culicidae  
247 (Diptera) selection of humans, chickens and rabbits in three different environments in  
248 the province of Chaco, Argentina. Mem Inst Oswaldo Cruz 108, 563-571.

249 Verloo, D., Magnus, E., Büscher, P., 2001. General expression of RoTat 1.2 variable antigen  
250 type in *Trypanosoma evansi* isolates from different origin. Veterinary parasitology 97,  
251 183-189.

252 Weiland, G., 1986. Species-specific serodiagnosis of equine piroplasma infections by means  
253 of complement fixation test (CFT), immunofluorescence (IIF), and enzyme-linked  
254 immunosorbent assay (ELISA). Veterinary parasitology 20, 43-48.

255

256

257

258 **Table 1.** Methods used for the serological diagnosis of equine diseases from the sera collected  
259 on a farm in the municipality of Laguna Limpia, province of Chaco, Argentina.

Disease	Serological method	Antigen source	Reference or provider	OIE Terrestrial Manual chapter
WNF	MNT <sup>a</sup>	WNV IS-98-ST1	AF481864	3.1.24.
EEE	MNT	NCPV689 strain H178/99	ICTV catalog number 0407041v	3.5.5.
VEE	MNT	Trinidad Donkey strain TC-83	L01443.1	3.5.5.
WEE	MNT	NCPV691, strain H160/99	ICTV catalog number 0407043v	3.5.5.
EVA	VNT <sup>b</sup>	EAV Bucyrus (ATCC VR - 796)	(Lazić et al., 2017)	3.5.10.
EIA	AGID <sup>c</sup>	Recombinant p26 protein	VMRD <sup>d</sup>	3.5.6.
Surra	IT <sup>e</sup>	<i>Trypanosoma evansi</i> Rotat 1.2	(Verloo et al., 2001)	3.1.21.
Equine piroplasmosis	CFT <sup>f</sup>	<i>Babesia caballi</i>	USDA <sup>g</sup>	3.5.8.
Glanders	CFT	<i>Burkholderia mallei</i> Ivan/NCTC10230 strain -	Ccpro GmbH, Germany	3.5.11.

260 <sup>a</sup> Virus-specific microneutralization tests.

261 <sup>b</sup> Virus neutralisation test.

262 <sup>c</sup> Agar gel immunodiffusion assay.

263 <sup>d</sup> Veterinary Medical Research & Development (Pullman, Washington, USA).

264 <sup>e</sup> Immune trypanolysis.

265 <sup>f</sup> Complement fixation test.

266 <sup>g</sup> United States Department of Agriculture (Ames, Iowa, USA).

267

268 **Figure captions**

269 **Figure 1.** Geographical situation of the Laguna Limpia farm (province of Chaco, Argentina)  
270 where the horse samples were collected.

271

272 **Figure 2.** Illustration of the sampled horse herd, clinical signs and evidence of  
273 haematophagous vectors. Photographs of the herd during the serum sampling campaign (A  
274 and B). The clinical signs observed included weight loss (C) and nasal discharge (D). The  
275 evidenced haematophagous vectors included ticks (E) and vampire bats (*Desmodus rotundus*),  
276 as shown by the fresh bite (F) on the neck of one of the horses.

277

278 **Figure 3.** Results of the serological tests performed on the sera collected from 20 horses on a  
279 farm in the province of Chaco, northern Argentina. (A) Percentage of positive results obtained  
280 in serological tests for equine infectious anaemia (EIA), West Nile fever (WNF), surra, equine  
281 piroplasmosis, glanders, equine viral arteritis (EVA), Eastern equine encephalomyelitis  
282 (EEE), Venezuelan equine encephalitis (VEE) and Western equine encephalomyelitis (WEE).  
283 (B) Frequency distribution of the number of seropositive results obtained for each infection  
284 per horse.

285

286

287 **Supporting information**

288 Table S1. Data recorded for 20 horses on a farm located in the province of Chaco, northern  
289 Argentina.

290



## Argentina



## Chaco Province







## Serological evidence of equine infectious anemia, West Nile fever, surra and equine piroplasmosis in a herd of horses in northern Argentina

Laurent Hébert<sup>1</sup>, Gonzalo Polledo<sup>2</sup>, Fanny Lecouturier<sup>1</sup>, Mariana Giorgi<sup>2</sup>, Cécile Beck<sup>3</sup>, Steeve Lowenski<sup>3</sup>, Karine Laroucau<sup>4</sup>, Philippe Büscher<sup>5</sup>, Aymeric Hans<sup>1</sup>, Teótimo Becù<sup>2</sup>

<sup>1</sup> Unité PhEED, Laboratoire de santé animale, site de Normandie, ANSES, RD675, 14430, Goustranville, France.

<sup>2</sup> Laboratorio de Diagnóstico, Clinica Equina SRL, Alem 1698 - Capitan Sarmiento, Buenos Aires, Argentina.

<sup>3</sup> UMR 1161 Virologie, Laboratoire de santé animale, site de Maisons-Alfort, ANSES, INRAE, ENVA, Maisons-Alfort, France.

<sup>4</sup> Unité Zoonoses bactériennes, Laboratoire de santé animale, site de Maisons-Alfort, ANSES, 94700 Maisons-Alfort, France.

<sup>5</sup> Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium.

This study describes, for the first time, horses simultaneously seropositive for equine infectious anaemia, West Nile fever, surra and equine piroplasmosis in northern Argentina.

