

Introducing Grasslands into Crop Rotations, a Way to Restore Microbiodiversity and Soil Functions

Wassila Riah-Anglet, Elodie Cusset, Rémi Chaussod, Criquet Stéven, Marie-Paule Norini, Nathalie Cheviron, Christian Mougin, Karine Laval, Xavier Latour, Marc Legras, et al.

► To cite this version:

Wassila Riah-Anglet, Elodie Cusset, Rémi Chaussod, Criquet Stéven, Marie-Paule Norini, et al.. Introducing Grasslands into Crop Rotations, a Way to Restore Microbiodiversity and Soil Functions. Agriculture, 2021, 11 (10), pp.909. 10.3390/agriculture11100909. hal-03354447v2

HAL Id: hal-03354447 https://hal.inrae.fr/hal-03354447v2

Submitted on 12 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License





Article Introducing Grasslands into Crop Rotations, a Way to Restore Microbiodiversity and Soil Functions

Wassila Riah-Anglet ^{1,2,*}, Elodie Cusset ^{1,2}, Rémi Chaussod ³, Stéven Criquet ⁴, Marie-Paule Norini ^{1,5}, Nathalie Cheviron⁶, Christian Mougin⁶, Karine Laval^{1,2}, Xavier Latour^{2,7}, Marc Legras^{1,2} and Isabelle Trinsoutrot-Gattin ^{1,2}

- 1 UniLaSalle, AGHYLE Research Unit UP 2018.C101, Rouen Team, 76134 Mont-Saint Aignan, France; elodie.cusset@unilasalle.fr (E.C.); mp.norini@brgm.fr (M.-P.N.); Karine.LAVAL@unilasalle.fr (K.L.); marc.LEGRAS@unilasalle.fr (M.L.); Isabelle.GATTIN@unilasalle.fr (I.T.-G.)
- 2 Research Federations NORVEGE Fed4277, Normandy University, F-76821 Mont-Saint-Aignan, France; xavier.latour@univ-rouen.fr 3
 - SEMSE, 2 Chemin du Lavoir, 21310 Viévigne, France; remi.chaussod@semse.fr
- 4 Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE, Campus Etoile St Jérôme, Av Escadrille Normandie Niémen-case 441-13397 Cedex 20, F-13397 Marseille, France; steven.criquet@imbe.fr 5
- BRGM, BP 36009, 45060 Orléans Cedex 2, France
- 6 Université Paris-Saclay, INRAE, AgroParisTech, UMR 1402 ECOSYS, Platform Biochem-Env, F-78026 Ver-sailles, France; nathalie.cheviron@inrae.fr (N.C.); christian.mougin@inrae.fr (C.M.)
- 7 Laboratory of Microbiology Signals and Microenvironment (LMSM/CBSA), University of Rouen Normandy, 55 rue Saint-Germain, F-27000 Evreux, France
- Correspondence: wassila.riah-anglet@unilasalle.fr

Abstract: The aims of this study were to investigate (i) the influence of aging grassland in the recovery of soil state by the comparison of permanent grassland, two restored grasslands, two temporary grasslands, and a continuous crop in the same pedoclimatic conditions, (ii) the extent and the persistence of the potential changes following a grassland/or cropland phase. We hypothesized that the level of microbial communities and enzyme activities could achieve a profile close to that of permanent grassland after the introduction of grassland for a few years in crop rotations. Soil biophysicochemical properties were studied. Our results indicated that the abundance of microbial communities and enzyme activities were positively correlated to soil C and N contents and negatively correlated to soil pH. The changes in microbial abundance level were strongly linked to the changes in functional level when grasslands are introduced into crop rotations. We also showed that a continuous crop regime had a stronger legacy on the soil biota and functions. By contrast, the legacy of a grassland regime changed quickly when the grassland regime is interrupted by recent culture events. A grassland regime enabled the restoration of functions after more than five cumulative years in the grassland regime.

Keywords: enzyme activities; microbial communities restoration; crop rotation; grassland; function level; grassland/cropland legacy

1. Introduction

During the last decade, nutrients deficiency has been considered as the main cause of poor productivity and crop failure. A study of the current trends in agronomic practices has suggested that the soil nutrients deficiency is further aggravated by continued use of high-yielding crop varieties, intensive cropping patterns, and the use of relatively poor fertilizers [1]. Soils are a non-renewable resource at a human time scale, and some adverse effects of degradative processes on soil quality are irreversible [2]. Thus, soils need to be recognized and valued for their productive capacities as well as their contribution to food security and the maintenance of key ecosystem services driven by soil biodiversity [3]. Among soil biodiversity, microorganisms are particularly abundant, given that just one



Citation: Riah-Anglet, W.; Cusset, E.; Chaussod, R.; Criquet, S.; Norini, M.-P.; Cheviron, N.; Mougin, C.; Laval, K.; Latour, X.; Legras, M.; et al. Introducing Grasslands into Crop Rotations, a Way to Restore Microbiodiversity and Soil Functions. Agriculture 2021, 11, 909. https:// doi.org/10.3390/agriculture11100909

Academic Editors: Purushothaman Chirakkuzhyil Abhilash, Pradeep Kumar Dubey, Eric C. Brevik, Rahul Datta, Othmane Merah and Suresh Babu

Received: 25 June 2021 Accepted: 16 September 2021 Published: 23 September 2021

Publisher's Note: MDPI stavs neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

gram of soil contains as many as 10^{10} – 10^{11} bacteria, including 6000–bacterial species and up to 200 m of fungal hyphae [4]. They represent the unseen majority and comprise a large portion of the genetic diversity of soil [4,5]. Bacteria and fungi communities play key roles in soil ecosystems because they constitute a major part of the biomass contributing to various biogeochemical cycles [6–8]. These soil microorganisms play a key role in the decomposition and mineralization of organic matter (OM) [9,10]. Indeed, they control and produce most of the enzymes involved in the breakdown of OM and influence soil carbon (C), nitrogen (N), phosphorus (P), and sulfur (S) cycling. The abundance and the diversity of soil microbial communities are also important in plant health, soil fertility, and wider ecosystem functioning [11–15].

Agricultural practices can affect directly and indirectly soil microbial communities and their functions [16,17]. Amongst these impacts, reduction in microbial diversity is often viewed as major threats to the future [18,19]. Today, agricultural practices should tend to sustainable agriculture to guarantee or improve crop yields and soil quality, thus limiting greenhouse gases emissions [20]. Among sustainable practices, farmers are reconsidering an old method of adding grasslands within arable crop rotations. This type of farming may be a way to reconcile agricultural productivity with concerns about environmental quality [21,22] by integrating crop-livestock farming systems [23]. Alternation of grassland and cropland has been indicated as more efficient than permanent cropping systems in terms of organic C sequestration, nutrient availability, and increasing biodiversity of the agrosystems [24–26]. Grasslands have been described as resilient soils, although soil cultivation results in well-documented losses of OM and soil structure [27–29].

The establishment of grasslands on previously cultivated soils can result in the recovery of some key soil properties and might act positively on soil microbial communities [30–34]. The analysis of soil microbial communities in cultivated soil and grassland found higher bacterial diversity in grassland as compared to highly disturbed cultivated soils [34–36]. McKinley et al. [37], Bach et al. [38] also documented that stress Phospholipid Fatty Acids (PLFAs) ratios indicators (saturated: monounsaturated and iso: anteiso) were highest in cultivated soils and decreased in restored grasslands. Authors explained that the reductions in these ratios during soil restoration by the alleviation of nutrient stress as grassland established on formerly cultivated soils, likely as a result of increased C substrate availability in the soil. The analysis of microbial communities showed differences in microbial abundance and diversity in response to time since grassland implementation [34,36,39,40]. Plassart et al. [41] also documented that a strong relationship was found between fungal genetic diversity and the aging of grasslands. Furthermore, various studies have reported changes in the soil functions on sites with an arable cropping history after which a grassland was established [32,41–43]. In fact, adding grasslands within arable crop rotations increase C mineralization, potential catabolic activity, microbial respiration, and soil N transformations (i.e., net N mineralization rates and net nitrification rates), indicating a strong activity of microorganisms [30,41–43].

The intimate connection between soil state, microbial communities, and enzyme activities emphasizes the importance of considering the role of microbial communities during soil restoration by the introduction of grasslands into crop rotations. Therefore, the objectives of this study were to investigate (i) the influence of aging grassland in the recovery of soil state by the comparison of different grassland regimes between permanent grassland and conventional crop in the same pedoclimatic context, (ii) the extent and the persistence of the potential changes of microbial communities and enzyme activities following a grassland/or cropland phase. We hypothesized that microbial communities could achieve an abundance level and functional state close to that of permanent grassland after the introduction of grassland for a few years into crop rotations. To achieve our objective, we analyzed (i) physicochemical properties, (ii) microbial community (indicators of total, bacterial and fungal abundance), and (iii) soil functions (14 enzyme activities involved in C, N, P, and S cycles) during the adding of grassland into crop rotations.

2. Materials and Methods

2.1. Soil Samples

The experimental site is located in northwestern France, in the Normandy area (at Yvetot, 35 km northwest from Rouen), under temperate oceanic climate (+10 °C mean annual temperature and 850 mm annual precipitation). This site was used in several previous studies and is considered as a reference site in the Normandy area [41,44–47] as well as in France [48–50].

Six plots with different land use and grassland regimes were studied. Their land uses, cropping history, and cumulative years under grasslands are presented, respectively, in Tables 1 and 2 for the 20 years before the sampling date in 2010. The permanent grassland (PG) is considered as a reference plot and has not been changed during at least the last 20 years. In contrast, continuous crop (CC) has a history of long-term cropping (wheat, maize, flax, or beet) with intensive management (plowing/fertilization). Two restored grasslands (RGI and RGII) characterized by grassland establishment after a cropping period of at least 8 years. These 2 restored grasslands differed significantly in the number of cumulative years under grassland. RGII cumulated 11 years of grassland, while RGI cumulated 5 years under grassland. We can also note that these 2 plots were replanted at the same time and have been managed in the same way since 2008. Temporary grasslands (TG and TC) were characterized by an alternating grassland period of 4 to 8 years and a corn/wheat rotation. In 2010, TC was sown with wheat after a previous corn crop, which is often used in the region as the head of rotation. The 2 temporary grasslands were managed differently in 2010. The plot TG is the oldest grassland after plot PG. The 6 plots were developed on the same native Luvisol, silty loam soil, within similar pedological and climatic contexts (Table 2).

Plots	Abbreviations	Clay (g/kg)	Silt (g/kg)	Sand (g/kg)	Agricultural Practices	Land Use at the Sampling Time	Plots Area (ha)	Cumulative Years under Grassland since 20 Years
Permanent grassland	PG ¹	163	633	204	Permanent grassland no tillage	Grassland	3	20 years
Postored grasslands	RGI ²	141	665	194	Restored grassland	Grassland	2	5.5 years
Restored grasslands	RGII ³	126	665	210	Restored grassland	Grassland	3	11 years
Temporary grasslands	TC ⁴	141	656	203	Temporary grassland, switched to crop in 2010	Wheat crop	3	14 years
leniporary grassianas	TG ⁵	147	651	202	Temporary grassland	Grassland	3	14 years
Continuous crop	CC ⁶	133	671	196	Continuous crop and fertilization	Wheat crop	4	0 year

Table 1. Descripti	on of the st	udied plots.
--------------------	--------------	--------------

¹ Permanent Grassland (PG): 20 years of pasture, permanent cover and no tillage. ² Continuous Crop (CC): permanent arable cropping. ³ Restored Grassland I (RGI): cumulate 5.5 years under grassland over a period of 20 years. ⁴ Restored Grassland II (RGII): cumulate 11 years under grassland over a period of 20 years. ⁵ Temporary Grassland (TG): cumulate 14 years under grassland over a period of 20 years. ⁶ Temporary crop (TC): this treatment is comparable to TG, but it switched to culture in the year of sampling.

Tab	le 2.	Descr	iption	of	the	cropp	ing	history	7 of t	he p	olots	since	1991.
-----	-------	-------	--------	----	-----	-------	-----	---------	--------	------	-------	-------	-------

Plots	Permanent Grassland	Restored Grasslands		Temporary	Continuous Crop	
Years	PG	RGI	RGII	TC	TG	CC
2010	grassland	grassland	grassland	Triticum aestivum	grassland	Triticum aestivum
2009	grassland	grassland	grassland	Zea mays	grassland	Zea mays
2008	grassland	Triticum aestivum	Triticum aestivum	grassland	grassland	Triticum aestivum
2007	grassland	Zea mays	Vicia faba	grassland	grassland	Zea mays
2006	grassland	Triticum aestivum	Linum usitatissimum	grassland	grassland	Triticum aestivum
2005	grassland	Zea mays	Triticum aestivum	grassland	Triticum aestivum	Linum usitatissimum
2004	grassland	Triticum aestivum	Zea mays	grassland	Zea mays	Triticum aestivum
2003	grassland	Zea mays	grassland	Triticum aestivum	grassland	Beta vulgaris
2002	grassland	grassland	grassland	Zea mays	grassland	Triticum aestivum
2001	grassland	grassland	grassland	grassland	grassland	Zea mays
2000	grassland	grassland	grassland	grassland	Triticum aestivum	Triticum aestivum

Plots	Permanent Grassland	Restored G	rasslands	Temporary	Temporary Grasslands		
Years	PG	RGI	RGII	TC	TG	CC	
1999	grassland	Pisum sativum	grassland	grassland	Zea mays	Beta vulgaris	
1998	grassland	Triticum aestivum	grassland	grassland	grassland	Triticum aestivum	
1997	grassland	Zea mays	grassland	grassland	grassland	Linum usitatissimum	
1996	grassland	Triticum aestivum	grassland	grassland	grassland	-	
1995	grassland	Zea mays	grassland	grassland	grassland	-	
1994	grassland	Linum usitatissimum	Triticum aestivum	grassland	grassland	-	
1993	grassland	Zea mays	Zea mays	Triticum aestivum	grassland	-	
1992	grassland	Triticum aestivum	Triticum aestivum	Zea mays	Triticum aestivum	-	
1991	grassland	Zea mays	Pisum sativum	grassland	Zea mays	-	

Table 2. Cont.

The sampling protocol followed the recommendations of the RMQS BioDiv national program [51], as described in Peres et al. [48] and Trap et al. [49]. For each plot, we designed a square with 10-m sides (10 m × 10 m) subdivided into 4 equal squares (5 m × 5 m) [49]. In March 2010, 12 soil cores were randomly collected in each sampling area, using an 8 cm diameter auger on the first 15 cm soil depth to form one composite sample. Four composite soil samples were, therefore, available for each plot (Figure S1). The soil samples were sieved at 2 mm and stored at 4 °C prior to analysis. Biological analyses were performed on the freshly sieved soils within 24 h after sampling, except for PLFAs and ergosterol. Indeed, for these 2 parameters, fresh soil samples were homogenized, sieved at 2 mm, and frozen at -80 °C until analyses. The main physicochemical characteristics (i.e., total carbon, total nitrogen, pH water, CEC, and P₂O₅) were measured on each individual air-dried soil sample, according to the normalized methods by the Soil Analysis Laboratory (LAS, INRA, Arras, France).

2.2. Microbial Biomass C Measurement

Soil total microbial biomass C (MBC) was determined using the chloroform fumigation extraction method according to Wu et al. [52] and Jenkinson et al. [53]. Briefly, soluble organic carbon was extracted with a 0.05 N K₂SO₄ solution from chloroform-fumigated (16 h) and non-fumigated soil samples (30 g, with 4 replicates). After centrifugation (10 min $3000 \times g$), the soluble organic carbon in the extracts was measured by persulfate-UV oxidation with a Dohrman DC-80 Total Organic Carbon Analyzer (Rosemount Analytical, Santa Clara, California, USA). MBC was calculated as a difference in the C content in the fumigated and non-fumigated samples (EC) using the k_{EC} coefficient (MBC = EC/ k_{EC}). $k_{\text{EC}} = 0.45$ was used to calculate MBC.

2.3. Phospholipid Fatty Acids (PLFAs) Analysis

After freeze-drying, PLFA were extracted using a modified method of Bligh and Dyer [54–56]. Fatty acids were extracted from 2 g of soil by a single-phase mixture of chloroform-methanol-pH 4 citrate buffer (1:2:0.8, v/v/v) shaken at 300 rpm for 1.5 h and centrifuged 15 min at 1500 rpm. The supernatant was retained, and the soil was re-extracted as before. Phase splitting in the combined supernatants was obtained by adding citrate buffer and chloroform (overnight separation). The CHCl₃ layer was dried under N₂ at ambient temperature, re-dissolved with chloroform, and purified on silica cartridges (chromabond[®] 3 mL/500 mg SiOH, Macherey and Nagel, Düren, Germany). The methanolic fraction containing phospholipids was evaporated under N₂. Before analysis by gas chromatography-mass spectrometry (GC-MS), PLFAs had to be transformed into their less polar fatty acid methyl ester (FAME) derivatives. The derivatization was performed online in GC injector by tri methyl sulfonium hydroxide (TMSH) [57]. FAMEs were analyzed with GC/MS (4000 GC/MS, Varian, Inc., Walnut Creek, CA, USA) equipped with a BPX70 column (60 m, 0.25 mm i.d., 0.25 mm df., SGE). The FAME identification and quantification were performed using as standards: 37 component FAME mix from Supelco,

methyl nonadecanoate (19:0, used as internal standard) from Fluka and Br1 Mix, methyl-13-methyl tetradecanoate (i15:0), methyl 15-methyl hexadecanoate (i17:0), methyl vaccinate (18:1w7c), methyl cis-9, 10-methyleneoctadecanoate (cy19:0) from Larodan, Solna, Sweden. The PLFAs i14:0, i15:0, a15:0, i16:0, i17:0 and a17:0 were chosen to represent Gram-positive bacteria while $16:1\omega7c$, $17:1\omega7c$ and cy19:0 were used as an indicator of Gram-negative bacteria [58,59]. The PLFAs $18:1\omega9c$ and $18:2\omega6c$ were designated as representatives of fungi [56,60].

2.4. Ergosterol Analysis

Two extraction methods were used for the determination of ergosterol. A modified method of Montgomery et al. [61] was used for the alkaline ergosterol extraction (total ergosterol, i.e., free and esterified ergosterol). Briefly, 1 g of fresh soil was mixed with 8 mL of methanol and 2 mL of 2 M NaOH. The extraction was assisted by a microwave oven (MarsX[®],CEM, Mathews, NC, USA), under the following conditions: 2×15 s with 15 min cooling intermission (2450 MHz, 495 W output). After the treatment, the soils and solvents of each reactor were recovered by 2×4 mL of MeOH. For the step of liquid extraction, we added 3×4 mL of pentane, and each addition was shaken by vortex for 10 s and allowed to separate the 2 layers. The combined pentane extracts were evaporated and taken to dryness under a stream of N_2 gas. The residues were then made up to 2 mL in methanol, vortexed, and filtered at 0.2 µm nylon Acrodisc® filter (Gelman Sciences, Ann Arbour, MI, USA). The second ergosterol extraction method consisted of the non-alkaline ergosterol extraction (free ergosterol) using the Gong et al. [62] modified protocol. Briefly, 4 g of fresh soil was combined with acid-washed glass beads (4 g of 400–800 µm) and 6 mL methanol. After shaking (10 s and then shaken for 1 h at 320 rpm), the soil-methanol mixture was centrifuged (10 min at 8000 rpm) and the supernatant was then filtered (0.2 µm nylon Acrodisc[®] filter. For both extraction methods, 4 replicates were performed for each plot. Prior to HPLC quantification, ergosterol was separated by a RP using Lichrospher C18, connected to a Varian Prostar 230 pump and a Prostar 330 PAD detector, Merck KGaA, Darmstadt, Germany). Samples (20 µL) were injected and eluted (with 90% acetonitrile 10% MeOH at a flow rate of 1.2 mL mn⁻¹). Ergosterol was detected at 282 nm with a retention time of 3.2 min.

Ergosterol and PLFA could be both considered as bioindicators of fungal biomass but since they corresponded to different lipid classes, they could be affected variously by agricultural practices [63]. Moreover, the 18S rDNA amplification was also used to address the abundance of fungi in soils. These 3 methods might seem redundant, but since they each have their own strengths and weakness, they should be considered as complementary [64].

2.5. Total DNA Extraction and Quantification

Nucleic acids were extracted from 0.5 g of soil using a FastDNA SPIN Kit for soil (MP-Biomedicals, Santa Ana, CA, USA) according to the instructions of the manufacturer. Four replicates were performed for each plot. DNA was quantified by a fluorimetric measurement using Hoeschst fluorochrome at 360/460 nm excitation/emission wavelengths with a Fluorescent DNA Quantitation Kit (Biorad, Hercules, CA, USA).

2.6. Real-Time PCR Amplification

The 18S rDNA amplifications for fungal biomass estimation by 18S rDNA realtime qPCR were carried out with a total volume of 50 μ L. The qPCR mix was prepared as follows: 10 ng of soil microbial DNA, 25 pmol of each primer [FU18S1 5'-GGAAACTCACCAGGTCCAGA-3' and Nu-SSU-1536 5'-ATTGCAATGCYCTATCCCCA-3' [65]], 25 μ L of qPCR Mastermix for SYBR Green I Master mix (Roche, Basel, Switzerland) and 2.5 mg mL⁻¹ BSA (GeneON Bioscience, Ludwigshafen, Germany). A dilution series containing known amounts of *Fusarium graminearum* genomic DNA (assuming that it was representative of the total fungal genomes) was used as the standard for the quantification of the sample dsDNA. The amplification protocol (40 cycles of PCR, 20 s at 95 °C, 30 s at 62 °C, and 30 s at 72 °C) was performed using LightCycler 480 real-time PCR system (Roche, Basel, Switzerland). The efficiency of the qPCR ranged from 93% to 98%.

The 16S rDNA amplifications for bacterial biomass estimation by 16S rDNA realtime PCR were carried out under the same conditions of the 18S rDNA PCR except for the primers [63f 5'-CAGGCCTAACA CATGCAAGTC-3' [66] and BU16S4 5' CTGCTGCCTCCCGTAGG-3' derived from 341F [67]] and the amplification protocol (40 s at 95 °C, 45 s at 64 °C and 30 s at 72 °C). The efficiency of the qPCR ranged from 98% to 102%. A dilution series of Pseudomonas aeruginosa (assuming that it is representative of the bacterial genomes). DNA was used as the standard, and the results were expressed in μg of bacterial and or fungal dsDNA per µg of dry soil. Two independent qPCR assays were performed for the bacterial and fungal dsDNA estimation. Four replicates were performed for each plot.

2.7. Enzyme Assays

A total of 14 enzyme activities involved in C, N, P, and S nutrient cycling: Deshydrogenase (DEH) [68], Fluorescein diacetate hydrolase (FDA) [69], β -Glalactosidase (GAL) [70], β -Glucosidase (GLU) [49], Cellulase (CEL) [49], Laccase (LAC) [71], Lipase (LIP) [72], Xylanase (XYL) [73], Arylamidase (ARYLN) [74], N-acetylglusosaminidase (NAG) [49], Urease (URE) [70], Arylsulphatase (ARYLS) [70], Acid phosphatase (ACP), and Alkalin phosphatse (AKP) [49], were quantified by spectrophotometry, using commercial substrate analogs for each enzyme (Table 3). Four replicates were performed for each plot. Enzyme activities were expressed as an international enzyme unit (nmole of hydrolyzed or oxidized substrate min⁻¹ g drv soil⁻¹).

Table 3. Enzymes and substrates used for enzyme activity assays.

Enzymes	Abbreviations	E.C. Numbers	Substrates (Buffer pH and Concentration)
Global enzymatic activities	DELL		
Dehydrogenase	DEH	1.1.1.1	2,3,5 triphenyltetrazolium chloride (pH 7; 0.12 M)
Fluorescein diacetate hydrolase	FDA		Fluorescein diacetate (pH 7.6; 1000 μ g mL ⁻¹)
C cycle enzymatic activities			
β-galactosidase	GAL	3.2.1.23	<i>p</i> -NP-β-D-galactopyranoside (pH 7; 0.02 M)
β-glucosidase	GLU	3.2.1.21	p -NP- β -D-glucopyranoside (pH 6; 50 mM)
Cellulase	CEL	3.2.1.4	p -NP- β -D-cellobioside (pH 6; 10 mM)
Laccase	LAC	1.10.3.2	2,2'-azino-bis-(3 ethylbenzothiazoline-6-sulfonate) (pH 5; 0.5 mM)
Lipase	LIP	3.1.1.3	<i>p</i> -NP-palmitate (pH 6.5; 1 mM)
N-acetylglucosaminidase	NAG	3.2.1.30	<i>p</i> -NP-N-acetylglucosaminide (pH 6; 10 mM)
Xylanase	XYL	3.2.1.8	Xylan (pH 5.5, 12 g L^{-1})
N cycle enzymatic activities			
Arylamidase	ARYLN	3.4.11.2	L-leucine b-naphtylamide (pH 8; 2 mM)
Urease	URE	3.5.1.5	Urea (pH 7; 0.05 mM)
S cycle enzymatic activities			
Arylsulfatase	ARYLS	3.1.6.1	<i>p</i> -NP-sulfate (pH7; 25 mM)
P cycle enzymatic activities			
Acid phosphatase	ACP	3.1.3.2	<i>p</i> -NP-phosphate (pH 5; 50 mM)
Alkaline phosphatase	AKP	3.1.3.1	p-NP-phosphate (pH 9; 50 mM)

E.C: Enzyme Commission numbers.NP: nitrophenyl.

2.8. Statistical Data Analysis

Statistical differences among the plots on physicochemical parameters (total C, total N, pH_{water}, CEC, P₂O₅), microbial abundance (MBC, total, bacterial and fungal PLFAs, total and free ergosterol, total dsDNA, bacterial and fungal dsDNA) and enzyme activities (GAL, GLU, CEL, LAC, LIP, XYL, ARYLN, NAG, URE, ARYLS, ACP, AKP, DEH, and FDA) were compared by multiple comparisons of means using the non-parametric post hoc test ("nparcomp" package in R), with a significance level of p < 0.05. The calculations of the correlations between the different parameters were carried out using a Spearman's rank correlation procedure, and the significance level of these correlations were tested

using a pairwise *t*-test. To examine the relationships between different physicochemical characteristics and both microbial abundance and enzyme activities, multiple linear regression was carried out. Two principal component analyses (PCAs) were performed on the data of abundance of microbial communities and enzyme activities. For each PCA, we collected the coordinates on the first and second axes for each sample and calculated Euclidean distances. Euclidean distances were calculated using the permanent grassland (PG) as the reference of the more abundant and highest diversified microbial communities. The Euclidean distances are commonly used to analyze the magnitude of modification of abundance/composition and functional profiles of soil microbial communities [17,75–78]. The significance of the difference between plots was evaluated using the Kruskal–Wallis test. The calculations were performed using R [79].

3. Results

3.1. Soil Physicochemical Characteristics

The different plots of the field experiment showed the same texture of silty loam soil but showed different physicochemical characteristics (Table 4). Across the six plots, the total C and N contents ranged between 10.76–15.74 g kg⁻¹ DW _{soil} and 1.08–2.46 g kg⁻¹ DW _{soil}, respectively. The total C and N contents differed significantly between PG and CC plots; the highest values were obtained for PG plot when the lowest ones were obtained for CC plot. PG plot also differed from the other plots.

Table 4.	Physicoc	hemical o	characteristics	of p	lots.
----------	----------	-----------	-----------------	------	-------

Plots	Permanent Grassland	Restored Grasslands		Temporary Grasslands		Continuous Crop
	PG	RGI	RGII	тс	TG	CC
Total C (g kg ⁻¹)	$25.74 \pm (0.96)$ ^a	$11.11 \pm (0.34)$ ^d	$14.30 \pm (0.22)$ ^c	$13.89 \pm (0.53)$ ^c	$18.60 \pm (0.81)^{\text{ b}}$	$10.67 \pm (0.34)$ ^d
Total N ($g kg^{-1}$)	$2.46 \pm (0.07)$ ^a	$1.14 \pm (0.03)^{ m d}$	$1.41 \pm (0.02)$ ^c	$1.44\pm(0.06)$ c	$1.81 \pm (0.06)^{ ext{ b}}$	$1.08 \pm (0.04)$ ^d
C/N ratio	$10.47\pm(0.18)$ $^{\mathrm{a}}$	$9.76 \pm (0.06)^{ ext{ b}}$	$10.15 \pm (0.11)$ ^a	$9.61 \pm (0.18)$ ^b	$10.25 \pm (0.09)$ ^a	$9.85 \pm (0.13)^{ m b}$
Moisture (%)	$13.83 \pm (0.55)$ ^a	$10.53 \pm (0.29)$ ^c	$9.58 \pm (0.19)^{ m d}$	$10.45 \pm (0.25)^{\text{ b}}$	$11.78 \pm (0.56)^{\text{ b}}$	$9.35 \pm (0.25)^{ ext{ d}}$
pH _{water}	$5.46 \pm (0.05)^{ ext{ b}}$	$5.57 \pm (0.05)^{\text{ b}}$	$6.04 \pm (0.05)$ ^c	$6.27 \pm (0.05)^{a}$	$5.51 \pm (0.05)^{\text{ b}}$	$6.43 \pm (0.05)^{a}$
CEC (cmol ⁺ kg ⁻¹)	$8.08 \pm (0.26)$ ^a	$5.46 \pm (0.37)^{ ext{ d}}$	$6.92 \pm (0.31)^{ ext{ b}}$	$7.76 \pm (0.05)$ ^c	$7.29 \pm (0.20)^{ ext{ b}}$	$7.04 \pm (0.30)^{ ext{ b}}$
$P_2O_5 (g kg^{-1})$	$0.13\pm(0.03)~^{\mathrm{bc}}$	$0.09 \pm (0.01)^{\text{d}}$	$0.13\pm(0.01)~^{\rm c}$	$0.15\pm(0.01)$ ^{ab}	$0.19\pm(0.02)$ $^{\mathrm{ab}}$	$0.20\pm(0.02)$ $^{\rm a}$

Mean \pm (SD). Different letters indicate significant differences among plots (p < 0.05, n = 4).

Based on the total C and N contents, plots can be classified in the following order PG > TG > RGII \approx TC > RGI \approx CC. The C/N ratio of the six plots was roughly the same, with an average of ten. Concerning moisture, the mean value observed was about 10 g kg⁻¹ (Table 4). Moisture differed significantly between the PG plot, which was the most humid soil and CC plot, which was the less humid plot. PG and CC plots differed significantly from other plots. For pH_{water}, only a slight effect was observed, and three groups can be distinguished from less to greater acid plot: TC \approx CC > RGII > PG \approx RGI \approx TG. Cation Exchange Capacity (CEC) was higher for PG, while the lowest was obtained for RGI. The P₂O₅ was partly affected by land uses, the highest value was obtained for CC, which was not different from the temporary grasslands (TG and TC) while the lowest value was obtained for PG.

3.2. Microbial Communities' Abundance

For the study of microbial communities, we have chosen to analyze total, bacterial, and fungal abundances by using different and complementary methods through the analysis of MBC, cell-bound lipid (total, bacterial and fungal PLFAs), total, and free ergosterol and nucleic acids (total ds DNA, bacterial, and fungal ds DNA). The results are presented in Table 5. Considering all these parameters, the highest values were recorded for the PG plot, which was significantly different from the other plots, except for the free ergosterol content, which was not different between PG and TG. Conversely, the lowest values were observed for the CC and, in most cases, in the TC Plot.

Plots	Permanent Grassland	ent Grassland Restored Grasslands		Temporary	Continuous Crop	
Abundance of Microbial Communities	PG	RGI	RGII	тс	TG	CC
Total microbial biomass Total dsDNA (μ g g ⁻¹) Total microbial PLFA (nmol g ⁻¹) Microbial biomass carbon (mg C kg ⁻¹)	$\begin{array}{l} 25.82 \pm (4.37)^{\;a} \\ 717.05 \pm (70.2)^{\;a} \\ 987.78 \pm (88.4)^{\;a} \end{array}$	$\begin{array}{c} 12.22 \pm (3.24) \ ^{c} \\ 184.26 \pm (4.00) \ ^{c} \\ 250.52 \pm (13.4) \ ^{d} \end{array}$	$\begin{array}{c} 16.49 \pm (0.28) \ ^{\rm b} \\ 259.41 \pm (29.1) \ ^{\rm b} \\ 348.50 \pm (6.03) \ ^{\rm c} \end{array}$	$9.76 \pm (1.79)^{b}$ $102.66 \pm (31.3)^{d}$ $339.32 \pm (23.3)^{c}$	$\begin{array}{l} 15.70 \pm (3.62) \ ^{bc} \\ 262.33 \pm (84.5) \ ^{bcd} \\ 570.32 \pm (63.1) \ ^{b} \end{array}$	$\begin{array}{c} 13.24 \pm (3.11) \ ^{b} \\ 141.33 \pm (26.8) \ ^{d} \\ 234.39 \pm (16.8) \ ^{d} \end{array}$
Bacterial biomass Bacterial dsDNA ($\mu g g^{-1}$) Total bacterial PLFA (nmol g^{-1})	$6.91 \pm (1.70)^{a}$ 348.88 $\pm (29.7)^{a}$	$3.30 \pm (1.12)^{ m bc}$ 72.66 $\pm (0.82)^{ m c}$	$3.96 \pm (0.50)^{\text{ b}}$ $105.34 \pm (13.4)^{\text{ b}}$	$2.91 \pm (0.39)$ ^c $41.62 \pm (17.0)$ ^d	$3.28 \pm (1.7) {}^{ m abc}$ $110.76 \pm (41.7) {}^{ m bcd}$	$\begin{array}{l} 2.93 \pm (1.20) {}^{\rm bc} \\ 53.78 \pm (9.79) {}^{\rm d} \end{array}$
Fungal biomass Fungal dsDNA (μg g ⁻¹) Total fungal PLFA (nmol g ⁻¹) Total ergosterol (μg g ⁻¹) Free ergosterol (μg g ⁻¹)	$\begin{array}{c} 1.78 \pm (0.25)\ ^{a} \\ 22.68 \pm (5.21)\ ^{a} \\ 2.08 \pm (0.60)\ ^{a} \\ 1.28 \pm (0.23)\ ^{a} \end{array}$	$\begin{array}{l} 1.55 \pm (0.84) \ ^{abc} \\ 9.95 \pm (0.33) \ ^{c} \\ 0.99 \pm (0.22) \ ^{b} \\ 0.77 \pm (0.08) \ ^{b} \end{array}$	$\begin{array}{l} 1.34 \pm (0.19) \ ^{ab} \\ 13.10 \pm (1.57) \ ^{b} \\ 1.21 \pm (0.21) \ ^{b} \\ 0.75 \pm (0.45) \ ^{abc} \end{array}$	$\begin{array}{c} 0.65 \pm (0.12) \ ^{\rm c} \\ 1.95 \pm (0.72) \ ^{\rm e} \\ 0.82 \pm (0.06) \ ^{\rm b} \\ 0.48 \pm (0.04) \ ^{\rm c} \end{array}$	$\begin{array}{c} 1.13 \pm (0.58) \ ^{abc} \\ 12.04 \pm (4.44) \ ^{bc} \\ 3.12 \pm (0.68) \ ^{a} \\ 1.06 \pm (0.10) \ ^{a} \end{array}$	$\begin{array}{l} 0.84 \pm (0.29) \ ^{bc} \\ 5.12 \pm (1.10) \ ^{d} \\ 0.65 \pm (0.22) \ ^{b} \\ 0.51 \pm (0.04) \ ^{c} \end{array}$

Table 5. Total, bacterial, and fungal biomasses of plots.

Mean \pm (SD). Different letters indicate significant differences among plots (p < 0.05, n = 4).

To reduce the dimensionality of the data set, a PCA was performed to compare the plots (Figure 1). The first axis (PC1) and the second axis (PC2) explained 54.8% and 17.5% of the total variability, respectively. PCA clearly illustrated how the data set of the microbial communities classified the plots. The distribution of the plots along the first axis showed that the RGI, RGII, and TG plots were clustered and occupied an intermediate position between the two contrasted land uses. Indeed, this cluster was placed in the middle between grassland (PG) and croplands (CC and TC), which occupied the most extreme positions of this "PCA gradient".





3.3. Enzyme Activities

To evaluate the functions of microbial communities, 14 enzyme activities involved in different biogeochemical cycles were measured: for C cycle (GAL, GLU, CEL, LAC, LIP and XYL), for N cycle (ARYLN, NAG, and URE), for S cycle (ARYLS), for P cycle (ACP and AKP), and two global enzymes (DEH and FDA). The PG plot showed the highest values of all studied enzyme activities, and it was in most cases significantly different from the other plots except for AKP and GLU (Table 6). CC plot showed weak values of enzyme activities in comparison to the others plots. Overall, the enzyme activity values of the other treatments were intermediate and ranged between the PG and CC activity values. The pattern of variation of enzyme activities in the TG plot was similar to that of PG, while in

Plots	Permanent Grassland	Restored C	Grasslands	Temporary	Grasslands	Continuous Crop
Enzyme Activities (nmole min ⁻¹ g _{dry soil} ⁻¹)	PG	RGI	RGII	тс	TG	СС
Global enzyme						
DEH	$0.20 \pm (0.10)$ ^a	$0.04 \pm (0.02)$ ^b	$0.04 \pm (0.03)$ ^{bc}	$0.04 \pm (0.03)$ ^{bc}	$0.08 \pm (0.04)$ $^{ m ab}$	$0.00 \pm (0.01)$ ^c
FDA	$3.00 \pm (0.30)$ ^a	$1.56 \pm (0.04)$ ^b	$2.56\pm(0.54)$ ^a	$1.30\pm(0.84)$ ^b	$2.12\pm(0.46)$ ^a	$1.08 \pm (0.25)^{\text{ b}}$
C cycle enzymes						
GAL	$4.65 \pm (0.28)^{a}$	$2.07 \pm (0.28)^{b}$	$2.20 \pm (0.36)^{b}$	$1.48 \pm (0.12)$ ^c	$3.42 \pm (0.36)^{\text{d}}$	$1.35 \pm (0.16)$ ^e
GLU	$9.65 \pm (0.81)^{ m b}$	$3.36 \pm (0.52)^{e}$	$7.97 \pm (0.44)$ ^c	$5.94 \pm (0.67)^{ m d}$	$12.10 \pm (0.94)$ ^a	$5.36 \pm (0.65)^{\text{d}}$
CEL	$1.96 \pm (0.24)^{a}$	$0.77 \pm (0.23)^{\text{ b}}$	$0.94 \pm (0.14)$ ^b	$0.81 \pm (0.16)$ ^b	$1.75 \pm (0.26)^{a}$	$0.65 \pm (0.09)$ ^b
LAC	$8.03 \pm (1.35)$ ^a	$5.80 \pm (1.47)^{ab}$	$2.57 \pm (1.30)$ ^c	$1.41 \pm (0.58)$ ^c	$3.53 \pm (1.88)^{bc}$	$2.73 \pm (0.63)$ ^c
LIP	$2.82 \pm (0.26)^{a}$	$1.68 \pm (0.16)^{b}$	$1.46 \pm (0.32)^{\text{b}}$	$1.06 \pm (0.16)$ ^c	$1.72 \pm (0.17)^{\text{ b}}$	$0.74 \pm (0.19)$ ^c
XYL	$0.08 \pm (0.01)^{a}$	$0.01 \pm (0.01)$ ^c	$0.01 \pm (0.00)$ ^c	$0.02 \pm (0.01)^{ m bc}$	$0.04 \pm (0.01)^{\text{ b}}$	$0.00\pm(0.01)~^{\rm c}$
N cycle enzymes						
ARYLN	$3.94 \pm (0.39)$ ^a	$1.32 \pm (0.09)$ ^c	$1.66 \pm (0.08)$ ^b	$1.49 \pm (0.11)$ ^c	$1.90 \pm (0.32)$ ^b	$2.04 \pm (0.24)$ ^b
NAG	$5.78 \pm (0.72)^{a}$	$2.03 \pm (0.41)^{ m cd}$	$2.70 \pm (0.20)^{\text{d}}$	$1.97 \pm (0.64)$ ^{cd}	$4.20 \pm (0.46)^{b}$	$1.36 \pm (0.31)$ ^c
URE	$79.43 \pm (10.03)^{a}$	$25.84\pm(2.46)$ $^{\rm d}$	$34.63\pm(2.01)$ $^{\rm c}$	30.01 \pm (3.57) $^{\mathrm{cd}}$	54.71 \pm (3.32) ^b	21.18 \pm (0.36) $^{ m e}$
S cycle enzyme						
ARYLS	$7.27\pm(0.23)$ ^a	$2.70 \pm (0.11)^{\text{ b}}$	$3.32\pm(0.22)^{\rm c}$	$2.70 \pm (0.10)$ ^b	$5.48 \pm (0.33)^{\text{ d}}$	$1.83 \pm$ (0.24) $^{\rm e}$
P cycle enzymes						
ACP	$42.72 \pm (1.25)^{a}$	$10.73 \pm (7.23)^{\text{ e}}$	$20.86 \pm (1.92)$ ^c	$17.74 \pm (0.91)^{ m d}$	$30.60 \pm (2.29)^{b}$	$10.81 \pm (0.90)$ ^e
AKP	$4.30 \pm (1.13)^{\text{ c}}$	$2.65 \pm (0.39)$ ^d	$7.90 \pm (0.56)^{a}$	$3.52 \pm (1.11)^{\text{ cd}}$	$5.24 \pm (1.17)^{\rm \ cb}$	$6.15 \pm (0.48)$ ^b

most cases, enzyme activities values in the CC plot were very close to those observed in RGI and TC.

	Table 6.	Enzyme	activities	of plot	s.
--	----------	--------	------------	---------	----

Mean \pm (SD). Different letters indicate significant differences among plots (p < 0.05, n = 4).

The comparison of the enzyme activities among the plots using PCA is presented in Figure 2. The first axis (PC1) and the second axis (PC2) explained 73.2% and 11.6% of the total variability, respectively. The plots were ranked as follows: PG > TG > RGII > TC \approx RGI > CC based on the first axis. The distribution of the plots along the first axis reflects the duration in numbers of years cumulated under grassland. The RGI, TC, and CC plots were almost aggregated. Interestingly, the plots CC and TC were under culture at the time of sampling, while RGI was under a grassland regime but had a history of five years of continuous crop. Conversely, TC, which was closed to CC on the "PCA gradient", was under continuous crop at the time of sampling but after 14 years cumulated under grassland. In this Figure, the separation of the plots is slightly different in comparison to those observed in Figure 1 for abundance parameters of microbial communities.



Figure 2. Principal component analysis and correlation circle of plots enzyme activities.

3.4. Relationship between Soil Physicochemical Characteristics, Microbial Communities, and Enzyme Activities

To test which physicochemical characteristics (total C, total N, pH_{water} , CEC, P_2O_5) shape microbial community and enzyme activities, Spearman's rank correlation was used (Figure 3). The results showed that with the exception of bacterial dsDNA and fungal PLFA, all the variables were positively correlated with the total C (r \ge 0.58, p < 0.001) and N (r \geq 0.51, p < 0.01) contents and were negatively correlated with pH_{water} (r \geq -0.55, p < 0.001). There was significant correlation between the CEC and total biomass (r ≥ 0.66 , p < 0.0001) and bacterial biomass (r = 0.43, p < 0.01), while fungal PLFA was strongly negatively correlated with CEC (r ≥ -0.81 , p < 0.0001) and in lesser extent with P₂O₅ $(r \ge -0.48, p < 0.01)$. Concerning enzyme activities, except for LAC and AKP activities, all enzyme activities were positively correlated with the total C (r \geq 0.55, p < 0.001) and total N ($r \ge 0.51$, p < 0.001) contents (except for ARYLN and AKP). In most cases, enzyme activities were highly negatively correlated with pH_{water}, (r ≥ -0.53 , *p* < 0.001, except for ARYLN and AKP). A positive relationship was shown between enzyme activities and CEC $(r \ge 0.50, p < 0.01)$, except for GAL, LAC, LIP, and AKP. Beyond LIP and ARYLN activities, which were positively correlated to P_2O_5 , no correlations were shown between P_2O_5 and any of the measured enzymes.



Figure 3. Correlation matrix between physicochemical parameters, microbial abundance, and enzyme activities. Red and blue color indicate statistically significant negative and positive correlations, respectively.

We further found that, except for AKP activity and in most cases LAC activity, all enzymes involved in C, N, S, and P cycling were strongly correlated with each other. For instance, the GAL and CEL activities were strongly correlated with the NAG, URE, ARYLS, and ACP activities (Figure 3). Among all soil parameters, the strongest correlations were

obtained for C and N contents and microbial biomass carbon and NAG, XYL, URE ARYLS, and ACP activities ($r \ge 0.90$, p < 0.0001)

3.5. Relationships between Abundance of Microbial Communities and Enzyme Activities Levels Following the Introduction of Grassland into Crop Rotation

To explore the changes in abundance and functional profiles of microbial communities following the introduction of grassland into crop rotation, Euclidean distances were calculated as described in Section 2. These distances were used to assess the magnitude of the changes on the abundance of microbial communities and the functions profiles in comparison to the reference plot (PG) (Figure 4).



Figure 4. Changes on the levels of the abundance of microbial communities and enzymes activities expressed by Euclidean distance using the PG soil state as a reference state. Different letters indicate a significant difference between plots using Kruskal–Wallis test (p < 0.05, n = 4). Upper letters indicate a significant difference in the abundance profile, and lowercase letters indicate a significant difference in enzyme profile.

The highest the Euclidean distance is, the highest is the difference between two plots. For example, the large distances between PG and CC (distance \approx 10) for the functional profile indicated that the cultivation had a large impact on enzyme activities. The analyses of distances showed significant changes in the abundance and functional profiles of microbial communities between PG and TC plot and between PG and CC plot. While these profiles in TG are closer to those of PG. The plots TC and TG have cumulated 14 years in grassland, but they differ in the land use at sampling day because the TC plot was re-cultivated the year of sampling (2010). Euclidean distance results showed that cultivation led to a significant change in soil microbial communities. The changes in the abundance and functions between PG and RGI, as well as between PG and RGII, were intermediate between those observed for PG/TC and PG/TG (Figure 4). Correlation analysis showed a positive correlation (r = 0.74, *p* < 0.0001, Figure S2) between the changes of microbial abundance profile and the changes on enzyme activities profile when grassland is introduced into crop rotations.

4. Discussion

The restoration of the physicochemical properties and biological characteristics of soil has become of interest to preserve the soil state and promote sustainable management [80]. In accordance with our expectations, the sustainable soil management, including the implementation of grassland in crop rotations, influenced the soil properties, microbial communities, and enzyme activities, but the variations depended on the number of cumulative years under grasslands of each plot and the type of land use at the time of sampling.

4.1. Influence of Soil Characteristics in Shaping the Whole Microbial Community and *Enzyme Activities*

This study aimed to assess how agricultural managements influenced the role of soil characteristics in shaping the belowground microbial communities and their enzyme activities. Microbial communities were associated with soil physicochemical characteristics determined by C and N content and soil pH. Indeed, when physicochemical characteristics are considered together as a grouping factor (Table S1), they explained between 55% and 97% of the variation in microbial communities and between 64% and 98% of the variation in enzyme activities (p < 0.0001), respectively. In our study, C and N contents were in relation with the cumulative years under grassland, whereas soil pH appeared to be in relation with land use (grassland or cropland) at the sampling time (wheat crop in TC, CC, Figure S3). Overall, all groups of microorganisms (bacteria and fungi) assessed using different methods were strongly related to soil properties, which help explaining the differences observed between plots communities. This is unsurprising as many edaphic conditions are co-correlated, and several studies are now beginning to shed light on the composition of different microbial communities and relationships with edaphic properties across various soils [17,81,82]. The effects of land-use change on soil conditions, which drive microbial community composition, was particularly demonstrated through this study, which investigated different grassland regimes between continuous grassland and continuous crop. Continuous cropping without OM restitution is known to deplete soil C and N contents [22,83]. On the opposite, permanent grassland contributes to C storage through both reduced soil disturbance and slow degradation dynamics of OM [26,84,85], with positive effects on microbial communities [17,60]. More generally, shifts in soil organic status resulting from land use (plant residues amounts and qualities, root exudates, etc.) induce modifications in microbial communities [86], confirming the relationships between land use and soil biota [87]. Furthermore, a stronger effect of soil pH on the microbial communities was already observed by Bissett et al. [88] and Kuramae et al. [89]. In our study, soil pH was negatively correlated with all the variables that describe microbial communities. Combining soil analyses with soil microbial communities revealed that the microbial communities are non-randomly distributed across soils but determined by environmental factors. More specifically, soil properties, notably the C and N contents and pH, appeared to be the main drivers of soil microbial community composition and distribution at the European scale [86,90–92].

Soil enzymes have been reported as useful bioindicators of soil-state because they provide information on the soil's ability to perform biogeochemical reactions [93]. The measured enzyme activities are involved in the decomposition of different substrates with varying complexity related to the C, N, P, and S cycles [94]. The highest enzyme activities were recorded in PG plots and the lowest in CC plots, while intermediate activities were observed in the plots that represented intermediate situations. We investigated enzyme activities in connection with soil chemical properties. Overall, the majority of the enzyme activities were negatively correlated with the soil pH but positively correlated with C and N contents. Several studies showed the same relationships between C and N contents and enzymes activities [95–99] because C and N cycles were also strongly coupled in soils [100–103]. Concerning AKP activity, several authors showed a positive correlation between C and N content and AKP activity [104,105]. This was not the case in our study because AKP was not correlated with any of the soil characteristics. However, a large study conducted by Margalef et al. [106] showed that climatic conditions were the first predictive variables of AKP variation in soil and to a lesser extent N content and soil pH. The lack of correlation between LAC activity and C and N contents could be due to that LAC is strongly linked to OM quality than quantity [107]. This activity is also specifically produced following exposure to xenobiotics, unlike hydrolases, which are constitutively expressed [44,108]. Indeed, the mechanism by which soil microbial communities maintain homeostasis in a fluctuating resource environment is based on regulating enzyme expression to optimize the responsiveness of substrate acquisition [109,110], confirming that OM

is the most driver of enzyme activities [100,111,112]. In our study, enzymes activities were significantly affected by soil pH; this result agrees with several studies showing that pH is an important factor in describing enzyme activities across soils [113,114]. Various soil enzymes have different optimal soil pH ranges. This is not always reflected by the association between the enzyme activities and pH because soil pH can affect the enzyme activities indirectly by changing nutrient availability, the soil OM content, and the composition of the microbial communities [59,98,115,116].

4.2. Relationships between Microbial Communities and Soil Enzyme Activities

We further found that the enzymes involved in C, N, S, and P cycling were strongly correlated with each other (Figure 3), which reflected the close link between soil biogeochemical cycles [103,117]. The study conducted by Ullah et al. [99] showed notably that the activities of C-cycling enzymes had a direct positive relationship with the activities of N-cycling enzymes, and soil C availability increased N-cycling enzymes. Beyond the relationships showed between enzyme activities involved in the different biogeochemical cycles of C, N, P, and S, our study highlighted strong relationships between microbial communities' level and functional profile in soils. Indeed, the changes in the abundance of microbial communities were strongly linked to the changes in enzymes activities profile under the introduction of grassland into crop rotations. Furthermore, our study also confirmed the strong functional redundancy of soil microbial communities [99,118,119]. Indeed, several enzymes were linked to both bacterial and fungal abundance at the same time (Figure 3). This functional redundancy is common for basic mediated processes by hydrolases activities and far less redundant for processes mediated by oxidase activities (LAC). Indeed, LAC activity, which is involved in the mineralization of recalcitrant organic pollutants, was only linked to fungal biomass underling that "rare" or "less redundant" functions are mediated by specific groups of microorganisms [108,119]. In our study, the variation of AKP activity was not explained by any measured parameters, and it was not correlated to them. We suggested that this enzyme activity could be linked to stocking density when plots are in grasslands. Indeed, some authors explained that phosphatase activity increased with grazing intensity [120,121].

4.3. Transition of Microbial Abundance and Functional States Following the Introduction of Grassland into Crop Rotation

Several studies have emphasized the modification in microbial communities, and enzyme activities in soil under different land uses [17,122,123]. The analysis of the introduction of grassland into crop rotation expressed by Euclidean distances revealed that the cumulative years in grassland were the primary factor that determined the abundance and functional states of microbial communities. Permanent grassland and continuous crop are two types of land use that have distinct effects on the soil biota [83,85,124–126]. In our study, the intermediate conditions between these two contrasted land uses showed that the level of microbial abundance of TG tended to return to the level of grassland (PG). Indeed, excluding TC plot, which was cultivated at the time of sampling, Euclidean distances showed that microbial abundance and functional states tended to resiliency with an increasing number of years cumulated in grassland. This observation is consistent with those of Van Eekeren et al. [32], who found that soil OM, earthworm abundance, soil structure, and major functions of the soil biota are restored after crop conversion to grassland. In our study, the restoration of the abundance of microbial communities and enzyme activities at a level equivalent to grassland required more than five years without disturbance (TG). In the meantime, the restoration of the soil biota and its functions in the grassland phase was only temporary due to the following crop phase, which can explain the behavior of RGI and RGII. In other words, a continuous crop regime apparently imposes a stronger legacy on microbial communities and enzyme activities (RGI), in contrast with the after-effect of a grassland regime, which appears to be readily "labile" when a grassland regime is interrupted by cultivation (TC). Most of the studies giving information about the potential legacy's effects of grassland/cropland rotation systems on soil biophysicochemical

properties and yields showed controversial findings [83,85,124,127]. Such practices, linked to grassland or cropland phases, can have "positive or negative legacy" effects resulting from the interactions between several factors such as (i) soil properties, (ii) the duration and the applied management under the grassland phase, (iii) the management practices applied for crop phase [32,83,85,125,128,129]. While most studies have focused on the evaluation of soil biophysicochemical parameters when introducing grasslands into crop rotations, only few studies have addressed the effect of such practices on soil ecosystem services. Indeed, recent studies showed a stronger beneficial grassland legacy effect on soil structure maintenance and biodiversity conservation. By contrast, water and pathogen regulation and forage production were not affected by the legacy of grassland during the rotation [130,131].

Our results suggest that the introduction of grassland into crop rotations can restore the microbial communities and functional soil states. A continuous culture regime has a stronger legacy on the abundance of microbial communities (RGI). By contrast, the legacy of the grassland regime changes quickly when the grassland regime is interrupted by recent culture events (TC). A grassland regime enables resilience by the restoration of functions. However, as observed for TG, more than five cumulative years under grassland are required to observe this resilience. Our results highlighted the strong relationships between microbial communities and enzyme activities and the strong functional redundancy of microbial communities that could explain the observed resiliency. Finally, including grasslands in crop rotations has strong consequences for the microbial communities and the sustainability of their functionality. Such biological studies could help to optimize crop rotations (e.g., frequency and duration of temporary leys) for both agricultural productivity and ecosystem services. This work confirms the potential benefits offered by the introduction of grassland into cropping systems and could be largely used in the context of diversification of agricultural systems.

Supplementary Materials: The following are available online at https://www.mdpi.com/article/ 10.3390/agriculture11100909/s1, Figure S1: Plots positions in the area of the study (A) and sampling protocol (B), Figure S2: Correlation between microbial abundance and enzyme activities profiles, Figure S3: Principal component analysis and correlation circle of plots physicochemical, Table S1: Physicochemical characteristics explaining the variation of the abundance of microbial communities and enzyme activities by multiple regression analysis.

Author Contributions: W.R.-A.: experiment execution, presentation of results and original draft manuscript preparation; E.C.: data analysis; M.-P.N.: data analysis and contribution on the original draft manuscript preparation; R.C., N.C., S.C., and C.M.: data acquisition, contribution on the original draft manuscript preparation; review, and editing; M.L.: data acquisition and field study supervision; X.L.: review and editing; I.T.-G. and K.L.: Identification of the research topic, resources, review, editing and supervision. All authors have read and agreed to the published version of the manuscript.

Funding: This research is a part of the French national "Bioindicators Programme" that was funded by the French Agency for the Environment and Energy Management (ADEME). Grant numbers, N° 0875C0074.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Please find the site address to look for details concerning the program and data portal: https://ecobiosoil.univ-rennes1.fr/ademe-bioindicateur/english/db.php.

Acknowledgments: We would like to thank the members of the Bioindicator program: A. Bispo, L Galsomies and C. Grand (moderators from ADEME), G. Pérès (Université Rennes 1, UMR CNRS EcoBio and Agrocampus Ouest, UMR INRA SAS Rennes) for coordinating the program and L. Rougé (Université Rennes 1, UMR CNRS EcoBio) for coordinating the sampling campaign and for database management. The authors acknowledge the lycée agricole d'Yvetot. Biochem-Env

(https://doi.org/10.15454/HA6V6Y) is a platform of the Investment d'Avenir infrastructure AnaEE-France, overseen by the French National Research Agency (ANR-11-INBS-0001).

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Hamdy, A.; Aly, A. Land Degradation, Agriculture Productivity and Food Security. In Proceedings of the Fifth International Scientific Agricultural Symposium "Agrosym 2014", Jahorina, Bosnia and Herzegovina, 23–26 October 2014; pp. 708–717.
- Eswaran, H.; Lal, R.; Reich, P.F. Land degradation: An overview. In *Response to Land Degradation*; CRC Press: Boca Raton, FL, USA, 2001; ISBN 978-0-429-18795-7.
- Eisenhauer, N.; Barnes, A.D.; Cesarz, S.; Craven, D.; Ferlian, O.; Gottschall, F.; Hines, J.; Sendek, A.; Siebert, J.; Thakur, M.P.; et al. Biodiversity–Ecosystem Function Experiments Reveal the Mechanisms Underlying the Consequences of Biodiversity Change in Real World Ecosystems. J. Veg. Sci. 2016, 27, 1061–1070. [CrossRef]
- 4. Heijden, M.G.A.V.D.; Bardgett, R.D.; Straalen, N.M.V. The Unseen Majority: Soil Microbes as Drivers of Plant Diversity and Productivity in Terrestrial Ecosystems. *Ecol. Lett.* **2008**, *11*, 296–310. [CrossRef]
- Sogin, M.L.; Morrison, H.G.; Huber, J.A.; Welch, D.M.; Huse, S.M.; Neal, P.R.; Arrieta, J.M.; Herndl, G.J. Microbial Diversity in the Deep Sea and the Underexplored "Rare Biosphere". Proc. Natl. Acad. Sci. USA 2006, 103, 12115–12120. [CrossRef] [PubMed]
- Kirk, J.L.; Beaudette, L.A.; Hart, M.; Moutoglis, P.; Klironomos, J.N.; Lee, H.; Trevors, J.T. Methods of Studying Soil Microbial Diversity. J. Microbiol. Methods 2004, 58, 169–188. [CrossRef]
- Gans, J.; Wolinsky, M.; Dunbar, J. Computational Improvements Reveal Great Bacterial Diversity and High Metal Toxicity in Soil. Science 2005, 309, 1387–1390. [CrossRef]
- 8. Rusch, A.; Amend, J.P. Order-Specific 16S RRNA-Targeted Oligonucleotide Probes for (Hyper)Thermophilic Archaea and Bacteria. *Extremophiles* **2004**, *8*, 357–366. [CrossRef] [PubMed]
- 9. Nannipieri, P.; Ascher, J.; Ceccherini, M.T.; Landi, L.; Pietramellara, G.; Renella, G. Microbial Diversity and Soil Functions. *Eur. J. Soil Sci.* 2003, *54*, 655–670. [CrossRef]
- 10. Coonan, E.C.; Kirkby, C.A.; Kirkegaard, J.A.; Amidy, M.R.; Strong, C.L.; Richardson, A.E. Microorganisms and Nutrient Stoichiometry as Mediators of Soil Organic Matter Dynamics. *Nutr. Cycl. Agroecosyst.* **2020**, *117*, 273–298. [CrossRef]
- 11. Zak, D.R.; Holmes, W.E.; White, D.C.; Peacock, A.D.; Tilman, D. Plant Diversity, Soil Microbial Communities, and Ecosystem Function: Are There Any Links? *Ecology* 2003, *84*, 2042–2050. [CrossRef]
- 12. Chaparro, J.M.; Sheflin, A.M.; Manter, D.K.; Vivanco, J.M. Manipulating the Soil Microbiome to Increase Soil Health and Plant Fertility. *Biol. Fertil. Soils* 2012, *48*, 489–499. [CrossRef]
- 13. Salles, J.F.; van Elsas, J.D.; van Veen, J.A. Effect of Agricultural Management Regime on Burkholderia Community Structure in Soil. *Microb. Ecol.* **2006**, *52*, 267–279. [CrossRef]
- Singh, B.K.; Quince, C.; Macdonald, C.A.; Khachane, A.; Thomas, N.; Al-Soud, W.A.; Sørensen, S.J.; He, Z.; White, D.; Sinclair, A.; et al. Loss of Microbial Diversity in Soils Is Coincident with Reductions in Some Specialized Functions. *Environ. Microbiol.* 2014, 16, 2408–2420. [CrossRef] [PubMed]
- 15. Guo, Y.; Luo, H.; Wang, L.; Xu, M.; Wan, Y.; Chou, M.; Shi, P.; Wei, G. Multifunctionality and Microbial Communities in Agricultural Soils Regulate the Dynamics of a Soil-Borne Pathogen. *Plant Soil* **2021**, *461*, 309–322. [CrossRef]
- 16. Bardgett, R. The Biology of Soil: A Community and Ecosystem Approach; OUP Oxford: Oxford, UK, 2005; ISBN 978-0-19-852503-5.
- 17. Riah-Anglet, W.; Trinsoutrot-Gattin, I.; Martin-Laurent, F.; Laroche-Ajzenberg, E.; Norini, M.-P.; Latour, X.; Laval, K. Soil Microbial Community Structure and Function Relationships: A Heat Stress Experiment. *Appl. Soil Ecol.* **2015**, *86*, 121–130. [CrossRef]
- 18. Hooper, D.U.; Dukes, J.S. Overyielding among Plant Functional Groups in a Long-Term Experiment. *Ecol. Lett.* **2004**, *7*, 95–105. [CrossRef]
- 19. Balvanera, P.; Pfisterer, A.B.; Buchmann, N.; He, J.-S.; Nakashizuka, T.; Raffaelli, D.; Schmid, B. Quantifying the Evidence for Biodiversity Effects on Ecosystem Functioning and Services. *Ecol. Lett.* **2006**, *9*, 1146–1156. [CrossRef] [PubMed]
- 20. Skinner, C.; Gattinger, A.; Krauss, M.; Krause, H.-M.; Mayer, J.; van der Heijden, M.G.A.; Mäder, P. The Impact of Long-Term Organic Farming on Soil-Derived Greenhouse Gas Emissions. *Sci. Rep.* **2019**, *9*, 1702. [CrossRef]
- 21. Franzluebbers, A.J.; Lemaire, G.; de Faccio Carvalho, P.C.; Sulc, R.M.; Dedieu, B. Toward Agricultural Sustainability through Integrated Crop-Livestock Systems: Environmental Outcomes. *Agric. Ecosyst. Environ.* **2014**, *190*, 1–3. [CrossRef]
- 22. Lemaire, G.; Franzluebbers, A.; de Faccio Carvalho, P.C.; Dedieu, B. Integrated Crop–Livestock Systems: Strategies to Achieve Synergy between Agricultural Production and Environmental Quality. *Agric. Ecosyst. Environ.* **2014**, *190*, 4–8. [CrossRef]
- 23. Russelle, M.P.; Entz, M.H.; Franzluebbers, A.J. Reconsidering Integrated Crop–Livestock Systems in North America. *Agron. J.* **2007**, *99*, 325–334. [CrossRef]
- 24. Lemaire, G.; Gastal, F.; Franzluebbers, A.; Chabbi, A. Grassland–Cropping Rotations: An Avenue for Agricultural Diversification to Reconcile High Production with Environmental Quality. *Environ. Manag.* **2015**, *56*, 1065–1077. [CrossRef]
- 25. Parras-Alcántara, L.; Díaz-Jaimes, L.; Lozano-García, B. Management Effects on Soil Organic Carbon Stock in Mediterranean Open Rangelands—Treeless Grasslands. *Land Degrad. Dev.* **2015**, *26*, 22–34. [CrossRef]
- 26. Rumpel, C.; Crème, A.; Ngo, P.T.; Velásquez, G.; Mora, M.L.; Chabbi, A. The Impact of Grassland Management on Biogeochemical Cycles Involving Carbon, Nitrogen and Phosphorus. *J. Soil Sci. Plant Nutr.* **2015**, *15*, 353–371. [CrossRef]
- 27. Lal, R. Soil Carbon Sequestration to Mitigate Climate Change. Geoderma 2004, 123, 1–22. [CrossRef]

- 28. Ussiri, D.A.N.; Lal, R. Long-Term Tillage Effects on Soil Carbon Storage and Carbon Dioxide Emissions in Continuous Corn Cropping System from an Alfisol in Ohio. *Soil Tillage Res.* **2009**, *104*, 39–47. [CrossRef]
- Dass, P.; Houlton, B.Z.; Wang, Y.; Warlind, D. Grasslands May Be More Reliable Carbon Sinks than Forests in California. *Environ. Res. Lett.* 2018, 13, 074027. [CrossRef]
- 30. Baer, S.G.; Kitchen, D.J.; Blair, J.M.; Rice, C.W. Changes in Ecosystem Structure and Function Along a Chronosequence of Restored Grasslands. *Ecol. Appl.* **2002**, *12*, 1688–1701. [CrossRef]
- 31. Garbeva, P.; van Elsas, J.D.; van Veen, J.A. Rhizosphere Microbial Community and Its Response to Plant Species and Soil History. *Plant Soil* **2008**, *302*, 19–32. [CrossRef]
- van Eekeren, N.; Bommelé, L.; Bloem, J.; Schouten, T.; Rutgers, M.; de Goede, R.; Reheul, D.; Brussaard, L. Soil Biological Quality after 36 Years of Ley-Arable Cropping, Permanent Grassland and Permanent Arable Cropping. *Appl. Soil Ecol.* 2008, 40, 432–446.
 [CrossRef]
- 33. Blair, J.; Nippert, J.; Briggs, J. Grassland Ecology. In *Ecology and the Environment*; Monson, R.K., Ed.; The Plant Sciences; Springer: New York, NY, USA, 2014; pp. 389–423, ISBN 978-1-4614-7501-9.
- Guillou, C.L.; Prévost-Bouré, N.C.; Karimi, B.; Akkal-Corfini, N.; Dequiedt, S.; Nowak, V.; Terrat, S.; Menasseri-Aubry, S.; Viaud, V.; Maron, P.-A.; et al. Tillage Intensity and Pasture in Rotation Effectively Shape Soil Microbial Communities at a Landscape Scale. *MicrobiologyOpen* 2019, 8, e00676. [CrossRef]
- Jangid, K.; Williams, M.A.; Franzluebbers, A.J.; Sanderlin, J.S.; Reeves, J.H.; Jenkins, M.B.; Endale, D.M.; Coleman, D.C.; Whitman, W.B. Relative Impacts of Land-Use, Management Intensity and Fertilization upon Soil Microbial Community Structure in Agricultural Systems. Soil Biol. Biochem. 2008, 40, 2843–2853. [CrossRef]
- 36. Jangid, K.; Williams, M.A.; Franzluebbers, A.J.; Blair, J.M.; Coleman, D.C.; Whitman, W.B. Development of Soil Microbial Communities during Tallgrass Prairie Restoration. *Soil Biol. Biochem.* **2010**, *42*, 302–312. [CrossRef]
- McKinley, V.L.; Peacock, A.D.; White, D.C. Microbial Community PLFA and PHB Responses to Ecosystem Restoration in Tallgrass Prairie Soils. Soil Biol. Biochem. 2005, 37, 1946–1958. [CrossRef]
- Bach, E.M.; Baer, S.G.; Meyer, C.K.; Six, J. Soil Texture Affects Soil Microbial and Structural Recovery during Grassland Restoration. Soil Biol. Biochem. 2010, 42, 2182–2191. [CrossRef]
- 39. Brye, K.R.; Slaton, N.A.; Savin, M.C.; Norman, R.J.; Miller, D.M. Short-Term Effects of Land Leveling on Soil Physical Properties and Microbial Biomass. *Soil Sci. Soc. Am. J.* 2003, *67*, 1405–1417. [CrossRef]
- 40. Bossio, D.A.; Fleck, J.A.; Scow, K.M.; Fujii, R. Alteration of Soil Microbial Communities and Water Quality in Restored Wetlands. *Soil Biol. Biochem.* **2006**, *38*, 1223–1233. [CrossRef]
- 41. Plassart, P.; Vinceslas, M.A.; Gangneux, C.; Mercier, A.; Barray, S.; Laval, K. Molecular and Functional Responses of Soil Microbial Communities under Grassland Restoration. *Agric. Ecosyst. Environ.* **2008**, 127, 286–293. [CrossRef]
- 42. Attard, E.; Le Roux, X.; Charrier, X.; Delfosse, O.; Guillaumaud, N.; Lemaire, G.; Recous, S. Delayed and Asymmetric Responses of Soil C Pools and N Fluxes to Grassland/Cropland Conversions. *Soil Biol. Biochem.* **2016**, *97*, 31–39. [CrossRef]
- 43. Wang, S.; Wilkes, A.; Zhang, Z.; Chang, X.; Lang, R.; Wang, Y.; Niu, H. Management and Land Use Change Effects on Soil Carbon in Northern China's Grasslands: A Synthesis. *Agric. Ecosyst. Environ.* **2011**, *142*, 329–340. [CrossRef]
- Lebrun, J.D.; Trinsoutrot-Gattin, I.; Vinceslas-Akpa, M.; Bailleul, C.; Brault, A.; Mougin, C.; Laval, K. Assessing Impacts of Copper on Soil Enzyme Activities in Regard to Their Natural Spatiotemporal Variation under Long-Term Different Land Uses. *Soil Biol. Biochem.* 2012, 49, 150–156. [CrossRef]
- 45. Beaumelle, L.; Lamy, I.; Cheviron, N.; Hedde, M. Is There a Relationship between Earthworm Energy Reserves and Metal Availability after Exposure to Field-Contaminated Soils? *Environ. Pollut.* **2014**, *191*, 182–189. [CrossRef]
- 46. Torres-Leguizamon, M.; Mathieu, J.; Decaëns, T.; Dupont, L. Genetic Structure of Earthworm Populations at a Regional Scale: Inferences from Mitochondrial and Microsatellite Molecular Markers in Aporrectodea Icterica (Savigny 1826). *PLoS ONE* **2014**, *9*, e101597. [CrossRef]
- 47. Coulibaly, S.F.M.; Winck, B.R.; Akpa-Vinceslas, M.; Mignot, L.; Legras, M.; Forey, E.; Chauvat, M. Functional Assemblages of Collembola Determine Soil Microbial Communities and Associated Functions. *Front. Environ. Sci.* **2019**, *7*, 52. [CrossRef]
- Pérès, G.; Vandenbulcke, F.; Guernion, M.; Hedde, M.; Beguiristain, T.; Douay, F.; Houot, S.; Piron, D.; Richard, A.; Bispo, A.; et al. Earthworm Indicators as Tools for Soil Monitoring, Characterization and Risk Assessment. An Example from the National Bioindicator Programme (France). *Pedobiologia* 2011, 54, 77–87. [CrossRef]
- Trap, J.; Riah, W.; Akpa-Vinceslas, M.; Bailleul, C.; Laval, K.; Trinsoutrot-Gattin, I. Improved Effectiveness and Efficiency in Measuring Soil Enzymes as Universal Soil Quality Indicators Using Microplate Fluorimetry. *Soil Biol. Biochem.* 2012, 45, 98–101. [CrossRef]
- 50. Pauget, B.; Gimbert, F.; Coeurdassier, M.; Crini, N.; Pérès, G.; Faure, O.; Douay, F.; Hitmi, A.; Beguiristain, T.; Alaphilippe, A.; et al. Ranking Field Site Management Priorities According to Their Metal Transfer to Snails. *Ecol. Indic.* **2013**, *29*, 445–454. [CrossRef]
- 51. Cluzeau, D.; Pérès, G.; Guernion, M.; Chaussod, R.; Cortet, J.; Fargette, M.; Martin-Laurent, F.; Mateille, T.; Pernin, C.; Ponge, J.-F.; et al. Intégration de La Biodiversité Des Sols Dans Les Réseaux de Surveillance de La Qualité Des Sols: Exemple Du Programme Pilote à l'échelle Régionale, Le RMQS BioDiv. *Etude Gest. Sols* 2009, *16*, 187–201.
- 52. Wu, J.; Joergensen, R.G.; Pommerening, B.; Chaussod, R.; Brookes, P.C. Measurement of Soil Microbial Biomass C by Fumigation-Extraction—an Automated Procedure. *Soil Biol. Biochem.* **1990**, *22*, 1167–1169. [CrossRef]
- 53. Jenkinson, D.S.; Brookes, P.C.; Powlson, D.S. Measuring Soil Microbial Biomass. Soil Biol. Biochem. 2003, 36, 5–7. [CrossRef]

- 54. Bligh, E.G.; Dyer, W.J. A Rapid Method of Total Lipid Extraction and Purification. *Can. J. Biochem. Physiol.* **1959**, *37*, 911–917. [CrossRef]
- 55. Frostegård, Å.; Tunlid, A.; Bååth, E. Microbial Biomass Measured as Total Lipid Phosphate in Soils of Different Organic Content. *J. Methods* **1991**, *14*, 151–163. [CrossRef]
- Shahzad, T.; Chenu, C.; Repinçay, C.; Mougin, C.; Ollier, J.-L.; Fontaine, S. Plant Clipping Decelerates the Mineralization of Recalcitrant Soil Organic Matter under Multiple Grassland Species. *Soil Biol. Biochem.* 2012, 51, 73–80. [CrossRef]
- 57. Gómez-Brandón, M.; Lores, M.; Domínguez, J. Comparison of Extraction and Derivatization Methods for Fatty Acid Analysis in Solid Environmental Matrixes. *Anal. Bioanal. Chem.* **2008**, 392, 505–514. [CrossRef] [PubMed]
- 58. Wu, Y.; Ma, B.; Zhou, L.; Wang, H.; Xu, J.; Kemmitt, S.; Brookes, P.C. Changes in the Soil Microbial Community Structure with Latitude in Eastern China, Based on Phospholipid Fatty Acid Analysis. *Appl. Soil Ecol.* **2009**, *43*, 234–240. [CrossRef]
- Bowles, T.M.; Acosta-Martínez, V.; Calderón, F.; Jackson, L.E. Soil Enzyme Activities, Microbial Communities, and Carbon and Nitrogen Availability in Organic Agroecosystems across an Intensively-Managed Agricultural Landscape. *Soil Biol. Biochem.* 2014, 68, 252–262. [CrossRef]
- 60. Acosta-Martínez, V.; Mikha, M.M.; Vigil, M.F. Microbial Communities and Enzyme Activities in Soils under Alternative Crop Rotations Compared to Wheat–Fallow for the Central Great Plains. *Appl. Soil Ecol.* **2007**, *37*, 41–52. [CrossRef]
- 61. Montgomery, H.J.; Monreal, C.M.; Young, J.C.; Seifert, K.A. Determinination of Soil Fungal Biomass from Soil Ergosterol Analyses. *Soil Biol. Biochem.* **2000**, *32*, 1207–1217. [CrossRef]
- 62. Gong, P.; Guan, X.; Witter, E. A Rapid Method to Extract Ergosterol from Soil by Physical Disruption. *Appl. Soil Ecol.* 2001, 17, 285–289. [CrossRef]
- Gorka, S.; Canarini, A.; Imai, B.; Teischinger, G.; Darcy, S.; Kaiser, C. A unified protocol for the high-throughput measurement of PLFA, NLFA, GLFA and Sterols from Soil, 2020. In Proceedings of the EGU General Assembly 2020, EGU2020-17791, Online, 4–8 May 2020; p. 64.
- 64. Frostegård, Å.; Tunlid, A.; Bååth, E. Use and Misuse of PLFA Measurements in Soils. *Soil Biol. Biochem.* **2011**, 43, 1621–1625. [CrossRef]
- 65. Borneman, J.; Hartin, R.J. PCR Primers That Amplify Fungal RRNA Genes from Environmental Samples. *Appl. Environ. Microbiol* **2000**, *66*, 4356–4360. [CrossRef]
- Marchesi, J.R.; Sato, T.; Weightman, A.J.; Martin, T.A.; Fry, J.C.; Hiom, S.J.; Dymock, D.; Wade, W.G. Design and Evaluation of Useful Bacterium-Specific PCR Primers That Amplify Genes Coding for Bacterial 16S RRNA. *Appl. Environ. Microbiol* 1998, 64, 795–799. [CrossRef]
- 67. Muyzer, G.; de Waal, E.C.; Uitterlinden, A.G. Profiling of Complex Microbial Populations by Denaturing Gradient Gel Electrophoresis Analysis of Polymerase Chain Reaction-Amplified Genes Coding for 16S RRNA. *Appl. Environ. Microbiol* **1993**, *59*, 695–700. [CrossRef]
- 68. Schaefer, R. DEHYDROGENASE ACTIVITY AS A MEASURE OF THE GLOBAL BIOLOGICAL ACTIVITY OF SOIL. Ann. Inst. Pasteur (Paris) 1963, 105, 326–331.
- 69. Adam, G.; Duncan, H. Development of a Sensitive and Rapid Method for the Measurement of Total Microbial Activity Using Fluorescein Diacetate (FDA) in a Range of Soils. *Soil Biol. Biochem.* **2001**, *33*, 943–951. [CrossRef]
- de Santiago-Martín, A.; Cheviron, N.; Quintana, J.R.; González, C.; Lafuente, A.L.; Mougin, C. Metal Contamination Disturbs Biochemical and Microbial Properties of Calcareous Agricultural Soils of the Mediterranean Area. *Arch. Environ. Contam. Toxicol.* 2013, 64, 388–398. [CrossRef] [PubMed]
- 71. Schultz, A.; Jonas, U.; Hammer, E.; Schauer, F. Dehalogenation of Chlorinated Hydroxybiphenyls by Fungal Laccase. *Appl. Environ. Microbiol.* **2001**, *67*, 4377–4381. [CrossRef] [PubMed]
- 72. Gupta, R.; Gupta, N.; Rathi, P. Bacterial Lipases: An Overview of Production, Purification and Biochemical Properties. *Appl. Microbiol. Biotechnol.* **2004**, *64*, 763–781. [CrossRef]
- Schinner, F.; von Mersi, W. Xylanase-, CM-Cellulase- and Invertase Activity in Soil: An Improved Method. Soil Biol. Biochem. 1990, 22, 511–515. [CrossRef]
- 74. Acosta-Martínez, V.; Tabatabai, M.A. Enzyme Activities in a Limed Agricultural Soil. Biol. Fertil. Soils 2000, 31, 85–91. [CrossRef]
- 75. Ranjard, L.; Nowak, V.; Echairi, A.; Faloya, V.; Chaussod, R. The Dynamics of Soil Bacterial Community Structure in Response to Yearly Repeated Agricultural Copper Treatments. *Res. Microbiol.* **2008**, *159*, 251–254. [CrossRef]
- 76. Constancias, F.; Saby, N.P.A.; Terrat, S.; Dequiedt, S.; Horrigue, W.; Nowak, V.; Guillemin, J.-P.; Biju-Duval, L.; Prévost-Bouré, N.C.; Ranjard, L. Contrasting Spatial Patterns and Ecological Attributes of Soil Bacterial and Archaeal Taxa across a Landscape. *MicrobiologyOpen* 2015, 4, 518–531. [CrossRef] [PubMed]
- Carbonetto, B.; Rascovan, N.; Álvarez, R.; Mentaberry, A.; Vázquez, M.P. Structure, Composition and Metagenomic Profile of Soil Microbiomes Associated to Agricultural Land Use and Tillage Systems in Argentine Pampas. *PLoS ONE* 2014, 9, e99949. [CrossRef] [PubMed]
- 78. Riah-Anglet, W.; Trinsoutrot-Gattin, I.; Norini, M.-P.; Gauthier, A.; Latour, X.; Laval, K. Initial State of Soil Microbial Communities Determines Their Stress Response. *J. Environ. Chem. Eng.* **2018**, *6*, 5470–5480. [CrossRef]
- 79. Ihaka, R.; Gentleman, R.R.; Development Core Team. *R: A Language and Environment for Statistical Computing, 1996;* R Foundation for Statistical Computing: Vienna, Austria, 2009. Available online: http://www.R-project.org (accessed on 16 October 2009).
- 80. Lal, R. Restoring Soil Quality to Mitigate Soil Degradation. Sustainability 2015, 7, 5875–5895. [CrossRef]

- 81. Griffiths, R.I.; Thomson, B.C.; James, P.; Bell, T.; Bailey, M.; Whiteley, A.S. The Bacterial Biogeography of British Soils. *Environ. Microbiol.* **2011**, *13*, 1642–1654. [CrossRef]
- 82. Tian, Q.; Jiang, Y.; Tang, Y.; Wu, Y.; Tang, Z.; Liu, F. Soil PH and Organic Carbon Properties Drive Soil Bacterial Communities in Surface and Deep Layers Along an Elevational Gradient. *Front. Microbiol.* **2021**, *12*, 646124. [CrossRef] [PubMed]
- Panettieri, M.; Guigue, J.; Chemidlin Prevost-Bouré, N.; Thévenot, M.; Lévêque, J.; Le Guillou, C.; Maron, P.-A.; Santoni, A.-L.; Ranjard, L.; Mounier, S.; et al. Grassland-Cropland Rotation Cycles in Crop-Livestock Farming Systems Regulate Priming Effect Potential in Soils through Modulation of Microbial Communities, Composition of Soil Organic Matter and Abiotic Soil Properties. *Agric. Ecosyst. Environ.* 2020, 299, 106973. [CrossRef]
- Knops, J.M.H.; Tilman, D. Dynamics of Soil Nitrogen and Carbon Accumulation for 61 Years after Agricultural Abandonment. Ecology 2000, 81, 88–98. [CrossRef]
- Panettieri, M.; Rumpel, C.; Dignac, M.-F.; Chabbi, A. Does Grassland Introduction into Cropping Cycles Affect Carbon Dynamics through Changes of Allocation of Soil Organic Matter within Aggregate Fractions? *Sci. Total Environ.* 2017, 576, 251–263. [CrossRef]
- Lauber, C.L.; Strickland, M.S.; Bradford, M.A.; Fierer, N. The Influence of Soil Properties on the Structure of Bacterial and Fungal Communities across Land-Use Types. Soil Biol. Biochem. 2008, 40, 2407–2415. [CrossRef]
- Köhl, L.; Oehl, F.; van der Heijden, M.G.A. Agricultural Practices Indirectly Influence Plant Productivity and Ecosystem Services through Effects on Soil Biota. *Ecol. Appl.* 2014, 24, 1842–1853. [CrossRef]
- 88. Bissett, A.; Richardson, A.E.; Baker, G.; Thrall, P.H. Long-Term Land Use Effects on Soil Microbial Community Structure and Function. *Appl. Soil Ecol.* **2011**, *51*, 66–78. [CrossRef]
- 89. Kuramae, E.E.; Yergeau, E.; Wong, L.C.; Pijl, A.S.; van Veen, J.A.; Kowalchuk, G.A. Soil Characteristics More Strongly Influence Soil Bacterial Communities than Land-Use Type. *FEMS Microbiol. Ecol.* **2012**, *79*, 12–24. [CrossRef]
- Wakelin, S.A.; Macdonald, L.M.; Rogers, S.L.; Gregg, A.L.; Bolger, T.P.; Baldock, J.A. Habitat Selective Factors Influencing the Structural Composition and Functional Capacity of Microbial Communities in Agricultural Soils. *Soil Biol. Biochem.* 2008, 40, 803–813. [CrossRef]
- 91. Rousk, J.; Bååth, E.; Brookes, P.C.; Lauber, C.L.; Lozupone, C.; Caporaso, J.G.; Knight, R.; Fierer, N. Soil Bacterial and Fungal Communities across a PH Gradient in an Arable Soil. *ISME J.* **2010**, *4*, 1340–1351. [CrossRef]
- Plassart, P.; Prévost-Bouré, N.C.; Uroz, S.; Dequiedt, S.; Stone, D.; Creamer, R.; Griffiths, R.I.; Bailey, M.J.; Ranjard, L.; Lemanceau, P. Soil Parameters, Land Use, and Geographical Distance Drive Soil Bacterial Communities along a European Transect. *Sci. Rep.* 2019, *9*, 605. [CrossRef] [PubMed]
- 93. Almeida, F.; Wolf, J.M.; Casadevall, A. Virulence-Associated Enzymes of Cryptococcus Neoformans. *Eukaryot. Cell* 2015, 14, 1173–1185. [CrossRef] [PubMed]
- 94. Sinsabaugh, R.L.; Follstad Shah, J.J. Ecoenzymatic Stoichiometry and Ecological Theory. *Annu. Rev. Ecol. Evol. Syst.* 2012, 43, 313–343. [CrossRef]
- Sinsabaugh, R.L.; Gallo, M.E.; Lauber, C.; Waldrop, M.P.; Zak, D.R. Extracellular Enzyme Activities and Soil Organic Matter Dynamics for Northern Hardwood Forests Receiving Simulated Nitrogen Deposition. *Biogeochemistry* 2005, 75, 201–215. [CrossRef]
- 96. Piotrowska, A.; Wilczewski, E. Effects of Catch Crops Cultivated for Green Manure and Mineral Nitrogen Fertilization on Soil Enzyme Activities and Chemical Properties. *Geoderma* **2012**, *189–190*, *72–80*. [CrossRef]
- Cattaneo, F.; Di Gennaro, P.; Barbanti, L.; Giovannini, C.; Labra, M.; Moreno, B.; Benitez, E.; Marzadori, C. Perennial Energy Cropping Systems Affect Soil Enzyme Activities and Bacterial Community Structure in a South European Agricultural Area. *Appl. Soil Ecol.* 2014, *84*, 213–222. [CrossRef]
- 98. Hendriksen, N.B.; Creamer, R.E.; Stone, D.; Winding, A. Soil Exo-Enzyme Activities across Europe—The Influence of Climate, Land-Use and Soil Properties. *Appl. Soil Ecol.* **2016**, *97*, 44–48. [CrossRef]
- Ullah, S.; Ai, C.; Huang, S.; Zhang, J.; Jia, L.; Ma, J.; Zhou, W.; He, P. The Responses of Extracellular Enzyme Activities and Microbial Community Composition under Nitrogen Addition in an Upland Soil. *PLoS ONE* 2019, 14, e0223026. [CrossRef] [PubMed]
- Manzoni, S.; Jackson, R.B.; Trofymow, J.A.; Porporato, A. The Global Stoichiometry of Litter Nitrogen Mineralization. *Science* 2008, 321, 684–686. [CrossRef] [PubMed]
- 101. Kuzyakov, Y. Priming Effects: Interactions between Living and Dead Organic Matter. *Soil Biol. Biochem.* **2010**, *42*, 1363–1371. [CrossRef]
- 102. Soussana, J.-F.; Lemaire, G. Coupling Carbon and Nitrogen Cycles for Environmentally Sustainable Intensification of Grasslands and Crop-Livestock Systems. *Agric. Ecosyst. Environ.* **2014**, *190*, 9–17. [CrossRef]
- 103. Han, G.; Tang, Y.; Liu, M.; Van Zwieten, L.; Yang, X.; Yu, C.; Wang, H.; Song, Z. Carbon-Nitrogen Isotope Coupling of Soil Organic Matter in a Karst Region under Land Use Change, Southwest China. *Agric. Ecosyst. Environ.* 2020, 301, 107027. [CrossRef]
- 104. Aon, M.; Colaneri, A., II. Temporal and Spatial Evolution of Enzymatic Activities and Physico-Chemical Properties in an Agricultural Soil. *Appl. Soil Ecol.* 2001, *18*, 255–270. [CrossRef]
- Bonmati, M.; Ceccanti, B.; Nanniperi, P. Spatial Variability of Phosphatase, Urease, Protease, Organic Carbon and Total Nitrogen in Soil. *Soil Biol. Biochem.* 1991, 23, 391–396. [CrossRef]

- 106. Margalef, O.; Sardans, J.; Fernández-Martínez, M.; Molowny-Horas, R.; Janssens, I.A.; Ciais, P.; Goll, D.; Richter, A.; Obersteiner, M.; Asensio, D.; et al. Global Patterns of Phosphatase Activity in Natural Soils. *Sci. Rep.* 2017, 7, 1337. [CrossRef]
- 107. Qasemian, L.; Guiral, D.; Farnet, A.-M. How Do Microlocal Environmental Variations Affect Microbial Activities of a Pinus Halepensis Litter in a Mediterranean Coastal Area? *Sci. Total Environ.* **2014**, *496*, 198–205. [CrossRef] [PubMed]
- Lebrun, J.D.; Demont-Caulet, N.; Cheviron, N.; Laval, K.; Trinsoutrot-Gattin, I.; Mougin, C. Oxidoreductases Provide a More Generic Response to Metallic Stressors (Cu and Cd) than Hydrolases in Soil Fungi: New Ecotoxicological Insights. *Environ. Sci. Pollut. Res.* 2016, 23, 3036–3041. [CrossRef] [PubMed]
- Burns, R.G.; DeForest, J.L.; Marxsen, J.; Sinsabaugh, R.L.; Stromberger, M.E.; Wallenstein, M.D.; Weintraub, M.N.; Zoppini, A. Soil Enzymes in a Changing Environment: Current Knowledge and Future Directions. *Soil Biol. Biochem.* 2013, 58, 216–234. [CrossRef]
- 110. Sinsabaugh, R.L.; Belnap, J.; Findlay, S.G.; Shah, J.J.F.; Hill, B.H.; Kuehn, K.A.; Kuske, C.R.; Litvak, M.E.; Martinez, N.G.; Moorhead, D.L.; et al. Extracellular Enzyme Kinetics Scale with Resource Availability. *Biogeochemistry* 2014, 121, 287–304. [CrossRef]
- Kivlin, S.N.; Treseder, K.K. Soil Extracellular Enzyme Activities Correspond with Abiotic Factors More than Fungal Community Composition. *Biogeochemistry* 2014, 117, 23–37. [CrossRef]
- 112. Talbot, J.M.; Bruns, T.D.; Taylor, J.W.; Smith, D.P.; Branco, S.; Glassman, S.I.; Erlandson, S.; Vilgalys, R.; Liao, H.-L.; Smith, M.E.; et al. Endemism and Functional Convergence across the North American Soil Mycobiome. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 6341–6346. [CrossRef]
- 113. Cao, H.; Chen, R.; Wang, L.; Jiang, L.; Yang, F.; Zheng, S.; Wang, G.; Lin, X. Soil PH, Total Phosphorus, Climate and Distance Are the Major Factors Influencing Microbial Activity at a Regional Spatial Scale. *Sci. Rep.* **2016**, *6*, 25815. [CrossRef]
- 114. Shi, B.; Zhang, J.; Wang, C.; Ma, J.; Sun, W. Responses of Hydrolytic Enzyme Activities in Saline-Alkaline Soil to Mixed Inorganic and Organic Nitrogen Addition. *Sci. Rep.* **2018**, *8*, 4543. [CrossRef]
- 115. Sinsabaugh, R.L.; Lauber, C.L.; Weintraub, M.N.; Ahmed, B.; Allison, S.D.; Crenshaw, C.; Contosta, A.R.; Cusack, D.; Frey, S.; Gallo, M.E.; et al. Stoichiometry of Soil Enzyme Activity at Global Scale. *Ecol. Lett.* **2008**, *11*, 1252–1264. [CrossRef]
- 116. Boeddinghaus, R.S.; Nunan, N.; Berner, D.; Marhan, S.; Kandeler, E. Do General Spatial Relationships for Microbial Biomass and Soil Enzyme Activities Exist in Temperate Grassland Soils? *Soil Biol. Biochem.* **2015**, *88*, 430–440. [CrossRef]
- 117. Dai, X.; Zhou, W.; Liu, G.; Liang, G.; He, P.; Liu, Z. Soil C/N and PH Together as a Comprehensive Indicator for Evaluating the Effects of Organic Substitution Management in Subtropical Paddy Fields after Application of High-Quality Amendments. *Geoderma* 2019, 337, 1116–1125. [CrossRef]
- 118. Grządziel, J. Functional Redundancy of Soil Microbiota–Does More Always Mean Better? Pol. J. Soil Sci. 2017, 50, 75. [CrossRef]
- 119. Jia, Y.; Whalen, J.K. A New Perspective on Functional Redundancy and Phylogenetic Niche Conservatism in Soil Microbial Communities. *Pedosphere* 2020, 30, 18–24. [CrossRef]
- 120. Shariff, A.R.; Biondini, M.E.; Grygiel, C.E. Grazing Intensity Effects on Litter Decomposition and Soil Nitrogen Mineralization. *J. Range Manag.* **1994**, *47*, 444. [CrossRef]
- 121. Esch, E.H.; Hernández, D.L.; Pasari, J.R.; Kantor, R.S.G.; Selmants, P.C. Response of Soil Microbial Activity to Grazing, Nitrogen Deposition, and Exotic Cover in a Serpentine Grassland. *Plant Soil* **2013**, *366*, 671–682. [CrossRef]
- 122. Waldrop, M.P.; Balser, T.C.; Firestone, M.K. Linking Microbial Community Composition to Function in a Tropical Soil. *Soil Biol. Biochem.* **2000**, *32*, 1837–1846. [CrossRef]
- 123. Zhen, Z.; Wang, S.; Luo, S.; Ren, L.; Liang, Y.; Yang, R.; Li, Y.; Zhang, Y.; Deng, S.; Zou, L.; et al. Significant Impacts of Both Total Amount and Availability of Heavy Metals on the Functions and Assembly of Soil Microbial Communities in Different Land Use Patterns. *Front. Microbiol.* 2019, 10, 2293. [CrossRef] [PubMed]
- Postma-Blaauw, M.B.; de Goede, R.G.M.; Bloem, J.; Faber, J.H.; Brussaard, L. Soil Biota Community Structure and Abundance under Agricultural Intensification and Extensification. *Ecology* 2010, *91*, 460–473. [CrossRef] [PubMed]
- Crotty, F.V.; Fychan, R.; Sanderson, R.; Rhymes, J.R.; Bourdin, F.; Scullion, J.; Marley, C.L. Understanding the Legacy Effect of Previous Forage Crop and Tillage Management on Soil Biology, after Conversion to an Arable Crop Rotation. *Soil Biol. Biochem.* 2016, 103, 241–252. [CrossRef]
- 126. Crème, A.; Rumpel, C.; Le Roux, X.; Romian, A.; Lan, T.; Chabbi, A. Ley Grassland under Temperate Climate Had a Legacy Effect on Soil Organic Matter Quantity, Biogeochemical Signature and Microbial Activities. Soil Biol. Biochem. 2018, 122, 203–210. [CrossRef]
- 127. Sirimarco, X.; Barral, M.P.; Villarino, S.H.; Laterra, P. Water Regulation by Grasslands: A Global Meta-Analysis. *Ecohydrology* **2018**, *11*, 1934. [CrossRef]
- Christensen, B.T.; Rasmussen, J.; Eriksen, J.; Hansen, E.M. Soil Carbon Storage and Yields of Spring Barley Following Grass Leys of Different Age. *Eur. J. Agron.* 2009, 31, 29–35. [CrossRef]
- Carter, D.L.; Blair, J.M. High Richness and Dense Seeding Enhance Grassland Restoration Establishment but Have Little Effect on Drought Response. *Ecol. Appl.* 2012, 22, 1308–1319. [CrossRef] [PubMed]
- 130. Martin, G.; Durand, J.-L.; Duru, M.; Gastal, F.; Julier, B.; Litrico, I.; Louarn, G.; Médiène, S.; Moreau, D.; Valentin-Morison, M.; et al. Role of Ley Pastures in Tomorrow's Cropping Systems. A Review. *Agron. Sustain. Dev.* **2020**, *40*, 17. [CrossRef]
- Hoeffner, K.; Hotte, H.; Cluzeau, D.; Charrier, X.; Gastal, F.; Pérès, G. Effects of Temporary Grassland Introduction into Annual Crop Rotations and Nitrogen Fertilisation on Earthworm Communities and Forage Production. *Appl. Soil Ecol.* 2021, 162, 103893. [CrossRef]