



HAL
open science

Meloxicam administration in the management of postoperative pain and inflammation associated with caesarean section in beef heifers: Evaluation of reproductive parameters

Vincent Mauffré, Thomas Cardot, Guillaume Belbis, Vincent Plassard, Fabienne F. Constant, Sandrine Bernard, Nicolas Roch, Arnaud Bohy, Nicolas Nehlig, Andrew A. A. Ponter, et al.

► To cite this version:

Vincent Mauffré, Thomas Cardot, Guillaume Belbis, Vincent Plassard, Fabienne F. Constant, et al.. Meloxicam administration in the management of postoperative pain and inflammation associated with caesarean section in beef heifers: Evaluation of reproductive parameters. *Theriogenology*, 2021, 175, pp.148-154. 10.1016/j.theriogenology.2021.09.005 . hal-03357780

HAL Id: hal-03357780

<https://hal.inrae.fr/hal-03357780v1>

Submitted on 24 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



HAL
open science

Meloxicam administration in the management of postoperative pain and inflammation associated with caesarean section in beef heifers: Evaluation of reproductive parameters

Alexander Bolotin, Anne Aucouturier, Alexei Sorokin, Elena Bidnenko

► To cite this version:

Alexander Bolotin, Anne Aucouturier, Alexei Sorokin, Elena Bidnenko. Meloxicam administration in the management of postoperative pain and inflammation associated with caesarean section in beef heifers: Evaluation of reproductive parameters. *Theriogenology*, 2021, 175 (1), pp.148-154. 10.1016/j.theriogenology.2021.09.005 . hal-03343070

HAL Id: hal-03343070

<https://hal.inrae.fr/hal-03343070>

Submitted on 16 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

REVISED

1

2 **Meloxicam administration in the management of**
3 **postoperative pain and inflammation associated with**
4 **caesarean section in beef heifers: evaluation of**
5 **reproductive parameters**

6

7 Vincent Mauffré ^{a,b}, Thomas Cardot ^b, Guillaume Belbis ^b, Vincent Plassard ^b,
8 Fabienne Constant ^{a,b}, Sandrine Bernard ^c, Nicolas Roch ^c, Arnaud Bohy ^c, Nicolas
9 Nehlig ^d, Andrew Ponter ^{a,b}, Bénédicte Grimard ^{a,b}, Laurence Guilbert-Julien ^e

10

11 ^a Université Paris-Saclay, UVSQ, INRAE, BREED, 78350, Jouy-en-Josas, France

12 ^b Ecole Nationale Vétérinaire d'Alfort, 94700, Maisons-Alfort, France

13 ^c Groupement Technique Vétérinaire Bourgogne-Franche-Comté, GTV BFC, 10 la
14 Platière, 71150 Fontaines, France

15 ^d Boehringer Ingelheim Animal Health, 29 avenue Tony Garnier, 69007 Lyon, France

16 ^e Laboratoire National de Contrôle des Reproducteurs, 13 rue Jouet, 94700 Maisons-
17 Alfort, France

18

19 Corresponding author: Vincent Mauffré, e-mail address: vincent.mauffre@vet-alfort.fr

20

21 **Abstract**

22 Post-operative pain and inflammation are normal physiological reactions to
23 caesarean section. Their management in cattle have rarely been investigated. This
24 surgical procedure negatively affects reproductive function with, for example, a
25 reduction in fertility resulting in an increase in calving interval. In this multicenter
26 clinical trial, the objective was to evaluate the impact on reproductive performance of
27 meloxicam injected before caesarean section to manage post-operative pain and
28 inflammation. Meloxicam is a non-steroidal anti-inflammatory drug. One hundred and
29 twenty-seven Charolais heifers ($n = 127$) were recruited from 47 farms in six French

30 veterinary practices in the Burgundy region. The heifers underwent a non-elective
31 standardized caesarean section operation. Heifers were randomly assigned to one of
32 two groups: meloxicam ($n = 66$), intravenous meloxicam injection before surgery, or
33 control ($n = 61$). Reproductive performance and health information were recorded
34 from the time of the caesarean section to the next calving or to culling. In our study,
35 meloxicam administration before caesarean section had no effect on the incidence of
36 retained placenta (18.2% of treated vs 25.0% of control cows, $p=0.35$). The
37 pregnancy rate was higher in treated than in control cows (83.1% vs 67.8%, $p=0.04$
38 after multivariate analysis) and a survival analysis showed that the median calving
39 interval was 35 days shorter in the meloxicam ($t_{50\%}=417$ days) compared to the
40 control group ($t_{50\%}=452$ days, $p=0.05$). A trend was also observed for culling rate to
41 be lower in treated (4.7%) compared to control cows (13.3%, $p=0.09$). In conclusion,
42 this study suggests that there is a beneficial effect of meloxicam administration
43 before caesarean section on reproductive performance in Charolais heifers.
44

45 **Keywords**

46 Meloxicam; C-section; Non-Steroidal Anti-Inflammatory drug; reproduction; beef
47 heifers; pain management.
48

49 **1. Introduction**

50 The livestock industry needs to take it into consideration the increase in public
51 concern regarding animal welfare and should evaluate the benefits of new farming
52 practices [1]. Managing pain in farm animals is an integral part of animal welfare.
53 However, a major difficulty lies in the perception and the evaluation of pain in animals
54 [2,3]. Veterinarians and farmers generally agree on the nature and sources of pain in
55 ruminants, but there is less consensus concerning the perception of its intensity and
56 the need for its management [4]. A commission of the French National Research
57 Institute for Agriculture, Food and Environment (INRAE) recommended to minimize
58 animal pain in farms using a “3S” approach: “*suppress, substitute or soothe*” [5].
59 Although some painful procedures can be “*suppressed*” or “*substituted*”, there is
60 currently no alternative to caesarean section (C-section) in many cases of dystocia

61 (foetal-pelvic disproportion, uterine torsion, complicated breech presentation...). C-
62 section is more painful than natural delivery [6]. Despite being performed often on
63 farms, C-section leads to visceral and somatic pain which have consequences on
64 post-surgical recovery [7]. The pain is responsible for a decrease in physical activity
65 and is often associated with decreased in feed intake [6], exacerbating the early
66 lactation energy deficit frequently observed in the postpartum period, which often
67 results in a decrease in fertility [8-11].

68 The use of anti-inflammatory drugs, steroidal (AIS) or non-steroidal (NSAID), in
69 the peripartum period has rarely been studied and the few results available are often
70 inconclusive (for reviews [12,13]). Effectiveness would appear to depend on the
71 molecules used and the interval between the onset of the painful act (calving or C-
72 section) and treatment [14-16]. Flunixin meglumine (NSAID), administered within 24
73 hours of delivery [17] or during C-section [18], is associated with increased
74 prevalence of retained placenta. Conversely, the administration of carprofen (NSAID)
75 increased feed intake in the days following calving and increased long-term milk
76 production [19]. In another study, the use of meloxicam (NSAID) during C-section
77 decreased pain indicators [20]. In addition, calves born to dams treated with
78 meloxicam prior to C-section spent more time sucking and had higher serum
79 immunoglobulin G (IgG) levels, indicating better transfer of passive immunity (serum
80 IgG content > 15 g/L) [21].

81 Finally, a recent clinical study showed an improvement in reproductive
82 performance in animals when mastitis was treated with meloxicam [22]. Based on
83 this study a simulation model showed that the management of inflammation as an
84 integral part of mastitis treatment during the first 120 days postpartum could also
85 have economic benefits [23].

86 Based on these observations, managing both pain and inflammation during C-
87 section may be associated with an improvement in reproductive performance.
88 Therefore, in this study we aimed to investigate reproductive performance (retained
89 foetal membranes rate, pregnancy rate, calving interval, culling rate) in beef heifers
90 receiving a non-steroidal anti-inflammatory treatment (meloxicam) prior to C-section.

91 **2. Material and methods**

92 This field study was performed in the Burgundy region (France) from December
93 2015 to September 2017. All procedures carried out in the present study were
94 approved by the Ethical Committee in Clinical Research of the National Veterinary
95 School of Alfort (France) under protocol # 2018-12-07.

96

97 **2.1. Animals**

98 This study was conducted on Charolais heifers (beef cattle) and focused on non-
99 elective caesarean deliveries following dystocia. The animals included in the study
100 were about three years old, nulliparous and from farms with a standard calving
101 interval of less than 400 days and for which veterinary and animal husbandry records
102 were available. In addition, the heifers had to be free from bovine viral diarrhoea virus
103 (BVDV) and infectious bovine rhinotracheitis (IBR) virus, no known infertility problems
104 and with a body condition score (BCS) between 2 and 4 (scale 1-5). Exclusion criteria
105 were: C-sections that resulted in post-operative complications (uterine prolapse,
106 metritis, peritonitis) or peroperative anomalies (tearing of the uterus, failed surgical
107 procedure), since these factors are known to have a negative impact on fertility. A
108 total of 127 heifers from 47 farms were included in the study.

109 **2.2. Study design**

110 The objective of this multicenter field trial with randomized clinical cases was to
111 monitor the fertility of beef heifers after treatment with meloxicam (a NSAID, $n = 66$)
112 or control (without a NSAID, $n = 61$) in the management of pain and inflammation
113 during C-section. The surgery under farm conditions was performed by seventeen
114 veterinarian bovine obstetricians from six different veterinary clinics. For each
115 investigator, the two experimental groups were randomly assigned. At the time of
116 each C-section, the veterinarian opened an envelope indicating the group
117 (meloxicam or control group) and the surgical protocol to be followed. A control visit
118 was conducted the following day to check for placenta expulsion and to monitor the
119 general condition of the cow. After the C-section, natural mating was used to initiate a
120 new gestation. Cows were followed over a 520-day period at the end of which the
121 different parameters of interest were recorded (cf. 2.4 data).

122 **2.3. Surgery**

123 In order to limit bias, the surgical protocol was standardized using a consensus
124 on the technique to perform a bovine C-section published in 2007 by a French
125 technical veterinary association [24].

126 In the control group, heifers only received local anesthesia. In the meloxicam
127 group, in addition to local anesthesia, heifers received 0.5 mg meloxicam /kg
128 liveweight(Metacam®, Boehringer Ingelheim Animal Health France, Lyon, France)
129 intravenously (jugular vein) before beginning the surgery. For each procedure
130 (meloxicam and control groups), premedication was performed using 10 ml of
131 clenbuterol (Planipart®, Boehringer Ingelheim Animal Health France, Lyon, France)
132 to induce tocolysis to facilitate manipulation and exteriorization of the gravid uterine
133 horn. Procaine was used for local anesthesia (Procamidol®, Axience S.A.S., Pantin,
134 France) and the surgical approach was carried out on the left flank of a standing
135 animal. The flank had been previously sheared or shaved. The surgeon wore sterile
136 gloves and used sterilized equipment. After its exteriorization, the uterus was opened
137 on its large curve with a single-use scalpel. After calf removal, the uterus was sutured
138 with two separate continuous suture patterns, at least one of which was inverted
139 (Lembert or Cushing pattern), using a round needle and absorbable synthetic
140 threads. Before closing the uterus, 1 g of amoxicillin (intra-uterine bolus, Clamoxyl®,
141 Zoetis, Malakoff, France) was placed in the lumen of the uterus. Each veterinarian
142 closed the laparotomy incision using the technique that he/she was familiar with.
143 After surgery, a broad-spectrum antibiotic therapy (penicillin/dihydrostreptomycin)
144 was initiated for at least four days.

145 **2.4. Data**

146 Several parameters were recorded to evaluate reproductive performance: the
147 incidence of retained placenta (RP), pregnancy rate after surgery, calving interval
148 (calving in the year n+1, extracted from BDIVET, a French national database) and
149 culling rate. RP was defined as the presence of foetal membranes in the uterus by
150 visual or vaginal examination more than 24 hours after calving [25]. Pregnancy
151 diagnosis was performed by ultrasound (after 30 days post-mating) or by transrectal
152 palpation (after 60 days post-mating).

153 Parameters known to influence reproductive performance were also recorded:
154 season of heifer birth (autumn, winter, spring), age and body condition score (BCS)
155 at C-section, season of calving and surgery records (size of the uterine incision,
156 length of surgery defined as the time from the start of the preparation of the animal
157 for surgery - including premedication - to the end of the cutaneous suture, first uterine
158 suture pattern: puncturing or inverting).

159

160 **2.5. Statistical analyses**

161 RP and the first uterine suture pattern were defined as binary variables. Each
162 remaining quantitative variable was transformed into a qualitative variable. Three
163 classes of variable were created using arbitrarily thresholds in order to obtain a
164 relatively balanced distribution of animals among the classes (Table 1). Data were
165 entered in Microsoft Office Excel 2016 and imported into SAS® Studio 3.8 (SAS®
166 University Edition) and GraphPad Prism® software (version 9.0.0 for Windows,
167 GraphPad Software, San Diego, California USA, www.graphpad.com).

168

169 **2.5.1. Univariate analysis**

170 Meloxicam and control groups were compared using the Chi-square test for
171 qualitative variables (RP rate and culling rate).

172

173 **2.5.2. Multivariate analysis**

174 A treatment effect on pregnancy rate and calving interval was investigated using
175 multivariate models to take into account the variables known to influence the
176 reproductive performance of primiparous cows (BCS at C-section, birth period, age at
177 C-section, month of C-section, retained placenta, surgical technique). A treatment
178 effect on the incidence of retained placenta and culling rate could not be tested by
179 multivariate analysis because there were too few observations in the different classes
180 for these two variables.

181 Univariate analysis was performed to assess the relationship between
182 explicative variables, pregnancy rate and calving interval (CI) comparing the
183 percentage of pregnant cows or mean CI for the different levels of the explicative

184 variable (Chi square test for the first parameter, T test or ANOVA for the second). All
185 the variables associated with pregnancy rate at the threshold of 20% were introduced
186 in the multivariate logistic regression models together with the treatment effect
187 (GLIMMIX procedure of SAS® Studio). A backward stepwise elimination of non-
188 associated ($p>0.10$) variables was performed to develop the models. The model with
189 the lowest Akaike's Information Criterion was retained.

190 The same approach was used to investigate the association between
191 explicative variables and CI in linear multivariate models (MIXED procedure of SAS®
192 Studio).

193

194 **2.5.3. Survival analysis**

195 A survival analysis was performed to investigate the interval between C-section
196 and the next calving or culling. The advantage of this analysis was that it included all
197 cows that were involved in the study, including those that did not calve after
198 breeding.

199 The estimation of the survival functions was carried out using the Kaplan-Meier
200 method. The log-rank test was used to compare the two survival curves.

201

202 **3. Results**

203 **3.1. Descriptive analysis**

204 In the present study, the predominant causes of dystocia were foetomaternal
205 disproportion and incomplete dilation of the vulva or cervix. The other causes were
206 irreducible uterine torsion (one case) and a bad foetal position that could not be
207 corrected by obstetrical manipulation (two breech presentations and three
208 uncomplicated posterior presentations).

209 The main characteristics of our sample are listed in Table 1. The variables
210 recorded before surgery («heifer birth season», «age and body condition score C-
211 section», «C-section period») and during surgery («size of the uterine incision»,
212 «length of surgery», «first uterine suture pattern») were not different between the
213 meloxicam and control groups.

214

215 **3.2. Meloxicam effect on reproductive performance**

216 Of the 127 heifers included in the study, 126 were used to examine the variable
217 «retained placenta» (one cow died after the intervention), 124 for «pregnancy rate»
218 (two cows culled before breeding) and 124 for «culling rate» (three died during the
219 study). The prevalence of RP after C-section was 21.4% (27/126), the pregnancy rate
220 was 75.8% (94/124), the calving interval (CI) was 412±40 days (mean±sd) and the
221 culling rate was 9.7% (12/124).

222 The main effects of meloxicam administration before C-section on reproductive
223 performance are summarized in Table 2. The incidence of retained placenta was not
224 significantly different between the meloxicam and the control group. The culling rate
225 tended ($p=0.09$) to be lower in the meloxicam compared to the control group.

226 In the multivariate analysis, six variables were related to pregnancy rate at the
227 threshold of 20% («treatment», «heifer birth season», «C-section period», «BCS»,
228 «size of the uterine incision», «length of surgery»). In the best multivariate model,
229 pregnancy rate was higher ($p<0.05$) in the heifers that received meloxicam prior to C-
230 section compared to controls. Three variables were related to CI at the threshold of
231 20% («treatment», « heifer birth season», «age at C-section»). After bias correction
232 by multivariate analysis, treatment effect on CI was not significant (meloxicam,
233 406.6±6.7 vs control, 417.5±8.0, $p=0.20$).

234 Finally, figure 1 shows the Kaplan-Meier survival plot (days from C- section to
235 next calving) for heifers receiving meloxicam or control before surgery. The median
236 survival time (i.e. median calving interval) was 35 days shorter in the treatment group
237 ($t_{50\%}=417$ days) compared to the control group ($t_{50\%}=452$ days, $p=0.05$).

239 **4. Discussion**

240 The objective of this field trial was to evaluate the impact of pre-C-section
241 administration of a NSAID (meloxicam), in pain and inflammation management, on
242 the fertility of Charolais beef heifers. The effects of meloxicam administration
243 preceding C-section were evaluated by comparison with a group of control animals
244 that only received the local anesthesia.

245 This study was conducted during the 2016 breeding season on Charolais
246 heifers in the Burgundy region, one of the main breeding areas for this breed in

247 France [26]. Meloxicam and control groups showed similar results for parameters
248 such as «heifer birth season», «age and body condition at C-section», and «C-
249 section period» (Table 1). These results are consistent with the data collected across
250 France for Charolais heifers during the same period. In our study, age at first calving
251 (C-section) was between 31 and 39 months old and represented 84% of calvings for
252 this breed [26]. Two calving periods are usually described in Charolais heifers:
253 «autumn calving» and «winter calving» [26]. Since C-sections took place from
254 December to May, this study focused on «winter calving». This calving period is
255 known to be associated with lower reproductive performance due to a delay in the
256 resumption of ovarian cyclicity 60 to 70 days after parturition (23-65% of cows with
257 normal ovarian cyclicity) compared to «autumn calving» cows (70-80% of cows with
258 normal ovarian cyclicity) [27-30]. At first calving, the recommended body condition
259 score (BCS) for beef heifers is 2.75 to 3 (scale 1-5) [27,31]. Heifers with BCS of 2.5
260 or less at calving are more likely to experience a delay in the resumption of ovarian
261 cyclicity, related to negative energy balance. This results in a prolonged calving
262 interval [32]. In comparison, heifers with a high BCS (>3.5) at calving have an
263 increased risk of dystocia [31]. In our study, dystocia was used as an inclusion
264 criterion necessitating a C-section. With less than 15% of the heifers having a BCS of
265 2.5 or less (scale 1-5), it can therefore be assumed that the impact of negative
266 energy balance on our data is limited.

267 In 2010, a survey of 710 bovine veterinary practitioners in Europe listed the
268 differences in performing a C-section [33,34]. In order to avoid potential biases
269 related to the use of different surgical techniques in our study, the surgical protocol
270 was standardized according to the recommendations of the French technical
271 veterinary association [24]. As a result of procedure standardization, the only
272 variation factors related to surgery were: «the size of the uterine incision», «the
273 nature of the first uterine suture pattern» and «the length of the procedure». The form
274 of the first uterine suture pattern was not dictated in the surgical procedure, but the
275 majority of veterinarians (>85%) chose a simple continuous suture pattern. The
276 average length of surgery was 31 ± 7 minutes, which is much shorter than the average
277 time (54 ± 12 minutes) reported in Europe in 2010 [34]. In the geographical area of our
278 study, C-section is a very common practice for heifers: veterinarians routinely
279 perform C-sections and farmers have become accustomed to preparing the heifer

280 (restraint, hair removal, washing of the surgical area). This preparation time, which
281 can represent 30% of the total time for the procedure [34], was not included in the
282 present study, which may explain the shorter length for the procedure observed in the
283 present experiment. Finally, no difference was observed for these variation factors
284 between the two experimental groups.

285 Following C-section, the animals were monitored over a 520-day period during
286 which reproductive performance (retained placenta rate, gestation rate, calving rate,
287 calving interval and culling rate) was recorded. C-section is known to impair fertility,
288 resulting in a lower conception and calving rates, and increased calving interval.
289 Compared to a normal calving population, the decrease in pregnancy rates due to C-
290 section ranged from 15% to 27% [35,36]: previous studies reported conception rates
291 of 48-80% [10,36,37] and calving rates of 41-52% [35,38]. The higher pregnancy rate
292 (75.8%) observed in this experiment may be explained by the fact that the animals in
293 our study were heifers intended for replacement when the other studies involved
294 animals with varying parities including multiparous cows, whose fertility is known to
295 be lower compared to heifers.

296 A reduction in fertility results in a longer calving interval. In the present study,
297 the mean calving interval for the Charolais primiparous cows after C-section was
298 412 ± 40 days. These data are consistent with the results of a study which analyzed
299 111,871 calvings and reported a calving interval of 426 days for Charolais cows with
300 C-section [38]. In contrast, in France, for the same breeding season (2016), the
301 calving interval in the Charolais breed was reported to be 396 ± 60 days [26]. C-
302 section is associated with an increase in the incidence of retained placenta and
303 culling rate for infertility. In the present experiment, the overall incidence of retained
304 placenta was 21% which is higher than the incidence encountered in elective
305 caesarean section in Belgian Blue cows (3.5%) [39] but consistent with the incidence
306 reported (26-35%) in other studies [9,40-42]. The overall culling rate for infertility in
307 the present study was 9.7%, similar to that previously reported in beef cattle (9%)
308 [40].

309 The aim of anti-inflammatory treatment with meloxicam before C-section is to
310 reduce inflammation in the reproductive tract following surgery and to decrease the
311 pain associated with the surgery. The mechanisms of pain go well beyond
312 nociception and include cortical integration of negative emotions associated with pain

313 as well as complex interplay of excitatory and inhibitory pathways [43]. Due to the
314 complexity of the pain network and its multiphase kinetics, pain management
315 requires multimodal analgesia, i.e. the use of complementary molecules capable of
316 combating different aspects of pain genesis [43,44]. Local anesthetics are powerful
317 molecules for suppressing nociception and as such are commonly used in surgery
318 [44,45]. However, after a single injection, their short-term action makes them
319 unsuitable to alleviate post-operative pain and they are ineffective in preventing the
320 development of post-incisional inflammation and the resulting peripheral hyperalgesia
321 [43]. Opioids and NSAIDs are effective analgesics, particularly for post-operative
322 pain. While opioids powerfully strengthen inhibitory pathways in the central nervous
323 system, NSAIDs suppress inflammation-induced peripheral hyperalgesia due to
324 nociceptor sensitization by inflammation mediators [44,45]. However, C-section
325 analgesia still often only includes the use of local anesthetics [46], this does not
326 achieve the goals of pain management. In human medicine, caesarean analgesia
327 usually includes combinations of local anesthetics, NSAIDs and opioids [47,48].
328 Opioids are not approved for use in food-producing animals, but meloxicam is a long-
329 acting NSAID approved in the European Union for use in livestock. Finally, the pre-
330 emptive use of local anesthetics and meloxicam in combination has been shown to
331 be effective in reducing the pain and distress in cattle associated with dehorning,
332 castration [45,46,49] and C-section [20].

333 Despite these observations on pain management, the contrasting results of
334 studies investigating the impact on fertility of the administration of anti-inflammatory
335 drugs around calving have long hindered their use in obstetrics. For instance,
336 ketoprofen administered just after or within 24 hours of calving had no effect on
337 fertility [50]. Similar results were observed with carprofen administered within three
338 weeks of parturition [16]. Studies on flunixin meglumine given after calving even
339 reported an increase in postpartum disorders (retained placenta, metritis) that
340 affected reproductive performance [17,18]. These studies differ from the present
341 study because of the type of drug used and the timing of the NSAID administration in
342 relation to the onset of the painful procedure. It would appear that the beneficial
343 effects are more pronounced when analgesia is implemented prior to the surgery
344 [51], which was the case in our study. Our data show that meloxicam prior to surgery
345 increased pregnancy rate by 15% and shortened median survival time (i.e. median

346 calving interval) by 35 days compared to control. The survival analysis, contrarily to
347 the comparison of CI averages, takes into account the cows which did not calve after
348 breeding and shows that more cows were likely to calve in a shorter time period after
349 C-section in the meloxicam than the control group. These positive effects on
350 reproduction may be related to peroperative pain management, which results in
351 improved comfort in animals after C-section [20] and better food intake in the days
352 following surgery [19], thus limiting the adverse effects on reproduction [8].

353 Furthermore, while some studies reported an increased incidence of RP when
354 using flunixin meglumine during the peripartum period [17,18], previous experiments
355 using meloxicam did not appear to have a similar effect on the incidence RP [52,53].
356 The present experiment confirmed the results of the latter studies.

357 Finally, among the long-term benefits, a decreased risk of culling for infertility
358 was reported when using meloxicam at calving [54] and flunixin during C-section [9].
359 This helps to reduce cow replacement rate and thus improves the longevity of the
360 animals. Although the difference is not significant, the culling rate in our study was
361 lower in the meloxicam group.

362

363 **5. Conclusions**

364 Our study suggests that administering meloxicam before C-section in order to
365 limit pain and inflammation does not compromise subsequent reproductive
366 performance in beef heifers. On the contrary, the results of the present study indicate
367 that the administration of meloxicam prior to C-section is associated with increased
368 pregnancy rate and tends to shorten the calving interval, with no increase in the risk
369 of retained placenta. These results are consistent with existing data. However, further
370 studies are required to confirm these using a larger group of animals and other
371 breeds, both beef and dairy cows.

372

373 **Source of funding**

374 This study was funded by Boehringer Ingelheim Animal Health (Lyon, France) and
375 the French technical veterinary association, section Bourgogne-Franche-Comté

376 (GTVR-BFC, Groupement Technique Vétérinaire Régional – Bourgogne-Franche-
377 Comté, Fontaines, France).

378 **Authors' contributions**

379 **Vincent Mauffré:** formal analysis, supervision, validation, writing – original draft,

380 **Thomas Cardot:** investigation, formal analysis, visualization

381 **Guillaume Belbis:** conceptualization, validation

382 **Vincent Plassard:** writing – review & editing

383 **Fabienne Constant:** writing – review & editing

384 **Sandrine Bernard:** resources

385 **Nicolas Roch:** conceptualization, funding acquisition

386 **Arnaud Bohy:** conceptualization, funding acquisition

387 **Nicolas Nehlig:** resources

388 **Andrew Ponter:** writing – review & editing

389 **Bénédicte Grimard:** formal analysis, writing – review & editing

390 **Laurence Guilbert-Julien:** conceptualization, funding acquisition, supervision,
391 validation, writing – review & editing

392 **Declaration of interest**

393 The study was funded by Boehringer Ingelheim Animal Health (Lyon, France).

394 **Acknowledgements**

395 The authors would like to thank the Boehringer Ingelheim Company, for making this
396 study possible, as well as the members of the GTV Bourgogne-Franche-Comté and
397 the farmers involved for their participation and implication in this field study. The
398 authors would also like to thank Fanny Pilot-Stork (Ecole Nationale Vétérinaire
399 d'Alfort, France) for her expertise on physiology of pain and its management.

400 **References**

401 [1] Ohl F, van der Staay FJ. Animal welfare: At the interface between science and
402 society. *The Veterinary Journal* 2012;192:13–9.
403 <https://doi.org/10.1016/j.tvjl.2011.05.019>.

- 404 [2] Livingston A. Pain and Analgesia in Domestic Animals. In: Cunningham F, Elliott
405 J, Lees P, editors. *Comparative and Veterinary Pharmacology*, Berlin, Heidelberg:
406 Springer Berlin Heidelberg; 2010, p. 159–89. [https://doi.org/10.1007/978-3-642-](https://doi.org/10.1007/978-3-642-10324-7_7)
407 [10324-7_7](https://doi.org/10.1007/978-3-642-10324-7_7).
- 408 [3] Mellor DJ, Stafford KJ. Physiological and behavioural assessment of pain in
409 ruminants: principles and caveats. *Altern Lab Anim* 2004;32 Suppl 1A:267–71.
- 410 [4] Thomsen PT, Anneberg I, Herskin MS. Differences in attitudes of farmers and
411 veterinarians towards pain in dairy cows. *The Veterinary Journal* 2012;194:94–7.
412 <https://doi.org/10.1016/j.tvjl.2012.02.025>.
- 413 [5] Guatteo R, Levionnois O, Fournier D, Guémené D, Latouche K, Leterrier C, et al.
414 Minimising pain in farm animals: the 3S approach – ‘Suppress, Substitute,
415 Soothe.’ *Animal* 2012;6:1261–74. <https://doi.org/10.1017/s1751731112000262>.
- 416 [6] Kolkman, Aerts, Vervaecke H, Vicca J, Vandelook J, De Kruif A, et al.
417 Assessment of differences in some indicators of pain in double muscled belgian
418 blue cows following naturally calving vs caesarean section. *Reproduction in*
419 *Domestic Animals* 2010;45:160–7. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0531.2008.01295.x)
420 [0531.2008.01295.x](https://doi.org/10.1111/j.1439-0531.2008.01295.x).
- 421 [7] Walker KA, Duffield TF, Weary DM. Identifying and preventing pain during and
422 after surgery in farm animals. *Applied Animal Behaviour Science* 2011;135:259–
423 65. <https://doi.org/10.1016/j.applanim.2011.10.021>.
- 424 [8] Roche JR, Burke CR, Crookenden MA, Heiser A, Loor JL, Meier S, et al. Fertility
425 and the transition dairy cow. *Reprod Fertil Dev* 2018;30:85–100.
- 426 [9] Lyons NA, Karvountzis S, Knight-Jones TJD. Aspects of bovine caesarean
427 section associated with calf mortality, dam survival and subsequent fertility. *The*
428 *Veterinary Journal* 2013;197:342–50. <https://doi.org/10.1016/j.tvjl.2013.01.010>.
- 429 [10] Ducrot C, Cimarosti I, Bugnard F, Luquet F. Calving effects on French beef-
430 cow fertility. *Preventive Veterinary Medicine* 1994;19:129–36.
431 [https://doi.org/10.1016/0167-5877\(94\)90045-0](https://doi.org/10.1016/0167-5877(94)90045-0).
- 432 [11] Barkema HW, Schukken YH, Guard CL, Brand A, van der Weyden GC.
433 Fertility, production and culling following cesarean section in dairy cattle.
434 *Theriogenology* 1992;38:589–99. [https://doi.org/10.1016/0093-691X\(92\)90022-J](https://doi.org/10.1016/0093-691X(92)90022-J).

- 435 [12] Laven R, Chambers P, Stafford K. Using non-steroidal anti-inflammatory drugs
436 around calving: Maximizing comfort, productivity and fertility. *The Veterinary*
437 *Journal* 2012;192:8–12. <https://doi.org/10.1016/j.tvjl.2011.10.023>.
- 438 [13] Chastant-Maillard S. Anti-inflammatoires et reproduction chez la vache. *Le*
439 *Point Vétérinaire* 2018;49:120–7.
- 440 [14] Newby NC, Pearl DL, LeBlanc SJ, Leslie KE, von Keyserlingk MAG, Duffield
441 TF. Effects of meloxicam on milk production, behavior, and feed intake in dairy
442 cows following assisted calving. *Journal of Dairy Science* 2013;96:3682–8.
443 <https://doi.org/10.3168/jds.2012-6214>.
- 444 [15] Mainau E, Cuevas A, Ruiz-de-la-Torre JL, Abbeloos E, Manteca X. Effect of
445 meloxicam administration after calving on milk production, acute phase proteins,
446 and behavior in dairy cows. *Journal of Veterinary Behavior: Clinical Applications*
447 *and Research* 2014;9:357–63. <https://doi.org/10.1016/j.jveb.2014.07.007>.
- 448 [16] Meier S, Priest NV, Burke CR, Kay JK, McDougall S, Mitchell MD, et al.
449 Treatment with a nonsteroidal antiinflammatory drug after calving did not improve
450 milk production, health, or reproduction parameters in pasture-grazed dairy cows.
451 *Journal of Dairy Science* 2014;97:2932–43. [https://doi.org/10.3168/jds.2013-](https://doi.org/10.3168/jds.2013-7838)
452 [7838](https://doi.org/10.3168/jds.2013-7838).
- 453 [17] Newby NC, Leslie KE, Dingwell HDP, Kelton DF, Weary DM, Neuder L, et al.
454 The effects of periparturient administration of flunixin meglumine on the health
455 and production of dairy cattle. *Journal of Dairy Science* 2017;100:582–7.
456 <https://doi.org/10.3168/jds.2016-11747>.
- 457 [18] Waelchli RO, Thun R, Stocker H. Effect of flunixin meglumine on placental
458 expulsion in dairy cattle after a caesarean. *Veterinary Record* 1999;144:702–3.
459 <https://doi.org/10.1136/vr.144.25.702>.
- 460 [19] Stilwell G, Schubert H, Broom DM. Short communication: Effects of analgesic
461 use postcalving on cow welfare and production. *Journal of Dairy Science*
462 2014;97:888–91. <https://doi.org/10.3168/jds.2013-7100>.
- 463 [20] Barrier AC, Coombs TM, Dwyer CM, Haskell MJ, Goby L. Administration of a
464 NSAID (meloxicam) affects lying behaviour after caesarean section in beef cows.
465 *Applied Animal Behaviour Science* 2014;155:28–33.
466 <https://doi.org/10.1016/j.applanim.2014.02.015>.

- 467 [21] Guatteo R, Lesort C, Durand D, Touzot-Jourde G. Impact of meloxicam
468 administration in cows prior to caesarean on the efficacy of transfer of passive
469 immunity in calves. Proceedings of the European Buiatrics Forum 2015, Rome,
470 Italy: 2015, p. 107.
- 471 [22] McDougall S, Abbeloos E, Piepers S, Rao AS, Astiz S, van Werven T, et al.
472 Addition of meloxicam to the treatment of clinical mastitis improves subsequent
473 reproductive performance. Journal of Dairy Science 2016;99:2026–42.
474 <https://doi.org/10.3168/jds.2015-9615>.
- 475 [23] van Soest FJS, Abbeloos E, McDougall S, Hogeveen H. Addition of
476 meloxicam to the treatment of bovine clinical mastitis results in a net economic
477 benefit to the dairy farmer. Journal of Dairy Science 2018;101:3387–97.
478 <https://doi.org/10.3168/jds.2017-12869>.
- 479 [24] Bohy A. Consensus sur la césarienne bovine : résultats et débats. Recueil des
480 Journées Nationales 2007, Nantes: SNGTV; 2007, p. 1097–101.
- 481 [25] Kelton DF, Lissemore KD, Martin RE. Recommendations for Recording and
482 Calculating the Incidence of Selected Clinical Diseases of Dairy Cattle. Journal of
483 Dairy Science 1998;81:2502–9. [https://doi.org/10.3168/jds.S0022-](https://doi.org/10.3168/jds.S0022-0302(98)70142-0)
484 [0302\(98\)70142-0](https://doi.org/10.3168/jds.S0022-0302(98)70142-0).
- 485 [26] Guerrier J, Leudet O. Résultats du contrôle des performances bovins
486 allaitants. France - Campagne 2016. Institut de l'Élevage; 2017.
- 487 [27] Grimard B, Agabriel J, Chambon G, Chanvallon A, Constant F, Chastant-
488 Maillard S. Particularités de la reproduction des vaches allaitantes de races
489 françaises. INRA Productions Animales 2017;30:125–38.
- 490 [28] Grimard B, Humblot P, Mialot JP, Ponter AA, Sauvart D, Thibier M. Facteurs
491 de variation de la durée de l'anoestrus postpartum et de la fertilité à l'oestrus
492 induit chez la vache allaitante : importance du niveau d'apport énergétique.
493 Rencontres Recherches Ruminants, vol. 1, Paris, France: 1994, p. 249–52.
- 494 [29] Mialot JP, Ponsart C, Gipoulou Ch, Bihoreau JL, Roux ME, Deletang F. The
495 fertility of autumn calving suckler beef cows is increased by the addition of
496 prostaglandin to progesterone and eCG estrus synchronization treatment.
497 Theriogenology 1998;49:1353–63. [https://doi.org/10.1016/S0093-](https://doi.org/10.1016/S0093-691X(98)00082-X)
498 [691X\(98\)00082-X](https://doi.org/10.1016/S0093-691X(98)00082-X).

- 499 [30] Mialot JP, Constant F, Dezaux P, Grimard B, Deletang F, Ponter AA. Estrus
500 synchronization in beef cows: comparison between GnRH+PGF2 α +GnRH and
501 PRID+PGF2 α +eCG. *Theriogenology* 2003;60:319–30.
502 [https://doi.org/10.1016/S0093-691X\(02\)01371-7](https://doi.org/10.1016/S0093-691X(02)01371-7).
- 503 [31] Diskin MG, Kenny DA. Managing the reproductive performance of beef cows.
504 *Theriogenology* 2016;86:379–87.
505 <https://doi.org/10.1016/j.theriogenology.2016.04.052>.
- 506 [32] Stagg K, Diskin MG, Sreenan JM, Roche JF. Follicular development in long-
507 term anoestrous suckler beef cows fed two levels of energy postpartum. *Animal*
508 *Reproduction Science* 1995;38:49–61. [https://doi.org/10.1016/0378-](https://doi.org/10.1016/0378-4320(94)01354-O)
509 [4320\(94\)01354-O](https://doi.org/10.1016/0378-4320(94)01354-O).
- 510 [33] Hanzen C, Théron L, Detilleux J. Réalisation de la césarienne dans l'espèce
511 bovine en Europe : l'intervention et ses conséquences. *Bulletin des GTV*
512 2011;61–72.
- 513 [34] Hanzen C, Théron L, Detilleux J. Modalités de réalisation de la césarienne en
514 Europe. *Bulletin Des GTV* 2011;15–28.
- 515 [35] Patterson DJ, Bellows RA, Burfening PJ. Effects of caesarean section,
516 retained placenta and vaginal or uterine prolapse on subsequent fertility in beef
517 cattle. *Journal of Animal Science* 1981;53:916–21.
518 <https://doi.org/10.2527/jas1981.534916x>.
- 519 [36] Vermunt JJ, Parkinson TJ, Noakes DE. Defects of Presentation, Position and
520 Posture in Livestock: Delivery by Caesarean Hysterotomy. *Veterinary*
521 *Reproduction and Obstetrics*. 10th edition, Elsevier; 2019, p. 291–314.
- 522 [37] Boucoumont D, Lecuyer B, Rosentiehl D, Tisserand R, Troccon B, Ouillier R.
523 Effect of caesarean section on fertility of cows. Comparative study of catgut and a
524 polyglycolic acid thread. *Le Point Vétérinaire* 1978;8:15–9.
- 525 [38] Coutard J-P, Ménard M, Benoteau G. Reproduction des troupeaux allaitants
526 dans les Pays de la Loire: facteurs de variation des performances. *Rencontres*
527 *Recherches Ruminants*, vol. 14, Paris, France: 2007, p. 359–62.
- 528 [39] Hanzen C. Etude des facteurs de risque de l'infertilité et des pathologies
529 puerpérales et du postpartum chez la vache laitière et chez la vache viandeuse.
530 Thèse d'agrégation de l'enseignement supérieur. Université de Liège, 1994,
531 172p.

- 532 [40] Cattell JH, Dobson H. A survey of caesarean operations on cattle in general
533 veterinary practice. *Veterinary Record* 1990;16:395–9.
- 534 [41] Laven RA, Peters AR. Bovine retained placenta: aetiology, pathogenesis and
535 economic loss. *Veterinary Record* 1996;139:465–71.
536 <https://doi.org/10.1136/vr.139.19.465>.
- 537 [42] Eiler H, Fecteau KA. Retained placenta. In: Threlfall RS, Youngquistwalter R,
538 editors. *Current therapy in large animal theriogenology*. 2nd ed., Saint Louis:
539 W.B. Saunders; 2007, p. 345–54. [https://doi.org/10.1016/B978-072169323-](https://doi.org/10.1016/B978-072169323-1.50048-9)
540 [1.50048-9](https://doi.org/10.1016/B978-072169323-1.50048-9).
- 541 [43] Lemke KA, Creighton CM. Analgesia for Anesthetized Patients. *Topics in*
542 *Companion Animal Medicine* 2010;25:70–82.
543 <https://doi.org/10.1053/j.tcam.2009.12.003>.
- 544 [44] Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative
545 recovery. *The Lancet* 2003;362:1921–8. [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(03)14966-5)
546 [6736\(03\)14966-5](https://doi.org/10.1016/S0140-6736(03)14966-5).
- 547 [45] Coetzee JF. A Review of Analgesic Compounds Used in Food Animals in the
548 United States. *Veterinary Clinics of North America: Food Animal Practice*
549 2013;29:11–28. <https://doi.org/10.1016/j.cvfa.2012.11.008>.
- 550 [46] Anderson DE, Edmondson MA. Prevention and Management of Surgical Pain
551 in Cattle. *Veterinary Clinics of North America: Food Animal Practice*
552 2013;29:157–84. <https://doi.org/10.1016/j.cvfa.2012.11.006>.
- 553 [47] Carvalho B, Butwick AJ. Postcesarean delivery analgesia. *Best Practice &*
554 *Research Clinical Anaesthesiology* 2017;31:69–79.
555 <https://doi.org/10.1016/j.bpa.2017.01.003>.
- 556 [48] Kerai S, Saxena KN, Taneja B. Post-caesarean analgesia: What is new?
557 *Indian J Anaesth* 2017;61:200–14. https://doi.org/10.4103/ija.IJA_313_16.
- 558 [49] Coetzee JF. Assessment and Management of Pain Associated with Castration
559 in Cattle. *Veterinary Clinics of North America: Food Animal Practice* 2013;29:75–
560 101. <https://doi.org/10.1016/j.cvfa.2012.11.002>.
- 561 [50] Richards BD, Black DH, Christley RM, Royal MD, Smith RF, Dobson H.
562 Effects of the administration of ketoprofen at parturition on the milk yield and
563 fertility of Holstein-Friesian cattle. *Veterinary Record* 2009;165:102–6.
564 <https://doi.org/10.1136/vetrec.165.4.102>.

- 565 [51] Anderson DE, Muir WW. Pain Management in Cattle. *Veterinary Clinics of*
566 *North America: Food Animal Practice* 2005;21:623–35.
567 <https://doi.org/10.1016/j.cvfa.2005.07.002>.
- 568 [52] Newby NC, Renaud D, Tremblay R, Duffield TF. Evaluation of the effects of
569 treating dairy cows with meloxicam at calving on retained fetal membranes risk.
570 *Can Vet J* 2014;55:1196–9.
- 571 [53] Swartz TH, Schramm HH, Bewley JM, Wood CM, Leslie KE, Petersson-Wolfe
572 CS. Meloxicam administration either prior to or after parturition: Effects on
573 behavior, health, and production in dairy cows. *Journal of Dairy Science*
574 2018;101:10151–67. <https://doi.org/10.3168/jds.2018-14657>.
- 575 [54] Shock DA, Renaud DL, Roche SM, Poliquin R, Thomson R, Olson ME.
576 Evaluating the impact of meloxicam oral suspension administered at parturition
577 on subsequent production, health, and culling in dairy cows: A randomized clinical
578 field trial. *PLOS ONE* 2018;13:e0209236.
579 <https://doi.org/10.1371/journal.pone.0209236>.
580

581 **Table 1. Comparison of meloxicam and control groups.**

582 Charolais heifers (n = 127) underwent a non-elective standardized caesarean section operation. Heifers were randomly assigned to

583 one of two groups: meloxicam (n = 66), intravenous meloxicam injection before surgery, or control (n = 61).

584

Variable	Meloxicam group		Control group		Total		p-value
	n (/66)	%	n (/61)	%	n (/127)	%	
Heifer birth season							
<i>Autumn (12/21/09 to 12/20/12)</i>	12	18%	9	15%	21	17%	<i>0.74</i>
<i>Winter (12/21/12 to 13/14/03)</i>	42	64%	42	69%	84	66%	
<i>Spring (13/15/03 to 13/21/06)</i>	11	16%	8	13%	19	15%	
<i>Missing data</i>	1	2%	2	3%	3	2%	
C-section period							
<i>December 2015</i>	25	38%	22	36%	47	37%	<i>0.11</i>
<i>January 2016</i>	33	50%	23	38%	56	44%	
<i>February-May 2016</i>	8	12%	16	26%	24	19%	
Age at C-section							
<i>31-33 months</i>	8	16%	5	13%	13	10%	<i>0.78</i>
<i>34-35 months</i>	25	74%	24	74%	49	39%	
<i>36-39 months</i>	32	8%	30	10%	62	49%	
<i>Missing data</i>	1	2%	2	3%	3	2%	
Body Condition Score at C-section ^a							
<i>< 3</i>	8	12%	9	15%	17	13%	<i>0.30</i>
<i>3</i>	23	35%	28	46%	51	40%	
<i>> 3</i>	35	53%	24	39%	59	46%	

Size of the uterine incision							
20-29 cm	8	12%	12	20%	20	16%	0.29
30-39 cm	51	77%	46	75%	97	76%	
40-49 cm	7	11%	3	5%	10	8%	
1 st uterine suture pattern type							
Simple continuous	59	89%	52	85%	111	87%	0.48
Inverting (Cushing or Lembert)	7	11%	9	15%	16	13%	
Length of surgery							
< 30 minutes	18	27%	14	23%	32	25%	0.25
30-35 minutes	36	55%	22	36%	58	46%	
> 35 minutes	10	15%	2	3%	12	9%	
Missing data	2	3%	23	38%	25	20%	
Average length (minutes) \pm SD	32 \pm 8		29 \pm 6		31 \pm 7		

585

^a scale 1-5

586

^b defined as the time from the start of the preparation of the animal for surgery - including premedication - to the end of the cutaneous suture

587 **Table 2. Comparison of reproductive performance for**
 588 **meloxicam and control groups.**

589 Charolais heifers (n = 127) underwent a non-elective standardized caesarean section
 590 operation. Heifers were randomly assigned to one of two groups: meloxicam (n = 66),
 591 intravenous meloxicam injection before surgery, or control (n = 61).

592


Reproductive parameters	Meloxicam group	Control group	p-value
Retained placenta (%) ^a	18.2 (12/66)	25.0 (15/60)	0.35
Pregnancy rate (%) ^b	83.1 (54/65)	67.8 (40/59)	0.04
Calving Interval (days) (mean ± standard error)	406.6 ±6.7	417.5 ±8.0	0.20
Culling rate (%)	4.7 (3/64)	13.3 (8/60)	0.09

593 ^a retained placenta was defined as the presence of foetal membranes in the uterus by visual or vaginal
 594 examination more than 24 hours after calving


595 ^b pregnancy diagnosis was performed by ultrasound (after 30 days post-mating) or by transrectal
 596 palpation (after 60 days post-mating)

597 **Figure 1. Kaplan-Meier survival plot (days from caesarean**
598 **section to next calving) for Charolais heifers (n = 127)**
599 **receiving meloxicam before the caesarean section**
600 **(meloxicam group, n = 66) or only local anesthesia (control**
601 **group, n = 61).**

602 The dashed line indicates the median survival time, i.e the calving interval
603 (meloxicam group, t=417 days; control group, t=452 days, p=0.050 - log-rank test)
604 and the dots the censored animals (subjects who left the study, or the study ended
605 before calving occurred). Each animal was followed for a 520-days period.



606
607



— Meloxicam group

— Control group