



HAL
open science

Classification of wine grape cultivars according to susceptibility to *Botrytis cinerea* in Chile and France: effects of fruit maturity and cluster compactness

Carolina Pañitrur-de La Fuente, Héctor Valdés-Gómez, Jean Roudet, Mauricio Lolas, César Acevedo-Opazo, Marc Fermaud

► To cite this version:

Carolina Pañitrur-de La Fuente, Héctor Valdés-Gómez, Jean Roudet, Mauricio Lolas, César Acevedo-Opazo, et al.. Classification of wine grape cultivars according to susceptibility to *Botrytis cinerea* in Chile and France: effects of fruit maturity and cluster compactness. 17. International Botrytis Symposium, Oct 2016, Santa Cruz, Chile. hal-03364677

HAL Id: hal-03364677

<https://hal.inrae.fr/hal-03364677>

Submitted on 4 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



CLASSIFICATION OF WINE GRAPE CULTIVARS ACCORDING TO SUSCEPTIBILITY TO *Botrytis cinerea* IN CHILE AND FRANCE: EFFECTS OF FRUIT MATURITY AND CLUSTER COMPACTNESS

Carolina Pañitrur-de la Fuente¹, Héctor Valdés-Gómez², Jean Roudet³, Mauricio Lolas¹, César Acevedo-Opazo¹, Marc Fermaud³

¹Universidad de Talca, Facultad de Ciencias Agrarias, Talca, Chile;

²Pontificia Universidad Católica de Chile, Departamento de Fruticultura y Enología, Santiago, Chile;

³INRA, UMR 1065 Santé et Agroécologie du Vignoble (SAVE) – ISVV, Centre INRA de Bordeaux, France.

XVII INTERNATIONAL BOTRYTIS SYMPOSIUM

**23rd – 28th October, 2016; Hotel Santa Cruz Plaza,
Santa Cruz, Valle de Colchagua, Chile.**



Yield

Quality



INTEGRATED PEST MANAGEMENT



**Human health
Environment**



PRESENTATION OUTLINE

- INTRODUCTION:
Botrytis bunch rot of grapes: Damage and Control
- CLASSIFICATION OF WINE GRAPE CULTIVARS :
According to susceptibility to *Botrytis cinerea* in Chile and France
- EFFECTS OF FRUIT MATURITY AND CLUSTER COMPACTNESS:
On the ranking of susceptibility to *Botrytis cinerea*
- CONCLUSIONS AND PERSPECTIVES



INTRODUCTION

Botrytis bunch rot of grapes (BBR): Damage and control



XVII INTERNATIONAL BOTRYTIS SYMPOSIUM

23rd – 28th October, 2016; Hotel Santa Cruz Plaza,
Santa Cruz, Valle de Colchagua, Chile.

Anuncio de nuevas lluvias para los próximos

Agraz situación para los productores de uva en Chile: producción cayó 30% y pérdidas superan los US\$220 millones



esperado

Tweet

por EL MOSTRADOR MERCADOS | 11 mayo 2016



Esta tarde la Sociedad Nacional de Agricultura (SNA) comunicó el complejo momento que atraviesan los productores de uva en el país.

El gremio, que calificó la vendimia de quienes trabajan la uva vinífera como una situación "muy compleja", apunta a las fuertes lluvias que golpearon a la zona central de Chile durante abril, así como los bajos precios internacionales de la uva.

Dicha situación habría dejado al proceso de recolección de la uva con una fuerte caída en la producción de 30%, lo que equivaldría a unos 400 millones menos de litros de vino para esta cosecha.


La uva vinífera habría sido golpeada por las intensas lluvias que cayeron en la zona central de Chile durante abril, así como los bajos precios internacionales, de acuerdo a la Sociedad Nacional de Agricultura.

De acuerdo a la SNA, la producción final de vinos para este 2016 sería de sólo unos 800 millones de litros, lo que traería pérdidas a la industria por sobre los US\$220 millones para los productores del

Especialmente preocupante es el caso de la botrytis, pues esta tiende a dejarse fruto. por lo que todavía el efecto que pueda tener está por verse.

Apur
podría
de ser
fruta.
Crédit

Botrytis bunch rot of grapes







GRAPEVINE CULTIVAR



Genetic and Phenotypic traits:

- ✓ Clusters compactness
- ✓ Morphological, anatomical and chemical features of the berry skin



T° : 15-25 ° C
 RH: >90% por 15 h



GRAPEVINE CULTIVAR

Essential management indicator in Integrated Pest Management (IPM)

Classification according to their susceptibility to BBR in literature

Based mostly on professional experience, rather than experimental data

Few studies compared it under the same environmental and management conditions

No study has compared it in contrasting climatic and cultural conditions (eg. Northern vs southern hemisphere)





MAIN OBJECTIVE:

To classify and compare the susceptibility to *B. cinerea* between different wine grape cvs in two contrasting climatic and cultivation conditions

SECONDARY OBJECTIVE:

To relate the fruit maturity with the resulting susceptibility ranking to *B. cinerea*



CLASSIFICATION OF WINE GRAPE CULTIVARS

*According to Susceptibility to Botrytis cinerea in Chile
and France*



XVII INTERNATIONAL BOTRYTIS SYMPOSIUM

23rd – 28th October, 2016; Hotel Santa Cruz Plaza,
Santa Cruz, Valle de Colchagua, Chile.

Classification of wine grape cultivars

Materials and Methods

- ❖ Chile (Maule Region) and France (Bordeaux Area)
- ❖ Seasons 2011 to 2015
- ❖ 3 grapevine collections

- ✓ 1 Grapevine collection “Panguilemo”
- ✓ 19 cultivars
- ✓ 2 Seasons (2013-14 and 2014-15)

- ✓ 2 Grapevine collections: “Tour Blanche” (TB) and “Grande Ferrade” (GF)
- ✓ 33 cultivars
- ✓ 3 Seasons (2011,2012 and 2014)



13 COMMON CULTIVARS

Cab. Franc
Cab. Sauvignon
Chardonnay
Cot
Gewürztraminer
Grenache noir
Merlot
Mourvèdre
Petit Verdot
Pinot noir
Roussanne
Sauvignon blanc
Syrah



Classification of wine grape cultivars

Materials and Methods

❖ EXPERIMENTAL CONDITIONS

Property	France		Chile
	Tour Blanche	Grande Ferrade	Panguilemo
Experimental Period	2011, 2012, 2014	2011	2013-14, 2014-15
Vineyard planting year	1995	2009	2006
Rootstock	3309	SO4	Own-rooted
Location (WGS84)	44°32' N, 0°21' W	44°47' N, 0°34' W	35°22' S, 71°36' W
Spacing (m x m)	1.8 x 0.9	1.8 x 1.0	2.0 x 1.0
Trellis/Pruning system		VSPSystem ^a / Two-bilateral	
Irrigation system	Non-irrigated	Non-irrigated	Drip irrigation (one dropper per plant with a flow rate of 4 L / h)



Classification of wine grape cultivars

Materials and Methods

❖ EVALUATIONS:

- ✓ Climatic characterization
- ✓ Disease susceptibility assessment
- ✓ Statistical analyses



Classification of wine grape cultivars

Materials and Methods

❖ EVALUATIONS:

- ✓ **Climatic characterization** →
- ✓ Disease susceptibility assessment
- ✓ Statistical analyses

- ✓ Automatic weather station
- ✓ Temperature, Relative Humidity and precipitation
- ✓ Second season in Chile: vines were moistened.



Classification of wine grape cultivars

Materials and Methods

❖ EVALUATIONS:

✓ Climatic characterization

✓ **Disease susceptibility assessment** →

✓ Statistical analyses

- ✓ Incidence and Severity at harvest (22°Brix)
- ✓ Susceptibility index (SI)



Highly Resistant (HR)=0-2.5%;
Resistant (R)=2.51-10%;
Intermediate (I)=10.1-25%;
Susceptible (S)=25.1-50%
Highly Susceptible (HS)= 50.1-100%.



$$SI = \frac{\text{Severity (\%)} \text{ for cultivar in question}}{\text{Highest severity (\%)} \text{ recorded in the season and in the most rotted cultivar}} \times 100$$



Classification of wine grape cultivars

Materials and Methods

❖ EVALUATIONS:

- ✓ Climatic characterization
- ✓ Disease susceptibility assessment

✓ **Statistical analyses** →

- ✓ ANOVA using the PROC GLM procedure; test (LSD) (severity and incidence data)
- ✓ Cluster Analysis (severity data)
- ✓ Box Plot Analysis (Susceptibility Index data)



Results

✓ Climatic characterization

FRANCE

humid and temperate conditions



Favored the growth and development of *B. cinerea*

CHILE

Dry and temperate conditions



Not conducive to disease development




Fig. 1. Monthly means for rainfall (mm) in France (a) and Chile (b), and mean air temperature (°C) in France (c) and Chile (d) during all seasons. The horizontal dotted line in (c) and (d) represents the mean air temperature (°C) in each season. Bud= Budbreak; Flo= Flowering; Ver= Veraison; Har= Harvest.



Results

✓ Disease susceptibility assessment






Fig. 2. Cluster classification of cultivars in France in the sites “Tour Blanche” (a) and both “Grande Ferrade and “Tour Blanche” (b), according to their severity values.



Results

✓ Disease susceptibility assessment





Fig. 3. Cluster classification of cultivars in Chile according to their severity values.



Results

RANKING SUSCEPTIBILITY

✓ Disease susceptibility assessment




Fig.2. Box-Plot of cultivars according to the Susceptibility Index. Where HR= Highly Resistant; R= Resistant; I= Intermediate; S= Susceptible; HS = Highly Susceptible. The vertical line in each box and the red cross, represents the median and the mean value of SI.


Results

RANKING SUSCEPTIBILITY

✓ Disease susceptibility assessment

Cultivar	a	b	c	d	e	f	g	h	Mean lit.	Our res.	Sd lit.	Sd res.
Grenache Noir	4	3	-	-	4	-	3	4	4	0	0.5	1.0
Cabernet Franc	3	-	-	-	-	-	4	1	3	0	1.5	1.6
Petit Verdot	0-1	-	-	-	-	-	1	1	1	0	0.3	1.6
Cabernet Sauvignon	2	-	0	1	1	0	1	1	1	0	0.7	1.2
Mourvèdre	-	-	-	-	-	-	1	-	1	1	-	0.5
Merlot	3	-	-	-	-	-	3	3	3	1	0	1.5
Syrah	2	-	1	3	3	-	-	2	2	1	0.8	1.2
Cot	3	-	-	-	-	-	3	3	3	1	0	1.5
Roussanne	4	-	-	-	-	-	-	4	4	2	0	1.2
Chardonnay	4	-	2	2	3	-	3	3	3	3	0.8	1.2
Pinot Noir	3	4	2	3	4	-	-	3	3	3	0.8	1.3
Gewürztraminer	4	-	-	-	-	-	1	4	3	4	1.7	0
Sauvignon Blanc	4	-	4	3	4	-	1	4	3	4	1.2	0.5

a= Dubos (2002), b=Dry and Gregory (1990), c= Orffer (1979), d=Jackson and Schuster (1987), e=Robinson (1986), f=Marois et al. (1992), g= Galet (1988), h= ACTA (1980); 0= highly resistant, 1= resistant, 2= intermediate, 3= susceptible, 4= highly susceptible; Mean lit= Mean of literature source, Our res= Resultats of our study; Sd lit= standard deviation of literature sources, Sd res= estándar deviation of our results.



EFFECTS OF FRUIT MATURITY AND CLUSTER COMPACTNESS

*On the ranking of susceptibility to *Botrytis cinerea*.*



XVII INTERNATIONAL BOTRYTIS SYMPOSIUM

23rd – 28th October, 2016; Hotel Santa Cruz Plaza,
Santa Cruz, Valle de Colchagua, Chile.

Effects of fruit maturity and cluster compactness

Materials and Methods

❖ EVALUATIONS:

- ✓ Fruit Maturity
- ✓ Cluster compactness
- ✓ Statistical analyses



Effects of fruit maturity and cluster compactness

Materials and Methods

❖ EVALUATIONS:

- ✓ Fruit Maturity → ✓ Maturity index (F_{Mat})
- ✓ Cluster compactness
- ✓ Statistical analyses

$$F_{Mat} = F_{B.c \text{ assessment}} - F_{veraison}$$

$F_{B.c \text{ assessment}}$ = Timing of the *B. cinerea* assessment
 $F_{veraison}$ = Timing of veraison for each cultivar
(GFV model of *Parker et al 2011, 2013*)

Critical degree-day sum (above 0°C) calculated from the 60th (France) and 242th (Chile) day of the year to the dates of *B. cinerea* assessment and veraison.

In order to remove the effect of each season

$$F_{Mat_adj} = F_{Mat} \text{ for each cultivar} - F_{Mat} \text{ Petit Verdot}$$



Effects of fruit maturity and cluster compactness

Materials and Methods

❖ EVALUATIONS:

- ✓ Fruit Maturity
- ✓ Cluster compactness
- ✓ Statistical analyses

→
✓ 40 clusters per cultivar
✓ visual scale from 1 to 6




Fig. 3. Visual scale of compactness of cluster.



Effects of fruit maturity and cluster compactness

Materials and Methods

❖ EVALUATIONS:

- ✓ Fruit Maturity
- ✓ Cluster compactness
- ✓ Statistical analyses →

- ✓ Correlation analysis
- ✓ nonlinear model



Effects of fruit maturity and cluster compactness

Results




Fig.4. Relationship between Maturity of cultivars (F Mat) and Susceptibility to BBR (SI), assessment at different dates, in France and Chile.

1) Fruit maturity




Fig.5. Relationship between Maturity of cultivars (F Mat_adj) and Susceptibility to BBR (SI) in both sites: France (a) and Chile (b) during all study seasons.



Effects of fruit maturity and cluster compactness

Results




Fig.6. Relationship between Cluster compactness and Susceptibility to BBR (SI) in France and Chile.

2) Cluster compactness




Fig.7. Relationship between Cluster compactness (CI) and Susceptibility to BBR (SI) in both sites: France (a) and Chile (b) during all study seasons.



Conclusions and perspectives

- ✓ The classification of the wine cultivars according to their susceptibility to *B. cinerea* was similar in both countries, despite the contrasting climatic conditions and management practices.
- ✓ Sauvignon Blanc and Gewürztraminer were the highest susceptible cultivars, whereas Petit Verdot, Cabernet Sauvignon, Mourvèdre and Syrah were rather resistant or highly resistant (in accordance to previous studies).
- ✓ For other cvs. evaluated (Grenache Noir, Cabernet Franc, Merlot, Cot and Roussanne), their ranking differed to some extent from the literature.
- ✓ This difference is presumably caused by variations in agronomic and/or environmental conditions under which the field experiments were carried out (eg. clone, rootstock, cluster compactness related to flower abortion).



Conclusions and perspectives

- ✓ Fruit maturity and cluster compactness, depending significantly on the cultivar, proved to be very important factors governing susceptibility to *B. cinerea*.
- ✓ The susceptibility classification of cultivars remains a key parameter in decision support systems and then, both variables cluster compactness and fruit maturity could be used to support this classification.
- ✓ Further investigation should be conducted to better understand the relationships between susceptibility to *B. cinerea* and other variables (e.g. clone, vigor, rootstock), in order to develop new IPM strategies



CONICYT Doctoral Fellowship 2013 N°
21130505





Thanks for your attention!