



The complete mitochondrial genome of the Basidiomycete fungus *Pleurotus cornucopiae* (Paulet) Rolland

Li-Ming Xu, D. D. Hinsinger, Guo-Feng Jiang

► To cite this version:

Li-Ming Xu, D. D. Hinsinger, Guo-Feng Jiang. The complete mitochondrial genome of the Basidiomycete fungus *Pleurotus cornucopiae* (Paulet) Rolland. *Mitochondrial DNA Part B Resources*, 2018, 3 (1), pp.73-75. 10.1080/23802359.2017.1422405 . hal-03370666

HAL Id: hal-03370666

<https://hal.inrae.fr/hal-03370666>

Submitted on 4 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

MITOGENOME ANNOUNCEMENT



The complete mitochondrial genome of the Basidiomycete fungus *Pleurotus cornucopiae* (Paulet) Rolland

Li-Ming Xu^a, Damien Daniel Hinsinger^b and Guo-Feng Jiang^{b,c} 

^aBiology Institute, Guangxi Academy of Sciences, Nanning, Guangxi, PR China; ^bBiodiversity Genomics Team, Plant Ecophysiology & Evolution Group, Guangxi Key Laboratory of Forest Ecology and Conservation, College of Forestry, Guangxi University, Nanning, Guangxi, PR China; ^cState Key Laboratory for Conservation and Utilization of Subtropical Agro-Bioresources, Guangxi University, Nanning, Guangxi, PR China

ABSTRACT

Pleurotus cornucopiae is a commercial edible and medicinal fungus. Herein, we determined and analyzed its complete mitochondrial genome. The mitogenome length was 72,134 bp with a GC content of 26.7%, contained 14 conserved protein coding genes, two rRNA genes (*rnl* and *rns*), ribosomal protein gene *rps3* and 24 tRNA genes. *Pleurotus cornucopiae* has a similar gene content and gene order of the mitogenome as *P. ostreatus* and *P. eryngii*. A phylogenetic analysis based on complete mitogenome in related fungi showed that *P. cornucopiae* is a member of the order Agaricales, forming a clade with *P. ostreatus* and *P. eryngii*, with *P. ostreatus* as a sister taxa. The mitochondrial genome sequence of *P. cornucopiae* appeared a promising tool for further studies of the taxonomy and evolution of Pleurotaceae and Agaricales.

ARTICLE HISTORY

Received 16 December 2017
Accepted 26 December 2017

KEYWORDS



Fungi; mushroom; *Pleurotus cornucopiae*; mitogenome; evolution

The genus *Pleurotus*, belonging to the family Pleurotaceae (Agaricales, Agaricomycetes, Basidiomycota), is one of the most diverse group of cultivated mushrooms in the world. The *Pleurotus* species establish a wide range of interactions with other organisms, and are efficient bioconverters of lignocellulosic residues into human food (Tsuneda and Thorn 1995; Philippoussis 2009). The biodiversity of *Pleurotus* remains highly investigated and recently, molecular studies have increase our knowledge about intra and inter-specific heterogeneity of the genus *Pleurotus* using both ribosomal and mitochondrial DNA (Li and Yao 2004; Wang et al. 2008; Yang et al. 2016; Chaudhary and John 2017). The edible fungus *Pleurotus cornucopiae* is emerging as an important species in the genus, with isolation of medicinal metabolites from the fruiting body (Wang et al. 2012). The genetic diversity and population structure of *P. cornucopiae* is well known (Iraçabal and Labarere 1994; Zervakis et al. 1994; Shnyreva et al. 1996; Zhang et al. 2007; Zhou et al. 2009; Adebayo et al. 2016); however, no complete mitogenome is available to date for the species. Here, we report the complete mitogenome of *P. cornucopiae* (GenBank accession MG652610) to provide new genetic resources and performed a phylogenetic analysis of related taxa of this species.

The strain SWS_15 of *Pleurotus cornucopiae* is maintained in the Biology Institute, Guangxi Academy of Sciences (Nanning, Guangxi, PR China). Total genomic DNA was extracted as previously described (Xu et al. 2017). Library construction and sequencing were processed by Novogene (Beijing, China), according to the Illumina HiSeqX-ten system

manufacturer instructions (Illumina, San Diego, CA). The mitogenome of *Pleurotus cornucopiae* was *de novo* assembled using ORG.Asm v0.2.05 (<https://pythonhosted.org/ORG.Asm/>) followed by manual curation in Geneious R9 v9.1.6 (Biomatters Ltd, Auckland, New Zealand) as described previously (Hinsinger and Strijk 2016; Jiang et al. 2016; Xu et al. 2017). Genome annotation and phylogenetic analysis were performed with DOGMA (<http://dogma.cccb.utexas.edu/index.html>) and Phym1 3.1 (Guindon et al. 2010), respectively. Maximum-likelihood (ML) tree was constructed including 10 available mitogenomes of Agaricales and *Heterobasidion irregulare* as an outgroup (Figure 1).

The complete mitogenome of *Pleurotus cornucopiae* was 72,134 bp in length with a GC content of 26.7%. The mitogenome contained 14 conserved protein coding genes, 2 rRNA genes (*rnl* and *rns*), the ribosomal protein gene *rps3* and 24 tRNA genes. The 14 conserved protein coding genes respectively encoded the seven ubiquinone reductase subunits of NADH (*nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, *nad5* and *nad6*), three cytochrome oxidase subunits (*cox1*, *cox2* and *cox3*), three ATP synthase subunits (*atp6*, *atp8* and *atp9*) and the apocytochrome b (*cob*). The 24 tRNA genes ranged in size from 71 bp to 93 bp, and covered all 20 standard amino acids. Phylogenetic analysis based on ML showed with high support that *Pleurotus cornucopiae* is a member of Agaricales and grouping with *P. ostreatus* and *P. eryngii*, and closely related to *P. ostreatus* (Figure 1). This result is consistent to the previous study based on SSU rDNA or LSU rDNA analyses (Gao et al. 2008; Chaudhary and John 2017), but differs from the

CONTACT Guo-Feng Jiang  gfiang@gxu.edu.cn  Plant Ecophysiology & Evolution Group, Guangxi Key Laboratory of Forest Ecology and Conservation, College of Forestry, Guangxi University, DaXueDongLu 100, Nanning, Guangxi 530005, PR China

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. ML phylogenetic tree of the 10 available mitogenomes of Agaricales in GenBank, plus the mitogenome of *Pleurotus cornucopiae*. The tree is rooted with *Heterobasidion irregulare*. Bootstraps values (1000 replicates) are shown at the nodes. Scale in substitution per site.

result based on ITS data (Shnyreva and Shnyreva 2015; He et al. 2016). The complete mitochondrial genome of *P. cornucopiae* provides the community a useful bioresource that will help to delineate taxonomic units in *Pleurotus*, and will be of interest for further applied researches by selecting the most suitable species for biotechnological and nutritional interest.

Acknowledgements

The authors would like to acknowledge the technical staff of the Biology Institute, Guangxi Academy of Sciences for their assistance in access to the *P. cornucopiae* strain SWS_15.

Disclosure statement

The authors report no conflicts of interest.

Funding

This work was supported by grants from the Natural Science Foundation of Guangxi Province of China (2014GXNSFBA118075 and 2015GXNSF AA139085) to G. -F. Jiang and D.D. Hinsinger (Guangxi University) and the Natural Science Foundation of Guangxi Province of China (2015GXNSFAA139087) to L. -M. Xu.

ORCID

Guo-Feng Jiang  <http://orcid.org/0000-0002-3221-8608>

References

- Adebayo EA, Oloke KJK, Azeez MA, Ayandele AA, Majolagbe ON. 2016. Compatibility study using hybridization procedure among *Pleurotus* genotypes and authentication by enzyme expression and ITS of rDNA. *Russian Agric Sci.* 42:423–430.
- Chaudhary MM, John P. 2017. Morphological and molecular characterization of oyster mushroom (*Pleurotus cystidiosus*). *Int J Curr Microbiol Appl Sci.* 6:246–250.
- Gao S, Huang CY, Chen Q, Bian YB, Zhang JX. 2008. Phylogenetic relationship of *Pleurotus* species based on nuclear large subunit ribosomal DNA sequences. *J Plant Genet Resour.* 9:328–334.
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol.* 59:307–321.
- He XL, Wu B, Li Q, Peng WH, Huang ZQ, Gan BC. 2016. Phylogenetic relationship of two popular edible *Pleurotus* in China, Bailinggu (*P. eryngii* var. *tuoliensis*) and Xingbaogu (*P. eryngii*), determined by ITS, RPB2 and EF1 α sequences. *Mol Biol Rep.* 43:573–582.
- Hinsinger DD, Strijk JS. 2016. Toward phylogenomics of Lauraceae: the complete chloroplast genome sequence of *Litsea glutinosa* (Lauraceae), an invasive tree species on Indian and Pacific Ocean islands. *Plant Gene.* 9:71–79.
- Iraçabal B, Labarere J. 1994. Restriction site and length polymorphism of the rDNA unit in the cultivated basidiomycete *Pleurotus cornucopiae*. *Theoret Appl Genet.* 88:824–830.
- Jiang GF, Hinsinger DD, Strijk JS. 2016. Comparison of intraspecific, inter-specific and intergeneric chloroplast diversity in Cycads. *Sci Rep.* 6:31473
- Li XL, Yao YJ. 2004. Assembling phylogenetic tree of based on 28S rDNA sequencing. *Mycosystema.* 23:345–350.
- Philippoussis AN. 2009. Biotechnology for agro-industrial residues utilisation: production of mushrooms using agro-industrial residues as substrates. Netherlands: Springer; p. 163–196.
- Shnyreva AV, Lomov AA, Mednikov BM, Dyakov JT. 1996. Oyster cap fungi (*Pleurotus* spp.) species and kinds identification using mating and molecular markers. *Mikologiya I Fitopatol.* 30:37–44.
- Shnyreva AA, Shnyreva AV. 2015. Phylogenetic analysis of *Pleurotus* species. *Russian J Genet.* 51:148–157.
- Tsuneda A, Thorn RG. 1995. Interactions of wood decay fungi with other microorganisms, with emphasis on the degradation of cell walls. *Can J Bot.* 73:1325–1333.
- Wang SJ, Bao L, Han JJ, Wang QX, Yang XL, Wen HA, Guo LD, Li SJ, Zhao F, Liu HW. 2012. Pleurospiroketals A – E, Perhydrobenzannulated 5,5-Spiroketals Sesquiterpenes from the edible mushroom *Pleurotus cornucopiae*. *J Natural Prod.* 76:45–50.
- Wang Y, Zeng FY, Hon CC, Zhang YZ, Leung FCC. 2008. The mitochondrial genome of the Basidiomycete fungus *Pleurotus ostreatus* (oyster mushroom). *FEMS Microbiol Lett.* 280:34–41.
- Xu LM, Hinsinger DD, Jiang GF. 2017. The complete mitochondrial genome of the *Agrocybe aegerita*, an edible mushroom. *Mitochondrial DNA Part A.* 2:791–792.

- Yang RH, Li Y, Li CH, Xu JP, Bao DP. 2016. The complete mitochondrial genome of the Basidiomycete edible fungus *Pleurotus eryngii*. Mitochondrial DNA Part B. 1:772–774.
- Zervakis G, Sourdís J, Balis C. 1994. Genetic variability and systematics of eleven *Pleurotus* species based on isozyme analysis. Mycol Res. 98:329–341.
- Zhang JX, Huang CY, Guan GP, Li HP, Zhang RY, Hu QX. 2007. Inter simple sequence repeat analysis for *Pleurotus cornucopiae*. Mycosystema. 26:115–121.
- Zhou J, Guo Y, Jia DH, Tan W, Zhang XP, Zheng LY, Huang ZQ. 2009. Hybrid strains of *Pleurotus cornucopiae* identified by ISSR. Southwest China J Agric Sci. 22:1694–1698.