



HAL
open science

IMPACT OF NON-STARCH POLYSACCHARIDES ON THE TEXTURAL BEHAVIOR OF PROCESSED YAM

Dutheil de La Rochère, Dominique Rinaldo, Agnès Rolland-Sabaté

► **To cite this version:**

Dutheil de La Rochère, Dominique Rinaldo, Agnès Rolland-Sabaté. IMPACT OF NON-STARCH POLYSACCHARIDES ON THE TEXTURAL BEHAVIOR OF PROCESSED YAM. EPNOE, Oct 2021, Nantes, France. hal-03372189

HAL Id: hal-03372189

<https://hal.inrae.fr/hal-03372189v1>

Submitted on 9 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IMPACT OF NON-STARCH POLYSACCHARIDES ON THE TEXTURAL BEHAVIOR OF PROCESSED YAM

Dutheil de la Rochère A.*^a, Rinaldo D.^b, Rolland-Sabaté A.^a

Congrès EPNOE, novembre 2021

^a INRAE, Avignon University, UMR 408 SQPOV, F-84000, Avignon, France

^b INRAE, UR 1321 ASTRO, Domaine de Duclos, F-97170 Petit-Bourg, Guadeloupe

* alienor.dutheil-de-la-rochere@inrae.fr

Abstract

Roots, tubers and bananas (RTB) products such as yam are staple food in Africa and the some countries of the Caribbean. Developers and breeders have worked for years to improve their yield and/or composition. However, they did not take into account consumer preferences such as cooking ability. Even ancient varieties highly appreciated by producers and consumers¹ have been evaluated for their yield and not for their quality traits including cooking ability.. Among the sensory criteria, texture is a primary quality factor for yam products, and it depends on both the initial characteristics of raw material and processing techniques ^{2,3}. In highly amylaceous products such as yam, the impact of starch on textural properties has been heavily explored. However, it cannot fully explain the texture and the cooking behaviour of these crops. Non-starch polysaccharides are known to be involved in the texture of non-starchy products such as apples⁴. Yam tuber hardness or mealiness after boiling has been hypothesized to be due to cell wall (CW) thickening, permeability changes and cell adhesion ^{2,3}. However there is little knowledge on the link between the texture of raw and boiled yam, and the relationship with cell wall structure and composition still remains to be detailed.

The aim of this work was to determine the CW polysaccharides (CWP) composition of yam tubers to evaluate the relationship between CWP, cooking ability and textural properties. To do so, cultivars and pedoclimatic conditions were varied to obtain yams with different tissue structure. Their cooking ability was evaluated (cooking time, texture). CWP were then extracted from five cultivars of *Dioscorea spp.* raw and steamed tubers, chosen for their highly-contrasted cooking behavior. A specific CW extraction procedure, involving an alcohol insoluble solids (AIS) extraction followed by an enzymatic starch removal procedure, was developed. Starch, lignin, CWP and pectin composition, acetylation and methylation degree were then determined by means of chromatographic and spectroscopic techniques (GC-FID, GC-MS).

AIS content in fresh tubers varies from 24% to 33%. The AIS contains 66% to 85% starch, representing 50 to 70% of the dry matter whereas cell walls weigh about 2%. Raw yam CWP contain 15 to 20% galacturonic acid and 20 to 25% galactose, which correlates with data reported previously for cassava roots⁵. Composition varies according to genotype, soil and cooking process. Steamed yam CWP show a decrease in galacturonic acid content (8 to 12%), which is consistent with β -elimination induced by the cooking process. This characterization of yam CWP in regard to cooking ability is the first of its kind. It gives an overview of West Indies yam composition and opens the way to a better understanding of yam quality traits for ancient varieties but also for the development of new hybrids. Many new varieties of yams, although resistant to diseases and pests have serious problems of acceptability and adoption by the targeted consumers. It is therefore important to provide breeders and developers with tools for the creation and selection of hybrids based on the expected sensory quality.

[1] Penet V, Cornet D, Blazy J - M, Alleyne A, Barthe E, Bussière F, Guyader S, Pavis C and Pétro, *Frontiers in Plant Science* 7:1962.

[2] Otegbayo B., Aina J., Sakyi-Dawson E., Bokanga M. & Asiedu R. *Journal of Texture Studies*, 36, 324-332, 2005.

- [3] Akissoe N., Mestres C., Handschin S., Gibert O., Hounhouigan J. & Nago M., *LWT - Food Science and Technology*, 44, 321-329, 2011.
- [4] Buergy A., Rolland-Sabaté A., Leca A. & Renard C.M.G.C. *Food Hydrocolloids*, 107, 105962, 2020.
- [5] Ngea G. L. N., Guillon F., Ngang J. J. E., Bonnin E., Bouchet B. & Saulnier L., *Food Chemistry*, 213, 402-409, 2016.