



**HAL**  
open science

## New virulent bacteriophages isolated from the surface of Epoisses cheese infect ripening bacteria

Thomas Paillet, Julien Lossouarn, Inès Pedros, Cédric Midoux, Olivier Rué,  
Marie-Agnès Petit, Eric Dugat-Bony

### ► To cite this version:

Thomas Paillet, Julien Lossouarn, Inès Pedros, Cédric Midoux, Olivier Rué, et al.. New virulent bacteriophages isolated from the surface of Epoisses cheese infect ripening bacteria. Sixth symposium of the French PHAGE Network, Oct 2021, Roscoff, France. hal-03441698

**HAL Id: hal-03441698**

**<https://hal.inrae.fr/hal-03441698v1>**

Submitted on 22 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# New virulent bacteriophages isolated from the surface of Epoisses cheese infect ripening bacteria

Thomas Paillet<sup>1</sup>, Julien Lossouarn<sup>2</sup>, Inès Pedros<sup>1</sup>, Cédric Midoux<sup>3</sup>, Olivier Rué<sup>3</sup>, Marie-Agnès Petit<sup>2</sup>, Eric Dugat-Bony<sup>1</sup>

<sup>1</sup>Université Paris-Saclay, INRAE, AgroParisTech, UMR SayFood, F-78850, Thiverval-Grignon, France; <sup>2</sup>Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, F-78352, Jouy-en-Josas, France; <sup>3</sup>Université Paris Saclay, INRAE, MaIAGE, 78350 Jouy-en-Josas, France



## Context



Bacteriophages are known to be major drivers of microbial communities in several ecosystems. In cheese, phages infecting lactic acid bacteria (LAB) starter cultures are well described. However, very little is known about those infecting ripening bacteria and their impact on the microbial successions observed during cheese maturation [1]. Recently, we used a viral metagenomics approach combined with transmission electron microscopy (TEM) to highlight the viral diversity present on the surface of Epoisses cheese [2]. Another group was also able to isolate and characterize several phages infecting the very common ripening culture *Brevibacterium aurantiacum* from smear and washed-rind cheeses or their production environment [3]. In order to evaluate the possible ecological role of phages present on the cheese surface, an effort was made to isolate and characterize some representative phages from Epoisses cheese.

## Materials & Methods


A collection of almost 200 bacterial isolates was retrieved from the surface of three Epoisses cheese and identified by sequencing the 16S rRNA gene. In parallel, the viral fraction was also purified according to the protocol described previously [2] containing the following steps: centrifugation (5,000 × g for 45 min), filtration (0.22µm), PEG precipitation and chloroform treatment. This viral particle pool was used to infect the collection of bacterial isolates through spot assays. Phages forming clear plaques were then isolated and purified three times. After purification, phages went through different characterization steps:

- Sequencing and annotation of complete phage genomes: after DNA extraction, long-reads were produced by a MinION device (Oxford Nanopore) and short reads by the NovaSeq platform (Illumina). An assembly was obtained from long-reads with Tricycler [4] and polished with short reads using Pilon [5]. The annotation was performed using RAST [6], supplemented with additional functional predictions using CDD [7], HHPred [8] and PHROGs [9].
- Morphological analysis of the virions by TEM
- Host range evaluation by spot assays against a range of reference strains corresponding to the same species as the original host as well as other species from the same genus.

## Isolation of bacteria and phages from Epoisses rind




- Glutamicibacter φ Voltaire**  
Host species: *Glutamicibacter arilaitensis*
- Psychrobacter φ d'Alembert**  
Host species: *Psychrobacter aquimaris*
- Leuconostoc φ Diderot**  
Host species: *Leuconostoc pseudomesenteroides*
- Brevibacterium φ Rousseau**  
Host species: *Brevibacterium aurantiacum*



Four virulent phages were isolated from the cheese rind. They infect four different bacterial species: *G. arilaitensis* and *B. aurantiacum* which are often used as ripening culture in the manufacture of washed-rind cheeses, *L. pseudomesenteroides* which is a non-starter lactic acid bacteria (NSLAB) and *P. aquimaris* which is frequently detected as a dominant species in washed-rind cheeses and is generally considered as an endogenous species.

## Sequencing, assembly and annotation





Tab1. Genomic information about the four isolated phages.

| Phage                      | Genome size (kb) | ORFs | Terminal Repeat (bp) | Best Blast hit (NCBI_viruses)                    |
|----------------------------|------------------|------|----------------------|--|
| Glutamicibacter φ Voltaire | 18               | 28   | 175                  | Brevibacterium phage Cantare 83,33% id 1% cov    |
| Psychrobacter φ d'Alembert | 92               | 158  | 5207                 | Vibrio phage vB_VhaM_VH-8 83,95% id 34% cov      |
| Leuconostoc φ Diderot      | 27               | 43   | /                    | Leuconostoc phage PhiLN03 98,20% id 98% cov      |
| Brevibacterium φ Rousseau  | 40               | 71   | /                    | Siphoviridae sp. Isolate ctmmc7 75,54% id 0% cov |

Following the latest proposed taxonomic criteria [10], where viruses with >70% nucleotide identity over the full genome belong to the same genus and viruses with >95% nucleotide identity belong to the same species, isolated phages may represent new genera except for Leuconostoc phage Diderot that belongs to the same species than PhiLN03.

## Morphology (TEM)




Tab2. Dimensions and morphologies of isolated phages

| Phage                      | Capsid (nm) | Tail (nm) | Morphotype |
|----------------------------|-------------|-----------|------------|
| Glutamicibacter φ Voltaire | 47          | 30        | Podophage  |
| Psychrobacter φ d'Alembert | 88          | 113       | Myophage   |
| Leuconostoc φ Diderot      | 57          | 141       | Siphophage |
| Brevibacterium φ Rousseau  | 62          | 177       | Siphophage |

Average sizes were obtained through the measure of 5 virions. They all belong to the *Caudoviricetes* class.

## Host range evaluation (Spot assay)



Tab3. Host range of isolated phages

| Phage                      | Sensitive species/ Tested species (same genus as the host) | Sensitive isolates/ Tested isolates (same species as the host) |                    |
|----------------------------|--|--|--------------------|
|                            |  | Isolated from Epoisses   | From other sources |
| Glutamicibacter φ Voltaire | 1/4  | 9/13   | 0/4                |
| Psychrobacter φ d'Alembert | 1/5  | 3/10   | 0/5                |
| Leuconostoc φ Diderot      | 2/4  | 7/9  | 0/1                |
| Brevibacterium φ Rousseau  | 1/6  | 1/2  | 0/16               |

Between 17 and 25 strains were tested in triplicate for each phage.

## Conclusions

New virulent phages infecting ripening bacteria were isolated from Epoisses cheese rind, and characterized. These steps allowed to confirm the originality of their genomes, as three of them may represent new viral genera. Their very narrow host range give first clues of their ecological impact in cheese ecosystem. New metaviromics analyses should allow to answer the question of the persistence of phages in Epoisses cheese.

## Acknowledgments

T.P. is the recipient of a doctoral fellowship from the French Ministry of Higher Education, Research and Innovation (MESRI) and the MICA department of INRAE. E.D-B is member of the knowledge hub SYSTEMIC, a joint action of JPI HDHL, JPI-OCEANS and FACCE-JPI launched under the ERA-NET ERA-HDHL (n° 696295).

## References

- Paillet, T. & Dugat-Bony, E. Bacteriophage ecology of fermented foods: anything new under the sun? *Current Opinion in Food Science* 40, 102–111 (2021).
- Dugat-Bony, E. et al. Viral metagenomic analysis of the cheese surface: A comparative study of rapid procedures for extracting viral particles. *Food Microbiology* 85, 103278 (2020).
- De Melo, A. G. de, Rousseau, G. M., Tremblay, D. M., Labrie, S. J. & Moineau, S. DNA tandem repeats contribute to the genetic diversity of *Brevibacterium aurantiacum* phages. *Environmental Microbiology* n/a, (2020).
- Wick, R. R. et al. Tricycler: consensus long-read assemblies for bacterial genomes. 2021.07.04.451066 <https://www.biorxiv.org/content/10.1101/2021.07.04.451066v1> (2021) doi:10.1101/2021.07.04.451066.
- Walker, B. J. et al. Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly Improvement. *PLOS ONE* 9, e112963 (2014).
- Aziz, R. K. et al. The RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9, 75 (2008).
- Lu, S. et al. CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res* 48, D265–D268 (2020).
- Zimmermann, L. et al. A Completely Reimplemented MPI Bioinformatics Toolkit with a New HHPred Server at its Core. *Journal of Molecular Biology* 430, 2237–2243 (2018).
- Terzian, P. et al. PHROG: families of prokaryotic virus proteins clustered using remote homology. *NAR Genomics and Bioinformatics* 3, (2021).
- Turner, D., Kropinski, A. M. & Adriaenssens, E. M. A Roadmap for Genome-Based Phage Taxonomy. *Viruses* 13, 506 (2021).