



HAL
open science

Heat treatment of concentrated milk protein system affect enzymatic coagulation properties

François Martin, Luisa Azevedo-Scudeller, Arnaud Paul, Romain Jeantet,
Gaëlle Tanguy, Serge Mejean, Cécile Le Floch-Fouéré

► **To cite this version:**

François Martin, Luisa Azevedo-Scudeller, Arnaud Paul, Romain Jeantet, Gaëlle Tanguy, et al.. Heat treatment of concentrated milk protein system affect enzymatic coagulation properties. Journées Plénières du GDR SLAMM (Solliciter LA Matière Molle), INRAE/CNRS, Nov 2021, Biarritz, France. hal-03451314

HAL Id: hal-03451314

<https://hal.inrae.fr/hal-03451314v1>

Submitted on 26 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License



l'institut Agro
agriculture • alimentation • environnement



Heat treatment of concentrated milk protein system affect enzymatic coagulation properties

François Martin^{1 2}, Luisa Azevedo-Scudeller³, Arnaud Paul⁴, Romain Jeantet¹,
Gaëlle Tanguy¹, Serge Mejean¹, Cécile Le Floch-Fouéré¹

1 STLO, INRAE, Institut Agro, 35042 Rennes, France

2 CNIEL, Paris, France

3 UMET, INRAE, PIHM Team, 59651 Villeneuve d'Ascq

4 Nancy Université, LIBio-ENSAIA-INPL, 54505 Vandoeuvre les Nancy, France

3 Novembre 2021, Biarritz

Email adress : francois.martin@inrae.fr



Context




- High added value products

- Expanding market

- Many Properties

- Coagulation
- Setting agent
- Emulsifier...



- Many outlets

- Infant formulas
- Cheese making
- Bakery...



Highly concentrated protein product with increasingly complex physico-chemistry

Process key stage : Heat treatment




More knowledge needed for controlling functionalities



Impact of heat treatments on highly concentrated dairy protein systems ?

➤ Scientific Background

Heat treatment of protein solution :
what happens in the case of milk?



Process : Temperature /
duration of heat
treatment

Physicochemical
conditions : pH, ionic
strength, whey
protein/Caseins ratio...

(Smits and Van Brouwershaven
.1980 ; Singh and Fox, 1985,1987 ;
Anema and Klostermeyer, 1997 ;
Oldfield et al. 2000 ; Anema and
Li. 2003, Anema, 2009 ; O'Connell
and Fox. 2003 ; Singh 2004)

UMET

l'institut Agro
agriculture • alimentation • environnement




LIBio
Laboratoire Français de Démolition



➤ Scientific Background

Heat treatment of protein solution :
what happens in the case of milk?



(Smits and Van Brouwershaven .1980 ; Singh and Fox, 1985,1987 ; Anema and Klostermeyer, 1997 ; Oldfield et al. 2000 ; Anema and Li. 2003, Anema, 2009 ; O'Connell and Fox. 2003 ; Singh 2004)

UMET

l'institut Agro
agriculture • alimentation • environnement




LIBio
Laboratoire Français des Dommages et des Pertes




➤ Scientific Background

Heat treatment of dairy solution with different protein concentration :

Heat treatment of milk



Heat treatment of highly concentrated dairy protein systems



- [Protein] : 33 g/L
- Ionic strength : 80 mM
- [Lactose] : 50 g/L



- [Protein] : 100-200 g/L
- Ionic strength : 150 - 300 mM
- [Lactose] : 2- 4 g/L

ie : Solution of dairy protein isolate

Many studies

(Smits and Van Brouwershaven .1980 ; Singh and Fox, 1985,1987 ; Anema and Klostermeyer, 1997 ; Oldfield et al. 2000 ; Anema and Li. 2003, Anema, 2009 ; O'Connell and Fox. 2003 ; Singh 2004)



Only parcellar understanding





➤ Research questions

Heat treatment of highly concentrated dairy protein systems


1/ Impact on heat-induced WP/Casein aggregation mechanisms ?

- Increase of collision probability
- Reaching of close packing of protein
 - ❖ Increasing formation of aggregates ?
 - ❖ Changes in whey protein/k-casein interaction ?
 - ❖ Changes in spatial location of protein complexes formed




Experimental strategy

➤ Experimental strategy



➤ Results : Protein denaturation / aggregation



HT After / 2HT :

- Highest whey protein denaturation level
- 2 times more than **Before** (ie 16% -> 30 %)

HT After / 2HT :

Very similar whey protein denaturation level (ie 27 and 30 %)


Literature data

Denaturation level : 30/35 % ->
Heat treatment 80°C / 6 min

(Giroux et al. 2020)

Protein concentration > parameter accelerating the whey protein denaturation / aggregation

➤ Results :Protein denaturation / aggregation




Type of aggregates formed ?



Quantitative analysis on the distribution of denatured whey proteins over WP aggregates and WP associated with the casein micelles


Coupled enzymatic and acid protein fractionation

(Noh et al.1989 ; Vasbinder et al. 2003)





➤ Results : Protein denaturation / aggregation

pH Heat treatment	Before	6.77	/	6.77
	After	/	6.68	6.70



Literature data

~ 6.70/6.75 -> 80 % solubles aggregates
(Anema et al.2003)




High protein concentration system

Different mechanism of k casein /whey protein association



➤ Result : Enzymatic coagulation properties



↗ protein aggregates formation
↳ ↗ gel time

In agreement with milk based studies


K-casein/Whey protein complexes disrupt hydrolyzed casein micelle aggregation

- Steric hindrance
- Electrostatic repulsion

(Kethireddipalli et al. 2015; Vasbinder et al. 2003 ; Donato et Guyomarc'h. 2009)



➤ Result : Enzymatic coagulation properties



Changes in K-casein/WP complexes ?



Mechanisms involved ?

Investigation of :

- Para casein micelles aggregation time
- Firming time


➤ Result : Enzymatic coagulation properties

Investigation of :


- Para casein micelles aggregation time
- Firming time

Using a light backscattering technique (Payne and Castillo. 2007 ; Bauland and al. 2020)

Time (s)



Shortest para-micelles aggregation time



Longest firming time



- Large number of K casein/whey protein complexes on micelles

- steric hindrance

2 HT -> preliminary hydrolyse of K-casein

Time to get 80% of micelles hydrolyse and start their aggregation is reduced

➤ Result : Enzymatic coagulation properties



✓ **HT After / 2 HT** : Whey proteins aggregates and k-casein/WP complexes disrupt gel reorganisation and generate very weak gels

(Giroux et al. 2015 ; Perreault et al. 2017)

✓ **HT Before** : Same gel strength as **Control** -> amount of whey proteins aggregates and k-casein/WP complexes seems to be insufficient to disrupt gel reorganisation



Aggregates quantity effect /
threshold effect

➤ Take home message

Heat treatment of highly concentrated dairy protein systems :

1/ Impact on heat-induced WP/Casein aggregation mechanisms ?

- Increase denaturation/aggregation of whey protein at 20 % (w/w) protein
- Drive K-casein/whey protein association toward major micellar aggregates

2/ Impact on the coagulation properties of the resulting heat-treated solution ?

- 2 Heat treatments -> generate K-casein hydrolysis -> reduce para-micelles aggregation time
- K-casein/whey protein complexes generated during heat treatment increase firming time, and lead to the formation of weak gels
- When only few aggregates are present in the bulk, gels produced have the same firmness than control gels



INRAE



➤ *Thanks*

Any questions ?