



HAL
open science

Modelling mixed crop-livestock farms for supporting farmers' strategic reflections: The CLIFS approach

Pierre-Yves Le Gal, Nadine Andrieu, Guillaume Bruelle, Patrick Dugué, Claude Monteil, Charles-Henri Moulin, Eric A. Penot, Julie J. Ryschawy

► To cite this version:

Pierre-Yves Le Gal, Nadine Andrieu, Guillaume Bruelle, Patrick Dugué, Claude Monteil, et al.. Modelling mixed crop-livestock farms for supporting farmers' strategic reflections: The CLIFS approach. Computers and Electronics in Agriculture, 2022, 192, 10.1016/j.compag.2021.106570 . hal-03460502

HAL Id: hal-03460502

<https://hal.inrae.fr/hal-03460502>

Submitted on 8 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Modelling mixed crop-livestock farms for supporting farmers' strategic reflections: the CLIFS**
2 **approach**

3
4 Le Gal P.-Y.^{1,2}, Andrieu N.^{1,2}, Bruelle G.^{1,2}, Dugué P.^{1,2}, Monteil C.^{3,4}, Moulin C.-H.^{5,6}, Penot E.^{1,2},
5 Ryschawy J.^{3,7}

6
7 ¹ CIRAD, UMR INNOVATION, F-34398 Montpellier, France

8 ² Innovation, Univ Montpellier, CIRAD, INRA, Institut Agro, Montpellier, France

9 ³ Université de Toulouse, INPT-ENSAT, F-31326 Castanet Tolosan

10 ⁴ INRAE, UMR DYNAFOR, F-31326 Castanet Tolosan

11 ⁵ Institut Agro, UMR SELMET, 2 Place Viala, 34060 Montpellier, France

12 ⁶ INRAE, UMR SELMET, 2 Place Viala, 34060 Montpellier, France

13 ⁷ INRAE, UMR AGIR, F-31326 Castanet Tolosan

14 Corresponding author: pierre-yves.le_gal@cirad.fr

15
16
17 **Abstract**

18
19 Integrated mixed crop-livestock farming systems help improve farm sustainability by securing
20 agricultural incomes through the diversification of productions and by enhancing farm autonomy
21 regarding agricultural inputs. Based on interactions between crop and livestock productions within
22 farms, such systems are complex to manage and to model. A one-to-one support methodology was
23 developed to help farmers in their strategic thinking regarding the future of their farms in a redesign
24 perspective. This methodology includes a three-step scenario process and is based on a spreadsheet
25 simulation tool called CLIFS (Crop Livestock Farm Simulator). CLIFS makes it possible to build
26 scenarios of the evolution of a farm and assess them *ex-ante* by calculating several balances at the
27 farm level (staple food, forage, manure) and their effects on the farm's economic results. The support
28 process has been tested in several African and South American contexts and with French suckling
29 cattle farms. The diversity of production contexts and issues addressed during the design process led
30 to the development of a generic tool that can be applied easily to a large range of situations. A
31 detailed description of the approach and the tool, with an illustration based on a Malagasy dairy farm,
32 are presented here. Farmers appreciate the support process because it addresses their own questions

33 within the context of their own farms. The process must now be transferred to advisory structures to
34 assess its relevance in a professional context.

35

36 **Key words:** design support, simulation, scenario, advisory

37

38

39 **1 Introduction**

40

41 The integration of mixed crop and livestock production on farms or within rural territories helps to
42 improve the sustainability of agricultural production (Herrero et al., 2010; Moraine et al., 2016).
43 Diversified production enables farmers to secure their incomes to cope with market fluctuations and
44 climate change while promoting biodiversity in certain modes of management (Kok et al., 2020).
45 Biomass exchanges between plant and animal enterprises help reduce the purchase of inputs and
46 increase farm autonomy (Bonaudo et al., 2014). In developed countries where the reintroduction of
47 these complex production systems is under debate (Schut et al., 2021), the survival of mixed crop-
48 livestock farms is being challenged by the specialisation of production between regions and a reduced
49 agricultural labour force (Ryschawy et al., 2013). In emerging and developing countries, some of these
50 mixed farms are benefiting from booming livestock sectors, particularly dairy and poultry (Sraïri et al.,
51 2013), but others are facing a decrease in available land, water resources and soil fertility due to
52 increasing population pressures (Kidron et al., 2010).

53

54 When mixed crop-livestock farmers wish to reconfigure their production systems to overcome certain
55 constraints, strengthen their autonomy with regard to inputs or respond to market demand, they must
56 consider the future interactions of all of their farm's production enterprises. The viability of their
57 projects depends on the balance between the land, forage and organic manure resources available
58 and the needs generated by their production objectives. This reflection may require the support of an
59 agricultural advisor which goes beyond conducting economic assessments (Penot, 2012), drawing
60 comparisons with farm types that are more or less similar to the farmer's specific situation (Titonnell et
61 al., 2009) or using technical management tools such as livestock ration calculators (FAO, 2016).

62

63 The use of calculation tools that evaluate *ex ante* the consequences of a given farm configuration on a
64 farm's technical-economic performance has proven useful in guiding farmers' reflections about future
65 directions (Semporé et al., 2015; Colnago et al., 2021). Most modelling work related to this activity has
66 a research objective, both in the design of the models and tools used, and in the intended uses (Le
67 Gal et al., 2011a). Many tools therefore provide a representation of a farm based on biotechnical
68 models, coupled with decision modules allowing the advantages of an innovation or a given farm
69 configuration to be tested (Snow et al., 2014). Optimisation models also are used for this purpose on
70 typical farms, for example to assess the benefits of conservation agriculture systems (Alary et al.,
71 2016) or the integration of production activities (Mosnier et al., 2017). Rule-based models aim to
72 reproduce farmers' decision-making processes (Vayssières et al., 2009), but are often difficult to use,
73 and do not always capture the complexity of real situations.

74

75 As these tools are based on a detailed representation of a mixed crop-livestock farm's operations, they
76 are difficult to understand by a farmer and to use for supporting his/her reflections. The use of
77 simulation tools representing these operations in a simplified but realistic way is one way to achieve
78 this objective (Le Gal et al., 2013). In the domain of mixed crop-livestock farming, these tools take
79 various forms, from calculation tools that compare the biomass produced by pastures or crops with the
80 biomass ingested by livestock (Lurette et al., 2013), up to board games that concretely represent
81 forage calendars, sometimes supplemented by spreadsheets (Martin et al., 2011). These tools are
82 often specialised for a certain type of livestock system (Machado et al., 2010; Parsons et al., 2011), or
83 a certain production context (Andrieu and Nogueira, 2010; Lisson et al., 2010). Despite their
84 operational objectives, they often use crop and livestock models that require certain data, the
85 availability of which varies with the work context, to be validated in the specific conditions of a farm.

86

87 This article presents an approach that aims to help farmers to think about the future direction of their
88 farms in terms of introducing or expanding crop and livestock production activities, or of introducing
89 technical innovations impacting all or part of their farms' operations. It relies on the use of a simulation
90 tool named CLIFS (Crop Livestock Farm Simulator) which is devoted to mixed crop-livestock farms.
91 The approach is specifically intended to be used with farmers and transferred to farm advisors. Earlier
92 versions of CLIFS were tested on samples of 2 to 20 family farms in a range of contexts, and in both

93 tropical and temperate environments (Le Gal et al., 2013; Ryschawy et al., 2014; Semporé et al.
94 2016). This article presents the current version, developed in Microsoft Excel®, which can be
95 downloaded freely at the following address: <https://doi.org/10.18167/DVN1/NZHWQQ>. A User Manual
96 (UM) provides a full description of the software and its equations to which the reader can refer for
97 additional information (Le Gal, 2021; <https://doi.org/10.18167/agritrop/00577>). The use of the
98 approach is then illustrated with the case of a dairy farm in Madagascar. In the discussion section, the
99 approach is assessed from the point of view of researchers and farmers, and its potential use is
100 explored.

101

102 **2 Empirical and methodological background**

103

104 **2.1 A set of seven case studies**

105

106 The design and development of the CLIFS approach drew from a set of case studies conducted over
107 ten years (from 2004 to 2013) on seven contrasting sites which differed in terms of the mixed crop-
108 livestock farming systems in place and the issues addressed (Table 1). This diversity gives the
109 approach and the application their generic character.

110

111 The case studies were distinguished by the type of livestock involved in the farming system: dairy
112 farming (five cases), mixed farming (one case) and suckler farming (one case), possibly combined
113 with a monogastric enterprise (pig, poultry). The size of the farms studied varied greatly, from very
114 small dairy farms in Morocco and Madagascar, to herds and cultivated areas comprising several
115 dozen heads and hectares. The forage systems encountered generally combined several sources of
116 biomass used differently depending on the time of year, including open natural pastures (Burkina
117 Faso, Madagascar), grazed cultivated grasslands (Brazil and France), forage crops cut and distributed
118 green (Morocco, Madagascar, Peru) or stored after silage or haymaking. Crop residues were used
119 widely in Burkina Faso and Peru.

120

121 These forage crops generally were combined on the farms with food and cash crops, which were more
122 or less diverse depending on the case study. Apart from the French case, the small family farms met

123 all or part of their cereal needs with their own production (wheat, rice or maize depending on the
124 case). They also were part of marketing channels through which surplus foodstuffs and specific crops
125 such as sugar beet in Morocco or cotton in Burkina Faso were sold. The crops could be irrigated or
126 rain-fed, with different yield potentials and periods of biomass availability. For example, irrigated alfalfa
127 in Morocco allowed dairy herds to be fed all year round, whereas cultivated rainfed grasslands in
128 Brazil and France could only be used during the period when grass was growing.

129

130 The issues addressed differed according to the type of livestock sector, the availability of biophysical
131 and socioeconomic resources, and the climate context. On dairy farms that were part of a dynamic
132 sector (Brazil, Morocco, Peru, Madagascar Highlands), most farmers were considering how to
133 increase their production, either by increasing the size of their herd or by increasing the productivity
134 per cow. These questions could be combined with more specific problems, such as reducing the
135 consumption of irrigation water in Morocco and Peru by diversifying forage resources, or improving soil
136 fertility in Burkina Faso and the Lake Alaotra region of Madagascar. In the view of local farmers, crop-
137 livestock integration and the introduction of cover crops combining soil protection and forage use (the
138 Madagascar case) were key to the long-term stability of agricultural production. In the French case,
139 the mixed crop-livestock farms studied were looking for solutions to ensure their sustainability in an
140 economic environment encouraging specialisation, such as reducing production costs through forage
141 autonomy or diversifying livestock sales outlets.

142 **Table 1.** Main characteristics of the seven case studies.

143

Country	Brazil	Burkina-Faso	France	Madagascar	Madagascar	Morocco	Peru
Location	Cerrados	Cotton belt	South-West	Lake Alaotra	Highlands	Tadla plain	Andean valley
Period of study	2009-2011	2007-2011	2013	2009-2013	2012	2004-2008	2011-2013
Involved farm number	6	24	2 + one group of 15 farmers	6	8	12	10
Production system	Rainfed crops Dairy cattle	Mixed crop-livestock	Rainfed crops Suckler cattle	Irrigated/rainfed crops x Dairy cattle	Irrigated/rainfed crops x Dairy cattle	Irrigated crops Dairy cattle	Irrigated crops Dairy cattle
Farm size	10-30 cows over 15-30 ha	1-3 cows 4 – 25 ha	43-50 cows over 85-130 ha	1-3 cows over 3-8 ha	2-11 cows over 1-24 ha	5-6 cows over 2-30 ha	3-65 cows over 1-60 ha
Livestock	Dairy cows	Oxen Suckler cows Fattening cattle Fattening sheep	Suckler cows Fattening cattle	Dairy cows Pigs, poultry	Dairy cows Pigs, poultry	Dairy cows	Dairy cows
Forage system ¹	<i>Brachiaria dec.</i> Sugarcane <u>Maize silage</u>	<u>Crop residues</u> (cereals, cowpea) <u>Mucuna pruriens</u> <i>Natural pasture</i>	<i>Cultivated pasture</i> <u>Grass/hay</u> <u>Maize</u>	Brachiaria ruz. ¹ Styloxanthès guia. ¹ Vetch ¹ Wild grass <u>Rice straw</u>	Oat Pennisetum kizosi Ray-grass Forage maize <u>Rice straw</u>	Alfalfa Berseem <u>Silage maize</u> <u>Alfalfa hay</u> <u>Wheat straw</u>	Alfalfa Oat-vetch <u>Maize stems</u>
Main other crops	Maize	Maize	Winter wheat	Irrigated/rainfed	Irrigated/rainfed	Winter wheat	Maize

	Rice	Cotton	Soya	rice	rice	Sugar beet	Potato
	Beans		Sunflower	Maize	Maize+Bean Sweet potato	Vegetables	Vegetables
Raised issues	Milk production increase	Crop-livestock integration	Forage autonomy Diversification of livestock product	Cover crop introduction	Increase of forage availability	Milk production increase Water consumption reduction	Milk production increase Forage diversification
References	Le Gal et al., 2013	Semporé et al., 2016	Ryschawy et al., 2014	Douhard, 2010 Foussat, 2011	Mouret, 2012	Le Gal et al., 2011b	Biens and Le Gal, 2012

144 ¹ Normal font: cut green and provided in troughs Underline font: stored and provided in troughs *Italic font*: grazed by herd

145

146 **2.2 A generic view of mixed crop-livestock farms**

147

148 The design of CLIFS is based on a generic representation of mixed crop-livestock farms (Figure 1)
149 that was derived from observations made in the seven study areas. This representation is organised
150 around exchanges between crop activities, which produce biomass on the farm, and livestock
151 activities, which consume biomass and produce organic manure that can be used on the crops. The
152 herd can be composed of ruminant and monogastric animals. For ruminants, a distinction is made
153 between (i) breeding females producing milk that is sold or consumed by their young, (ii) animals kept
154 for renewal, savings, traction, and reproduction, and (iii) animals that are fattened and sold.
155 Monogastrics are distinguished by their function: breeder or producer of meat and eggs. All of these
156 animals produce manure, either directly on the fields while they are consuming the biomass available
157 there, or while stabled in livestock buildings or pens. In the latter case, the manure can be mixed with
158 litter from crop residues to form organic fertiliser. This can then be spread on the cultivated fields. This
159 process provides more control over the quantity and quality of the organic manure than the practice of
160 keeping animals on the fields (Blanchard et al., 2014).

161

162 The crops potentially present on the farm are grouped into three categories: (i) crops to feed the
163 farmer's family, the surplus of which can be marketed once family needs have been met. Some crops
164 such as grain maize also can be fed to animals, both as the main feed for monogastric animals or as a
165 supplement for ruminants; (ii) crops intended solely or mainly to be sold, such as cotton and
166 groundnuts, but which could contribute to the family's diet; (iii) grassland and forage crops directly
167 intended to feed ruminants; when there is a surplus in relation to the herd's needs, these may be
168 marketed. This biomass can be distributed in three forms: green, hay or silage. This on-farm forage
169 resource is supplemented by crop residues from food crops, such as cereal straw, and from marketed
170 crops, such as groundnut leaves. All of these crops can potentially receive organic manure produced
171 by the farm's herd.

172

173 The management of crop and livestock mobilises the workforce, family and hired labour, which is
174 allocated by the farm head throughout the year to the various agricultural activities according to needs.
175 While crop management is determined by crop cycles, seasons and technical practices, livestock

176 farming involves both seasonal work and routine work such as milking reproductive females, tasks that
177 are repeated every day for all or part of the year (Hostiou and Dedieu, 2012). Mixed crop-livestock
178 farms therefore represent a complex situation in terms of work organisation.


179

180 The farm is open to its environment, with which it interacts to (i) generate income from the sale of its
181 products according to its marketing strategy, and (ii) obtain goods and services for the farm: inputs for
182 crop (seeds, fertilisers, pesticides, mechanised services) and livestock activities (fattening animals,
183 food supplements, veterinary interventions); seasonal labour to supplement the permanent labour
184 force at certain times of the year; pastures and natural biomass outside the farm on which animals can
185 be fed directly or after mowing and distribution.

186

187 Mixed crop-livestock farmers must therefore make strategic and tactical choices concerning (i) the
188 nature of the crop and livestock production they wish to develop in relation to the desired degree of
189 specialisation vs diversification, (ii) the coordinated sizing of the crop and livestock enterprises with
190 regard to the desired degree of autonomy in terms of animal feed and crop fertilizers from outside the
191 farm, (iii) the way in which the crops and the herd will be managed in order to attain a given production
192 objective. The timing of biological cycles is an important element to consider in these choices,
193 including crop cycles according to the seasons and irrigation options, which determine in particular the
194 moment when biomass is available for animals and whether or not it must be stored as hay or silage
195 for later use, and animal breeding cycles which may or may not be synchronised with previous ones.
196 The internal degree of integration between crops and livestock, considered as one of the levers
197 leading towards more agroecological production systems (Ryschawy et al, 2017), depends on this set
198 of interlocking decisions.

199



200

201 **Figure 1.** Generic representation of the components and flows between components of mixed crop-
 202 livestock farms.

203

204 **2.3 A three-step support process**

205

206 CLIFS is first and foremost an individual support tool that aims to provide farmers elements to consider
 207 and assess when considering a medium to long-term development project for their farms. Its main
 208 objective consists of supporting farmers' reflections in an exploratory and iterative way rather than
 209 defining precisely the project content and how it will be implemented. Indeed, the strategic evolution of
 210 a farm includes many interacting elements that are difficult to comprehensively address, especially
 211 since the future of some of these elements remains uncertain. This approach has three steps based
 212 on the (i) design, (ii) simulation and (iii) assessment of successive scenarios (Figure 2). Each scenario
 213 corresponds to a configuration of the farm that is simulated over one year.

214

215 The Initial Scenario (IS) is based on an analysis made with the farmer about the current situation on
 216 his/her farm. This analysis allows the advisor to better understand the farmer's objectives and

217 strategies, to characterise the farm's structure, operations and performance, and to calibrate certain
218 CLIFS input variables that are difficult to access, such as pasture productivity. Depending on the data
219 available, it may take several loops to arrive at a representation of the farm that the farmer thinks to be
220 consistent with reality. This representation is then considered to be valid for the next steps in the
221 process. This validation stage is important to ensure that the rest of the process is based on
222 knowledge that is shared by the farmer and the advisor, and to enable the farmer to understand the
223 structure of CLIFS and the calculations made.

224

225 The next step is to build a coherent and balanced Project Scenario (PS) corresponding to choices as
226 to how the farm might evolve. CLIFS highlights possible imbalances between supply and demand for
227 resources, for example between the size and productivity of a dairy herd and the forage resources
228 generated by the cropping pattern planned. At the end of this step, which may include several
229 intermediate scenarios if necessary, farmers have a more precise and concrete idea of their project
230 and the consequences on their production and economic results.

231

232 During the third step, Alternative Scenarios (AS) are developed by the farmer and the advisor based
233 on the results of the PS and proposed changes or technical innovations considered to be potentially
234 interesting. If the farmer is satisfied with the PS, this last step is optional. Nonetheless, it opens up the
235 field of possibilities and enriches the thinking of both participants, as the farmer may ultimately prefer
236 one of these AS. It also makes it possible to assess *ex ante* the relevance of an innovation at the level
237 of a given farm, and to evaluate the variability of scenario results according to changes in prices or
238 yields, for instance.


239

240 The loop may be run through as often as needed to test new ideas or to assess the sensitivity of a
241 farm configuration to variables such as product and input prices or variations in yields according to the
242 climate year. As CLIFS is a static tool, these analyses involve multiplying scenarios that vary the
243 values given to these variables. This procedure allows the user to maintain control over the changes
244 made during the support process. Once completed, the process can be evaluated with the farmer to
245 assess what the user has learned from it (Matthews et al., 2010). The approach does not include
246 support for the eventual implementation of the preferred scenario. In some cases, however, we

247 returned to the farms a few years later to discuss with the farmers what they finally had achieved, as in
 248 the case presented in this article.

249
 250 CLIFS can be used to address a wide range of farm issues and projects, such as the choice and
 251 resizing of livestock enterprises to increase milk production, the choice of a forage system for self-
 252 sufficiency, the introduction of innovations such as catch crops or the partial use of cover crop
 253 biomass, and the analysis of a farm's sensitivity to climatic and economic shocks.

254



255
 256

257 **Figure 2.** Organization of the farmer support approach in three main loops.

258
 259

260 **3 CLIFS description**

261

262 **3.1 Overview**

263

264 CLIFS translates the generic representation of a mixed crop-livestock farm (Figure 1) into a
 265 spreadsheet-based simulation tool (Microsoft Excel®). The format makes it usable and accessible to a
 266 wide range of users. The overall structure of the tool, along with its calculation procedures and output
 267 variables, have been designed so that the farmers involved can understand them while also providing
 268 a representation that closely matches the farms which the farmers can validate. This led to limiting the
 269 number of variables to be characterized on each farm and to excluding the integration of crop and
 270 livestock models, which are furthermore often unavailable in many contexts.

271
 272 The only equations related to biological processes concern: (i) for all ruminants, excreta production as
 273 a function of their live weight to calculate the supply of organic manure as a function of herd structure
 274 and management (Table 5); (ii) for cattle only, energy and nitrogen requirements for animal
 275 maintenance, gestation of breeding females, and production of milk and meat per head (Table 2) in
 276 order to link these productions to the rations distributed, with the choice between two calculation
 277 systems independent of the user's working language: French (INRA, 2007) and American (NRC,
 278 2001). The user can nevertheless inactivate these calculations of milk and meat production if required
 279 data are unavailable or another feed system than those proposed is being used. The other
 280 calculations only use the four mathematical operators.

281

282 **Table 2.** Equations of energy and protein requirements for a lactating cow according to the feed value
 283 system

		INRA	NRC
Daily maintenance	Energy	$5 + ((LW_b - 600) \div 100) \times 0.6$	$0.079 \times LW_b^{0.75}$
	Protein ($NIMR_b$)	$395 + ((LW_b - 600) \div 100) \times 50$	$3.8 \times LW_b^{0.75}$
Pregnancy	Energy	$0.00072 \times CBW_b \times e^{(0.116 \times g \times 4.33)}$	$g < 7 : 0$ $g > 6 : (0.00318 \times DAF_g - 0.0352) \times (CBW_b \div 45) \div 0.218$
	Protein	$g < 7 : 0$ $g = 7 : 0.18 \times NIMR_b$ $g = 8 : 0.33 \times NIMR_b$ $g = 9 : 0.50 \times NIMR_b$	$g < 8 : 0$ $g > 7 : ((0.69 \times DAF_g) - 69.2) \times (CBW_b \div 45) \div 0.33$
Production of 1 liter of milk	Energy	0.44	0.699
	Protein	48	0.05

284 With:

285 LW_b : Live weight of the average cow of batch b (kg; considered as constant throughout the year)

286 CBW_b : Calf Birth Weight for batch b (kg)

287 g : Month after fertilization [1,9]

288 DAF_g : Day after fertilization [from $g=1$ to $g=9$: 15; 45; 75; 105; 135; 165; 200; 230; 260]

289

290 For each farm configuration simulated, CLIFS calculates the resource supply-demand balances of
291 three key components of mixed crop-livestock farms, namely: (i) the annual balance of food and cash
292 production, between the supply per crop linked to its area and yield and its consumption by the family
293 and livestock (UM F.6); (ii) the monthly forage balance between the supply linked to the forage system
294 (area and yield per forage crop, crop residues) and the demand linked to the ruminants, which itself
295 depends on herd structure (numbers per type of animal), the diet over the 12 months of the year and
296 the reproductive strategy for female breeders (distribution of births over the 12 months of the year)
297 (UM F.3, F.4 and F.5 for green forage, hay and silage, and crop residues, respectively); (iii) the annual
298 balance of organic fertiliser based on the manure produced by stabled animals and the use of the
299 organic fertiliser on crop fields (UM F.7). In addition to these three balance sheets, the economic
300 results (variable and fixed costs, gross and net margins) corresponding to each farm configuration are
301 calculated (UM F.8). However, CLIFS does not include a labour balance sheet due to the time
302 required to estimate labour supply and demand with a precision that makes sense to farmers.

303

304 CLIFS is available in English, French, Spanish and Portuguese for names and titles that cannot be
305 modified by the user. However, users can enter in their own language the contents of the choice lists
306 linked to certain variables.

307

308 **3.2 Structure and operation**

309

310 *3.2.1 General structure*

311


312 CLIFS consists of a series of Excel® sheets grouped into three linked modules (Figure 3, and UM C.
313 for a full list of the sheets):

314 (i) Parameters (UM D.) with seven sheets grouping variables whose values are identical for a set of
315 farms, such as the characterisation of animals' feed and requirements;

316 (ii) Input variables, specific to each farm, with an "animal" sub-module of 16 sheets characterising the
317 dairy, growth and fattening ruminant enterprises and the batches of monogastric animals (UM E.1 to
318 E.8); and a "crop" sub-module of six sheets characterising the crop blocks making up the farm. Each
319 block is defined by the combination of one to two food, cash or forage crops, the technical practices


320 applied and the associated yields. It may or may not correspond to physical crop fields (UM E.9 to
 321 E.14). Also included is an economic sub-module of four sheets to enter prices of inputs and services,
 322 and prices of marketed products (UM E.15 to E.18);
 323 (iii) Output variables, grouping all of the balances and economic calculations resulting from the farm's
 324 sizing, technical choices and performance (UM F.) in 14 sheets;
 325 Input and output variables are linked through a set of calculations included in sheets not visible to the
 326 user (UM Eq.1 to Eq.22).

327
 328



329

Legend



330

331 **Figure 3.** CLIFS structure into three modules and links between modules.

332

333 Each batch of ruminants is represented as a sum of individuals homogeneous in weight, and female
334 breeders by a lactation curve. Each animal receives the same daily diet comprising a forage
335 component, which can be modified each month from January to December for breeding and growing
336 animals, and a concentrates component, distributed to breeding animals according to their monthly
337 stage of lactation (Figure 4). Fattening enterprises are defined by their duration and position in the
338 year, the number of animals concerned and a uniform daily forage+concentrates diet for each period
339 considered. CLIFS considers that the feed distributed or grazed is ingested completely by the average
340 animal per batch as long as the animal's intake capacity is not saturated. The user must decrease the
341 distributed quantities when the calculated forage saturation exceeds this capacity.

342

343 The structure of Excel® in sheets and tables limits the number of elements that can be characterised
344 per activity (UM B.3). The sizing was determined to (i) make it easier to read the tables on the screen,
345 and (ii) represent many different production systems, combining cattle, small ruminants and
346 monogastrics for livestock, and food, cash and forage crops (including permanent grasslands) in pure,
347 associated or catch crop form for crops.

348

349 CLIFS uses different time steps depending on the process represented (Table 3). These time steps
350 are those used by farmers to manage their production activities while being aggregated to limit the
351 amount of data needed to be entered. For example, diets are entered in kg per day, which can be
352 modified each month, i.e., 12 values to be entered per feed. Similarly, the units used vary according to
353 the variables (UM B.2). They are either specified in the row and column headings (e.g., kg/ha for crop

354 yields), or left to the user's choice (e.g., quantities of all crop inputs except organic fertilisers given in
 355 kg/ha). The monetary unit related to the economic variables is also left to the user's choice.

356

357 **Table 3.** Time steps used in CLIFS according to the modelled item.

Modelled item	Time step	Comments
Input variables		
Lactation curve (l/day)	Daily	One value entered every month over 12 months
Birth distribution	Monthly	From January to December
Diet		
Ruminants (kg/day)	Daily	
Lactating females		One value entered every calendar month for forage and every month after parturition for concentrates
Growing animals		One value entered every calendar month for forage and for concentrates
Fattening animals		Uniform for the whole fattening period
Monogastrics (kg/year)	Annual	Total quantity of concentrates distributed per batch
Manure production		
Total duration of stabling	Day	Between 1 and 365 days over the year
Daily duration of stabling	Hour	Between 1 and 24 h
Technical sequence per crop block	Crop cycle	From land preparation to harvest over the year
Forage yields (%)	Month	From January to December
Forage type (hay/silage)	Month	From January to December
Fixed costs	Annual	Total amount per cost
Result variables		
Milk production	Month	+ annual total
Green forage balance	Month	+ annual balance
Hay/Silage stock balance	Month	+ annual balance
Crop residue balance	Annual	
Food/Market balance	Annual	
Organic fertilizer balance	Annual	
Economic results	Annual	

358

359 For each change introduced into a scenario, the internal consistency of the data entered must be
360 verified since the mechanistic links between variables are limited to cattle feed, if the user wishes to
361 use this functionality, and to excreta production for ruminants. This vigilance is required in particular to
362 ensure consistency between technical practices and corresponding yields per crop block, and between
363 livestock diets and milk and meat production in the absence of equations (in the case of small
364 ruminants) or when the proposed equations (Table 2) are considered unsuitable for the case studied.
365 This alignment may be based on expert knowledge, whether provided by the farmers themselves, the
366 users of the tool or specialists in the field, or from crop models and rationing tools that are not part of
367 CLIFS as long as their area of validity covers the situation under study.

368

369 *3.2.2 Calculation of food-market crop balance*

370

371 On farms combining self-consumption, sales and herd feeding for certain crops such as maize,
372 farmers must reconcile different objectives: to cover family food needs, meet production targets related
373 to the herd, and generate a certain income. CLIFS therefore calculates the family's food needs for up
374 to two crops from the list defined by the user, based on the structure of the family and the annual
375 quantities needed to cover the needs of one person.

376

377 The food balance for the two selected crops corresponds to the difference between their total
378 production on the farm and the amount consumed by the family over the year. A negative value
379 generates an expense based on the market purchase price entered by the user. These amounts are
380 not deducted from the net margin of the scenario but their total is provided as an indicator of non-
381 achievement of the food self-sufficiency objective. A positive value leads to the calculation of a second
382 balance, which subtracts the total quantities consumed by the animals from this value. A negative
383 balance generates a purchase, a positive one, proceeds.

384

385 *3.2.3 Forage balance*

386

387 The forage balance makes it possible to assess the degree of a farm's forage self-sufficiency for each
388 scenario designed. Each forage is the subject of a calculation that subtracts the herd's total

389 consumption according to the diets applied in the different enterprises from the supply produced by
 390 the farm (see Table 4 with the example of green forage). For each forage crop, this supply
 391 corresponds to the sum of the {area x yield} of the blocks where it is grown pure or in association.
 392 Yields are entered in gross weight and not in dry matter to be consistent with farmers' practices
 393 regarding the quantification of diets distributed. The annual quantity produced is then distributed on a
 394 monthly basis using a percentage of the yield depending on the production dynamics of the different
 395 forage crops (growing all or part of the year like grassland or alfalfa, or without regrowth like forage
 396 maize). The user then specifies for each crop and each month how it is used: green (grazing on the
 397 field or distribution synchronised with the harvest), stored as hay, or silage.

398

399 **Table 4.** Equations used for calculating green forage balance

Balance component	Equation
Supply for forage f month m $Prod_GF_{f,m}$	$\sum_l Surf_GF_{f,l} \times Yield_{f,l} \times \%Yield_{f,m} \div \sum_{m=1}^{12} \%Yield_{f,m}$
Annual supply for forage f $Prod_GF_f$	$\sum_{m=1}^{12} Prod_GF_{f,m}$
Consumption of forage f month m $Q_GF_{f,m}$	$\sum_b Q_{f,m,b} \times Day_m \times n_b$
Annual consumption of forage f Q_GF_f	$\sum_{m=1}^{12} Q_GF_{f,m}$
Balance for forage f month m $Bal_GF_{f,m}$	$Q_GF_{f,m} - Prod_GF_{f,m}$
Annual balance of forage f	$Q_GF_f - Prod_GF_f$
Total annual deficit of forage f	$\sum_{m=1}^{12} Bal_GF_{f,m} \text{ with } Bal_GF_{f,m} < 0$

400 With:

$Surf_GF_{f,l}$	Area of green forage crop f block l (these blocks can bear up to two forage crops f)
$Yield_{f,l}$	Total gross yield of green forage f on crop block l (kg/ha)
$\%Yield_{f,m}$	Percentage of the total gross yield of forage f for month m
$Q_{f,m,b}$	Quantity of forage f provided daily per ruminant head during month m for batch b (kg green matter)
Day_m	number of days of month m during which animals are fed
n_b	Number of heads in each of the seven ruminant batches

401

402 The demand is calculated monthly for each forage based on the diets distributed in each enterprise,
 403 multiplied by the size of the enterprise (Fig.4 for an example of a diet in a dairy unit). A monthly
 404 balance is then calculated by comparing the supply and demand from January to December. For
 405 green forage, no carry-over from one month to the next is possible. For hay and silage, the monthly
 406 evolution of the stock is calculated, which makes it possible to pinpoint the date when a shortage is

407 possible. A positive balance, monthly for green forage or at the end of the annual cycle for stocks,
 408 reflects an imbalance between production and needs. However, surplus stocks can be sold if a market
 409 price exists. A negative balance triggers the purchase of forage, whatever its type, which is added to
 410 the farm's expenses if a market price exists. In the absence of a market, the scenario highlights a
 411 structural forage imbalance that must be corrected by reconfiguring demand (nature and size of the
 412 herd, diets) or supply (cropping pattern and yields). The user then has to evaluate these changes,
 413 including the choices and management of animal diets, by designing and simulating additional
 414 scenarios since CLIFS does not provide any optimisation algorithm to define a relevant farm
 415 configuration.

416
 417 Crop residues (straw, stover, tops) stored dry by the farm at harvest and then incorporated into diets
 418 are subject to a supply-demand balance only on an annual basis. The supply is calculated from the
 419 crop production to which a grain/residue ratio based on raw material is applied. The demand includes
 420 the quantities consumed by the ruminants and those incorporated into organic manure.

421

422 3.2.4 Manure balance

423

424 With CLIFS, it is possible to define up to 10 types of organic manure combining, in proportions chosen
 425 by the user, a type of excreta based on the animals present on the farm and a crop residue (e.g., rice
 426 or wheat straw) used as litter. The supply in gross weight of each type of manure is then calculated as
 427 a function of the time spent in stable stalls by each group of animals and the size of each group (Table
 428 5). Only the organic manure produced is considered to be available for return to the plots via crop
 429 fertilisation. The contribution of excrement deposited while animals are grazing on fields, which is still
 430 frequent in certain regions such as Burkina Faso, is not included in the quantities of manure provided
 431 because it is difficult for farmers to control.

432

433 **Table 5.** Equations used for calculating manure production by ruminant batches

Production component	Equation
Daily production of excreta by one animal of type <i>a</i> in batch <i>b</i> ¹	$LW_{a,b} \times 0.01 / DM_Exc_x$
$QD_Exc_{x,a,b}$	
Annual production of excreta <i>x</i> by one animal <i>a</i> in batch <i>b</i>	$QD_Exc_{x,a,b} \times Dur_Stab_{a,b} \times (HD_Stab_{a,b} \div 24)$

$$\frac{QY_Exc_{x,a,b}}{Prod_Manure_{u,b}} = \frac{QY_Exc_{x,a,b} \times n_man_{a,b} \times Conv_Exc_{x,b} \times ((100 - Loss_Exc_{u,b}) \div 100) \div ((100 - \%Exc_{x,u}) \div 100)}{Prod_Manure_{u,b}}$$

434 With:

- $LW_{a,b}$ Live weight of the average animal *a* of batch *b* (kg)
- DM_Exc_x Dry matter of excreta *x* (g/kg)
- $Dur_Stab_{a,b}$ Duration of stabling (days) of animal *a* in batch *b*
- $HD_Stab_{a,b}$ Daily duration of stabling (hours) animal *a* in batch *b*
- $n_man_{a,b}$ Number of heads of animal *a* in batch *b*
- $Conv_Exc_{x,b}$ Conversion rate for excreta *x* in batch *b*²
- $Loss_Exc_{u,b}$ Loss rate (%) of manure *u* in batch *b*
- $\%Exc_{x,u}$ Proportion of excreta *x* in manure *u* (%)³

435 ¹The coefficient '0.01' is based on the following assumptions:

- 436 - A ruminant eats 2.5% DM of its live weight
- 437 - The diet has an average digestibility of 60%
- 438 - The dried excreted quantities = $LW_{a,b} \times 0.025 \times (1 - 0.60)$

439 ²When excreta are processed before their use as fertilizer.

440 ³Considering that a given manure can consist of a mix of excreta and crop residue.

441

442 The quantities applied per hectare by type of organic manure are entered with all of the technical
 443 practices used on each crop block. These quantities come from farmer declarations, local standards,
 444 or research results. The balance is calculated for each type by subtracting the sum of the quantities
 445 applied to the blocks from the total quantity produced. A negative balance indicates too much demand
 446 for manure while a positive one indicates a surplus. If a market exists for the corresponding manures
 447 and prices have been entered these values trigger respectively a purchase or a sale.

448

449 3.2.5 Milk and meat production for the ruminant batches

450

451 Milk production is calculated monthly for each batch of breeding females by first defining with the
 452 farmer a production target based on the genetics of the herd, or depending on what the farmer
 453 believes he/she is able to produce from the farm's feed resources (UM E.4 and UM Eq.5 to Eq.13 for a
 454 detailed presentation of the procedure, the variables and the equations used). This objective is
 455 translated into a simplified lactation curve applied to all of the breeders in the batch according to their
 456 calving months. For cows only, the production linked to the diet ingested each month is then
 457 calculated from the feed values of the forage and concentrates concerned (Figure 4). Concentrate
 458 quantities are entered according to the lactation stage of the breeders and not the calendar month.
 459 This practice is in fact widely used in most of the dairy farms observed. The final production


460 considered each month is equal to the minimum of the production target and the productions linked to
 461 the diet's protein and energy content.

462

463 A graph is provided showing the production curves related to the target and those to the diet. In cattle,
 464 the comparison of these curves makes it possible to detect, month by month, phenomena of over or
 465 under-feeding that may have multiple causes. During IS characterization these include poor
 466 knowledge of the animals' requirements and of the dietary values of the inputs, an underestimation of
 467 the genetic potential of the herd, or a poor quantitative estimate of the diets actually provided.

468

469 The increase in live weight of fattening and growing cattle such as replacement heifers is calculated by
 470 looking up, in the tables entered, the average daily gain (ADG) corresponding to the total energy and
 471 nitrogen input of the diet as a function of the average live weight of the animal over the fattening
 472 period. The user can inactivate all the calculations linking diet and production in cattle by not filling in
 473 certain variables at the level of parameters or enterprises. The productions linked to targets are then
 474 used for the rest of the calculations, as in the case of sheep and goats.



475

476 **Figure 4.** Example of graphic user interface (entry of daily diets and calculation of milk production for
 477 the average lactating cow of batch 1).

478

479 3.2.6 Economic results

480

481 The calculations of the above balances and of milk and meat production provide the quantities of crop
482 and livestock products that can be sold or need to be purchased, as well as forage and concentrates.
483 Added to this are the other inputs needed for the crop blocks. These different quantities are valued
484 economically by multiplying them by the unit purchase and sale prices entered by the user. On this
485 basis, and after entering the amounts of fixed costs for the entire farm, CLIFS calculates the following
486 economic variables, both in total and per hectare: expenses, gross proceeds, total gross margins (in
487 total and per crop and livestock component), and net profits.

488

489 **4 Example of CLIFS implementation on a Malagasy dairy farm**

490

491 The use of CLIFS is illustrated with the case of a real dairy farm located in the Lake Alaotra area in
492 Madagascar. Support was given to this farm in 2011 under a research project aiming to develop
493 mulch-based cropping systems (MCS) combining a crop of interest such as maize with permanent
494 plant cover. This innovation is being promoted by local development and research structures for its
495 expected effects on soil fertility and in reducing the risk of erosion on sloping fields (Corbeels et al.,
496 2013). This example was chosen because the entire process was carried out with the farmer, with
497 scenarios assessing the benefits of this innovation on the farm's dairy performance by making the
498 most of the cover plants' forage potential.

499

500 **4.1 Farm structure and production**

501

502 BH's farm, located on the shores of Lake Alaotra (17°41'S; 48°27'E), combines rice and dairy
503 production on about six hectares. Irrigated rice fields with high yield potential (5.5 t/ha of paddy) cover
504 80% of the area. The harvest covers the family's rice needs and provides two-thirds of the farm's
505 income. The remaining 20% of the farm consists of a lowland field (0.20 ha) producing green forage
506 throughout the year (*Chloris gayana*), and two rainfed plots (0.20 ha in total) cultivated as pure forage
507 (*Bracharia ruziziensis* and *Stylosanthes guianensis*). A rainfed area of 0.80 ha has been left to lie
508 fallow as it is too degraded to be productive (Table 6). The dairy herd consists of three improved breed

509 cows, the target production estimated by BH is 2800 l/cow/lactation. Calving takes place in November-
510 December with peak production in January-February (Figure 5). The cows are fed from the forage
511 plots, supplemented throughout the year with natural grasses that are grazed or collected. This
512 collection of biomass requires the employment of five permanent paid employees, who also are
513 responsible for milking the cows, processing the milk into yoghurt, and marketing. The cows are
514 permanently stalled all year round, which allows an abundant production of manure (a mixture of dung
515 and rice straw produced on the rice fields) that generally is spread on the rice fields.

516

517 **4.2 Building the scenarios**

518

519 After introducing BH to the support process and the general structure of CLIFS, an IS was built based
520 on his current situation. This step revealed that the cows' diet did not allow them to reach the expected
521 peak production in the rainy season because the energy content of the forage was insufficient (Figure
522 5). The focus then turned to redesigning the forage system in relation to the herd feeding system by
523 integrating three objectives expressed by BH: (i) produce more milk in the dry season (May-
524 September) because this would earn a better return (1,200 Ar¹/l instead of 1,000 Ar/l the rest of the
525 year); (ii) reduce the manpower devoted to the collection of natural grasses by feeding the cows with
526 on-farm cultivated forage; (iii) improve degraded soils to increase forage production and diversify food
527 production. To achieve these objectives, BH considered (i) shifting calving to May and June, (ii) setting
528 up a 0.25 ha plot of vetch as a catch crop after irrigated rice, (iii) setting up on 0.60 ha of rainfed land a
529 crop sequence consisting of a maize/*Stylosanthes* combination for one year, followed by three years
530 of *Stylosanthes* alone, and (iv) diversifying food production with a plot of groundnuts (0.20 ha) and
531 cassava (0.10 ha) in rainfed conditions on soils improved by MCS. The tops of groundnut plants also
532 can be fed to dairy cows.

533

534 The project scenario (PS) was developed on this basis by seeking technical references on MCS from
535 local experts in the field (Husson et al., 2013). The biomass of *Stylosanthes* exportable each year for
536 the herd was estimated in such a way that a minimum of vegetation would be maintained on the

¹ 2800 Ariary = 1 € in 2011 when the study was conducted.

537 ground in order to manage fertility and control erosion (Naudin et al., 2015). The maize-*Stylosanthes*
538 system, practically divided into four equal subparts in rotation, was modelled to produce the same
539 amount of maize and forage each year.

540

541 After simulating the PS, a first alternative scenario AS1 was built to reach BH's objectives of forage
542 autonomy and increased production during the dry season. This scenario expanded the type of MCS
543 introduced on the farm to diversify the forage available for the herd, and transformed the entire
544 production of *Stylosanthes* into hay starting from May. This technique, which BH was not practicing but
545 did know, represented an additional innovation in relation to the farm's current system and its planned
546 evolution. A second alternative scenario, AS2, added to AS1 the distribution of a concentrate to the
547 dairy cows, composed of raw materials directly available on the farm (grain maize, cassava) or after
548 processing (rice bran and groundnut cake).

549

550 **4.3 Evaluation**

551

552 This first stage of the process confirmed BH's observation that there was a milk production deficit in
553 the dry season in the initial situation (IS). It also enabled him to verify that the model was able to
554 represent his current situation before moving on to represent his project. The PS shows a reduction in
555 the volume of natural grasses collected in the dry season, while the cows' demand for forage
556 increases due to the higher proportion of milk produced in the dry season (Table 7; Figure 5).
557 However, the permanent labour force could only be decreased by one person due to the volumes that
558 continue to be needed. Furthermore, the target milk production could not be reached at peak
559 production in June-July. AS1 made it possible to forgo natural grasses in the dry season, saving an
560 additional permanent labour position. However, peak production still did not reach the target because
561 the diet remained too low in energy content. Only AS2 allowed both objectives to be achieved through
562 the use of concentrates produced on the farm.

563

564 From an economic point of view, each scenario increased the net margin of the farm compared to the
565 initial situation, with the milk enterprise accounting for a higher share of income while nonetheless
566 remaining less than that of rice. The existence of potential forage surpluses, especially in the

567 alternative scenarios, raised the possibility of increasing the herd size, but this was not modelled
 568 during the support process with BH due to time constraints. In view of these results, BH was interested
 569 in the alternative scenarios, especially as he already had a building in which he could store hay.
 570 Discussions also focused on the benefits and feasibility of the MCS modelled, which are presented
 571 here under steady state conditions but which require an installation time of one year to establish the
 572 cover crop. This installation time was a constraint for BH because it meant that he would have to adapt
 573 how he fed his herd due to the lack of forage.

574

575 **4.4 Outcomes**

576

577 Two years after this intervention, a visit was made to BH to learn about the developments on his farm
 578 related or not to what had been discussed. The farm's rice orientation had been accentuated with the
 579 opportunity to acquire a new plot of land with good control of irrigation water. BH also had reduced his
 580 herd to two dairy cows, which were better fed than before thanks to the increased production of
 581 groundnut tops and the addition of concentrates, and had shifted calving to the dry season as
 582 planned. However, he had not implemented the MCS tested in the scenarios because he considered
 583 that they were too complicated and not well adapted to his production conditions. Natural grasses
 584 therefore continued to provide much of the forage biomass, supplemented by green *Bracharia r.* grown
 585 in rotation with maize and cassava.

586

587

588 **Table 6.** Characteristics of the four scenarios simulated with CLIFS on the BH case.

	BS	PS	AS1 & AS2
Permanent staff (n)	5	4	3
Lowland cropping pattern (ha)			
Monocropped Rice paddy	4.75	4.50	4.25
Rice+Vetch ²		0.25	0.25
Rice+Oat ²			0.25
Chloris gayana ¹	0.20	0.20	0.20

Rainfall cropping pattern (ha)			
Groundnut ^{5,8}		0.20	0.10
Cassava ^{7,8}		0.10	
Maïze ^{7,8}			0.10
Maïze//Stylosanthes ^{3,5,6,8,9}		0.60	
Groundnut//Stylosanthes ^{3,5,6,7,8,9}			0.40
Cassava//Stylosanthes ^{3,6,7,8,9}			0.30
Stylosanthes guianensis ^{3,9}	0.10		
Bracharia ruziziensis ^{3,9}	0.10		
Fallow	0.70		
Total (ha)	5.85	5.85	5.85
% Total forage area	6.8	10.7	13.7

589 Crop objectives:

590 ¹ provide balanced green forage throughout the year

591 ² provide nitrogen-rich green forage in the dry season (August-November)

592 ³ provide green forage in the rainy season and at the beginning of the dry season

593 ⁴ provide green forage in the dry season (June-November)

594 ⁵ provide nitrogen-rich dry forage (peanut tops)

595 ⁶ provide usable hay in the dry season

596 ⁷ provide the ingredients of the self-produced concentrate

597 ⁸ diversify food crops

598 ⁹ improve degraded soils

599

600 **Table 7.** Technical-economic outcomes of the four scenarios simulated.


	IS	PS	AS1 (without concentrate)	AS2 (with concentrate)
Marketable milk production (l)	6380	7050	7220	7760
% production in dry season	36	52	53	56
Average sale price (Ar/l)	1071	1104	1105	1113
Total natural grasses (t)	23.2	11.1	9.2	9.2

Of which dry season	10.2	6.5	0	0
% dry season	44	59	0	0
Excess green forage (t/year)				
Chloris g.	0.7	1.2	0.7	0.7
Vetch		0.0	0.8	0.8
Oat			2.2	2.2
Stylosanthes g.	0.3	0.9	0.0	0.0
Bracharia r.	0.2			
Expenses (1000 Ar)				
Permanent work force	1375	1100	825	825
Seasonal work force (harvest)	1730	1870	1880	1880
Intermediate consumption	660	1190	1450	1450
Concentrates	-	-	-	97
Total	3765	4160	4345	4442
Total net margin (1000 Ar)				
	19740	20880	21090	21720
% livestock ¹	35	37	38	39

601

602 ¹ 100* Livestock Gross Margin / Total Net Margin

603



604

605 **Figure 5.** Monthly milk yields as a function of the production target and diet distributed.

606

607

608 **5 Discussion**

609

610 **5.1 Characterizing scenarios**

611

612 CLIFS limits the use of mechanistic equations to excreta production for ruminants, and to the
613 relationship between feed and production for cattle. The latter equations are assumed to be valid for
614 any type of context where animals have a live weight of between 400 and 800 kg (INRA, 2007), and to
615 be sufficient for farm-level estimates that do not require the same degree of precision as a feed
616 rationing tool. Aside from these situations, or when the data needed for the calculations are
617 unavailable, users can switch to a simplified use of the simulator where the link between diet and
618 production is based on their own expertise and not on calculations.

619

620 This option, and the fact that CLIFS essentially only mobilizes data describing the farm's resources
621 and how these are used, provides sufficient flexibility for the software to be used in many contexts.
622 Users are thus freed from the constraints encountered by whole-farm models integrating biophysical
623 models which require data that are not systematically available or validated, which limits their use and
624 utility in supporting farmers (Jones et al., 2017). This flexibility appears well suited to non-research
625 contexts with real farms, in individual or group situations (Ryschawy et al., 2014), and not just typical
626 farms reconstructed for the purposes of a research study (Tittonell et al., 2009; Smith et al., 2020).

627

628 However, this flexibility calls for vigilance with regard to three points. The first concerns the quality of
629 the data describing the farm in the initial scenario, which may not be available or entirely accurate on
630 all of the farms likely to benefit from the approach. The construction of this scenario can therefore be
631 time-consuming when extensive discussions with the farmer are needed to arrive at a plausible
632 representation of his/her situation. The interactions between the components of the production system
633 and the technical and economic performances, which often are known or even recorded (e.g., milk
634 production marketed), make it possible to arrive at a result that farmers themselves can evaluate and
635 validate within a participatory modelling approach (Voinov and Bousquet, 2010).

636

637 The second point concerns how to determine the performance of crop and livestock enterprises as a
638 function of respectively crop and feed management sequences when equations linking them are

639 absent. These values have a direct impact on the balance sheets calculated and the resulting
640 economic results. They need to be considered for each technical change in a scenario, either with the
641 farmer concerned or with experts in the field, in order not to lead to outliers. These considerations can
642 include the effects of adverse climatic and health events to assess the capacity of the simulated
643 configurations to absorb such shocks.

644

645 The third point concerns the characterisation of the technical innovations integrated into the
646 prospective scenarios when the farmer is unfamiliar with them. The data needed to describe the
647 innovations are then obtained from scientists and experts in the field. However, their validity in the
648 case under study cannot be guaranteed. In such situations, it is useful to develop a set of scenarios in
649 which yields are varied within a range considered plausible in the context studied to assess the
650 variability of the resulting technical and economic results.

651

652 Unlike other whole-farm models (Pissonnier et al., 2017; McDonald et al., 2019), the balance between
653 supply and demand for labour has not been integrated into CLIFS due to the complexity of calculating
654 this at the scale of a farm where there are a multitude of tasks, some seasonal, some routine, multiple
655 possibilities of division of labour within the farm (Davison, 2019), and multiple viewpoints on this
656 question depending on the farmer involved. When faced with an increasing demand for labour, some
657 farmers may be willing to work more, while others may prefer to hire labour from outside the farm
658 (Hostiou et al., 2015). These contexts lead to the establishment of individual balances per worker
659 within the farm, requiring a large amount of data. This is why, rather than embarking on such
660 calculations, it is preferred to assess with the farmer the additional costs generated by a given
661 configuration involving significant changes in labour demand (hiring temporary workers or employment
662 of permanent staff) (Ryschawy et al., 2014).

663

664

665 **5.2 Contributions and limitations for farmers and advisors**

666

667 The individual farmer support approach was implemented in a similar way on all of the sites where it
668 was tested. A number of points emerged that were remarkably consistent across the farms

669 participating in the study. First, farmers particularly appreciate the fact that the work is carried out on
670 their own farms and not on a virtual case (Ryschawy et al., 2014). This point, which is related to the
671 individual nature of the support, is due to the unique character of each farm which follows its own
672 development trajectory, even if certain objectives or issues which arise are shared by others (Rose et
673 al., 2016). Nonetheless, the outcomes of individual cases can be used to fuel collective reflection, as
674 long as the group is used to working collectively and uses real farms in the discussions (Ryschawy et
675 al., 2014).

676

677 Farmers also appreciate the ability to integrate the different components of the farm in a holistic
678 approach which they can understand, and the realistic and concrete character of the simulated
679 scenarios. This last point is the result of both the individual nature of the support and the attention paid
680 to the validation of the scenarios by the farmers themselves. From this perspective, the transparent
681 structure of CLIFS is an advantage. As with any support process based on the analysis of scenarios
682 (Martin et al., 2011), the farmers emphasise the capacity of the process to compare different options
683 and their impacts in terms of the management of production factors and performance, which can go as
684 far as reorienting their initial project.

685

686 They also highlight the knowledge gained in certain technical areas, such as dairy cow diets or forage
687 quality, through discussions about the scenarios (Semporé et al., 2016), and an increased awareness
688 of the value of collecting data on farm activities and planning these activities, as proposed by certain
689 advisory methods (Faure et al., 2018). The existence of quantified data about their farms known by the
690 farmers facilitates the implementation of the approach. When such is not the case, the advisor must
691 spend more time characterising the initial scenario, but in so doing gains a deeper understanding of
692 the farm.

693

694 However, as demonstrated by the case presented here, the projects discussed are not necessarily
695 implemented. This may be because this step involves questions of feasibility that the approach did not
696 address (McCown, 2002), or because farmers encounter opportunities or hazards that lead them to
697 reconsider their choices. Discussions and simulations may consider some of these points, such as the
698 occurrence of an extreme weather event, potential market opportunities and price dynamics, but the

699 reality encountered later may be quite different. The added value of the approach thus lies rather in
700 the capacity to transfer and increase the knowledge of both farmers and advisors about possible future
701 options (Martin et al., 2011).

702

703 For the advisors, who to date have mainly consisted of researchers and students, this participatory
704 approach provides a framework that goes beyond qualitative approaches, with the quantified outputs
705 of the simulations feeding into discussions with the farmer(s) on a concrete basis. By placing farmers
706 in a position to react and reflect on the questions they are asked and the results they are presented,
707 they are led to a better understanding of their own objectives, strategies, constraints and knowledge.
708 The advisor is required to seek biotechnical references from experts, researchers and technicians,
709 both to configure CLIFS under local conditions and to define the scenarios. This presents
710 opportunities to discuss possible technical innovations in a given case, and to point out knowledge
711 gaps in certain areas. A lack of local references indeed may limit the range of possible scenarios.

712

713 **5.3 From the tool to its use**

714

715 CLIFS was designed with the explicit aim of transferring the approach and the tool to non-researchers.
716 To achieve this objective, a compromise had to be reached between over simplifying, which would
717 make the tool ineffective with regard to the reflections to be conducted on strategic issues, and over
718 complicating, which would make the tool unusable in a professional advisory context (Rose et al.,
719 2016). This compromise led to the structural choices concerning the static nature of the simulations
720 and the calculation of balances.

721

722 The different case studies have shown that the tool appeals to farmers because it is relevant to their
723 questions, helps them understand scenario outputs and is useful for their reflections (Matthews et al.,
724 2008). However, the inclusion of new actors, such as agricultural advisors, and the structures in which
725 they are part have not yet been tested. Such an inclusion raises new questions regarding the skills of
726 those involved, both technically and in terms of their ability to analyse, make proposals and interact
727 with farmers in order to carry out the process. These skills are not common in advisory structures,
728 even in well-staffed contexts such as France, where strategic farm advice remains rare (Capitaine et

729 al., 2013). This situation stems in part from the need to focus on individual farms due to their specific
730 features, which is both time-consuming and costly. Mixed crop-livestock farming adds to the difficulty,
731 as agricultural advisors are often specialised by production sector, with a distinct separation between
732 the crop and livestock worlds (Garrett et al., 2020).

733

734 However, given the challenges of transforming farms facing environmental issues and economic
735 uncertainties in the agricultural and food sectors, such advisory services are needed. Consequently,
736 the goal now is to test the use of the approach by agricultural advisors in their own work contexts and
737 in various forms, ranging from individual support to advising producer groups. It also can be used in
738 frameworks involving research, development structures and farmers in the co-design of innovative
739 mixed crop-livestock systems (Berthet et al., 2016). In the academic sector, this type of tool makes it
740 possible to assess *in silico* highly innovative production system configurations (Pissonnier et al., 2019)
741 and to make students aware of the complexity of these systems and how to think about their evolution.

742

743 **6 Conclusion**

744

745 The farmer support approach built around CLIFS software is specifically dedicated to the strategic
746 questions that mixed crop-livestock farmers ask themselves about their future. The various
747 experiments conducted with family farms in the tropics and in France show that the tool meets the
748 expectations of farmers and enables them to better understand possible changes in their production
749 systems. CLIFS was designed to adapt to a wide range of situations and geographical contexts while
750 remaining simple to use, particularly with regard to the variables to be filled in on each farm. Its
751 transfer to advisory structures remains to be tested, but this simplicity should be an asset in this effort.
752 The use of the tool also could evolve towards being linked with environmental assessment methods in
753 order to think about crop-livestock combinations on farms that limit environmental impacts and
754 improve certain components such as biodiversity.

755

756 **Acknowledgements**

757

758 We would like to express our thanks to the many producers, researchers, experts and students who
759 participated to the design, development and experimentation of the CLIFS approach in the various
760 studied contexts. We thank Grace Delobel for translating the paper into English. This work was
761 supported by the following research projects: SIRMA (Water Saving in Irrigation Systems in North
762 Africa), financed by French institutional co-operation and North African countries in Morocco,
763 PEPITES project (Ecological, Technical and Social Innovation Processes in Conservation Agriculture)
764 ANR-08-STRA-10 in Brazil and Madagascar, and MOUVE ANR-2010-STRA-005 in France, both
765 financed by the French National Research Agency, the TFESSD (Trust Fund for Environmentally &
766 Socially Sustainable Development) of the World Bank in Peru.

767

768 **References**

769

770 Alary, V., Corbeels, M., Affholder, F., Alvarez, S., Soria, A., Valadares Xavier, J.H., da Silva, F.A.M.,
771 Scopel E., 2016. Economic assessment of conservation agriculture options in mixed crop-livestock
772 systems in Brazil using farm modelling. *Agricultural Systems* 144, 33–45.
773 <http://dx.doi.org/10.1016/j.agsy.2016.01.008>

774

775 Andrieu, N., Nogueira, D.M., 2010. Modeling biomass flows at the farm level: A discussion support tool
776 for farmers. *Agronomy for Sustainable Development* 30, 505-513.
777 <http://dx.doi.org/10.1051/agro/2009047>

778

779 Berthet, E.T.A., Barnaud, C., Girard, N., Labatut, J., Martin, G., 2016. How to foster agroecological
780 innovations? A comparison of participatory design methods. *Journal of Environmental Planning and*
781 *Management* 59, 280-301. <https://doi.org/10.1080/09640568.2015.1009627>

782

783 Bienz, N., Le Gal, P.-Y., 2012. Cultivating Prospective Thinking: A Gateway into the Future for
784 Peruvian Dairy Farmers in the Mantaro Valley. *Experimenting a Support Approach Based on the Use*
785 *of Modelling Tools*. Cirad, Montpellier, France, 53 p. <https://agritrop.cirad.fr/579168/>

786

787 Blanchard, M., Coulibaly, K., Bognini, S., Dugué, P., Vall, E. 2014. Diversité de la qualité des engrais
788 organiques produits par les paysans d'Afrique de l'Ouest : quelles conséquences sur les
789 recommandations de fumure ?. *Biotechnologie, Agronomie, Société et Environnement*, 18, 512-523.
790 <https://popups.uliege.be/1780-4507/index.php?id=16864&file=1&pid=11654>
791

792 Bonaudo, T., Burlamaqui Bendahan, A., Sabatier, R., Ryschawy, J., Bellon, S., Leger, F., Magda, D.,
793 Tichit, M., 2014. Agroecological principles for the redesign of integrated crop–livestock systems.
794 *European Journal of Agronomy* 57, 43-51. <https://doi.org/10.1016/j.eja.2013.09.010>
795

796 Capitaine, M., Garnier, A., Jeanneaux, P., Pervanchon, F., Chabin, Y., Bletterie, N., de Torcy B., de
797 Framond, H., 2013. Accompagner la démarche de management stratégique de l'exploitation agricole,
798 *Économie rurale* 337, 75-90. <https://doi.org/10.4000/economierurale.4118>
799

800 Colnago, P., Rossing, W.A.H., Dogliotti, S., 2021. Closing sustainability gaps on family farms:
801 Combining on-farm co-innovation and model-based explorations. *Agricultural Systems* 188, 103017.
802 <https://doi.org/10.1016/j.agsy.2020.103017>
803

804 Corbeels, M., de Graaff, J., Ndah, T.H., Penot, E., Baudron, F., Naudin K., Andrieu, N., Chirat, G.,
805 Schuler, J., Nyagumbo, I., Rusinamhodzi, L., Traore, K., Dulla Mzoba, H., Solomon Adolwa, I., 2014.
806 Understanding the impact and adoption of conservation agriculture in Africa: A multi-scale analysis.
807 *Agriculture, Ecosystems & Environment* Volume 187, 155-170.
808 <https://doi.org/10.1016/j.agee.2013.10.011>
809

810 Davison, J. (Ed.), 2019. *Agriculture, women, and land: The African experience*. Routledge, New York,
811 286 p.
812

813 Douhard, F., 2010. Conception et expérimentation d'outils de simulation pour l'accompagnement
814 d'agro-éleveurs. Application dans la région du Lac Alaotra (Madagascar). Unpublished Master thesis,
815 SupAgro, Cirad, VetAgro Sup, 34 p. <https://agritrop.cirad.fr/557653/>
816

817 FAO, 2016 FAO ration formulation tool for dairy cows. FAO, Rome, Italy.
818 <https://www.feedipedia.org/content/fao-ration-formulation-tool-dairy-cows>
819

820 Faure, G., Toillier, A., Havard, M., Rebuffel, P., Moumouni, I., Gasselin, P., Tallon, H. 2018. Advice to
821 farms to facilitate innovation: between supervision and support. In: Faure, G., Chiffolleau, Y., Goulet,
822 F., Temple, L., Touzard, J.-M. (Eds.), Innovation and development in agricultural and food systems.
823 Quae, Versailles, pp.144-156.
824

825 Foussat, M.-C., 2011. Evaluation prospective de systèmes de production incluant des techniques
826 d'agriculture de conservation dans une démarche d'accompagnement d'agro-éleveurs. Application
827 dans la région du Lac Alaotra (Madagascar). Unpublished Master thesis SupAgro, Cirad, AgroCampus
828 Ouest, 47 p. <https://agritrop.cirad.fr/561604/>
829

830 Garrett, R. D., Ryschawy, J., Bell, L. W., Cortner, O., Ferreira, J, Garik, A.V.N., Gil, J. D. B., Klerkx, L.,
831 Moraine, M., Peterson, C. A., dos Reis, J.C., Valentim, J.F., 2020. Drivers of decoupling and
832 recoupling of crop and livestock systems at farm and territorial scales. *Ecology and Society* 25, 24.
833 <https://doi.org/10.5751/ES-11412-250124>
834

835 Herrero, M., Thornton, P. K., Notenbaert, A. M., Wood, S., Msangi, S., Freeman, H. A., Bossio, D.,
836 Dixon, J., Peters, M., van de Steeg, J., Lynam, J., Rao, P.P., Macmillan, S., Gerard, B., McDermott, J.,
837 Sere, C., Rosegrant, M., 2010. Smart Investments in Sustainable Food Production: Revisiting Mixed
838 Crop-Livestock Systems. *Science* 327, 822-825. <https://doi.org/10.1126/science.1183725>
839

840 Hostiou, N., Dedieu, B., 2012. A method for assessing work productivity and flexibility in livestock
841 farms. *Animal* 6, 852-862. <https://doi.org/10.1017/S1751731111002084>
842

843 Hostiou, N., Cialdella, N., Vasquez, V., Müller, A.G., Le Gal, P.-Y., 2015. Work organization on
844 smallholder dairy farms: a process unique to each farm. *Tropical Animal Health Production* 47, 1271-
845 1278. <http://dx.doi.org/10.1007/s11250-015-0859-7>
846

847 Husson, O. (ed.), Séguy, L. (ed.), Charpentier, H. (ed.), Rakotondramanana (ed.), 2013. Manuel
848 pratique du Semis direct sur couverture végétale permanente (SCV). Application à Madagascar.
849 GSDM-CIRAD, Antananarivo, Madagascar, 716 p.
850 [https://www.researchgate.net/publication/283259038_Manuel_pratique_du_Semis_direct_sur_Couvert](https://www.researchgate.net/publication/283259038_Manuel_pratique_du_Semis_direct_sur_Couverture_Vegetale_permanente_SCV_Application_a_Madagascar/link/562f65b508ae4742240aca6d/download)
851 [ure_Vegetale_permanente_SCV_Application_a_Madagascar/link/562f65b508ae4742240aca6d/downl](https://www.researchgate.net/publication/283259038_Manuel_pratique_du_Semis_direct_sur_Couverture_Vegetale_permanente_SCV_Application_a_Madagascar/link/562f65b508ae4742240aca6d/download)
852 [oad](https://www.researchgate.net/publication/283259038_Manuel_pratique_du_Semis_direct_sur_Couverture_Vegetale_permanente_SCV_Application_a_Madagascar/link/562f65b508ae4742240aca6d/download)
853
854 INRA (Ed.), 2007. Alimentation des bovins, ovins et caprins. Besoins des animaux - Valeur des
855 aliments. Tables INRA 2007. Quae Editions, Versailles, France. 307 p.
856
857 Jones, J.W., Antle, J.M., Basso, B., Boote, K.J., Conant, R.T., Foster, I., Godfray, H.C.J., Herrero, M.,
858 Howitt, R.E., Janssen, S., Keating, B.A., Munoz-Carpena, R., Porter, C.H., Rosenzweig, C., Wheeler,
859 T.R., 2017. Toward a new generation of agricultural system data, models, and knowledge products:
860 State of agricultural systems science. *Agricultural Systems* 155, 269-288.
861 <http://dx.doi.org/10.1016/j.agsy.2016.09.021>
862
863 Kidron, G. J., Karnieli, A., & Benenson, I., 2010. Degradation of soil fertility following cycles of cotton–
864 cereal cultivation in Mali, West Africa: A first approximation to the problem. *Soil and Tillage Research*,
865 106, 254-262. <https://doi.org/10.1016/j.still.2009.11.004>
866
867 Kok, A., de Olde, E.M., de Boer, I.J.M., Ripoll-Bosch, R., 2020. European biodiversity assessments in
868 livestock science: A review of research characteristics and indicators. *Ecological Indicators* 112,
869 105902. <https://doi.org/10.1016/j.ecolind.2019.105902>
870
871 Le Gal, P.-Y., 2021. CLIFS (Crop Livestock Farm Simulator). User manual. CIRAD, Montpellier,
872 France, 88 p. <https://doi.org/10.18167/agritrop/00577>
873
874 Le Gal, P.-Y., Dugué, P., Faure, G., Novak, S., 2011a. How does research address the design of
875 innovative agricultural production systems at the farm level? A review. *Agricultural Systems* 104, 714-
876 728. <http://dx.doi.org/10.1016/j.agsy.2011.07.007>

877

878 Le Gal, P.-Y., Andrieu, N., Dugué, P., Kuper, M., Sraïri, M.T., 2011b. Des outils de simulation pour
879 accompagner des agroéleveurs dans leurs réflexions stratégiques. *Cahiers Agriculture*, 20, 413-420.
880 <http://dx.doi.org/10.1684/agr.2011.0509>

881

882 Le Gal P.-Y., Bernard J., Moulin C.-H., 2013. Supporting strategic thinking of smallholder dairy farmers
883 using a whole farm simulation tool. *Tropical Animal Health Production* 45, 1119–1129.
884 <http://dx.doi.org/10.1007/s11250-012-0335-6>

885

886 Lisson, S., MacLeod, M., McDonald, C., Corfield, J., Pengelly, B., Wirajaswadi, L., Rahman, R.,
887 Bahar, S., Padjung, R., Razak, N., Puspadi, K., Dahlanuddin, Sutaryono, Y., Saenong, S., Panjaitan,
888 T., Hadiawati, L., Ash, A., Brennan, L., 2010. A participatory, farming systems approach to improving
889 Bali cattle production in the smallholder crop–livestock systems of Eastern Indonesia. *Agricultural*
890 *Systems* 103, 486-497. <http://dx.doi.org/10.1016/j.agsy.2010.05.002>

891

892 Lurette, A., Aubron, C., Moulin, C.-H., 2013. A simple model to assess the sensitivity of grassland
893 dairy systems to scenarios of seasonal biomass production variability. *Computers and Electronics in*
894 *Agriculture* 93, 27–36. <http://dx.doi.org/10.1016/j.compag.2013.01.008>

895

896 Machado, C.F., Morris, S.T., Hodgson, J., Arroquia, M.A., Mangudo, P.A., 2010. . A web-based model
897 for simulating whole-farm beef cattle systems. *Computers and Electronics in Agriculture* 74, 129–136.
898 <http://dx.doi.org/10.1016/j.compag.2010.07.007>

899

900 Martin, G., Felten, B., Duru, M., 2011. Forage rummy: A game to support the participatory design of
901 adapted livestock systems. *Environmental Modelling & Software* 26, 1442-1453.
902 <https://doi.org/10.1016/j.envsoft.2011.08.013>

903

904 Matthews, K.B., Schwarz, G., Buchan, K., Rivington, M., Miller, D., 2008. Wither agricultural DSS?
905 *Computers and Electronics in Agriculture* 61, 149–159. <https://doi.org/10.1016/j.compag.2007.11.001>

906

907 McCown, R.L., 2002. Locating agricultural decision support systems in the troubled past and socio-
908 technical complexity of 'models for management'. *Agricultural Systems* 74, 11–25.
909

910 McDonald, C.K., MacLeod, N. D., Lisson, S., Corfield, J.P., 2019. The Integrated Analysis Tool (IAT) –
911 A model for the evaluation of crop-livestock and socio-economic interventions in smallholder farming
912 systems. *Agricultural Systems* 176, 102659. <https://doi.org/10.1016/j.agsy.2019.102659>
913

914 Moraine, M., Grimaldi, J., Murgue, C., Duru, M., Therond, O., 2016. Co-design and assessment of
915 cropping systems for developing crop-livestock integration at the territory level. *Agricultural Systems*
916 147, 87-97. <http://dx.doi.org/10.1016/j.agsy.2016.06.002>
917

918 Mosnier, C., Duclos, A., Agabriel, J., Gac, A., 2017. Orfee: A bio-economic model to simulate
919 integrated and intensive management of mixed crop-livestock farms and their greenhouse gas
920 emissions. *Agricultural Systems* 157, 202-215. <https://doi.org/10.1016/j.agsy.2017.07.005>
921

922 Mouret, P., 2012. Evaluation prospective des stratégies d'évolution d'exploitations laitières dans la
923 région Vakinankaratra, Madagascar. Unpublished Master thesis, AgroParisTech, Cirad, 49 p.
924 <https://agritrop.cirad.fr/570246/>
925

926 Naudin, K., Bruelle, G., Salgado, P., Penot, E., Scopel, E., Lubbers, M., de Ridder, N., Giller, K.E.,
927 2015. Trade-offs around the use of biomass for livestock feed and soil cover in dairy farms in the
928 Alaotra lake region of Madagascar. *Agricultural Systems* 134, 36-47.
929 <http://dx.doi.org/10.1016/j.agsy.2014.03.003>
930

931 NRC, 2001. *Nutrient Requirements of Dairy Cattle*, 7th revised edition. National Academy of Sciences,
932 Washington, DC, USA. <https://doi.org/10.17226/9825>
933

934 Parsons, D., Nicholson, C.F., Blake, R.W., Ketterings, Q.M., Ramírez-Aviles, L., Fox, D.G., Tedeschi,
935 L.O., Cherney, J.H., 2011. Development and evaluation of an integrated simulation model for

936 assessing smallholder crop–livestock production in Yucatán, Mexico. *Agricultural Systems* 104, 1-12.
937 <http://dx.doi.org/10.1016/j.agsy.2010.07.006>
938

939 Penot, E. (Ed), 2012. *Exploitations agricoles, stratégies paysannes et politiques publiques. Les*
940 *apports du modèle Olympe*. Editions Quae, Versailles. 336 p.
941

942 Pissonnier S., Lavigne C., Le Gal P.-Y., 2017. A simulation tool to support the design of crop
943 management strategies in fruit tree farms. Application to the reduction of pesticide use. *Computers*
944 *and Electronics in Agriculture* 142(A), 260–272. <http://dx.doi.org/10.1016/j.compag.2017.09.002>
945

946 Pissonnier, S., Dufils, A., Le Gal, P.-Y., 2019. A methodology for redesigning agroecological radical
947 production systems at the farm level. *Agricultural Systems* 173, 161–171.
948 <https://doi.org/10.1016/j.agsy.2019.02.018>

949 Rose, D.C., Sutherland, W.J., Parker, C., Lobley, M., Winter, M., Morris, C., Twining, S., Foulkes, C.,
950 Amano, T., Dicks, L.V., 2016. Decision support tools for agriculture: Towards effective design and
951 delivery. *Agricultural Systems* 149, 165-174. <https://doi.org/10.1016/j.agsy.2016.09.009>
952

953 Ryschawy, J., Choisis, N., Choisis, J.-P., Gibon, A., 2013. Paths to last in mixed crop-livestock
954 farming: lessons from an assessment of farm trajectories of change. *Animal* 7, 673-681.
955 <https://doi.org/10.1017/S1751731112002091>
956

957 Ryschawy, J., Joannon, A., Choisis, J.-P., Gibon, A., Le Gal, P.-Y. , 2014. Participative assessment of
958 innovative scenarios for enhancing sustainability of French mixed crop-livestock farms. *Agricultural*
959 *Systems* 129: 1-8. <http://dx.doi.org/10.1016/j.agsy.2014.05.004>
960

961 Sempore, A.W., Andrieu, N., Nacro, H.B., Sedogo, M.P., Le Gal, P.-Y., 2015. Relevancy and role of
962 whole-farm models in supporting smallholder farmers in planning their agricultural season.
963 *Environmental Modelling & Software* 68, 147-155. <http://dx.doi.org/10.1016/j.envsoft.2015.02.015>
964

965 Sempore, A.W., Andrieu, N., Le Gal, P.-Y., Nacro, H.B., Sedogo, M.P., 2016. Supporting better crop-
966 livestock integration on small-scale West African farms: a simulation-based approach. *Agroecology*
967 *and Sustainable Food Systems* 40, 3–23. <http://dx.doi.org/10.1080/21683565.2015.1089966>
968

969 Schut, A.G.T., Cooledge, E.C., Moraine, M., Van de Ven, G.W.J., Jones, D.L., Chadwick, D.R., 2021.
970 Reintegration of crop-livestock systems in Europe: an overview. *Frontiers of Agricultural Science and*
971 *Engineering*, Online first. <https://doi.org/10.15302/J-FASE-2020373>
972

973 Smith, J., Nayak, D, Datta, A., Narkhede, W.N., Albanito, F., Balana, B., Bandyopadhyay, S.K., Black,
974 H., Boke, S., Brand, A., Byg, A., Dinato, M., Habte, M., Hallett, P. D., Lemma, T., Mekuria, W., Moges,
975 A., Muluneh, A., Novo, P., Rivington, M., Tefera, T., Vanni, E.M., Yakob, G., Phimister, E., 2020. A
976 systems model describing the impact of organic resource use on farming households in low to middle
977 income countries. *Agricultural Systems* 184, 102895. <https://doi.org/10.1016/j.agsy.2020.102895>
978

979 Snow, V.O., Rotz, C.A., Moore, A.D., Martin-Clouaire. R., Johnson, I.R., Hutchings, N.J., Eckard, R.J.
980 2014. The challenges – and some solutions – to process-based modelling of grazed agricultural
981 systems. *Environmental Modelling & Software* 62, 420-436.
982 <http://dx.doi.org/10.1016/j.envsoft.2014.03.009>
983

984 Sraïri, M.T., Benyoucef, M.T., Kraiem, K., 2013. The dairy chains in North Africa (Algeria, Morocco
985 and Tunisia): from self sufficiency options to food dependency? *SpringerPlus*, 2:162.
986 <http://www.springerplus.com/content/2/1/162>
987

988 Tiftonell, P., van Wijk, M.T, Herrero, M., Rufino, M.C., de Ridder, N., Giller, K.E., 2009. Beyond
989 resource constraints – Exploring the biophysical feasibility of options for the intensification of
990 smallholder crop-livestock systems in Vihiga district, Kenya. *Agricultural Systems* 101, 1-19.
991 <https://doi.org/10.1016/j.agsy.2009.02.003>
992

993 Vayssières, J., Bocquier, F., Lecomte, P., 2009. GAMEDE: a global activity model for evaluating the
994 sustainability of dairy enterprises. Part II – Interactive simulation of various management strategies

995 with diverse stakeholders. *Agricultural Systems* 101, 139–151.

996 <http://dx.doi.org/10.1016/j.agsy.2009.05.006>

997

998 Voinov, A., Bousquet, F., 2010. Modelling with stakeholders. *Environmental Modeling & Software*

999 25, 1268–1281. <http://dx.doi.org/10.1016/j.envsoft.2010.03.007>

1000