



HAL
open science

Exploring complementarities between modelling approaches that enable upscaling from plant community functioning to ecosystem services as a way to support agroecological transition

Noémie Gaudio, Gaëtan Louarn, Romain Barillot, Clémentine Meunier, Rémi Vezy, Marie Launay

► To cite this version:

Noémie Gaudio, Gaëtan Louarn, Romain Barillot, Clémentine Meunier, Rémi Vezy, et al.. Exploring complementarities between modelling approaches that enable upscaling from plant community functioning to ecosystem services as a way to support agroecological transition. *in silico Plants*, 2022, 4 (1), pp.1-13. 10.1093/insilicoplants/diab037/6449487 . hal-03470234v1

HAL Id: hal-03470234

<https://hal.inrae.fr/hal-03470234v1>

Submitted on 8 Dec 2021 (v1), last revised 1 Mar 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Exploring complementarities between modelling approaches that enable upscaling from plant community functioning to ecosystem services as a way to support agroecological transition

Noémie Gaudio¹, Gaëtan Louarn², Romain Barillot², Clémentine Meunier¹, Rémi Vezy³, Marie Launay^{4*}

¹ AGIR, Univ Toulouse, INRAE, Castanet-Tolosan, France

² UR P3F, INRAE, F-86600, Lusignan, France

³ SYSTEM, Univ Montpellier, CIRAD, INRA, IRD, Montpellier SupAgro, Montpellier, France

⁴ US AGROCLIM, INRAE, Avignon, France

*corresponding author: marie.launay@inrae.fr

Accepted Manuscript

© The Author(s) 2021. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Promoting plant diversity through crop mixtures is a mainstay of the agroecological transition. Modelling this transition requires considering both plant-plant interactions and plants' interactions with abiotic and biotic environments. Modelling crop mixtures enables designing ways to use plant diversity to provide ecosystem services, as long as they include crop management as input. A single modelling approach is not sufficient, however, and complementarities between models may be critical to consider the multiple processes and system components involved at different and relevant spatial and temporal scales. In this article, we present different modelling solutions implemented in a variety of examples to upscale models from local interactions to ecosystem services. We highlight that modelling solutions (i.e. coupling, metamodelling, inverse or hybrid modelling) are built according to modelling objectives (e.g. understand the relative contributions of primary ecological processes to crop mixtures, quantify impacts of the environment and agricultural practices, assess the resulting ecosystem services) rather than to the scales of integration. Many outcomes of multispecies agroecosystems remain to be explored, both experimentally and through the heuristic use of modelling. Combining models to address plant diversity and predict ecosystem services at different scales remains rare but is critical to support the spatial and temporal prediction of the many systems that could be designed.

Keywords: crop mixtures, process-based models, crop models, individual-based models, modelling synergies, pest regulation

1. Introduction

New models are frequently developed for specialists in a field to answer specific scientific questions, without much interaction with other disciplines in the initial stages. During the past decade, however, modellers have integrated knowledge from multiple disciplines (e.g. micro-meteorology, environmental physics, ecophysiology, ecology, soil science) to better represent interactions between processes within plants, between plants, and between plants and their environment (e.g. Gauthier *et al.* 2020). The result is a distinct diversity of modelling approaches that can be used to benefit the complementary properties and strengths of models (see e.g. Colbach *et al.* 2021; Pointurier *et al.* 2021). This knowledge and model sharing requires certain upstream steps that are necessary to render models more accessible, such as free licenses, open-source code, accessible software products, improved usability, extensive documentation and training sessions. These steps are especially important when setting up close collaboration between teams of modellers that include in-depth work on exchanges between models, such as the international modelling communities working on crop models, such as AgMip (“Agricultural Model Intercomparison and Improvement Project”; Rosenzweig *et al.* 2013) and MACSUR (“Modelling European Agriculture with Climate Change”; <https://macsur.eu/>). A good example of this desire to share and standardize practices is the study of Midingoyi *et al.* (2021) on the development of a meta-language to facilitate the exchange and reuse of crop-model components between modelling platforms.

The need to combine several modelling approaches, each with trade-offs in accuracy and generality, is crucial in all scientific disciplines and assumes that each model may improve understanding and predictions of ecosystem functioning. For instance, Evans *et al.* (2016) highlighted that the global models used to predict the geographic distribution of plant species throughout the world have low predictive power if they are not improved with process-based range models that predict impacts of environmental changes. Therefore, the need exists for accurate predictions of processes and more global and qualitative modelling approaches to understand an ecosystem, while also considering the feedback between different approaches, especially as the factors involved in ecosystem functioning are not necessarily the same for the spatial scales considered (Pearson and Dawson 2003; Xu *et al.*, 2021).

Building connections between modelling approaches is particularly crucial in the context of the current agroecological transition, which involves in-depth changes to agricultural practices, with more complex and diversified agroecosystems and a multifunctional view of agriculture (Caron *et al.* 2014; Duru *et al.* 2015; Gaba *et al.* 2015). Increasing plant diversity is a mainstay of the agroecological transition and the cornerstone for “biodiversity-based agriculture” (Duru *et al.* 2015), which depends on agrobiodiversity at field, farm and landscape scales (Kremen and Miles 2012;

Prieto *et al.* 2015; Tschardtke *et al.* 2021). In these types of agriculture, ecological processes are fundamental to agricultural production, which requires particular focus on production-ecology trade-offs (Sabatier and Mouysset 2018). More than ever, modelling synergies must be identified to enable upscaling from plant functioning (i.e. ecophysiological processes and plant-plant interactions) to ecosystem services to support the agroecological transition (Tixier *et al.* 2013).

To illustrate how these modelling synergies and complementarities are essential to better characterize biodiversity-based agriculture, we focus on modelling species and cultivar crop mixtures along the continuum of plant (and plant-plant interactions), field and farm scales. Each scale requires representing specific abiotic and biotic factors (Pearson and Dawson 2003; Peng *et al.* 2020), as well as ecosystem functions that support ecosystem services, including production and regulating services (Haines-Young and Potschin 2013). The nature, importance and level of expression of ecosystem functions also depend on the scale considered. While experiments have identified agronomic advantages of these diversified systems (e.g. Kiar *et al.* 2012; Beillouin *et al.* 2021), better understanding is needed about their agricultural management and especially about how to integrate them into cropping systems to attain the ecosystem services targeted. The diversity of ecosystem services targeted and the extent of the temporal and spatial scales at which these services are developed make modelling choices complex. Issues to consider include which processes the model should simulate and at what resolution, as well as which temporal scale (e.g. instantaneous, daily, crop-cycle, rotation or long-term) and spatial resolution (e.g. plant, field or landscape, along with its multiple cultivated and uncultivated components) to use to represent the multiple interactions of interest. These considerations suggest that a single modelling approach is not sufficient to meet these legitimate expectations. Moreover, how these issues are addressed depends on which stakeholders use the models.

In this opinion article, we advocate that complementarities and coupling of different modelling approaches are critical to consider the complex and diversified agroecosystems involved in the agroecological transition, as well as to upscale from the plant and/or field scales to the ecosystem services targeted in diversified agroecosystems. Using several examples, we demonstrate that the complementarity between individual-based models (including functional-structural plant models (FSPMs)), crop models and physical or more qualitative or statistical models, improves understanding and facilitates simulating the functioning of crop mixtures and the ecosystem services for which they are designed. These modelling complementarities are discussed through the lens of crop mixtures or are integrated at larger scales to address three important modelling challenges: to i) quantify and understand plant-plant interactions and their underlying processes, ii) represent

impacts of the environment and agricultural practices on the functioning of crop mixtures and iii) assess the ecosystem services provided by these heterogeneous covers (Figure 1).

2. Modelling plant-plant interactions in crop mixtures to assess ecosystem services at fine scales (plant and field)

Plant-plant interactions in crop mixtures are the foundation for the ecosystem services provided by diversified agroecosystems. Although these interactions can provide large-scale ecosystem services, for most processes they usually occur at a fine scale due to interactions between neighbouring plants or between plants and microorganisms. To illustrate this, we focus on complementarities between existing modelling approaches to simulate production and regulating services quantified at the plant and/or field scales.

2.1. Modelling plant-plant interactions to quantify underlying processes for production services

One widely known advantage of crop mixtures is their potential to achieve higher yields due to more efficient and complementary use of abiotic resources compared to that of sole crops, especially in a low-input context (Banik *et al.* 2006; Dhima *et al.* 2007; Gaudio *et al.* 2021; Hauggaard-Nielsen *et al.* 2008). Predicting effects of plant-plant interactions on the capture and use of abiotic resources is therefore crucial to assess the production services provided by crop mixtures.

From an ecophysiological viewpoint, some processes are particularly determinant in crop mixtures because they strongly influence functioning and performance, especially related to competition (for light, water and nutrients), complementarity (spatio-temporal and niche processes) and facilitation (Malézieux *et al.* 2009; Brooker *et al.* 2015). The main modelling approaches developed to study plant-plant interactions within crop mixtures are process-based models at plant and crop scales (Gaudio *et al.* 2019). We do not provide a detailed description here of how process-based modelling approaches simulate the processes that underlie these plant-plant interactions, as several recent reviews and integrated studies have done so and have described their strengths and weaknesses (e.g. Gaudio *et al.* (2019) for crop or individual process-based models, Evers *et al.* (2019) and Louarn and Song (2020) for FSPMs). When downscaling to quantify and understand the relative contributions of primary ecological processes in crop mixtures, individual-based models, especially FSPMs, are usually required, as the phenotype of individual plants emerges from interactions between the local environment they perceive and their functioning (Evers *et al.* 2019). Thus, some FSPMs can quantify ecological processes (Zhu *et al.* 2015; Faverjon *et al.* 2019) or assess the role of given traits involved in the performance or resource-use efficiency of crops (usually light or nitrogen; Barillot *et al.* 2014; Louarn *et al.* 2020). For instance, Zhu *et al.* (2015) used an FSPM to simulate

wheat-maize relay-intercropping and highlighted that plasticity is the main process involved in the higher light-use efficiency in intercrops than in sole crops.

Important processes highlighted by FSPMs can be summarized using functional relationships to include them in crop models (Escobar-Gutiérrez *et al.* 2009), which would enable understanding of plant-plant interactions and thus achieve upscaling. However, few studies have highlighted complementarities between these two modelling approaches. For instance, FSPMs can be used to evaluate simplified assumptions applied in upscaled crop models. Crop models can simulate light interception and partitioning differently, but the common way to represent light interception is to use the “turbid-medium approach” and Beer-Lambert law. However, the question remains as to whether this simplified approach is sufficiently accurate to simulate light partitioning among plants in crop mixtures. Barillot *et al.* (2011) addressed this question for three grass-legume mixtures (wheat-pea, fescue-alfalfa and fescue-clover) and compared light partitioning between the component species. They simulated detailed 3D representations of plants, coupled with a solar radiation model that followed the turbid-medium approach, with the plant canopy represented by one, two or ten layers. The results indicated that more detailed representation of the canopy (several layers or in 3D) improved the prediction of light partitioning in mixtures only slightly, thus validating the turbid-medium approach for estimating light competition at the canopy scale in crop models. Similarly, Pao *et al.* (2021) transformed a 1D light model using Beer-Lambert equation by estimating empirically the light extinction coefficient from the canopy geometry formalised by plant and row distances. The performance of this approach in combination with hourly-step time resolution of simulation was equivalent to a 3D light model using ray tracing, in a dynamic plant model predicting leaf-level photosynthetic acclimation and plant-level dry matter accumulation. This smart solution provided efficient estimation for long-term processes integrated over weeks. Another example exhibiting how emerging results can be simplified using FSPM demonstrated that a model input that is time-consuming to assess, the red: far-red ratio, was successfully replaced by a proxy computed from the leaf area of the upper ten leaves and the plant density (Kahlen and Stützel 2011).

The complex integration of local plant responses to light competition and plant structure considered in FSPMs is largely incompatible with the simple representation of plants in crop models. However, indirect connections can be identified using inverse modelling and the adaptive calibration of input parameters in crop models. For instance, these models frequently use the response function of crops to plant density to represent the competitive effect of neighbours in mixtures by calibrating a dominance ratio or an equivalent-density parameter (Brisson *et al.* 2008; Confalonieri 2014). These parameters depend strongly on environmental conditions and the identity of neighbouring species in

the mixture (Van Oijen *et al.* 2020). Their variations can be derived from simulated data produced by more detailed FSPMs, such as the relative-density responses of species (e.g. as illustrated by Louarn and Faverjon (2018) for contrasting legume species).

Interactions for belowground resources are not reflected to this extent in modelling platforms (Evers *et al.* 2019). Because the spatial distribution of resources and physiological characteristics of species drive the growth and plasticity of roots, they are important factors that influence the response to water or nutrient changes in intercropping (Yin *et al.* 2020). Results of intercropping studies that focus on how to regulate root systems through spatio-temporal variation in the water or nutrient supply are rarely reported. However, studies have focused on species-specific responses of root morphological plasticity as influenced by nutrient availability, showing greater plasticity in graminaceous species (e.g. maize) than in leguminous species (e.g. faba bean, chickpea) (Li *et al.* 2014). Other studies compared mixed cropping to segregated strip-intercropping, showing that using strip-intercropping to concentrate low C:N species increased N mineralization potential in the planting zone for the subsequent crop (Lowry and Brainard 2016). Finally, benefits of legume-based intercrops have been shown through direct plant-to-plant N transfer, depending on the physical co-location of the root systems and thus on the spatial arrangement of the two species (Johansen and Jensen 1996; He *et al.* 2009; Chapagain and Riseman 2014). These results indicate the need to model phenotypic plasticity of roots as a function of the distribution of resources in a predefined spatial arrangement of intercrops. However, simple and generic approaches that focus on belowground resources already exist. For instance, Bertrand *et al.* (2018) developed BISWAT, a crop model that simulates dynamics of water stress in plants in sole crops and crop mixtures. The model integrates and combines simple approaches to simulate the main processes in the system with a 2D representation of the plants and soil, radiation-use efficiency, total transpirable soil water content and a simple representation of root dynamics. The resulting model requires few data for parametrization and yet remains robust for simulating water-stress dynamics in a wide range of systems.

Compared to these ecophysiological approaches, ecological approaches are particularly relevant for studying and understanding the functioning of systems in which multiple heterogeneous populations, such as crop mixtures, interact. However, representing mechanistically the processes that interact within these systems and quantifying the resulting ecosystem services requires knowledge and conceptual frameworks that are well theorized in ecophysiology and agronomy. Thus, dialogue between these different disciplines – environmental physics, ecophysiology, agronomy and ecology – is crucial for modelling these agroecological systems (Evans *et al.* 2016; Brooker *et al.* 2021). In particular, the concept of “functional trait” commonly used in ecology and

recently used to characterize agrobiodiversity and ecosystem services (Wood *et al.* 2015) can be related to the parameters and variables in process-based models using the distinction between pattern and process traits developed by Voltaire *et al.* (2020). Unlike the strict definition of functional traits, which are measured independent of the environment (Violle *et al.* 2007), the authors argue that process traits (*a trait measured under environmental conditions fluctuating in time, which characterizes processes, that is, flows of material and energy in a given environment during a defined period of time*, e.g. growth rate or phenological stage duration) are also functional and crucial for parameterizing models. This distinction may provide a bridge between ecology and crop science, partly because it allows to discuss upon a common semantics linking pattern and process traits with input parameters and state variables which are used in process-based models.

2.2. Modelling plant-plant interactions to quantify the underlying processes for regulating services

Crop mixtures should promote regulating services as well as production services (Haines-Young and Potschin 2013). For instance, vegetation diversity at all spatial scales (intra- vs. inter-field) improves pest regulation in several ways. It increases the matrix of unfavourable habitats and thus limits pest dispersal (Fabre *et al.* 2012; Papaïx *et al.* 2014). The spatial heterogeneity of host plants can also restrict pest population dynamics, which slows the specialization process (Plantagenest *et al.* 2007). In the next modelling example, a splash dispersal model was coupled with a snapshot of virtual 3D canopies provided by an FSPM to understand fine processes involved in controlling rain-borne diseases in wheat-cultivar mixtures (Vidal *et al.* 2018, Figure 2).

By mixing susceptible and resistant cultivars, the habitat favourable to a pathogen is spatially fragmented, thus generating i) a "barrier" effect, related to the presence of resistant cultivars, and ii) a "dilution" effect, as the probability of an individual finding a favourable habitat is reduced proportionally to the reduction in density of the susceptible cultivar (Finckh *et al.* 2000). This latter effect can be reinforced by a difference in height between cultivars when the pathogen spreads from the bottom to the top of the plant. Vidal *et al.* (2018) showed that a wheat-cultivar mixture composed of a short resistant cultivar and a taller susceptible cultivar would result in a lower layer that mixes susceptible and resistant leaves, which may provide a strong barrier effect (Figure 2). In contrast, the upper part of the canopy would be less dense (containing only the taller susceptible cultivar), and the upper leaves would be protected by their increased distance from the inoculum source at the bottom (height effect) and the presence of resistant leaves in the lower part of the canopy (barrier effect). In this example, the coupled model clarified understanding of the mechanisms involved and identified height as a relevant architectural trait to reduce spore dispersal

when mixing cultivars. However, manipulating such models is extremely time-consuming. The idea again is to identify emerging results – response functions to plant architecture and cultivar resistance – and then introduce them in a simplified manner into models that simulate other important factors in epidemics, such as the microclimate in the canopy. The next step is to quantify the influence of these associations on regulating pest populations and limiting crop yield losses. Consequently, coupling process-based modelling approaches and food-web modelling could provide a promising path for upscaling (Tixier *et al.* 2013; Malard *et al.* 2020).

3. Representing impacts of the environment and agricultural practices when assessing ecosystem services provided by crop mixtures at the cropping-system scale

3.1. Representing impacts of external drivers on plant-plant interactions

As described in previous sections, understanding mechanisms of plant-plant interactions requires describing the plant environment in detail. Promoting the use of crop mixtures at the cropping-system scale (e.g. rotation, farm) requires considering effects of agricultural practices and environmental factors that influence plant-plant interactions when building and evaluating the ability of crop mixtures to provide one or more ecosystem services. Current FSPMs often do not consider this particular point as extensively as crop models due to their complex structure and the associated modelling costs (Louarn and Song 2020). This could be mitigated by borrowing the strengths of different approaches and developing hybrid modelling. However, the level of precision and degree of simplification required to consider plant-plant interactions are not necessarily the same and depend on the outputs targeted by the simulation study. Colas *et al.* (2021) illustrated this point by simplifying a complex individual-based model at the cropping-system scale to design effective strategies for weed control. They simplified light partitioning – which is usually represented with a 3D voxelized canopy in their mechanistic model (FlorSys, Colbach *et al.* 2014) – using a random-forest-based metamodelling approach to accelerate the simulations and enable interactive testing of many complex cropping systems with end users.

In crop models, agricultural practices influence the crop environment, such as soil fertility and water availability, which modifies the soil-climate context in which crop mixtures may adapt as a function of their complementarity and/or plasticity properties (Stöckle and Kemanian 2020). Crop models are thus able to supply inputs for FSPMs, i.e. quantified and dynamic descriptions of abiotic constraints

under which a crop mixture grows. For instance, the STICS crop model was used to simulate impacts of delayed sowing dates on plant-plant competition for light (Launay *et al.* 2009) and impacts of different levels of nitrogen fertilization on relative dominance (Corre-Hellou *et al.* 2009) in barley-pea intercrops.

3.2. Assessment of a given ecosystem service: pest regulation

Pests in agroecological cropping systems can be regulated by significantly increasing plant diversity in the field or landscape using arable crops and semi-natural or natural elements (Sirami *et al.* 2019). However, the effectiveness of these systems depends greatly on their spatial organization at the field scale (Landis *et al.* 2000). Collard *et al.* (2018) transposed the spatial concepts of landscape ecology to the field scale, assuming that proximity, edge length or aggregation could improve understanding of how the spatial organization of non-crop habitats might alter the predator effect and thus increase crop health. They used an individual-based and spatially explicit model to simulate individual behaviours of predators, such as the duration and frequency of visits to orchard crops. They tested several spatial organizations that varied in the clumping of non-crop habitats, the distance between crop and non-crop habitats, and the number of alternative favourable neighbouring non-crop habitats around the crop habitat. To assess pest regulation, however, this modelling approach now needs to include the dynamics of pests and their interactions with predators. The current version partly meets this aim using proxies such as visit duration and frequency, as well as the duration of the predator's absence from the crop.

A good example of a modelling solution that uses complementarities between models and that can represent heterogeneous canopies was built by coupling the individual-based model FlorSys (Colbach *et al.* 2014) with RSCone, a metamodel produced using the architectural root model ArchiSimple (Pagès *et al.* 2020), and the soil submodel of the STICS crop model (Brisson *et al.* 2008). While FlorSys simulates aboveground crop-weed canopies, the ArchiSimple metamodel represents the trophic connection between above- and belowground growth, and the STICS soil submodels represent soil structure and climate and their effects on root growth (Figure 3). Illustrating the issues faced by coupling models or modules with different time or space scales, conversions had to be done to make optimal days of RSCone compatible with thermal time in FlorSys. This smart solution thus generates outputs and proxies that can be used to assess contrasting ecosystem services such as crop grain yield; weed-caused yield loss; weed seed production (as a proxy for future yield loss) and weed-based trophic resources for domestic bees (as one example of weed benefits), resource uptake or striga risk (Pointurier *et al.* 2021). It was necessary to develop working and modelling assumptions as the aim of this multi-faceted model was to cover a wide range of flora (including

many contrasting annual species) and address multiple ecosystem services (Colbach *et al.* 2021). Simpler, empirical relationships were preferred for processes for which mechanistic representation would have required downscaling to the cellular or molecular scale. Because representation of individual plants had to be compatible with multi-annual and multi-field simulations of thousands of plants per field, detailed representations, such as used in FSPMs, were rejected in favour of individual-based modelling.

This research model was then used to identify agroecological mechanisms and provide decision support for farmers. This mechanistic and individual-based approach induces considerable algorithmic complexity and slow simulations; thus, using it in decision-support systems is time consuming, as it requires assigning many input variables and calibrating many parameters, particularly when simulating many diverse crops simultaneously. This is solved by aspects of metamodelling (Colas *et al.* 2020) that can identify potential changes to cropping systems that might improve their performance. However, a biophysical parent model is still required to provide biophysical explanations that farmers will accept (Colbach *et al.* 2021).

The authors reconstructed the functioning of a diversified agroecosystem by coupling models that could represent systems (the plant and its aerial and root structure, seeds, soil layers and their structure and microclimate) and mechanisms at similar scales. In particular, they integrated two aspects that are essential to understand and manage these types of agroecological systems: consideration of long-term processes (e.g. evolution of a seed bank) and impacts of management decisions on these processes and the targeted ecosystem services, with consequences that could occur over several years.

3.3. Assessment of a bundle of ecosystem services at the farm scale

Assessing the ecosystem services provided by diverse crop mixtures is challenging due to the many ecosystem services targeted by farmers and the diversity of crops to be investigated (Verret *et al.* 2020). Coupling models may be a promising solution to understand this complexity and diversity because it benefits from the strengths of diverse modelling approaches. However, predicting how management activities and changing future conditions will alter ecosystem services is rendered more complex by interactions (e.g. trade-offs, synergies) among multiple ecosystem services (Agudelo *et al.* 2020). More widespread use of process-based models to estimate ecosystem services could identify physiological processes, or even the traits, that influence interactions between ecosystem services. However, simulating the ecosystem services provided by crop mixtures requires representing their inclusion in crop rotations and long-term effects of the environment. This could

be achieved by combining the knowledge provided by process-based models and using more qualitative models based on farmers' expertise.

In agreement with this idea, Meunier *et al.* (2022) designed a serious game to help users (farmers or students) explore and assess a bundle of ecosystem services (i.e. cereal and legume grain yield, cereal protein content, potential nitrogen supply to the next crop, maintenance of soil structure and pest regulation) provided by a wide range of binary cereal–grain legume intercrops (Figure 4). The serious game encapsulates a modelling chain that they constructed from three modelling approaches:

- (i) STICS (Brisson *et al.* 2008) was used to simulate the potential and water-limited biomasses of the cereal and legume sole crops independently under a variety of soil-climate conditions and management practices.
- (ii) A statistical model built using R software (R Core Team 2018), using a field-trial database of cereal-legume intercrops and their corresponding sole crops, was used to correct these potential and water-limited biomasses into attainable (i.e. water and nutrient-limited) biomasses (Van Ittersum *et al.* 2013).
- (iii) A knowledge-based multi-attribute model built using DEXi software (Bohanec 2008) was used to turn attainable biomass into actual biomass considering pest damage and assessing pest-regulation services. Other multi-attribute models also enabled assessment of five more ecosystem services that result from the actual biomass of the cereal-legume intercrop at harvest and/or cropping-system features.

The serious game was designed to explore the ecosystem services provided by both common and less-common intercropping scenarios, and to promote debate and knowledge sharing among users.

4. Upscaling models from local interactions to ecosystem services: realities, opportunities and obstacles

4.1. Modelling solutions to benefit from model complementarities

From the examples listed above, different strategies can be identified to combine models at different scales and predict consequences of plant-plant interactions, from local responses up to ecosystem services at the cropping system and farm scales (Figure 5). Besides direct coupling of models, which is rarely feasible across all scales, we identified three particularly promising approaches to address this issue:

- 10 ● **Inverse modelling**, which connects models by identifying input parameters from simulated
11 data. This approach is common to many scientific disciplines (Evans *et al.* 2016) and uses
12 simulated datasets to determine parameter values from other models to supplement the
13 observed data available. Using simulated datasets to improve exchanges between models
14 and modellers is particularly valuable to facilitate parametrization of existing models, as
15 illustrated by the adaptive parametrization of density responses and dominance ratios in
16 crop models (individual-based model to a crop model, Van Oijen *et al.* 2020) or the definition
17 of input scenarios in serious games (crop model to a farm-management model; Meunier *et*
18 *al.* 2022).
- 19 ● **Metamodelling**, which connects models by developing a simpler model of outputs from a
20 more complex model (Jin *et al.* 2001). Defining such new models is a particularly interesting
21 way to simplify complex simulation models that have high computing costs into something
22 tractable and reusable in a particular domain using a more integrated approach. For
23 instance, this is illustrated by the integration of a root-morphogenesis metamodel in FlorSys
24 to represent root competition (from root FSPM to a cropping-system model, Pointurier *et al.*
25 2021). The approach has also been effective at scaling up local plant interactions over large
26 areas and representing vegetation dynamics by considering soil and landscape variability
27 (e.g. Moorcroft *et al.* 2001). Metamodelling can reduce the computing costs of complex
28 models by several orders of magnitude.
- 29 ● **Hybrid modelling**, which connects models by combining the strengths of existing models in a
30 new model (Louarn and Song 2020). The goal is to perform hierarchical modelling at multiple
31 scales by including only the level of detail required to represent the critical processes
32 involved in targeted outputs of the system (e.g. scaling-up a mechanistic model of dynamic
33 protein turnover from leaf to canopy level, to provide a physiological explanation of the
34 photosynthetic acclimation under various light availability and nitrogen supply environments;
35 Pao *et al.* 2019). A direct example is the re-use of complementary modules from existing
36 models in original modelling solutions (e.g. individual-based plant models with soil and
37 management modules from crop models; Faverjon *et al.* 2019). Merging knowledge can also
38 result in formalizing emergent properties of a complex model in simpler robust equations or
39 in validating a simplified formalism used in cropping-system models (Barillot *et al.* 2011).
40 Although not yet developed for agroecosystems, hybrid modelling could also consider
41 Bayesian approaches that have been effective at aggregating different types of models and
42 data, including those that concern consequences of plant-plant interactions in natural
43 systems (Pagel and Schurr 2012).

44

45 These three broad categories are not mutually exclusive and can be combined to build original
46 models across scales. Each can help illustrate the potential of process-based models to assess certain
47 key ecosystem services (related mainly to crop productivity, biogeochemical cycling and weed
48 control). Moreover, the temporal scale at which processes and interactions occur and ecosystem
49 services are built may require long-term simulations (Figure 5). Generally, when the spatial scale
50 increases (from plant-plant to the landscape), the temporal scale increases. However, modelling
51 could target more ambitious applications than those documented to date, such as more
52 comprehensive representation of environmental drivers (e.g. pests and pathogens, soil phosphorus
53 content, climate change) and greater detail in the relationships between plant diversity (crop, service
54 and weed plants) and biodiversity at other trophic levels in agroecosystems (pests and diseases).

55 4.2. Challenges and difficulties linked to modelling solutions

56 Reusing and coupling existing models faces several methodological and technical challenges. To be
57 effective, direct coupling and hybrid modelling often require developing specific adapters or new
58 model code. Too many inconsistencies between models, such as differences in temporal and spatial
59 resolutions, concepts and coupling variables, can make it more difficult to couple the models. The
60 coupling time step must be defined and be consistent with the time step of the interactions between
61 the simulated systems. This indicates that it may be necessary to increase (temporal upscaling) or
62 decrease (temporal downscaling) the time step of one of the coupled models; the latter assumes
63 knowing how to describe processes at a finer temporal resolution. Furthermore, the processes
64 considered can occur at different spatial scales (e.g. from field to watershed) depending on the type
65 of organisms and the factors involved, and can be influenced by multiple interactions. Modelling
66 platforms do not always have sufficient technical development to combine these contrasting
67 resolutions to describe systems and their functioning. Moreover, coupling models promotes dialogue
68 between disciplines (e.g. agronomy and hydrology) and thus requires agreeing on a common lexicon
69 or an ontology.

70 4.3. Modelling perspectives and opportunities

71 Many consequences of multispecies systems remain to be explored, both experimentally and
72 through modelling and theoretical studies. We advocate practicing both during the transition
73 towards more agroecological systems. Models cannot be developed without supporting data, and a
74 lack of reliable models hinders data analysis. This is particularly true regarding consequences of
75 plant-plant interactions, for which the magnitude and hierarchy of the major processes involved

76 remain hotly debated despite over 80 years of manipulative and observational studies (Brooker
77 2006; Weisser *et al.* 2017). This lack of understanding prevents identification of a consensual, much
78 less optimal, model structure. However, it also promotes the development of a variety of models to
79 test and benchmark interactions between mechanisms that act simultaneously (e.g. competition and
80 complementarity for resources, different forms of facilitation, physical and chemical signalling). In
81 this context, combining specific model developments with effective strategies to aggregate them
82 encourages parallel progress in key disciplinary issues (related to biophysical aspects and social
83 sciences in managed ecosystems), while still enabling integration of outputs relevant for predicting
84 ecosystem services at different scales.

85 Connecting data with models to develop diversified cropping systems provides an opportunity to
86 address issues involved in quantifying biodiversity-based services. As a part of managed ecosystems,
87 these services are scrutinized more closely than those in natural systems and benefit from
88 observation in agricultural networks (e.g. Lechenet *et al.* 2017), as well as developments in digital
89 agriculture that are increasingly used for diversified systems (Reboud 2019; Chen *et al.* 2019). They
90 also depend greatly on crops that have a long history of biological characterization and modelling
91 and are now benefiting from the early development of high-throughput information systems in
92 plants (Tardieu *et al.* 2017). Rich benchmarking datasets that cover multiple ecosystem services
93 rather than only productivity are increasingly available. These data remain rare, but they are required
94 to understand potential trade-offs between services and to identify inconsistent predictions across
95 scales (Schneider *et al.* 2014). Here, models are needed to go beyond the observational posture of
96 naturalists and quantitatively represent and analyse effects of plant diversity in a high number of
97 possible scenarios

98 To this end, the ability to predict consequences of within-field diversity at different spatial and
99 temporal scales is required in order to assess the overall interest of various diversification scenarios.
100 A general belief about natural ecosystems is that plant diversity alone provides the ecosystem
101 services targeted, and that increasing species and genetic diversity in cropping systems should be a
102 goal to provide multiple services. However, how and why a particular arrangement of practices, or a
103 given range of diversity, should be chosen largely remain to be solved. By definition, managed
104 agroecosystems have an economic purpose and often target particular marketable products. From a
105 farmer's perspective, diversification thus has advantages (resilience) and disadvantages (not all
106 species are equal from an economic viewpoint; more complex management). Our opinion is that
107 combining models that can represent plant diversity and predict ecosystem services at multiple

108 scales is critical to support the spatial and temporal prediction of the many systems that could be
109 designed.

110 **Concluding remarks**

111 Diversified agriculture points a clear route toward more sustainable systems able to provide a range
112 of services to the society beyond agricultural production. Exploring and evaluating the diversity of
113 possible solutions is by no means simple, and will require the combination of different approaches
114 relying on field experiments, farmers networking and new technological tools taking advantage of
115 heterogeneous sources of data. We support that plant, crop and cropping system models will be
116 among the key tools to help achieving this goal. We illustrated it here through the example of one
117 major option regarding crop diversification, the increase of within-field variability by mixing different
118 crops, and highlighted potential connection and complementarities in the range of models already at
119 hand. A proximate use of multi-scale modelling solutions could be to help explore numerically the
120 benefits and constraints of different diversification scenarios and address the “diversification
121 dilemma” of an almost infinite number of combinations to test. At the very least, it could help
122 focusing the experimental development efforts on the most promising solutions and limit the test in
123 field trials of non-beneficial systems (e.g. non-compatible diversity that does not provide the
124 expected services or whose costs decrease system resilience or farmers’ incomes). When mature and
125 more robust, an ultimate use of these models could also be to help quantify non-productive
126 ecosystem services. It is clear that intercropping and other diversification practices will not become
127 widespread without sufficient economic justification. Such models could be very useful to help
128 determine the added value of diversified systems. We are witnessing a renewal of interest for these
129 systems and anticipate that further developments of models in this area will be critical in the coming
130 years.

131
132 **Acknowledgements:** The authors are very grateful to Evelyne Costes and Kathrin Kahlen, who invited
133 them to reflect on this subject at the 9th International Conference on Functional-Structural Plant
134 Models in Hanover, Germany, from 5-9 October 2020. We also thank Michael and Michelle Corson
135 for their helpful comments and English revision.

136

137 **Contribution of authors**

- 138 - M.L., N.G. and G.L. conceived and wrote the manuscript
139 - R.B., C.M. and R.V. contributed ideas, supported the writing, read and commented on the
140 entire manuscript

141 REFERENCES

- 142 **Agudelo CAR, Bustos SLH, Moreno CAP. 2020.** Modeling interactions among multiple ecosystem services. A
143 critical review. *Ecological Modelling* **429**: 109103.
- 144 **Banik P, Midya A, Sarkar BK, Ghose SS. 2006.** Wheat and chickpea intercropping systems in an additive series
145 experiment: Advantages and weed smothering. *European Journal of Agronomy* **24**: 325–332.
- 146 **Barillot R, Escobar-Gutierrez AJ, Fournier C, Huynh P, Combes D. 2014.** Assessing the effects of architectural
147 variations on light partitioning within virtual wheat-pea mixtures. *Annals of Botany* **114**: 725–737.
- 148 **Barillot R, Louarn G, Escobar-Gutiérrez AJ, Huynh P, Combes D. 2011.** How good is the turbid medium-based
149 approach for accounting for light partitioning in contrasted grass–legume intercropping systems? *Annals of*
150 *Botany* **108**: 1013–1024.
- 151 **Beillouin D, Ben-Ari T, Malézieux E, Seufert V, Makowski D. 2021.** Positive but variable effects of crop
152 diversification on biodiversity and ecosystem services. *Global Change Biology* **27**: 4697–4710.
- 153 **Bertrand N, Roux S, Forey O, Guinet M, Wery J. 2018.** Simulating plant water stress dynamics in a wide range
154 of bi-specific agrosystems in a region using the BISWAT model. *European Journal of Agronomy* **99**: 116–128.
- 155 **Bohanec M. 2008.** *DEXi : Program for Multi-Attribute Decision Making User's Manual. Evaluation.*
- 156 **Brisson N, Launay M, Mary B, Beaudoin N. 2008.** *Conceptual Basis, Formalisations and Parameterization of the*
157 *STICS Crop Model.* Editions Quae.
- 158 **Brooker RW. 2006.** Plant–plant interactions and environmental change. *New Phytologist* **171**: 271–284.
- 159 **Brooker RW, Bennett AE, Cong W-F, et al. 2015.** Improving intercropping: a synthesis of research in agronomy,
160 plant physiology and ecology. *New Phytologist* **206**: 107–117.
- 161 **Brooker RW, George TS, Homulle Z, et al. 2021.** Facilitation and biodiversity–ecosystem function relationships
162 in crop production systems and their role in sustainable farming. *Journal of Ecology* **n/a**.
- 163 **Caron P, Biénabe E, Hainzelin E. 2014.** Making transition towards ecological intensification of agriculture a
164 reality: the gaps in and the role of scientific knowledge. *Current Opinion in Environmental Sustainability* **8**: 44–
165 52.
- 166 **Chapagain T, Riseman A. 2014.** Barley-pea intercropping: Effects on land productivity, carbon and nitrogen
167 transformations. *Field Crops Research* **166**: 18-25.
- 168 **Chen T-W, Cabrera-Bosquet L, Alvarez Prado S, et al. 2019.** Genetic and environmental dissection of biomass
169 accumulation in multi-genotype maize canopies. *Journal of Experimental Botany* **70**: 2523–2534.
- 170 **Colas F, Cordeau S, Granger S, et al. 2020.** Co-development of a decision support system for integrated weed
171 management: Contribution from future users. *European Journal of Agronomy* **114**: 126010.
- 172 **Colas F, Gauchi J-P, Villerd J, Colbach N. 2021.** Simplifying a complex computer model: Sensitivity analysis and
173 metamodelling of an 3D individual-based crop-weed canopy model. *Ecological Modelling* **454**: 109607.
- 174 **Colbach N, Colas F, Cordeau S, et al. 2021.** The FLORSYS crop-weed canopy model, a tool to investigate and
175 promote agroecological weed management. *Field Crops Research* **261**: 108006.

- 176 **Colbach N, Collard A, Guyot SHM, Meziere D, Munier-Jolain N. 2014.** Assessing innovative sowing patterns for
177 integrated weed management with a 3D crop:weed competition model. *European Journal of Agronomy* **53**: 74–
178 89.
- 179 **Collard B, Tixier P, Carval D, Lavigne C, Delattre T. 2018.** Spatial organisation of habitats in agricultural plots
180 affects per-capita predator effect on conservation biological control: An individual based modelling study.
181 *Ecological Modelling* **388**: 124–135.
- 182 **Confalonieri R. 2014.** CoSMo: A simple approach for reproducing plant community, dynamics using a single
183 instance of generic crop simulators. *Ecological Modelling* **286**: 1–10.
- 184 **Corre-Hellou G, Faure M, Launay M, Brisson N, Crozat Y. 2009.** Adaptation of the STICS intercrop model to
185 simulate crop growth and N accumulation in pea-barley intercrops. *Field Crops Research* **113**: 72–81.
- 186 **Dhima KV, Lithourgidis AS, Vasilakoglou IB, Dordas CA. 2007.** Competition indices of common vetch and cereal
187 intercrops in two seeding ratio. *Field Crops Research* **100**: 249–256.
- 188 **Duru M, Therond O, Fares M. 2015.** Designing agroecological transitions; A review. *Agronomy for Sustainable*
189 *Development* **35**: 1237–1257.
- 190 **Escobar-Gutiérrez AJ, Combes D, Rakocevic M, et al. 2009.** Functional relationships to estimate
191 Morphogenetically Active Radiation (MAR) from PAR and solar broadband irradiance measurements: The case
192 of a sorghum crop. *Agricultural and Forest Meteorology* **149**: 1244–1253.
- 193 **Evans MEK, Merow C, Record S, McMahon SM, Enquist BJ. 2016.** Towards Process-based Range Modeling of
194 Many Species. *Trends in Ecology & Evolution* **31**: 860–871.
- 195 **Evers JB, van der Werf W, Stomph TJ, Bastiaans L, Anten NPR. 2019.** Understanding and optimizing species
196 mixtures using functional-structural plant modelling. *Journal of experimental botany* **70**: 2381–2388.
- 197 **Fabre F, Rousseau E, Mailleret L, Moury B. 2012.** Durable strategies to deploy plant resistance in agricultural
198 landscapes. *New Phytologist* **193**: 1064–1075.
- 199 **Faverjon L, Escobar-Gutiérrez A, Litrico I, Julier B, Louarn G. 2019.** A generic individual-based model can
200 predict yield, nitrogen content, and species abundance in experimental grassland communities. *Journal of*
201 *Experimental Botany* **70**: 2491–2504.
- 202 **Finckh MR, Gacek ES, Goyeau H, et al. 2000.** Cereal variety and species mixtures in practice, with emphasis on
203 disease resistance. *Agronomie* **20**: 813–837.
- 204 **Gaba S, Lescourret F, Boudsocq S, et al. 2015.** Multiple cropping systems as drivers for providing multiple
205 ecosystem services: from concepts to design. *Agronomy for Sustainable Development* **35**: 607–623.
- 206 **Gaudio N, Escobar-Gutiérrez AJ, Casadebaig P, et al. 2019.** Current knowledge and future research
207 opportunities for modeling annual crop mixtures. A review. *Agronomy for Sustainable Development* **39**: 20.
- 208 **Gaudio N, Violle C, Gendreau X, et al. 2021.** Interspecific interactions regulate plant reproductive allometry in
209 cereal-legume intercropping systems. *Journal of Applied Ecology*. <https://doi.org/10.1111/1365-2664.13979>
- 210 **Gauthier M, Barillot R, Schneider A, et al. 2020.** A functional structural model of grass development based on
211 metabolic regulation and coordination rules. *Journal of Experimental Botany* **71**: 5454–5468.

- 212 **Haines-Young R, Potschin M. 2013.** *Common International Classification of Ecosystem Services (CICES):*
213 *Consultation on Version 4, August-December 2012 (Report to the European Environment Agency, revised*
214 *January 2013).*
- 215 **Hauggaard-Nielsen H, Jørnsgaard B, Kinane J, Jensen ES. 2008.** Grain legume–cereal intercropping: The
216 practical application of diversity, competition and facilitation in arable and organic cropping systems.
217 *Renewable Agriculture and Food Systems* **23**: 3–12.
- 218 **He XH, Xu MG, Qiu GY, Zhou JB. 2009.** Use of N-15 stable isotope to quantify nitrogen transfer between
219 mycorrhizal plants. *Journal of Plant Ecology* **2**: 107-118.
- 220 **van Ittersum MK, Cassman K, Grassini P, Wolf J, Tittone P, Hochman Z. 2013.** Yield gap analysis with local to
221 global relevance - a review. *Field Crops Research* **143**: 4–17.
- 222 **Jin R, Chen W, Simpson TW. 2001.** Comparative studies of metamodelling techniques under multiple modelling
223 criteria. *Structural and Multidisciplinary Optimization* **23**: 1–13.
- 224 **Johansen A, Jensen ES. 1996.** Transfer of N and P from intact or decomposing roots of pea to barley
225 interconnected by an arbuscular mycorrhizal fungus. *Soil Biology and Biochemistry* **28**: 73-81.
- 226 **Kahlen K, Stützel H. 2011.** Simplification of a light-based model for estimating final internode length
227 in greenhouse cucumber canopies. *Annals of Botany* **108**: 1055–1063.
- 228 **Kiaer LP, Skovgaard IM, Ostergard H. 2012.** Effects of inter-varietal diversity, biotic stresses and environmental
229 productivity on grain yield of spring barley variety mixtures. *Euphytica* **185**: 123–138.
- 230 **Kremen C, Miles A. 2012.** Ecosystem Services in Biologically Diversified versus Conventional Farming Systems:
231 Benefits, Externalities, and Trade-Offs. *Ecology and Society* **17**: 40.
- 232 **Landis DA, Wratten SD, Gurr GM. 2000.** Habitat Management to Conserve Natural Enemies of Arthropod Pests
233 in Agriculture. *Annual Review of Entomology* **45**: 175–201.
- 234 **Launay M, Brisson N, Satger S, et al. 2009.** Exploring options for managing strategies for pea-barley
235 intercropping using a modeling approach. *European Journal of Agronomy* **31**: 85–98.
- 236 **Lechenet M, Dessaint F, Py G, Makowski D, Munier-Jolain N. 2017.** Reducing pesticide use while preserving
237 crop productivity and profitability on arable farms. *Nature Plants* **3**: 17008.
- 238 **Li HB, Ma QH, Li HG, Zhang FS, Rengel Z, Shen JB. 2014.** Root morphological responses to localized nutrient
239 supply differ among crop species with contrasting root traits. *Plant and Soil* **376**: 151-163.
- 240 **Louarn G, Barillot R, Combes D, Escobar-Gutiérrez A. 2020.** Towards intercrop ideotypes: non-random trait
241 assembly can promote overyielding and stability of species proportion in simulated legume-based mixtures.
242 *Annals of Botany*.
- 243 **Louarn G, Faverjon L. 2018.** A generic individual-based model to simulate morphogenesis, C-N acquisition and
244 population dynamics in contrasting forage legumes. *Annals of Botany* **121**: 875–896.
- 245 **Louarn G, Song Y. 2020.** Two decades of functional–structural plant modelling: now addressing fundamental
246 questions in systems biology and predictive ecology. *Annals of Botany* **126**: 501–509.

- 247 **Lowry CJ, Brainard DC. 2016.** Strip-Intercropping of Rye-Vetch Mixtures Affects Biomass, Carbon/Nitrogen
248 Ratio, and Spatial Distribution of Cover Crop Residue. *Agronomy Journal* **108**: 2433-2443.
- 249 **Malard JJ, Adamowski JF, Diaz MR, et al. 2020.** Agroecological food web modelling to evaluate and design
250 organic and conventional agricultural systems. *Ecological Modelling* **421**: 108961.
- 251 **Meunier C, Alletto L, Bedoussac L, et al. 2022.** A modelling chain combining soft and hard models to assess a
252 bundle of ecosystem services provided by a diversity of cereal-legume intercrops. *European Journal of*
253 *Agronomy* **132**: 126412.
- 254 **Midingoyi CA, Pradal C, Enders A, et al. 2021.** Crop2ML: An open-source multi-language modeling framework
255 for the exchange and reuse of crop model components. *Environmental Modelling & Software* **142**: 105055.
- 256 **Moorcroft PR, Hurtt GC, Pacala SW. 2001.** A Method for Scaling Vegetation Dynamics: The Ecosystem
257 Demography Model (ed). *Ecological Monographs* **71**: 557–586.
- 258 **Pagel J, Schurr FM. 2012.** Forecasting species ranges by statistical estimation of ecological niches and spatial
259 population dynamics. *Global Ecology and Biogeography* **21**: 293–304.
- 260 **Pagès L, Pointurier O, Moreau D, Voisin A-S, Colbach N. 2020.** Metamodelling a 3D architectural root-system
261 model to provide a simple model based on key processes and species functional groups. *Plant and Soil* **448** :
262 231-251.
- 263 **Pao YC, Chen TW, Moualeu-Ngangue DP, Stützel H. 2019.** Environmental triggers for photosynthetic
264 protein turnover determine the optimal nitrogen distribution and partitioning in the canopy. *Journal of*
265 *Experimental Botany* **70** : 2419-2434.
- 266 **Papaïx J, Touzeau S, Monod H, Lannou C. 2014.** Can epidemic control be achieved by altering landscape
267 connectivity in agricultural systems? *Ecological Modelling* **284**: 35–47.
- 268 **Pearson RG, Dawson TP. 2003.** Predicting the impacts of climate change on the distribution of species: are
269 bioclimate envelope models useful? *Global Ecology and Biogeography* **12**: 361–371.
- 270 **Peng B, Guan K, Tang J, et al. 2020.** Towards a multiscale crop modelling framework for climate change
271 adaptation assessment. *Nature Plants* **6**: 338–348.
- 272 **Pointurier O, Moreau D, Pagès L, Caneill J, Colbach N. 2021.** Individual-based 3D modelling of root systems in
273 heterogeneous plant canopies at the multiannual scale. Case study with a weed dynamics model. *Ecological*
274 *Modelling* **440**: 109376.
- 275 **Prieto I, Violle C, Barre P, Durand J-L, Ghesquiere M, Litrico I. 2015.** Complementary effects of species and
276 genetic diversity on productivity and stability of sown grasslands. *Nature Plants* **1**: 15033.
- 277 **R Core Team. 2018.** R: A language and environment for statistical computing. R Foundation for Statistical
278 Computing In: Vienna, Austria.
- 279 **Reboud X. 2019.** Le lien entre agriculture et biodiversité : ce qui a bougé sur le plan de la société, de
280 la réflexion entre 2008 et 2018. *Innovations Agronomiques* **75**: 1–14.
- 281 **Rosenzweig C, Jones JW, Hatfield JL, et al. 2013.** The Agricultural Model Intercomparison and Improvement
282 Project (AgMIP): Protocols and pilot studies. *Agricultural and Forest Meteorology* **170**: 166–182.

- 283 **Sabatier R, Mouysset L. 2018.** A robustness-based viewpoint on the production-ecology trade-off in
284 agroecosystems. *Agricultural Systems* **167**: 1–9.
- 285 **Schneider MK, Lüscher G, Jeanneret P, et al. 2014.** Gains to species diversity in organically farmed fields are
286 not propagated at the farm level. *Nature Communications* **5**: 4151.
- 287 **Sirami C, Gross N, Baillod AB, et al. 2019.** Increasing crop heterogeneity enhances multitrophic diversity across
288 agricultural regions. *Proceedings of the National Academy of Sciences* **116**: 16442–16447.
- 289 **Stöckle CO, Kemanian AR. 2020.** Can Crop Models Identify Critical Gaps in Genetics, Environment, and
290 Management Interactions? *Frontiers in Plant Science* **11**.
- 291 **Tardieu F, Cabrera-Bosquet L, Pridmore T, Bennett M. 2017.** Plant Phenomics, From Sensors to Knowledge.
292 *Current Biology* **27**: R770–R783.
- 293 **Tscharntke T, Grass I, Wanger TC, Westphal C, Batáry P. 2021.** Beyond organic farming – harnessing
294 biodiversity-friendly landscapes. *Trends in Ecology & Evolution*. <https://doi.org/10.1016/j.tree.2021.06.010>.
- 295 **Tixier P, Duyck P-F, Cote F-X, Caron-Lormier G, Malezieux E. 2013.** Food web-based simulation for
296 agroecology. *Agronomy for Sustainable Development* **33**: 663–670.
- 297 **Van Oijen M, Barcza Z, Confalonieri R, et al. 2020.** Incorporating Biodiversity into Biogeochemistry Models to
298 Improve Prediction of Ecosystem Services in Temperate Grasslands: Review and Roadmap. *Agronomy* **10**: 259.
- 299 **Verret V, Pelzer E, Bedoussac L, Jeuffroy M-H. 2020.** Tracking on-farm innovative practices to support crop
300 mixture design: The case of annual mixtures including a legume crop. *European Journal of Agronomy* **115**:
301 126018.
- 302 **Vidal T, Gigot C, de Vallavieille-Pope C, Huber L, Saint-Jean S. 2018.** Contrasting plant height can improve the
303 control of rain-borne diseases in wheat cultivar mixture: modelling splash dispersal in 3-D canopies. *Annals of*
304 *botany* **121**: 1299–1308.
- 305 **Violle C, Navas M-L, Vile D, et al. 2007.** Let the concept of trait be functional! *Oikos* **116**: 882–892.
- 306 **Volaire F, Gleason SM, Delzon S. 2020.** What do you mean “functional” in ecology? Patterns versus processes.
307 *Ecology and Evolution* **10**: 11875–11885.
- 308 **Weisser WW, Roscher C, Meyer ST, et al. 2017.** Biodiversity effects on ecosystem functioning in a 15-year
309 grassland experiment: Patterns, mechanisms, and open questions. *Basic and Applied Ecology* **23**: 1–73.
- 310 **Wood SA, Karp DS, DeClerck F, Kremen C, Naeem S, Palm CA. 2015.** Functional traits in agriculture:
311 agrobiodiversity and ecosystem services. *Trends in Ecology & Evolution* **30**: 531–539.
- 312 **Xu QN, Yang X, Yan Y, Wang SP, Loreau M, Jiang L. 2021.** Consistently positive effect of species diversity on
313 ecosystem, but not population, temporal stability. *Ecology Letters* **24**:2256-66
- 314 **Yin W, Chai Q, Zhao C, Yu A, Fan Z, Hu F, Fan H, Guo Y, Coulter JA. 2020.** Water utilization in intercropping: A
315 review. *Agricultural Water Management* **241**: 106335.
- 316 **Zhu J, van der Werf W, Anten NPR, Vos J, Evers JB. 2015.** The contribution of phenotypic plasticity to
317 complementary light capture in plant mixtures. *New Phytologist* **207**: 1213–1222.

318 Figure 1. Different features are required to upscale from plant-plant interactions (e.g. competition,
319 complementary, facilitation) to agroecosystems that provide ecosystem services. This combines the individual
320 (plant) model level with biophysical (abiotic and biotic environments) and technical levels. However, the
321 corollary of this upscaling is downscaling, in the sense that modellers and users may need to apply parsimony
322 to simulate these systems at large scales (at which only a few factors explain variability in performance)
323 (Pearson and Dawson 2003; Evans et al. 2016), or to identify an optimal system that responds to a set of
324 constraints and objectives at a given location. Adapted from Louarn and Song (2020).

325

326 Figure 2. Illustration of facilitation against pests resulting from mixing two wheat cultivars that vary in height
327 and resistance to pests. An aerial functional-structural plant model (FSPM) and a rain-splash model were
328 coupled to simulate facilitation (Vidal et al. 2018). Source of photographs: Sébastien Saint-Jean (INRAE, UMR
329 EcoSys).

330

331 Figure 3. Overview of the main processes that connect the FlorSys individual-based model (IBM, Colbach et al.
332 2014), which simulates the aboveground crop-weed canopy, to the RSConc metamodel (Pagès et al. 2020),
333 which simulates root growth, and the STICS crop model (Brisson et al. 2008), which provides soil structure and
334 climate and their effects on root growth. Adapted from Pointurier et al. (2021).

335

336 Figure 4. Introduction to the serious game “Interplay”, used to assess a bundle of ecosystem services provided
337 by including crop mixtures in a crop rotation. The game and overall structure of the modelling chain are
338 illustrated. Green boxes are examples of options selected in a designed intercropping scenario. The main steps
339 of the game and the variables of the modelling chain are in white, the dry matter (DM) of cereal biomass in the
340 scenario is in light orange, the DM of legume biomass is in green and additional variables that influence soil
341 structure are in dark orange.

342

343 Figure 5. Conceptual illustration of modelling solutions (i.e. inverse modelling, metamodelling, coupling and
344 hybrid modelling) used to simulate crop mixtures at different spatial scales (plant, field, farm and landscape),
345 which are characterized by contrasting processes and ecosystem services. Asterisks indicate that the multiple
346 spatial scales involve both short- and long-term simulations.

347


Figure 1

Modelling...

crop mixtures that provide ecosystem services




... requires modelling plant-plant & plant-environment interactions




AND provisioning & regulating services

Accepted




Choice of the genotypes = $f(\text{architecture, resistance})$

Aerial FSPM x physical model




Accepted

Figure 3




Accepted Manuscript

Figure 4



Accepted Manuscript

Figure 5



Accepted Manuscript